

# DAGGER CATEGORIES OF RELATIONS

MATTHEW DiMEGLIO

Joint work with Chris Heunen, Paolo Perrone and Dario Stein

EDINBURGH CATEGORY SEMINAR  
MARCH 2025

Bijection between contractions and  
relations in category of coisometries

Linear maps  $f:X \rightarrow Y$  between Hilbert spaces  
such that  $\|fx\| \leq \|x\|$  for all  $x \in X$

Bijection between contractions and  
relations in category of coisometries

Linear maps  $f:X \rightarrow Y$  between Hilbert spaces  
such that  $\|fx\| \leq \|x\|$  for all  $x \in X$

Bijection between contractions and  
relations in category of coisometries

Isomorphism classes of  
jointly monic spans

Linear maps  $f:X \rightarrow Y$  between Hilbert spaces  
such that  $\|fx\| \leq \|x\|$  for all  $x \in X$

Bijection between contractions and  
relations in category of coisometries

Isomorphism classes of  
jointly monic spans


Adjointable maps  $f:X \rightarrow Y$  such that  $ff^+ = 1$   
(orthogonal projections onto closed subspaces)

But category of coisometries  
is not regular

But category of coisometries  
is not regular

It does not have pullbacks





# DEFINITION

A  $t$ -category is a category with  
 $f^+: Y \rightarrow X$  for each  $f: X \rightarrow Y$  such that  
 $1^+ = 1$      $(gf)^+ = f^+g^+$      $f^{++} = f$

# DEFINITION

A  $t$ -category is a category with  
 $f^+: Y \rightarrow X$  for each  $f: X \rightarrow Y$  such that  
 $1^+ = 1$      $(gf)^+ = f^+g^+$      $f^{++} = f$

## EXAMPLES

Con

Hilbert spaces and contractions

$$\langle y | f x \rangle = \langle f^+ y | x \rangle$$

# DEFINITION

A  $t$ -category is a category with  
 $f^+: Y \rightarrow X$  for each  $f: X \rightarrow Y$  such that  
 $1^+ = 1$      $(gf)^+ = f^+g^+$      $f^{++} = f$

## EXAMPLES

### Con

Hilbert spaces and contractions

$$\langle y | f x \rangle = \langle f^+ y | x \rangle$$

### PInj

Sets and injective partial functions

$$y = f x \Leftrightarrow x = f^+ y$$

# DEFINITION

A  $t$ -category is a category with  $f^+: Y \rightarrow X$  for each  $f: X \rightarrow Y$  such that

$$1^+ = 1 \quad (gf)^+ = f^+ g^+ \quad f^{++} = f$$

## EXAMPLES

### Con

Hilbert spaces and contractions

$$\langle y | f x \rangle = \langle f^+ y | x \rangle$$

### Plnj

Sets and injective partial functions

$$y = f x \Leftrightarrow x = f^+ y$$

## FinPS

A finite probability space  $X$  is a finite set  $X$  equipped with a function  $P_x: X \rightarrow (0,1]$  such that  $\sum_{x \in X} P_x(x) = 1$ .

# DEFINITION

A  $t$ -category is a category with  $f^+: Y \rightarrow X$  for each  $f: X \rightarrow Y$  such that

$$1^+ = 1 \quad (gf)^+ = f^+ g^+ \quad f^{++} = f$$

## EXAMPLES

### Con

Hilbert spaces and contractions

$$\langle y | f x \rangle = \langle f^+ y | x \rangle$$

### PInj

Sets and injective partial functions

$$y = f x \Leftrightarrow x = f^+ y$$

# FinPS

A finite probability space  $X$  is a finite set  $X$  equipped with a function  $\mathbb{P}_x: X \rightarrow [0, 1]$  such that  $\sum_{x \in X} \mathbb{P}_x(x) = 1$ .

A stochastic map  $f: X \rightarrow Y$  is a function  $\mathbb{P}_f(-| -): Y \times X \rightarrow [0, 1]$  such that

$$\sum_{y \in Y} \mathbb{P}_f(y|x) = 1$$

$$\sum_{x \in X} \mathbb{P}_f(y|x) \mathbb{P}_x(x) = \mathbb{P}_Y(y)$$

# DEFINITION

A  $t$ -category is a category with  $f^+: Y \rightarrow X$  for each  $f: X \rightarrow Y$  such that  $1^+ = 1$ ,  $(gf)^+ = f^+g^+$ ,  $f^{++} = f$

## EXAMPLES

### Con

Hilbert spaces and contractions

$$\langle y | f x \rangle = \langle f^+ y | x \rangle$$

### Plnj

Sets and injective partial functions

$$y = f x \Leftrightarrow x = f^+ y$$

## FinPS

A finite probability space  $X$  is a finite set  $X$  equipped with a function  $\mathbb{P}_x: X \rightarrow (0,1]$  such that  $\sum_{x \in X} \mathbb{P}_x(x) = 1$ .

A stochastic map  $f: X \rightarrow Y$  is a function  $\mathbb{P}_f(-|-): Y \times X \rightarrow [0,1]$  such that

$$\sum_{y \in Y} \mathbb{P}_f(y|x) = 1$$

$$\sum_{x \in X} \mathbb{P}_f(y|x) \mathbb{P}_x(x) = \mathbb{P}_Y(y)$$

The Bayesian inverse  $f^+$  of  $f$  is given by

$$\mathbb{P}_f(y|x) \mathbb{P}_x(x) = \mathbb{P}_{f^+}(x|y) \mathbb{P}_Y(y)$$

# DEFINITION

A morphism  $f$  is coisometric if  $ff^+=1$ .

# DEFINITION

A morphism  $f$  is coisometric if  $ff^t=1$ .

Let  $\text{Coisometry}(C)$  be the wide  
subcategory of coisometries in  $C$ .


# DEFINITION

A morphism  $f$  is coisometric if  $ff^*=1$ .

Let  $\text{Coisometry}(C)$  be the wide subcategory of coisometries in  $C$ .

## EXAMPLES

Isometries in  $\text{Con}$  are inclusions of closed subspaces.




# DEFINITION

A morphism  $f$  is coisometric if  $ff^*=1$ .


Let **Coisometry**( $C$ ) be the wide subcategory of coisometries in  $C$ .

## EXAMPLES

Isometries in **Con** are inclusions of closed subspaces.



Isometries in **Plnj** are total.




# DEFINITION

A morphism  $f$  is coisometric if  $ff^*=1$ .


Let **Coisometry**( $C$ ) be the wide subcategory of coisometries in  $C$ .

## EXAMPLES

Isometries in **Con** are inclusions of closed subspaces.



Isometries in **Plnj** are total.



Coisometries in **FinPS** are deterministic

$$P_f(y|x) \in \{0, 1\}$$


# DEFINITION

A morphism  $f$  is coisometric if  $ff^*=1$ .

Let **Coisometry**( $C$ ) be the wide subcategory of coisometries in  $C$ .


## EXAMPLES

Isometries in **Con** are inclusions of closed subspaces.



Isometries in **Plnj** are total.

2



Coisometries in **FinPS** are deterministic

$$P_f(y|x) \in \{0, 1\}$$

Write  $f$  also for the underlying function

$$P_f(y|x) = \begin{cases} 1 & \text{if } y = fx \\ 0 & \text{otherwise} \end{cases}$$

# DEFINITION

A dilation of  $f: X \rightarrow Y$  is a span

$$X \xleftarrow{s_1} S \xrightarrow{s_2} Y$$

of coisometries such that  $f = s_2 s_1^+$ .

# DEFINITION

A dilation of  $f: X \rightarrow Y$  is a span

$$X \xleftarrow{s_1} S \xrightarrow{s_2} Y$$

of coisometries such that  $f = s_2 s_1^+$ .

A dilator of  $f$  is a terminal dilation of  $f$ .

# DEFINITION

A dilation of  $f: X \rightarrow Y$  is a span

$$X \xleftarrow{s_1} S \xrightarrow{s_2} Y$$

of coisometries such that  $f = s_2 s_1^+$ .

A dilator of  $f$  is a terminal dilation of  $f$ .

$$X \xleftarrow{p_1} X \boxplus_f Y \xrightarrow{p_2} Y$$

# DEFINITION

A dilation of  $f: X \rightarrow Y$  is a span

$$X \xleftarrow{s_1} S \xrightarrow{s_2} Y$$

of coisometries such that  $f = s_2 s_1^+$ .

A dilator of  $f$  is a terminal dilation of  $f$ .

$$\begin{array}{ccccc} & & S & & \\ & \swarrow s_1 & & \searrow s_2 & \\ X & \xleftarrow{p_1} & X \boxplus_f Y & \xrightarrow{p_2} & Y \end{array}$$


# DEFINITION

A dilation of  $f: X \rightarrow Y$  is a span

$$X \xleftarrow{s_1} S \xrightarrow{s_2} Y$$

of coisometries such that  $f = s_2 s_1^+$ .

A dilator of  $f$  is a terminal dilation of  $f$ .



# DEFINITION

A dilation of  $f: X \rightarrow Y$  is a span

$$X \xleftarrow{s_1} S \xrightarrow{s_2} Y$$

of coisometries such that  $f = s_2 s_1^*$ .

A dilator of  $f$  is a terminal dilation of  $f$ .

$$\begin{array}{ccccc} & & S & & \\ & \swarrow s_1 & \downarrow s & \searrow s_2 & \\ X & \xleftarrow{p_1} & X \oplus_f Y & \xrightarrow{p_2} & Y \end{array}$$

# EXAMPLES

Codilators  
in Con

$$\begin{array}{ccc} X \oplus Y & & \\ \uparrow & [\sqrt{1-f^*f} & f] \\ X & & \\ & & \uparrow [0 & 1] \\ & & Y \end{array}$$

# DEFINITION

A dilation of  $f: X \rightarrow Y$  is a span

$$X \xleftarrow{s_1} S \xrightarrow{s_2} Y$$

of coisometries such that  $f = s_2 s_1^+$ .

A dilator of  $f$  is a terminal dilation of  $f$ .

$$\begin{array}{ccccc} & & S & & \\ & \swarrow s_1 & \downarrow s & \searrow s_2 & \\ X & \xleftarrow{p_1} & X \boxplus_f Y & \xrightarrow{p_2} & Y \end{array}$$

# EXAMPLES

Codilators  
in **Con**

$$\begin{array}{ccc} & X \oplus Y & \\ s_1 \nearrow & & \swarrow s_2 \\ X & & Y \end{array}$$

# DEFINITION

A dilation of  $f: X \rightarrow Y$  is a span

$$X \xleftarrow{s_1} S \xrightarrow{s_2} Y$$

of coisometries such that  $f = s_2 s_1^+$ .

A dilator of  $f$  is a terminal dilation of  $f$ .

$$\begin{array}{ccccc} & & S & & \\ & \swarrow s_1 & \downarrow s & \searrow s_2 & \\ X & \xleftarrow{p_1} & X \boxplus_f Y & \xrightarrow{p_2} & Y \end{array}$$

# EXAMPLES

Codilators  
in **Con**

$$\begin{array}{ccccc} & & X \oplus Y & & \\ & \nearrow s_1 & \uparrow s_1 \vee s_2 & \searrow s_2 & \\ X & \longrightarrow & X \vee Y & \longleftarrow & Y \end{array}$$

# DEFINITION

A dilation of  $f: X \rightarrow Y$  is a span

$$X \xleftarrow{s_1} S \xrightarrow{s_2} Y$$

of coisometries such that  $f = s_2 s_1^+$ .

A dilator of  $f$  is a terminal dilation of  $f$ .

$$\begin{array}{ccccc} & & S & & \\ & \swarrow s_1 & \downarrow s & \searrow s_2 & \\ X & \xleftarrow{p_1} & X \boxplus_f Y & \xrightarrow{p_2} & Y \end{array}$$

# EXAMPLES

Codilators  
in **Con**

$$\begin{array}{ccccc} & & X \oplus Y & & \\ & \nearrow s_1 & \uparrow s_1 \vee s_2 & \searrow s_2 & \\ X & \longrightarrow & X \vee Y & \longleftarrow & Y \end{array}$$

$$X \vee Y = \overline{\text{Im } s_1 + \text{Im } s_2}$$

# DEFINITION

A dilation of  $f: X \rightarrow Y$  is a span

$$X \xleftarrow{s_1} S \xrightarrow{s_2} Y$$

of coisometries such that  $f = s_2 s_1^+$ .

A dilator of  $f$  is a terminal dilation of  $f$ .

$$\begin{array}{ccccc} & & S & & \\ & \swarrow s_1 & \downarrow s & \searrow s_2 & \\ X & \xleftarrow{p_1} & X \boxplus_f Y & \xrightarrow{p_2} & Y \end{array}$$

# EXAMPLES

Codilators  
in Con

$$\begin{array}{ccccc} & & X \oplus Y & & \\ & \nearrow s_1 & \uparrow s_1 \vee s_2 & \searrow s_2 & \\ X & \longrightarrow & X \vee Y & \longleftarrow & Y \end{array}$$

$$X \vee Y = \overline{\text{Im } s_1 + \text{Im } s_2}$$

# DEFINITION

A dilation of  $f: X \rightarrow Y$  is a span

$$X \xleftarrow{s_1} S \xrightarrow{s_2} Y$$

of coisometries such that  $f = s_2 s_1^+$ .

A dilator of  $f$  is a terminal dilation of  $f$ .

$$\begin{array}{ccccc} & & S & & \\ & \swarrow s_1 & \downarrow s & \searrow s_2 & \\ X & \xleftarrow{p_1} & X \boxplus_f Y & \xrightarrow{p_2} & Y \end{array}$$

# EXAMPLES

Codilators  
in **Con**

$$\begin{array}{ccccc} & & X \oplus Y & & \\ & \nearrow s_1 & \uparrow s_1 \vee s_2 & \searrow s_2 & \\ X & \longrightarrow & X \vee Y & \longleftarrow & Y \end{array}$$

$$X \vee Y = \overline{\text{Im } s_1 + \text{Im } s_2}$$

Codilators  
in **Plnj**

$$\begin{array}{ccc} (X \setminus \text{Dom } f) \sqcup Y & & \\ \nearrow & & \searrow \\ X & & Y \end{array}$$

# DEFINITION

A dilation of  $f: X \rightarrow Y$  is a span

$$X \xleftarrow{s_1} S \xrightarrow{s_2} Y$$

of coisometries such that  $f = s_2 s_1^+$ .

A dilator of  $f$  is a terminal dilation of  $f$ .

$$\begin{array}{ccccc} & & S & & \\ & \swarrow s_1 & \downarrow s & \searrow s_2 & \\ X & \xleftarrow{p_1} & X \boxplus_f Y & \xrightarrow{p_2} & Y \end{array}$$

# EXAMPLES

Codilators  
in **Con**

$$\begin{array}{ccccc} & & X \oplus Y & & \\ & \nearrow s_1 & \uparrow s_1 \vee s_2 & \searrow s_2 & \\ X & \longrightarrow & X \vee Y & \longleftarrow & Y \end{array}$$

$$X \vee Y = \overline{\text{Im } s_1 + \text{Im } s_2}$$

Codilators  
in **Plnj**

$$\begin{array}{ccccc} & & (X \setminus \text{Dom } f) \sqcup Y & & \\ & \nearrow & \downarrow & \swarrow & \\ X & \longrightarrow & \text{Dom } f & \xrightarrow{f|_{\text{Dom } f}} & Y \end{array}$$


# DEFINITION

A dilation of  $f: X \rightarrow Y$  is a span

$$X \xleftarrow{s_1} S \xrightarrow{s_2} Y$$

of coisometries such that  $f = s_2 s_1^+$ .

A dilator of  $f$  is a terminal dilation of  $f$ .



# EXAMPLES


# DEFINITION

A dilation of  $f: X \rightarrow Y$  is a span

$$X \xleftarrow{s_1} S \xrightarrow{s_2} Y$$

of coisometries such that  $f = s_2 s_1^+$ .

A dilator of  $f$  is a terminal dilation of  $f$ .



# EXAMPLES

Dilators  
in FinPS

$$X \xleftarrow{p_1} X \boxplus_f Y \xrightarrow{p_2} Y$$


# DEFINITION

A dilation of  $f: X \rightarrow Y$  is a span

$$X \xleftarrow{s_1} S \xrightarrow{s_2} Y$$

of coisometries such that  $f = s_2 s_1^+$ .

A dilator of  $f$  is a terminal dilation of  $f$ .



# EXAMPLES

Dilators  
in **FinPS**

$$X \boxplus_f Y = \{(x, y) \in X \times Y : P_f(y|x) \neq 0\}$$

$$X \xleftarrow{p_1} X \boxplus_f Y \xrightarrow{p_2} Y$$


# DEFINITION

A dilation of  $f: X \rightarrow Y$  is a span

$$X \xleftarrow{s_1} S \xrightarrow{s_2} Y$$

of coisometries such that  $f = s_2 s_1^+$ .

A dilator of  $f$  is a terminal dilation of  $f$ .



# EXAMPLES

Dilators  
in **FinPS**

$$X \boxplus_f Y = \{(x, y) \in X \times Y : P_f(y|x) \neq 0\}$$

$$P_{x \boxplus_f Y}(x, y) = P_f(y|x)P_x(x)$$

$$X \xleftarrow{p_1} X \boxplus_f Y \xrightarrow{p_2} Y$$


# DEFINITION

A dilation of  $f: X \rightarrow Y$  is a span

$$X \xleftarrow{s_1} S \xrightarrow{s_2} Y$$

of coisometries such that  $f = s_2 s_1^+$ .

A dilator of  $f$  is a terminal dilation of  $f$ .



# EXAMPLES

Dilators  
in **FinPS**

$$X \boxplus_f Y = \{(x, y) \in X \times Y : P_f(y|x) \neq 0\}$$


$$P_{x \boxplus_f Y}(x, y) = P_f(y|x)P_x(x)$$

$$X \xleftarrow{p_1} X \boxplus_f Y \xrightarrow{p_2} Y$$

$$p_1(x, y) = x \quad p_2(x, y) = y$$


Remnants of dilators in *Coisometry*(C)

# Remnants of dilators in Coisometry(C)




factorisation of spans

# Remnants of dilators in Coisometry(C)




factorisation of spans




“pullbacks”

# Remnants of dilators in Coisometry(C)



factorisation of spans




“pullbacks”

# DEFINITION

An **independence category** is a category equipped with a predicate  $\perp$  on commuting squares such that

# DEFINITION



An **independence category** is a category equipped with a predicate  $\perp$  on commuting squares such that



Inspired by Simpson “Equivalence and Independence in Atomic Sheaf Logic”



# DEFINITION

An **independence category** is a category equipped with a predicate  $\perp$  on commuting squares such that





# DEFINITION

An **independence category** is a category equipped with a predicate  $\perp$  on commuting squares such that



# DEFINITION

An independence category is a category equipped with a predicate  $\perp$  on commuting squares such that



# LEMMA

If  $\mathbf{C}$  is a  $t$ -category, then

$\text{Coisometry}(\mathbf{C})$  is an independence category where

$$\begin{array}{ccc} & \xrightarrow{a} & \\ f \downarrow & \perp & \downarrow g \\ & \xrightarrow{s} & \end{array} \quad \text{if} \quad \begin{aligned} ga &= sf \\ af^+ &= g^+s \end{aligned}$$

# LEMMA


If  $\mathbf{C}$  is a  $t$ -category, then  
 $\mathbf{Coisometry}(\mathbf{C})$  is an independence  
 category where

$$\begin{array}{ccc} & \xrightarrow{a} & \\ f \downarrow & \perp & \downarrow g \\ & \xrightarrow{s} & \end{array} \quad \text{if} \quad \begin{aligned} ga &= sf \\ af^+ &= g^+s \end{aligned}$$

# EXAMPLES

In  $\mathbf{Isometry}(\mathbf{Con})$ ,  
 $\perp$  captures relative orthogonality.

$$\begin{array}{ccccc} (0, y, z) & \longleftrightarrow & (y, z) & & \\ (x, 0, z) & \mathbb{R}^3 & \longleftarrow & \mathbb{R}^2 & (0, z) \\ \uparrow & & \uparrow & & \uparrow \\ (x, z) & \mathbb{R}^2 & \longleftarrow & \mathbb{R} & z \\ (0, z) & & \longleftrightarrow & & z \end{array}$$



# LEMMA

If  $\mathbf{C}$  is a  $t$ -category, then



$\text{Coisometry}(\mathbf{C})$  is an independence category where

$$\begin{array}{ccc} & \xrightarrow{a} & \\ f \downarrow & \perp & \downarrow g \\ & \xrightarrow{s} & \end{array} \quad \text{if} \quad \begin{array}{l} ga = sf \\ af^+ = g^+s \end{array}$$

# EXAMPLES

In  $\text{Isometry}(\text{Plnj})$ ,

$\perp$  captures relative disjointness.



# LEMMA

If  $C$  is a  $t$ -category, then

$\text{Coisometry}(C)$  is an independence category where

$$\begin{array}{ccc} & \xrightarrow{a} & \\ f \downarrow & \perp & \downarrow g \\ & \xrightarrow{s} & \end{array} \quad \text{if} \quad \begin{aligned} ga &= sf \\ af^+ &= g^+s \end{aligned}$$

# EXAMPLES

In  $\text{Coisometry}(\text{FinPS})$ ,

$$\begin{array}{ccc} Z & \xrightarrow{g} & Y \\ f \downarrow & \perp & \downarrow t \\ X & \xrightarrow{s} & A \end{array}$$

$\Updownarrow$

$$sf = tg \text{ and}$$

$$\mathbb{P}_z(f^{-1}\{x\} \cap g^{-1}\{y\}) = \frac{\mathbb{P}_x(x)\mathbb{P}_Y(y)}{\mathbb{P}_A(a)}$$

for all  $x \in s^{-1}\{a\}$  and  $y \in t^{-1}\{a\}$

# DEFINITION

In an independence category, an independent pullback is a square

$$\begin{array}{ccc} & g & \\ f \downarrow & \perp & \downarrow t \\ & \perp & \\ s \rightarrow & & \end{array}$$

such that for all

$$\begin{array}{ccccc} & b & & & \\ a \downarrow & \perp & \downarrow t & & \\ & \perp & & & \\ r \rightarrow & \xrightarrow{s} & & & \end{array}$$

exists unique  $c$  such that

$$\begin{array}{ccc} c & \xrightarrow{\quad} & \\ a \downarrow & \perp & \downarrow f \\ & \perp & \\ r & \xrightarrow{\quad} & \end{array} \quad \text{and} \quad b = gc.$$

# DEFINITION

In an independence category, an independent pullback is a square

$$\begin{array}{ccc} & \xrightarrow{g} & \\ f \downarrow & \perp & \downarrow t \\ & \xrightarrow{s} & \end{array}$$

such that for all

$$\begin{array}{ccccc} & & b & & \\ & \searrow & \perp & \swarrow & \\ a \downarrow & & & & t \downarrow \\ & r \longrightarrow & s \longrightarrow & & \end{array}$$

exists unique  $c$  such that

$$\begin{array}{ccc} c & \xrightarrow{\quad} & \\ a \downarrow & \perp & \downarrow f \\ & \xrightarrow{r} & \end{array} \quad \text{and} \quad b = gc.$$

Weak independent pullbacks are similar, but with  $r = 1$ .

# DEFINITION

In an independence category, an independent pullback is a square

$$\begin{array}{ccc} & \xrightarrow{g} & \\ f \downarrow & \perp & \downarrow t \\ & \xrightarrow{s} & \end{array}$$

such that for all

$$\begin{array}{ccccc} & & b & & \\ & \xrightarrow{\quad} & & \xrightarrow{\quad} & \\ a \downarrow & \perp & & \downarrow t & \\ & \xrightarrow{r} & \xrightarrow{s} & & \end{array}$$

exists unique  $c$  such that

$$\begin{array}{ccc} c & \xrightarrow{\quad} & \\ a \downarrow & \perp & \downarrow f \\ & \xrightarrow{r} & \end{array} \quad \text{and} \quad b = gc.$$


Weak independent pullbacks are similar, but with  $r = 1$ .

## LEMMA


If  $\mathbf{C}$  is a  $t$ -category with dilators, then  $\mathbf{Coisometry}(\mathbf{C})$  has weak independent pullbacks.

Inspired by Simpson “Equivalence and Independence in Atomic Sheaf Logic”

# Remnants of dilators in Coisometry(C)




factorisation of spans




“pullbacks”

# Remnants of dilators in Coisometry(C)



factorisation of spans



“pullbacks”


# DEFINITION

A morphism  $e$  is **strong epic** if it is left orthogonal to the jointly monic spans.

(Non-standard definition)

# DEFINITION


A morphism  $e$  is **strong epic** if it is left orthogonal to the jointly monic spans.



(Non-standard definition)

# DEFINITION


A morphism  $e$  is **strong epic** if it is left orthogonal to the jointly monic spans.



(Non-standard definition)

# DEFINITION

A morphism  $e$  is **strong epic** if it is left orthogonal to the jointly monic spans.



(Non-standard definition)


# LEMMA

Let  $\mathbf{C}$  be a  $\mathbb{t}$ -category with dilators. Then

- (i) A span in  $\mathbf{Coisometry}(\mathbf{C})$  is jointly monic if and only if it is a dilator in  $\mathbf{C}$ .

# DEFINITION

A morphism  $e$  is **strong epic** if it is left orthogonal to the jointly monic spans.



(Non-standard definition)


# LEMMA

Let  $\mathbf{C}$  be a  $\mathbb{t}$ -category with dilators. Then

- (i) A span in  $\mathbf{Coisometry}(\mathbf{C})$  is jointly monic if and only if it is a dilator in  $\mathbf{C}$ .
- (ii) Every morphism in  $\mathbf{Coisometry}(\mathbf{C})$  is strong epic.

# DEFINITION

A morphism  $e$  is **strong epic** if it is left orthogonal to the jointly monic spans.



(Non-standard definition)

# LEMMA

Let  $\mathbf{C}$  be a  $\mathbb{t}$ -category with dilators. Then

- (i) A span in **Coisometry(C)** is jointly monic if and only if it is a dilator in  $\mathbf{C}$ .
- (ii) Every morphism in **Coisometry(C)** is strong epic.
- (iii) Every span in **Coisometry(C)** has a (strong epic, jointly monic) factorisation.

# DEFINITION

An independence category is regular-ish if

# DEFINITION

An independence category is **regular-ish** if

- (i) it has weak independent pullbacks,

# DEFINITION

An independence category is **regular-ish** if

(i) it has weak independent pullbacks,

(ii) every span factors as a strong epi followed by a jointly monic span,

# DEFINITION

An independence category is **regular-ish** if

(i) it has weak independent pullbacks,

(ii) every span factors as a strong epi  
followed by a jointly monic span,

(iii) every morphism is strong epic

# DEFINITION

An independence category is **regular-ish** if

- (i) it has weak independent pullbacks,  
every span factors as a strong epi
- (ii) followed by a jointly monic span,
- (iii) every morphism is strong epic

# THEOREM

Let  $\mathbf{C}$  be a  $t$ -category with dilators.  
Then  $\mathbf{Coisometry}(\mathbf{C})$  is regular-ish.

# DEFINITION

An independence category is **regular-ish** if

- (i) it has weak independent pullbacks,  
every span factors as a strong epi
- (ii) followed by a jointly monic span,
- (iii) every morphism is strong epic

# THEOREM

Let  $\mathbf{C}$  be a  $t$ -category with dilators.  
Then  $\mathbf{Coisometry}(\mathbf{C})$  is regular-ish.

# LEMMA

In a regular-ish independence category  
every weak independent pullback is  
an independent pullback.

# DEFINITION

Let  $D$  be a regular-ish independence category. Define  $\text{Rel}(D)$  as follows:

- objects are objects of  $D$
- morphisms are relations in (isomorphism classes of jointly monic spans)
- composition is by independent pullback and span factorisation

# DEFINITION

Let  $D$  be a regular-ish independence category. Define  $\text{Rel}(D)$  as follows:

- objects are objects of  $D$
- morphisms are relations in (isomorphism classes of jointly monic spans)
- composition is by independent pullback and span factorisation

# THEOREM

- $\text{Rel}(D)$  is a dagger category with dilators.
- $\text{Coisometry}(\text{Rel}(D)) \cong D$
- $\text{Rel}(\text{Coisometry}(C)) \cong C$

What is the connection between  
dilators and tabulators?

**Rel** is a poset-enriched  $+$ -category.

$$f \leq g \quad \text{implies} \quad f^+ \leq g^+$$

**Rel** is a poset-enriched  $+$ -category.

$$f \leq g \quad \text{implies} \quad f^+ \leq g^+$$

A partial map is a morphism  $f$  such that  $ff^+ \leq 1$ .

**Rel** is a poset-enriched  $+$ -category.

$$f \leq g \quad \text{implies} \quad f^+ \leq g^+$$

A partial map is a morphism  $f$  such that  $ff^+ \leq 1$ .

Let  $f: X \rightarrow Y$  be a relation.

**Rel** is a poset-enriched  $+$ -category.

$$f \leq g \quad \text{implies} \quad f^+ \leq g^+$$

A partial map is a morphism  $f$  such that  $ff^+ \leq 1$ .

Let  $f: X \rightarrow Y$  be a relation.


$$X \xleftarrow{[1 \ p_1 \ \perp]} X \sqcup f \sqcup Y \xrightarrow{\perp \ p_2 \ 1} Y$$

**Rel** is a poset-enriched  $+$ -category.

$$f \leq g \quad \text{implies} \quad f^+ \leq g^+$$

A partial map is a morphism  $f$  such that  $ff^+ \leq 1$ .

Let  $f: X \rightarrow Y$  be a relation.




**Rel** is a poset-enriched  $+$ -category.

$$f \leq g \quad \text{implies} \quad f^+ \leq g^+$$

A partial map is a morphism  $f$  such that  $ff^+ \leq 1$ .

Let  $f: X \rightarrow Y$  be a relation.




$\text{Rel}$  is a poset-enriched  $+$ -category.

$$f \leq g \quad \text{implies} \quad f^+ \leq g^+$$

A partial map is a morphism  $f$  such that  $ff^+ \leq 1$ .

Let  $f: X \rightarrow Y$  be a relation.




A morphism  $f: (X, x) \rightarrow (Y, y)$  in  $\text{Rel}_*$  is a relation  $f: X \rightarrow Y$  such that  $(x, y) \in f$ . 10

$\text{Rel}$  is a poset-enriched  $+$ -category.

$$f \leq g \quad \text{implies} \quad f^+ \leq g^+$$

A partial map is a morphism  $f$  such that  $ff^+ \leq 1$ .

Let  $f: X \rightarrow Y$  be a relation.



A morphism  $f: (X, x) \rightarrow (Y, y)$  in  $\text{Rel}_{*}$  is a relation  $f: X \rightarrow Y$  such that  $(x, y) \in f$ .


The tabulator of  $f: (X, x) \rightarrow (Y, y)$  is  $(X, x) \xleftarrow{P_1} (f, (x, y)) \xrightarrow{P_2} (Y, y)$

$\text{Rel}$  is a poset-enriched  $+$ -category.

$$f \leq g \quad \text{implies} \quad f^+ \leq g^+$$

A partial map is a morphism  $f$  such that  $ff^+ \leq 1$ .

Let  $f: X \rightarrow Y$  be a relation.



A morphism  $f: (X, x) \rightarrow (Y, y)$  in  $\text{Rel}_{*}$  is a relation  $f: X \rightarrow Y$  such that  $(x, y) \in f$ .

The tabulator of  $f: (X, x) \rightarrow (Y, y)$  is

$$(X, x) \xleftarrow{P_1} (f, (x, y)) \xrightarrow{P_2} (Y, y)$$


Consider  $\text{Rel} \rightarrow \text{Rel}_{*}$ :

$\text{Rel}$  is a poset-enriched  $+$ -category.

$$f \leq g \quad \text{implies} \quad f^+ \leq g^+$$

A partial map is a morphism  $f$  such that  $ff^+ \leq 1$ .

Let  $f: X \rightarrow Y$  be a relation.



A morphism  $f: (X, x) \rightarrow (Y, y)$  in  $\text{Rel}_*$  is a relation  $f: X \rightarrow Y$  such that  $(x, y) \in f$ .

The tabulator of  $f: (X, x) \rightarrow (Y, y)$  is

$$(X, x) \xleftarrow{P_1} (f, (x, y)) \xrightarrow{P_2} (Y, y)$$

Consider  $\text{Rel} \rightarrow \text{Rel}_*$ :


$$\begin{matrix} X \\ \Downarrow \\ (X \sqcup \{\ast\}, \ast) \end{matrix}$$

$\text{Rel}$  is a poset-enriched  $+$ -category.

$$f \leq g \quad \text{implies} \quad f^+ \leq g^+$$

A partial map is a morphism  $f$  such that  $ff^+ \leq 1$ .

Let  $f: X \rightarrow Y$  be a relation.



A morphism  $f: (X, x) \rightarrow (Y, y)$  in  $\text{Rel}_{*}$  is a relation  $f: X \rightarrow Y$  such that  $(x, y) \in f$ .

The tabulator of  $f: (X, x) \rightarrow (Y, y)$  is

$$(X, x) \xleftarrow{P_1} (f, (x, y)) \xrightarrow{P_2} (Y, y)$$

Consider  $\text{Rel} \rightarrow \text{Rel}_{*}$ :

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ \downarrow & & \downarrow \\ (X \sqcup \{\ast\}, \ast) & \xrightarrow{\left[ \begin{matrix} f & \{\ast\} \times Y \\ X \times \{\ast\} & \{(\ast, \ast)\} \end{matrix} \right]} & (Y \sqcup \{\ast\}, \ast) \end{array}$$

# PROJECTS

With Heunen, Perrone and Stein

---

m.dimeglio@ed.ac.uk

<https://mdimeglio.github.io>

# PROJECTS

With Heunen, Perrone and Stein

Characterise  
the category of  
Hilbert spaces  
and *coisometries*

---

m.dimeglio@ed.ac.uk

<https://mdimeglio.github.io>

All coisometries  
have a quantum  
interpretation

# PROJECTS

With Heunen, Perrone and Stein

Characterise  
the category of  
Hilbert spaces  
and coisometries

Not a  $\mathbb{C}$ -category

---

m.dimeglio@ed.ac.uk

<https://mdimeglio.github.io>

All coisometries  
have a quantum  
interpretation

# PROJECTS

With Heunen, Perrone and Stein

Characterise  
the category of  
Hilbert spaces  
and coisometries

Characterise  
a category of  
probability  
spaces

Not a  $\mathbb{C}$ -category

---

m.dimeglio@ed.ac.uk

<https://mdimeglio.github.io>

All coisometries  
have a quantum  
interpretation

# PROJECTS

With Heunen, Perrone and Stein

The first  
characterisation  
in categorical  
probability

Characterise  
the category of  
Hilbert spaces  
and coisometries

Characterise  
a category of  
probability  
spaces

Not a  $\mathbb{C}$ -category

Not Markov categories

---

m.dimeglio@ed.ac.uk

<https://mdimeglio.github.io>