

Lecture 20: Max-Flow Min-Cut

Jessica Sorrell

November 6, 2025

601.433/633 Introduction to Algorithms

Slides by Michael Dinitz

Introduction

Flow Network:

- ▶ Directed graph $G = (V, E)$
- ▶ Capacities $c : E \rightarrow \mathbb{R}_{\geq 0}$ (simplify notation: $c(x, y) = 0$ if $(x, y) \notin E$)
- ▶ Source $s \in V$, sink $t \in V$

Today: flows and cuts

- ▶ Flow: “sending stuff” from s to t
- ▶ Cut: separating t from s

Turn out to be very related!

Today: some algorithms but not efficient. Mostly structure. Better algorithms Tuesday.

Flows

Intuition: send “stuff” from s to t

- ▶ Water in a city water system, traffic along roads, trains along tracks, ...

Flows

Intuition: send “stuff” from s to t

- ▶ Water in a city water system, traffic along roads, trains along tracks, ...

Definition

An (s, t) -flow is a function $f : E \rightarrow \mathbb{R}_{\geq 0}$ such that

$$\sum_{u:(u,v) \in E} f(u,v) = \sum_{u:(v,u) \in E} f(v,u)$$

for all $v \in V \setminus \{s, t\}$. This constraint is known as *flow conservation*.

Flows

Intuition: send “stuff” from s to t

- ▶ Water in a city water system, traffic along roads, trains along tracks, ...

Definition

An (s, t) -flow is a function $f : E \rightarrow \mathbb{R}_{\geq 0}$ such that

$$\sum_{u:(u,v) \in E} f(u, v) = \sum_{u:(v,u) \in E} f(v, u)$$

for all $v \in V \setminus \{s, t\}$. This constraint is known as *flow conservation*.

Value of flow $|f|$: “total amount of stuff sent from s to t ”

Flows

Intuition: send “stuff” from s to t

- ▶ Water in a city water system, traffic along roads, trains along tracks, ...

Definition

An (s, t) -flow is a function $f : E \rightarrow \mathbb{R}_{\geq 0}$ such that

$$\sum_{u:(u,v) \in E} f(u, v) = \sum_{u:(v,u) \in E} f(v, u)$$

for all $v \in V \setminus \{s, t\}$. This constraint is known as *flow conservation*.

Value of flow $|f|$: “total amount of stuff sent from s to t ”

$$|f| = \sum_{u:(s,u) \in E} f(s, u) - \sum_{u:(u,s) \in E} f(u, s) = \sum_{u:(u,t) \in E} f(u, t) - \sum_{u:(t,u) \in E} f(t, u)$$

Feasible Flows

Capacity constraints: $\mathbf{0} \leq \mathbf{f}(u, v) \leq \mathbf{c}(u, v)$ for all $(u, v) \in V \times V$

Feasible Flows

Capacity constraints: $0 \leq f(u, v) \leq c(u, v)$ for all $(u, v) \in V \times V$

Definitions:


- ▶ An (s, t) -flow satisfying capacity constraints is a *feasible* flow.
- ▶ If $f(e) = c(e)$ then f *saturates* e .
- ▶ If $f(e) = 0$ then f *avoids* e .

Feasible Flows

Capacity constraints: $0 \leq f(u, v) \leq c(u, v)$ for all $(u, v) \in V \times V$

Definitions:

- ▶ An (s, t) -flow satisfying capacity constraints is a *feasible* flow.
- ▶ If $f(e) = c(e)$ then f *saturates* e .
- ▶ If $f(e) = 0$ then f *avoids* e .




An (s, t) -flow with value 10. Each edge is labeled with its flow/capacity.

Feasible Flows

Capacity constraints: $0 \leq f(u, v) \leq c(u, v)$ for all $(u, v) \in V \times V$

Definitions:

- ▶ An (s, t) -flow satisfying capacity constraints is a *feasible* flow.
- ▶ If $f(e) = c(e)$ then f *saturates* e .
- ▶ If $f(e) = 0$ then f *avoids* e .



An (s, t) -flow with value 10. Each edge is labeled with its flow/capacity.

Problem we'll talk about: find feasible flow of maximum value (max flow)

Cuts

Definition

- ▶ An (s, t) -cut is a partition of V into (S, \bar{S}) such that $s \in S, t \notin S$

Cuts

Definition

- ▶ An (s, t) -cut is a partition of V into (S, \bar{S}) such that $s \in S$, $t \notin S$
- ▶ The *capacity* of an (s, t) -cut (S, \bar{S}) is


$$\text{cap}(S, \bar{S}) = \sum_{(u,v) \in E: u \in S, v \in \bar{S}} c(u, v) = \sum_{u \in S} \sum_{v \in \bar{S}} c(u, v)$$

Cuts

Definition

- ▶ An **(s, t) -cut** is a partition of V into (S, \bar{S}) such that $s \in S$, $t \notin S$
- ▶ The **capacity** of an (s, t) -cut (S, \bar{S}) is

$$cap(S, \bar{S}) = \sum_{(u,v) \in E: u \in S, v \in \bar{S}} c(u, v) = \sum_{u \in S} \sum_{v \in \bar{S}} c(u, v)$$




Cuts

Definition

- ▶ An **(s, t) -cut** is a partition of V into (S, \bar{S}) such that $s \in S$, $t \notin S$
- ▶ The **capacity** of an (s, t) -cut (S, \bar{S}) is

$$cap(S, \bar{S}) = \sum_{(u,v) \in E: u \in S, v \in \bar{S}} c(u, v) = \sum_{u \in S} \sum_{v \in \bar{S}} c(u, v)$$



Problem we'll talk about: find (s, t) -cut of minimum capacity (min cut)

Warmup Theorem

Theorem

Let \mathbf{f} be a feasible (s, t) -flow, and let (S, \bar{S}) be an (s, t) -cut. Then $|\mathbf{f}| \leq \mathbf{cap}(S, \bar{S})$.

Warmup Theorem

Theorem

Let \mathbf{f} be a feasible (s, t) -flow, and let (S, \bar{S}) be an (s, t) -cut. Then $|\mathbf{f}| \leq \mathbf{cap}(S, \bar{S})$.

$$|\mathbf{f}| = \sum_{v \in V} \mathbf{f}(s, v) - \sum_{v \in V} \mathbf{f}(v, s) \quad (\text{definition})$$

Warmup Theorem

Theorem

Let \mathbf{f} be a feasible (s, t) -flow, and let (S, \bar{S}) be an (s, t) -cut. Then $|\mathbf{f}| \leq \mathbf{cap}(S, \bar{S})$.

$$|\mathbf{f}| = \sum_{v \in V} \mathbf{f}(s, v) - \sum_{v \in V} \mathbf{f}(v, s) \quad (\text{definition})$$

$$= \sum_{u \in S} \left(\sum_{v \in V} \mathbf{f}(u, v) - \sum_{v \in V} \mathbf{f}(v, u) \right) \quad (\text{flow conservation constraints})$$

Warmup Theorem

Theorem

Let \mathbf{f} be a feasible (s, t) -flow, and let $(\mathbf{S}, \bar{\mathbf{S}})$ be an (s, t) -cut. Then $|\mathbf{f}| \leq \mathbf{cap}(\mathbf{S}, \bar{\mathbf{S}})$.

$$|\mathbf{f}| = \sum_{v \in V} \mathbf{f}(s, v) - \sum_{v \in V} \mathbf{f}(v, s) \quad (\text{definition})$$

$$= \sum_{u \in \mathbf{S}} \left(\sum_{v \in V} \mathbf{f}(u, v) - \sum_{v \in V} \mathbf{f}(v, u) \right) \quad (\text{flow conservation constraints})$$

$$= \sum_{u \in \mathbf{S}} \left(\sum_{v \in \bar{\mathbf{S}}} \mathbf{f}(u, v) - \sum_{v \in \bar{\mathbf{S}}} \mathbf{f}(v, u) \right) \quad (\text{remove terms which cancel})$$

Warmup Theorem

Theorem

Let \mathbf{f} be a feasible (s, t) -flow, and let (S, \bar{S}) be an (s, t) -cut. Then $|\mathbf{f}| \leq \mathbf{cap}(S, \bar{S})$.

$$|\mathbf{f}| = \sum_{v \in V} \mathbf{f}(s, v) - \sum_{v \in V} \mathbf{f}(v, s) \quad (\text{definition})$$

$$= \sum_{u \in S} \left(\sum_{v \in V} \mathbf{f}(u, v) - \sum_{v \in V} \mathbf{f}(v, u) \right) \quad (\text{flow conservation constraints})$$

$$= \sum_{u \in S} \left(\sum_{v \in \bar{S}} \mathbf{f}(u, v) - \sum_{v \in \bar{S}} \mathbf{f}(v, u) \right) \quad (\text{remove terms which cancel})$$

$$\leq \sum_{u \in S} \sum_{v \in \bar{S}} \mathbf{f}(u, v) \quad (\text{flow is nonnegative})$$

Warmup Theorem

Theorem

Let \mathbf{f} be a feasible (s, t) -flow, and let (S, \bar{S}) be an (s, t) -cut. Then $|\mathbf{f}| \leq \text{cap}(S, \bar{S})$.

$$|\mathbf{f}| = \sum_{v \in V} \mathbf{f}(s, v) - \sum_{v \in V} \mathbf{f}(v, s) \quad (\text{definition})$$

$$= \sum_{u \in S} \left(\sum_{v \in V} \mathbf{f}(u, v) - \sum_{v \in V} \mathbf{f}(v, u) \right) \quad (\text{flow conservation constraints})$$

$$= \sum_{u \in S} \left(\sum_{v \in \bar{S}} \mathbf{f}(u, v) - \sum_{v \in \bar{S}} \mathbf{f}(v, u) \right) \quad (\text{remove terms which cancel})$$

$$\leq \sum_{u \in S} \sum_{v \in \bar{S}} \mathbf{f}(u, v) \quad (\text{flow is nonnegative})$$

$$\leq \sum_{u \in S} \sum_{v \in \bar{S}} c(u, v) = \text{cap}(S, \bar{S}) \quad (\text{flow is feasible})$$

Max-Flow Min-Cut

Corollary

If \mathbf{f} avoids every $\bar{\mathbf{S}} \rightarrow \mathbf{S}$ edge and saturates every $\mathbf{S} \rightarrow \bar{\mathbf{S}}$ edge, then \mathbf{f} is a maximum flow and $(\mathbf{S}, \bar{\mathbf{S}})$ is a minimum cut.

Max-Flow Min-Cut

Corollary

If \mathbf{f} avoids every $\bar{\mathbf{S}} \rightarrow \mathbf{S}$ edge and saturates every $\mathbf{S} \rightarrow \bar{\mathbf{S}}$ edge, then \mathbf{f} is a maximum flow and $(\mathbf{S}, \bar{\mathbf{S}})$ is a minimum cut.

Theorem (Max-Flow Min-Cut Theorem)

In any flow network, value of max (\mathbf{s}, \mathbf{t}) -flow = capacity of min (\mathbf{s}, \mathbf{t}) -cut.

Max-Flow Min-Cut

Corollary

If \mathbf{f} avoids every $\bar{\mathbf{S}} \rightarrow \mathbf{S}$ edge and saturates every $\mathbf{S} \rightarrow \bar{\mathbf{S}}$ edge, then \mathbf{f} is a maximum flow and $(\mathbf{S}, \bar{\mathbf{S}})$ is a minimum cut.

Theorem (Max-Flow Min-Cut Theorem)


In any flow network, value of max (\mathbf{s}, \mathbf{t}) -flow = capacity of min (\mathbf{s}, \mathbf{t}) -cut.

Spend rest of today proving this.

- ▶ Many different valid proofs.
- ▶ We'll see a classical proof which will naturally lead to algorithms for these problems.


One Direction

Cycles of length 2 will turn out to be annoying. Get rid of them.



One Direction

Cycles of length 2 will turn out to be annoying. Get rid of them.



- ▶ Doesn't change max-flow or min-cut
- ▶ Increases #edges by constant factor, # nodes to original # edges.

Residual

Let \mathbf{f} be feasible (\mathbf{s}, \mathbf{t}) -flow. Define *residual capacities*:

$$c_f(\mathbf{u}, \mathbf{v}) = \begin{cases} c(\mathbf{u}, \mathbf{v}) - \mathbf{f}(\mathbf{u}, \mathbf{v}) & \text{if } (\mathbf{u}, \mathbf{v}) \in E \\ 0 & \text{otherwise} \end{cases}$$

Residual

Let \mathbf{f} be feasible (\mathbf{s}, \mathbf{t}) -flow. Define *residual capacities*:



$$c_f(\mathbf{u}, \mathbf{v}) = \begin{cases} c(\mathbf{u}, \mathbf{v}) - \mathbf{f}(\mathbf{u}, \mathbf{v}) & \text{if } (\mathbf{u}, \mathbf{v}) \in E \\ \mathbf{f}(\mathbf{v}, \mathbf{u}) & \text{if } (\mathbf{v}, \mathbf{u}) \in E \\ 0 & \text{otherwise} \end{cases}$$

Residual

Let f be feasible (s, t) -flow. Define *residual capacities*:

$$c_f(u, v) = \begin{cases} c(u, v) - f(u, v) & \text{if } (u, v) \in E \\ f(v, u) & \text{if } (v, u) \in E \\ 0 & \text{otherwise} \end{cases}$$

Residual Graph: $G_f = (V, E_f)$ where $(u, v) \in E_f$ if $c_f(u, v) > 0$.



A flow f in a weighted graph G and the corresponding residual graph G_f .

Start of Proof

Let \mathbf{f} be a max (\mathbf{s}, \mathbf{t}) -flow with residual graph $\mathbf{G}_{\mathbf{f}}$.

Want to Show: There is a cut $(\mathbf{S}, \bar{\mathbf{S}})$ with $\mathbf{cap}(\mathbf{S}, \bar{\mathbf{S}}) = |\mathbf{f}|$.

Start of Proof

Let f be a max (s, t) -flow with residual graph G_f .

Want to Show: There is a cut (S, \bar{S}) with $cap(S, \bar{S}) = |f|$.

Case 1: There is no $s \rightarrow t$ path in G_f


Start of Proof

Let f be a max (s, t) -flow with residual graph G_f .

Want to Show: There is a cut (S, \bar{S}) with $\text{cap}(S, \bar{S}) = |f|$.

Case 1: There is no $s \rightarrow t$ path in G_f

Let $S = \{\text{vertices reachable from } s \text{ in } G_f\}$




Start of Proof

Let f be a max (s, t) -flow with residual graph G_f .

Want to Show: There is a cut (S, \bar{S}) with $\text{cap}(S, \bar{S}) = |f|$.

Case 1: There is no $s \rightarrow t$ path in G_f

Let $S = \{\text{vertices reachable from } s \text{ in } G_f\}$



► (S, \bar{S}) an (s, t) -cut. ✓


Start of Proof

Let f be a max (s, t) -flow with residual graph G_f .

Want to Show: There is a cut (S, \bar{S}) with $\text{cap}(S, \bar{S}) = |f|$.

Case 1: There is no $s \rightarrow t$ path in G_f

Let $S = \{\text{vertices reachable from } s \text{ in } G_f\}$



- ▶ (S, \bar{S}) an (s, t) -cut. ✓
- ▶ $c_f(a, b) =$


Start of Proof

Let f be a max (s, t) -flow with residual graph G_f .

Want to Show: There is a cut (S, \bar{S}) with $\text{cap}(S, \bar{S}) = |f|$.

Case 1: There is no $s \rightarrow t$ path in G_f

Let $S = \{\text{vertices reachable from } s \text{ in } G_f\}$



- ▶ (S, \bar{S}) an (s, t) -cut. ✓
- ▶ $c_f(a, b) = 0$


Start of Proof

Let f be a max (s, t) -flow with residual graph G_f .

Want to Show: There is a cut (S, \bar{S}) with $\text{cap}(S, \bar{S}) = |f|$.

Case 1: There is no $s \rightarrow t$ path in G_f

Let $S = \{\text{vertices reachable from } s \text{ in } G_f\}$



- ▶ (S, \bar{S}) an (s, t) -cut. ✓
- ▶ $c_f(a, b) = 0$
 - $\implies c(a, b) - f(a, b) = 0$
 - $\implies c(a, b) = f(a, b)$


Start of Proof

Let f be a max (s, t) -flow with residual graph G_f .

Want to Show: There is a cut (S, \bar{S}) with $\text{cap}(S, \bar{S}) = |f|$.

Case 1: There is no $s \rightarrow t$ path in G_f

Let $S = \{\text{vertices reachable from } s \text{ in } G_f\}$



- ▶ (S, \bar{S}) an (s, t) -cut. ✓
- ▶ $c_f(a, b) = 0$
 $\implies c(a, b) - f(a, b) = 0$
 $\implies c(a, b) = f(a, b)$
- ▶ $c_f(y, x) =$


Start of Proof

Let f be a max (s, t) -flow with residual graph G_f .

Want to Show: There is a cut (S, \bar{S}) with $\text{cap}(S, \bar{S}) = |f|$.

Case 1: There is no $s \rightarrow t$ path in G_f

Let $S = \{\text{vertices reachable from } s \text{ in } G_f\}$



- ▶ (S, \bar{S}) an (s, t) -cut. ✓
- ▶ $c_f(a, b) = 0$
 $\implies c(a, b) - f(a, b) = 0$
 $\implies c(a, b) = f(a, b)$
- ▶ $c_f(y, x) = 0$


Start of Proof

Let f be a max (s, t) -flow with residual graph G_f .

Want to Show: There is a cut (S, \bar{S}) with $\text{cap}(S, \bar{S}) = |f|$.

Case 1: There is no $s \rightarrow t$ path in G_f

Let $S = \{\text{vertices reachable from } s \text{ in } G_f\}$



- ▶ (S, \bar{S}) an (s, t) -cut. ✓
- ▶ $c_f(a, b) = 0$
 $\implies c(a, b) - f(a, b) = 0$
 $\implies c(a, b) = f(a, b)$
- ▶ $c_f(y, x) = 0$
 $\implies f(x, y) = 0$


Start of Proof

Let f be a max (s, t) -flow with residual graph G_f .

Want to Show: There is a cut (S, \bar{S}) with $\text{cap}(S, \bar{S}) = |f|$.

Case 1: There is no $s \rightarrow t$ path in G_f

Let $S = \{\text{vertices reachable from } s \text{ in } G_f\}$



- ▶ (S, \bar{S}) an (s, t) -cut. ✓
- ▶ $c_f(a, b) = 0$
 $\implies c(a, b) - f(a, b) = 0$
 $\implies c(a, b) = f(a, b)$
- ▶ $c_f(y, x) = 0$
 $\implies f(x, y) = 0$

f saturates $S \rightarrow \bar{S}$ edges, avoids $\bar{S} \rightarrow S$ edges $\implies \text{cap}(S, \bar{S}) = |f|$ by corollary

Case 2

Suppose \exists an $s \rightarrow t$ path P in G_f .

- ▶ Called an *augmenting path*

Idea: show that we can “push” more flow along P , so f not a max flow. Contradiction, can't be in this case.

Case 2



Suppose \exists an $s \rightarrow t$ path P in G_f .

- ▶ Called an *augmenting path*



Idea: show that we can “push” more flow along P , so f not a max flow. Contradiction, can't be in this case.

- ▶ Foreshadowing: augmenting path allows us to send more flow. Algorithm to increase flow!

Intuition



A flow f in a weighted graph G and the corresponding residual graph G_f .



An augmenting path in G_f with value $F = 5$ and the augmented flow f' .

Formalities

Let \mathbf{P} be (simple) augmenting path in \mathbf{G}_f . Let $\mathbf{F} = \min_{e \in \mathbf{P}} c_f(e)$.

Formalities

Let P be (simple) augmenting path in G_f . Let $F = \min_{e \in P} c_f(e)$.

Define new flow f' : for all $(u, v) \in E$, let

$$f'(u, v) = \begin{cases} f(u, v) + F & \text{if } (u, v) \text{ in } P \\ f(u, v) - F & \text{if } (v, u) \text{ in } P \\ f(u, v) & \text{otherwise} \end{cases}$$

Formalities

Let P be (simple) augmenting path in G_f . Let $F = \min_{e \in P} c_f(e)$.

Define new flow f' : for all $(u, v) \in E$, let

$$f'(u, v) = \begin{cases} f(u, v) + F & \text{if } (u, v) \text{ in } P \\ f(u, v) - F & \text{if } (v, u) \text{ in } P \\ f(u, v) & \text{otherwise} \end{cases}$$

Claim: f' is a feasible (s, t) -flow with $|f'| > |f|$.

Formalities

Let P be (simple) augmenting path in G_f . Let $F = \min_{e \in P} c_f(e)$.

Define new flow f' : for all $(u, v) \in E$, let

$$f'(u, v) = \begin{cases} f(u, v) + F & \text{if } (u, v) \text{ in } P \\ f(u, v) - F & \text{if } (v, u) \text{ in } P \\ f(u, v) & \text{otherwise} \end{cases}$$

Claim: f' is a feasible (s, t) -flow with $|f'| > |f|$.


Plan: prove (sketch) each subclaim individually

- ▶ $|f'| > |f|$
- ▶ f' an (s, t) -flow (flow conservation)
- ▶ f' feasible (obeys capacities)


$$|f'| > |f|$$

Consider first edge of P (out of s), say (s, v_1)

- ▶ If $(s, v_1) \in E$, then $f'(s, v_1) = f(s, v_1) + F$
- ▶ If $(v_1, s) \in E$ then $f'(v_1, s) = f(v_1, s) - F$




or




$$|f'| > |f|$$

Consider first edge of P (out of s), say (s, v_1)

- ▶ If $(s, v_1) \in E$, then $f'(s, v_1) = f(s, v_1) + F$
- ▶ If $(v_1, s) \in E$ then $f'(v_1, s) = f(v_1, s) - F$



or



$$|f'| = \sum_u f'(s, u) - \sum_u f'(u, s) = |f| + F > |f|$$

f' obeys flow conservation

Consider some $u \in V \setminus \{s, t\}$.

f' obeys flow conservation


Consider some $u \in V \setminus \{s, t\}$.

- ▶ If $u \notin P$, no change in flow at $u \implies$ still balanced.

f' obeys flow conservation

Consider some $u \in V \setminus \{s, t\}$.

- ▶ If $u \notin P$, no change in flow at $u \implies$ still balanced.
- ▶ If $u \in P$, four possibilities:



f' obeys capacity constraints

Let $(u, v) \in E$

f' obeys capacity constraints

Let $(u, v) \in E$

- ▶ If $(u, v), (v, u) \notin P$: $f'(u, v) = f(u, v) \leq c(u, v)$

f' obeys capacity constraints

Let $(u, v) \in E$

- ▶ If $(u, v), (v, u) \notin P$: $f'(u, v) = f(u, v) \leq c(u, v)$
- ▶ If $(u, v) \in P$:

f' obeys capacity constraints

Let $(u, v) \in E$

- ▶ If $(u, v), (v, u) \notin P$: $f'(u, v) = f(u, v) \leq c(u, v)$
- ▶ If $(u, v) \in P$:

$$\begin{aligned} f'(u, v) &= f(u, v) + F \\ &\leq f(u, v) + c_f(u, v) \\ &= f(u, v) + c(u, v) - f(u, v) \\ &= c(u, v) \end{aligned}$$

f' obeys capacity constraints

Let $(u, v) \in E$

► If $(u, v), (v, u) \notin P$: $f'(u, v) = f(u, v) \leq c(u, v)$

► If $(u, v) \in P$:

► If $(v, u) \in P$:

$$\begin{aligned} f'(u, v) &= f(u, v) + F \\ &\leq f(u, v) + c_f(u, v) \\ &= f(u, v) + c(u, v) - f(u, v) \\ &= c(u, v) \end{aligned}$$

f' obeys capacity constraints

Let $(u, v) \in E$

► If $(u, v), (v, u) \notin P$: $f'(u, v) = f(u, v) \leq c(u, v)$

► If $(u, v) \in P$:

$$\begin{aligned} f'(u, v) &= f(u, v) + F \\ &\leq f(u, v) + c_f(u, v) \\ &= f(u, v) + c(u, v) - f(u, v) \\ &= c(u, v) \end{aligned}$$

► If $(v, u) \in P$:

$$\begin{aligned} f'(u, v) &= f(u, v) - F \\ &\geq f(u, v) - c_f(v, u) \\ &= f(u, v) - f(u, v) \\ &= 0 \end{aligned}$$

Ford-Fulkerson Algorithm and Integrality

FF Algorithm

Obvious algorithm from previous proof: keep pushing flow!

```
 $f = \vec{0}$   
while( $\exists s \rightarrow t$  path  $P$  in  $G_f$ ) {  
     $F = \min_{e \in P} c_f(e)$   
    Push  $F$  flow along  $P$  to get new flow  $f'$   
     $f = f'$   
}  
return  $f$ 
```

FF Algorithm

Obvious algorithm from previous proof: keep pushing flow!

```
 $f = \vec{0}$   
while( $\exists s \rightarrow t$  path  $P$  in  $G_f$ ) {  
     $F = \min_{e \in P} c_f(e)$   
    Push  $F$  flow along  $P$  to get new flow  $f'$   
     $f = f'$   
}  
return  $f$  or  $\{v \in V : v \text{ reachable from } s \text{ in } G_f\}$ 
```












FF Algorithm

Obvious algorithm from previous proof: keep pushing flow!

```
 $f = \vec{0}$   
while( $\exists s \rightarrow t$  path  $P$  in  $G_f$ ) {  
     $F = \min_{e \in P} c_f(e)$   
    Push  $F$  flow along  $P$  to get new flow  $f'$   
     $f = f'$   
}  
return  $f$  or  $\{v \in V : v \text{ reachable from } s \text{ in } G_f\}$ 
```

Correctness: directly from previous proof

Example



Integrality

Corollary

If all capacities are integers, then there is a max flow such that the flow through every edge is an integer

Integrality

Corollary

If all capacities are integers, then there is a max flow such that the flow through every edge is an integer

Proof.

Induction on iterations of the Ford-Fulkerson algorithm: initially true, stays true \implies true at end. □

Running Time

Theorem


If all capacities are integers and the max flow value is F , Ford-Fulkerson takes time at most $O(F(m + n))$

Running Time

Theorem

If all capacities are integers and the max flow value is F , Ford-Fulkerson takes time at most $O(F(m + n))$

Finding path takes $O(m + n)$ time, increase flow by at least 1




A bad example for the Ford-Fulkerson algorithm.

Running Time

Theorem

If all capacities are integers and the max flow value is F , Ford-Fulkerson takes time at most $O(F(m + n))$

Finding path takes $O(m + n)$ time, increase flow by at least 1



Running time $\geq \#$ iterations.


A bad example for the Ford-Fulkerson algorithm.

Running Time

Theorem

If all capacities are integers and the max flow value is F , Ford-Fulkerson takes time at most $O(F(m + n))$

Finding path takes $O(m + n)$ time, increase flow by at least 1



Running time $\geq \#$ iterations.

This example:

- ▶ Running time: $\Omega(x)$


A bad example for the Ford-Fulkerson algorithm.

Running Time

Theorem

If all capacities are integers and the max flow value is F , Ford-Fulkerson takes time at most $O(F(m + n))$

Finding path takes $O(m + n)$ time, increase flow by at least 1



A bad example for the Ford-Fulkerson algorithm.

Running time $\geq \#$ iterations.

This example:


- ▶ Running time: $\Omega(x)$
- ▶ Input size $O(\log x) + O(1)$

Running Time

Theorem

If all capacities are integers and the max flow value is F , Ford-Fulkerson takes time at most $O(F(m + n))$

Finding path takes $O(m + n)$ time, increase flow by at least 1



A bad example for the Ford-Fulkerson algorithm.

Running time $\geq \#$ iterations.

This example:

- ▶ Running time: $\Omega(x)$
- ▶ Input size $O(\log x) + O(1)$

\implies Exponential time!