



# **CIS 635 - Knowledge Discovery & Data Mining**

- Linear to Polynomial Regression
- Model Regularization




# Plan

- LR to Polynomial Regression
- Regularization
  - Theory
  - Practical - Notebook presentation



## Non linear data/function

- Does this data points seem familiar matching a known function?




# Non linear data/function

- Does this data points seem familiar matching a known function?

- A Sinusoidal function

$$y(t) = A \sin(\omega t + \varphi) = A \sin(2\pi f t + \varphi)$$




# Non linear data/function

- Does this data points seem familiar matching a known function?

- A Sinusoidal function

$$y(t) = A \sin(\omega t + \varphi) = A \sin(2\pi f t + \varphi)$$





Clearly this is not a linear function; right?

# Non linear data/function

- Does this data points seem familiar matching a known function?
- Can we approximate this function using LR?

$$\hat{y} = \beta_0 + \beta_1 x$$



LR will not work; right?



## What no-linear functions we are aware of?


- Can you recall any nonlinear function you learned at your high school/colleges?

# What no-linear functions we are aware of?

- Can you recall any nonlinear function you learned at your high school/colleges?
- **Quadratic ( $x^2$ )**

$$f(x) = x^2, \quad f(x) = 2x^2, \quad f(x) = 5x^2.$$


What is the impact of changing the coefficient of  $x^2$  as we have done in these examples? One way to find out is to sketch the graphs of the functions.





# What no-linear functions we are aware of?

- Can you recall any nonlinear function you learned at your high school/colleges?
- **Cubic ( $x^3$ )**



Cubic functions of the form

$$y = x^3 + cx.$$

The graph of any cubic function is  
**similar** to such a curve.

# What no-linear functions we are aware of?

- Can you recall any nonlinear function you learned at your high school/colleges?
- Quadratic ( $x^2$ )
- Cubic ( $x^3$ )
- 

$$\hat{y} = \beta_0 + \beta_1 x$$

$$\hat{y} = \beta_0 + \beta_1 x + \beta_2 x^2$$

$$\hat{y} = \beta_0 + \beta_1 x + \beta_2 x^2 + \beta_3 x^3$$

$$\hat{y} = \beta_0 + \beta_1 x + \beta_2 x^2 + \beta_3 x^3 + \dots$$

Linear ( $x$ )

Quadratic ( $x^2$ )

Cubic ( $x^3$ )

# LR to Polynomial Regression





- Polynomial function
  - $M$  is the order/degree of polynomial ..

$$\hat{y} = \beta_0 + \beta_1 x$$

$$\hat{y} = \beta_0 + \beta_1 x + \beta_2 x^2$$

$$\hat{y} = \beta_0 + \beta_1 x + \beta_2 x^2 + \beta_3 x^3$$

$$\hat{y} = \beta_0 + \beta_1 x + \beta_2 x^2 + \beta_3 x^3 + \dots$$



# LR to Polynomial Regression





- Polynomial function
  - $M$  is the order/degree of polynomial ..
  - **Where to stop? What is the best  $M$ ?**

$$\hat{y} = \beta_0 + \beta_1 x$$

$$\hat{y} = \beta_0 + \beta_1 x + \beta_2 x^2$$

$$\hat{y} = \beta_0 + \beta_1 x + \beta_2 x^2 + \beta_3 x^3$$

$$\hat{y} = \beta_0 + \beta_1 x + \beta_2 x^2 + \beta_3 x^3 + \dots$$



# LR to Polynomial Regression

- Polynomial function
  - $M$  is the order/degree of polynomial ..
  - **Where to stop? What is the best  $M$ ?**





Good news is our gradient descent (iterative learning ) remains the same!

$$\hat{y} = \beta_0 + \beta_1 x$$

$$\hat{y} = \beta_0 + \beta_1 x + \beta_2 x^2$$

$$\hat{y} = \beta_0 + \beta_1 x + \beta_2 x^2 + \beta_3 x^3$$

$$\hat{y} = \beta_0 + \beta_1 x + \beta_2 x^2 + \beta_3 x^3 + \dots$$



# LR to Polynomial Regression

- Polynomial function
  - M is the order ..
  - **Where to stop? What is the best M?**





- Good news is our gradient descent (iterative learning) remains the same!
- You only need to change your objective function (from LR to Polynomial LR)

$$\hat{y} = \beta_0 + \beta_1 x$$


$$\hat{y} = \beta_0 + \beta_1 x + \beta_2 x^2$$

$$\hat{y} = \beta_0 + \beta_1 x + \beta_2 x^2 + \beta_3 x^3$$

$$\hat{y} = \beta_0 + \beta_1 x + \beta_2 x^2 + \beta_3 x^3 + \dots$$



# LR to Polynomial Regression


$$\begin{aligned}\hat{y} &= \beta_0 + \beta_1 x \\ \hat{y} &= \beta_0 + \beta_1 x + \beta_2 x^2 \\ \hat{y} &= \beta_0 + \beta_1 x + \beta_2 x^2 + \beta_2 x^3 \\ \hat{y} &= \beta_0 + \beta_1 x + \beta_2 x^2 + \beta_2 x^3 + \dots\end{aligned}$$

Model

$$\hat{y} = \beta_0 + \beta_1 x$$

$$\Theta = \{\beta_0, \beta_1\}$$

$$\epsilon = |\hat{y} - y|$$

Optimization function

$$E_{\Theta} = \frac{1}{2} \sum_{i=1}^N (\hat{y}_i - y_i)^2$$

$$\Theta^* = \operatorname{argmin}_{\Theta} E\{(x_i, y_i)\}_{i=1, \dots, N}$$

Our model got a little bigger: 2 params to  $M$  param



GPT

I know one of your tricks; get you soon!!



Our model yesterday



Our model got a little bigger: 2 params to  $M$  param



GPT

I know one of your tricks; get you soon!!




Our model today



# Regularization

# Regularization



So, essentially we are fitting a function; right?

Model

$$\hat{y} = \beta_0 + \beta_1 x$$

$$\Theta = \{\beta_0, \beta_1\}$$

Fitting Error

$$\epsilon = |\hat{y} - y|$$

Optimization function

$$E_{\Theta} = \frac{1}{2} \sum_{i=1}^N (\hat{y}_i - y_i)^2$$

$$\Theta^* = \operatorname{argmin}_{\Theta} E\{(x_i, y_i)\}_{i=1, \dots, N}$$

# Regularization

$$E(\mathbf{w}) = \frac{1}{2} \sum_{n=1}^N \{y(x_n, \mathbf{w}) - t_n\}^2$$

Model

$$\hat{y} = \beta_0 + \beta_1 x$$

$$\Theta = \{\beta_0, \beta_1\}$$

$$\epsilon = |\hat{y} - y|$$

Optimization function

$$E_{\Theta} = \frac{1}{2} \sum_{i=1}^N (\hat{y}_i - y_i)^2$$

$$\Theta^* = \operatorname{argmin}_{\Theta} E\{(x_i, y_i)\}_{i=1, \dots, N}$$

# Regularization

$$E(\mathbf{w}) = \frac{1}{2} \sum_{n=1}^N \{y(x_n, \mathbf{w}) - t_n\}^2$$

Essentially, the same formulation

Generally **ML** vs **Math** conventions

Model

$$\hat{y} = \beta_0 + \beta_1 x$$

$$\Theta = \{\beta_0, \beta_1\}$$

$$\epsilon = |\hat{y} - y|$$

Optimization function

$$E_{\Theta} = \frac{1}{2} \sum_{i=1}^N (\hat{y}_i - y_i)^2$$

$$\Theta^* = \operatorname{argmin}_{\Theta} E\{(x_i, y_i)\}_{i=1, \dots, N}$$

# Regularization

$$E(\mathbf{w}) = \frac{1}{2} \sum_{n=1}^N \{y(x_n, \mathbf{w}) - t_n\}^2$$

$x$  : scalar  
 $\mathbf{x}, \mathbf{x}$ : vector  
 $\mathbf{X}$ : Matrix

Model

$$\hat{y} = \beta_0 + \beta_1 x$$

$$\Theta = \{\beta_0, \beta_1\}$$

Essentially, the same formulation

Generally ML vs Math conventions

$$W^* = \operatorname{argmin}_W E\{(x_i, t_i)\}_{i=1, \dots, N}$$


$$\epsilon = |\hat{y} - y|$$

Optimization function

$$E_{\Theta} = \frac{1}{2} \sum_{i=1}^N (\hat{y}_i - y_i)^2$$

$$\Theta^* = \operatorname{argmin}_{\Theta} E\{(x_i, y_i)\}_{i=1, \dots, N}$$


# Regularization



**Table 1.1** Table of the coefficients  $w^*$  for polynomials of various order. Observe how the typical magnitude of the coefficients increases dramatically as the order of the polynomial increases.

	$M = 0$	$M = 1$	$M = 6$	$M = 9$
$w_0^*$	0.19	0.82	0.31	0.35
$w_1^*$		-1.27	7.99	232.37
$w_2^*$			-25.43	-5321.83
$w_3^*$			17.37	48568.31
$w_4^*$				-231639.30
$w_5^*$				640042.26
$w_6^*$				-1061800.52
$w_7^*$				1042400.18
$w_8^*$				-557682.99
$w_9^*$				125201.43

# Regularization




**Table 1.1** Table of the coefficients  $w^*$  for polynomials of various order. Observe how the typical magnitude of the coefficients increases dramatically as the order of the polynomial increases.

	$M = 0$	$M = 1$	$M = 6$	$M = 9$
$w_0^*$	0.19	0.82	0.31	0.35
$w_1^*$		-1.27	7.99	232.37
$w_2^*$			-25.43	-5321.83
$w_3^*$			17.37	48568.31
$w_4^*$				-231639.30
$w_5^*$				640042.26
$w_6^*$				-1061800.52
$w_7^*$				1042400.18
$w_8^*$				-557682.99
$w_9^*$				125201.43

Abs values  
Are increasing



# Regularization



$$E(\mathbf{w}) = \frac{1}{2} \sum_{n=1}^N \{y(x_n, \mathbf{w}) - t_n\}^2$$

Regularizer

$$\tilde{E}(\mathbf{w}) = \frac{1}{2} \sum_{n=1}^N \{y(x_n, \mathbf{w}) - t_n\}^2 + \frac{\lambda}{2} \|\mathbf{w}\|^2$$

# Regularization

$$E(\mathbf{w}) = \frac{1}{2} \sum_{n=1}^N \{y(x_n, \mathbf{w}) - t_n\}^2$$

Regularizer

$$\tilde{E}(\mathbf{w}) = \frac{1}{2} \sum_{n=1}^N \{y(x_n, \mathbf{w}) - t_n\}^2 + \frac{\lambda}{2} \|\mathbf{w}\|^2$$

**Table 1.1** Table of the coefficients  $\mathbf{w}^*$  for polynomials of various order. Observe how the typical magnitude of the coefficients increases dramatically as the order of the polynomial increases.


	$M = 0$	$M = 1$	$M = 6$	$M = 9$
$w_0^*$	0.19	0.82	0.31	0.35
$w_1^*$		-1.27	7.99	232.37
$w_2^*$			-25.43	-5321.83
$w_3^*$			17.37	48568.31
$w_4^*$				-231639.30
$w_5^*$				640042.26
$w_6^*$				-1061800.52
$w_7^*$				1042400.18
$w_8^*$				-557682.99
$w_9^*$				125201.43

Abs values  
Are increasing

# Regularization

$$E(\mathbf{w}) = \frac{1}{2} \sum_{n=1}^N \{y(x_n, \mathbf{w}) - t_n\}^2$$

$$\tilde{E}(\mathbf{w}) = \frac{1}{2} \sum_{n=1}^N \{y(x_n, \mathbf{w}) - t_n\}^2 + \frac{\lambda}{2} \|\mathbf{w}\|^2$$




**Table 1.1** Table of the coefficients  $\mathbf{w}^*$  for polynomials of various order. Observe how the typical magnitude of the coefficients increases dramatically as the order of the polynomial increases.

	$M = 0$	$M = 1$	$M = 6$	$M = 9$
$w_0^*$	0.19	0.82	0.31	0.35
$w_1^*$		-1.27	7.99	232.37
$w_2^*$			-25.43	-5321.83
$w_3^*$			17.37	48568.31
$w_4^*$				-231639.30
$w_5^*$				640042.26
$w_6^*$				-1061800.52
$w_7^*$				1042400.18
$w_8^*$				-557682.99
$w_9^*$				125201.43


Abs values  
Are increasing

# Regularization

$$E(\mathbf{w}) = \frac{1}{2} \sum_{n=1}^N \{y(x_n, \mathbf{w}) - t_n\}^2$$



$$\tilde{E}(\mathbf{w}) = \frac{1}{2} \sum_{n=1}^N \{y(x_n, \mathbf{w}) - t_n\}^2 + \frac{\lambda}{2} \|\mathbf{w}\|^2$$



**Table 1.1** Table of the coefficients  $\mathbf{w}^*$  for polynomials of various order. Observe how the typical magnitude of the coefficients increases dramatically as the order of the polynomial increases.


	$M = 0$	$M = 1$	$M = 6$	$M = 9$
$w_0^*$	0.19	0.82	0.31	0.35
$w_1^*$		-1.27	7.99	232.37
$w_2^*$			-25.43	-5321.83
$w_3^*$			17.37	48568.31
$w_4^*$				-231639.30
$w_5^*$				640042.26
$w_6^*$				-1061800.52
$w_7^*$				1042400.18
$w_8^*$				-557682.99
$w_9^*$				125201.43

Abs values  
Are increasing

# Regularization

$$E(\mathbf{w}) = \frac{1}{2} \sum_{n=1}^N \{y(x_n, \mathbf{w}) - t_n\}^2$$

$$\tilde{E}(\mathbf{w}) = \frac{1}{2} \sum_{n=1}^N \{y(x_n, \mathbf{w}) - t_n\}^2 + \frac{\lambda}{2} \|\mathbf{w}\|^2$$



**Table 1.1** Table of the coefficients  $\mathbf{w}^*$  for polynomials of various order. Observe how the typical magnitude of the coefficients increases dramatically as the order of the polynomial increases.

	$M = 0$	$M = 1$	$M = 6$	$M = 9$
$w_0^*$	0.19	0.82	0.31	0.35
$w_1^*$		-1.27	7.99	232.37
$w_2^*$			-25.43	-5321.83
$w_3^*$			17.37	48568.31
$w_4^*$				-231639.30
$w_5^*$				640042.26
$w_6^*$				-1061800.52
$w_7^*$				1042400.18
$w_8^*$				-557682.99
$w_9^*$				125201.43

*How to control this?*

# Regularization

$$E(\mathbf{w}) = \frac{1}{2} \sum_{n=1}^N \{y(x_n, \mathbf{w}) - t_n\}^2$$

$$\tilde{E}(\mathbf{w}) = \frac{1}{2} \sum_{n=1}^N \{y(x_n, \mathbf{w}) - t_n\}^2 + \frac{\lambda}{2} \|\mathbf{w}\|^2$$

**Table 1.1** Table of the coefficients  $\mathbf{w}^*$  for polynomials of various order. Observe how the typical magnitude of the coefficients increases dramatically as the order of the polynomial increases.


	$M = 0$	$M = 1$	$M = 6$	$M = 9$
$w_0^*$	0.19	0.82	0.31	0.35
$w_1^*$		-1.27	7.99	232.37
$w_2^*$			-25.43	-5321.83
$w_3^*$			17.37	48568.31
$w_4^*$				-231639.30
$w_5^*$				640042.26
$w_6^*$				-1061800.52
$w_7^*$				1042400.18
$w_8^*$				-557682.99
$w_9^*$				125201.43

Who to control this?

# Regularization

$$E(\mathbf{w}) = \frac{1}{2} \sum_{n=1}^N \{y(x_n, \mathbf{w}) - t_n\}^2$$

$$\tilde{E}(\mathbf{w}) = \frac{1}{2} \sum_{n=1}^N \{y(x_n, \mathbf{w}) - t_n\}^2 + \frac{\lambda}{2} \|\mathbf{w}\|^2$$




**Table 1.1** Table of the coefficients  $\mathbf{w}^*$  for polynomials of various order. Observe how the typical magnitude of the coefficients increases dramatically as the order of the polynomial increases.

	$M = 0$	$M = 1$	$M = 6$	$M = 9$
$w_0^*$	0.19	0.82	0.31	0.35
$w_1^*$		-1.27	7.99	232.37
$w_2^*$			-25.43	-5321.83
$w_3^*$			17.37	48568.31
$w_4^*$				-231639.30
$w_5^*$				640042.26
$w_6^*$				-1061800.52
$w_7^*$				1042400.18
$w_8^*$				-557682.99
$w_9^*$				125201.43

**Who to control this?**

Lambda is called the  
**Hyper Parameter** of this model


# Regularization



$$\tilde{E}(\mathbf{w}) = \frac{1}{2} \sum_{n=1}^N \{y(x_n, \mathbf{w}) - t_n\}^2 + \frac{\lambda}{2} \|\mathbf{w}\|^2$$


**Table 1.2** Table of the coefficients  $w^*$  for  $M = 9$  polynomials with various values for the regularization parameter  $\lambda$ . Note that  $\ln \lambda = -\infty$  corresponds to a model with no regularization, i.e., to the graph at the bottom right in Figure 1.4. We see that, as the value of  $\lambda$  increases, the typical magnitude of the coefficients gets smaller.

	$\ln \lambda = -\infty$	$\ln \lambda = -18$	$\ln \lambda = 0$
$w_0^*$	0.35	0.35	0.13
$w_1^*$	232.37	4.74	-0.05
$w_2^*$	-5321.83	-0.77	-0.06
$w_3^*$	48568.31	-31.97	-0.05
$w_4^*$	-231639.30	-3.89	-0.03
$w_5^*$	640042.26	55.28	-0.02
$w_6^*$	-1061800.52	41.32	-0.01
$w_7^*$	1042400.18	-45.95	-0.00
$w_8^*$	-557682.99	-91.53	0.00
$w_9^*$	125201.43	72.68	0.01





# Linear to Polynomial Regression + Regularization




Learned function is nonlinear

$$\begin{aligned}\hat{y} &= \beta_0 \\ \hat{y} &= \beta_0 + \beta_1 x \\ \hat{y} &= \beta_0 + \beta_1 x + \beta_2 x^2 \\ \hat{y} &= \beta_0 + \beta_1 x + \beta_2 x^2 + \beta_3 x^3 \\ \hat{y} &= \beta_0 + \beta_1 x + \beta_2 x^2 + \beta_3 x^3 + \dots\end{aligned}$$

$$\tilde{E}(\mathbf{w}) = \frac{1}{2} \sum_{n=1}^N \{y(x_n, \mathbf{w}) - t_n\}^2 + \frac{\lambda}{2} \|\mathbf{w}\|^2$$

# Linear to Polynomial Regression + Regularization



Learned function is **nonlinear**

$$\begin{aligned}\hat{y} &= \beta_0 \\ \hat{y} &= \beta_0 + \beta_1 x \\ \hat{y} &= \beta_0 + \beta_1 x + \beta_2 x^2 \\ \hat{y} &= \beta_0 + \beta_1 x + \beta_2 x^2 + \beta_3 x^3 \\ \hat{y} &= \beta_0 + \beta_1 x + \beta_2 x^2 + \beta_3 x^3 + \dots\end{aligned}$$

$$\tilde{E}(\mathbf{w}) = \frac{1}{2} \sum_{n=1}^N \{y(x_n, \mathbf{w}) - t_n\}^2 + \frac{\lambda}{2} \|\mathbf{w}\|^2$$

Model (still) **linear**