CIS 635 Knowledge Discovery & Data Mining

Predictive modeling: Classification: Metrics and Imbalanced Data

Accuracy =
$$\frac{\text{Nb of correct predictions}}{\text{Nb of (correct + incorrect) predictions}}$$

Accuray


• Is accuracy a good metric?

- Is accuracy a good metric?
- Not always

- Is accuracy a good metric?
- Not always
- Let's analyze the confusion matrix of our Breast cancer prediction problem
 - Accuray metric can be catastrophic

- Is accuracy a good metric?
- Not always
- Let's analyze the confusion matrix of our Breast cancer prediction problem
 - Accuray metric can be catastrophic
- What other metrics we may use?


$$Accuracy = \frac{PP + NN}{PP + NN + NP + PN}$$



Other important classification metrics


- Precision (also called **Positive Predictive Value**)
- Recall (also called Sensitivity)
- F1 Score


Precision (also called Positive Predictive Value)




• Recall (also called **Sensitivity**)

Recall =
$$\frac{PP}{PP + PN}$$







- Demonstration through a practical example
 - CC application

- How to deal with Data Imbalance Problems
 - Through Sampling Bias

- How to deal with Data Imbalance Problems
 - Through Sampling Bias
 - Undersampling
 - Oversampling

- How to deal with Data Imbalance Problems
 - Through Sampling Bias
 - Undersampling
 - Oversampling
 - Redefining model (loss function for an example)

- How to deal with Data Imbalance Problems
 - Through Sampling Bias
 - Undersampling
 - Oversampling
 - Redefining model (loss function for an example)
 - Applying Higher Penalty to data points from the less occurring class.

QA