



CIS 678 - Machine Learning

- Model Regularization (part 2)




Plan

- Regularization
 - Theory
 - Practical - Notebook presentation



Non linear data/function

- Does this data points seem familiar matching a known function?




Non linear data/function

- Does this data points seem familiar matching a known function?


- A Sinusoidal function

$$y(t) = A \sin(\omega t + \varphi) = A \sin(2\pi f t + \varphi)$$






Polynomial Regression with Regularization



$$\hat{y} = \beta_0$$


Polynomial Regression with Regularization



$$\hat{y} = \beta_0 + \beta_1 x$$


Polynomial Regression with Regularization

$$\hat{y} = \beta_0 + \beta_1 x + \beta_2 x^2 + \beta_3 x^3$$




Polynomial Regression with Regularization

$$\hat{y} = \beta_0 + \beta_1 x + \beta_2 x^2 + \beta_3 x^3 + \dots$$




Polynomial Regression with Regularization



$$\begin{aligned}\hat{y} &= \beta_0 \\ \hat{y} &= \beta_0 + \beta_1 x \\ \hat{y} &= \beta_0 + \beta_1 x + \beta_2 x^2 \\ \hat{y} &= \beta_0 + \beta_1 x + \beta_2 x^2 + \beta_3 x^3 \\ \hat{y} &= \beta_0 + \beta_1 x + \beta_2 x^2 + \beta_3 x^3 + \dots\end{aligned}$$


Polynomial Regression with Regularization



$$\begin{aligned}\hat{y} &= \beta_0 \\ \hat{y} &= \beta_0 + \beta_1 x \\ \hat{y} &= \beta_0 + \beta_1 x + \beta_2 x^2 \\ \hat{y} &= \beta_0 + \beta_1 x + \beta_2 x^2 + \beta_3 x^3 \\ \hat{y} &= \beta_0 + \beta_1 x + \beta_2 x^2 + \beta_3 x^3 + \dots\end{aligned}$$

$$\tilde{E}(\mathbf{w}) = \frac{1}{2} \sum_{n=1}^N \{y(x_n, \mathbf{w}) - t_n\}^2$$

Polynomial Regression with Regularization




$$\begin{aligned}\hat{y} &= \beta_0 \\ \hat{y} &= \beta_0 + \beta_1 x \\ \hat{y} &= \beta_0 + \beta_1 x + \beta_2 x^2 \\ \hat{y} &= \beta_0 + \beta_1 x + \beta_2 x^2 + \beta_3 x^3 \\ \hat{y} &= \beta_0 + \beta_1 x + \beta_2 x^2 + \beta_3 x^3 + \dots\end{aligned}$$

$$\tilde{E}(\mathbf{w}) = \frac{1}{2} \sum_{n=1}^N \{y(x_n, \mathbf{w}) - t_n\}^2$$

decreasing

Polynomial Regression with Regularization



$$\hat{y} = \beta_0$$

$$\hat{y} = \beta_0 + \beta_1 x$$

$$\hat{y} = \beta_0 + \beta_1 x + \beta_2 x^2$$

$$\hat{y} = \beta_0 + \beta_1 x + \beta_2 x^2 + \beta_3 x^3$$


$$\hat{y} = \beta_0 + \beta_1 x + \beta_2 x^2 + \beta_3 x^3 +$$

increasing

$$\tilde{E}(\mathbf{w}) = \frac{1}{2} \sum_{n=1}^N \{y(x_n, \mathbf{w}) - t_n\}^2 + \frac{1}{2} \|\mathbf{w}\|^2$$

decreasing

Polynomial Regression with Regularization



$$\hat{y} = \beta_0$$

$$\hat{y} = \beta_0 + \beta_1 x$$

$$\hat{y} = \beta_0 + \beta_1 x + \beta_2 x^2$$

$$\hat{y} = \beta_0 + \beta_1 x + \beta_2 x^2 + \beta_3 x^3$$


$$\hat{y} = \beta_0 + \beta_1 x + \beta_2 x^2 + \beta_3 x^3 + \beta_4 x^4$$

Loss function

Regularizer

$$\tilde{E}(\mathbf{w}) = \frac{1}{2} \sum_{n=1}^N \{y(x_n, \mathbf{w}) - t_n\}^2 + \frac{1}{2} \|\mathbf{w}\|^2$$

Polynomial Regression with Regularization



$$\hat{y} = \beta_0$$

$$\hat{y} = \beta_0 + \beta_1 x$$

$$\hat{y} = \beta_0 + \beta_1 x + \beta_2 x^2$$

$$\hat{y} = \beta_0 + \beta_1 x + \beta_2 x^2 + \beta_3 x^3$$

$$\hat{y} = \beta_0 + \beta_1 x + \beta_2 x^2 + \beta_3 x^3 + \beta_4 x^4$$

Loss function

Regularizer

$$\tilde{E}(\mathbf{w}) = \frac{1}{2} \sum_{n=1}^N \{y(x_n, \mathbf{w}) - t_n\}^2 + \frac{\lambda}{2} \|\mathbf{w}\|^2$$

Lambda is for balancing these two -
Hyper Parameter of this model



How to find the best HP (Lamda)?

Predictive modeling: [Regression \(diabetes\)](#)