



# **CIS 678 Machine Learning**

Introduction to Linear Algebra



# Basic Math - Concept of Vectors, and Vector Space



# Basic Math - Concept of Vectors

We are aware of Scalars: A person's

Height (1.72m)



# Basic Math - Concept of Vectors

We are aware of Scalars: A person's

Height (1.72m)

Weight (72kg)



# Basic Math - Concept of Vectors

We are aware of Scalars: A person's

Height (1.72m)

Weight (72kg)

Salary (100K)



# Basic Math - Concept of Vectors

We are aware of Scalars: A person's

Height (1.72m)

Weight (72kg)

Salary (100K)

....



# Basic Math - Concept of Vectors

A closed form definition of a person through some features

[Height (1.72m), Weight (72kg), Salary (100K)]



## Basic Math - Concept of Vectors

A closed form definition of a person through some features

- no explicit unit mentions

[1.72, 72, 100]





## Basic Math - Concept of Vectors

A closed form definition of a person through some features

- no explicit unit mentions

[1.72, 72, 100]

Is a vectorized representation of a person  
through some attributes: height, weight, salary



## Basic Math - Concept of Vectors

A closed form definition of a person through some features

- no explicit unit mentions

[1.72, 72, 100],  
[1.65, 70, 120],  
[1.81, 110, 90],  
...  
[1.45, 65, 130],

And here we are talking about a number of  
people through same features: height,  
weight, salary



# Formal definition of Vectors

# Basic Math - Concept of Vectors

## Formal definition of Vectors

### 1. Vectors

We begin by defining a mathematical abstraction known as a **vector space**. In linear algebra the fundamental concepts relate to the  **$n$ -tuples** and their algebraic properties.

**Definition:** An ordered  $n$ -tuple is considered as a sequence of  $n$  **terms**  $(a_1, a_2, \dots, a_n)$ , where  $n$  is a positive integer.

We see that an ordered  **$n$ -tuple** has **terms** whereas a set has **members**.

**Example:** A sequence (5) is called an ordered **1-tuple**. A **2-tuple**, for example (3, 6) (where 6 appears after 3) is called an ordered pair, and **3-tuple** is called an ordered triple. A sequence (9, 3, 4, 4, 1) is called an ordered **5-tuple**.

Let us denote the set of all ordered **1-tuples** of real numbers by  $\mathbb{R}$ . We will write for example  $(3.5) \in \mathbb{R}$ .

$$\mathbf{X} = [1.78, 72, 100]$$

$$\mathbf{x} = \begin{bmatrix} 1 \\ 3 \\ 4 \end{bmatrix} \quad \mathbf{x} = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix}$$



## Basic Math - Concept of Vectors

We are aware of Scalars: A person's height, weight, salary

Physics vector: velocity (scalar value + direction)

Algebraic vector (in general): Common representation of an entity (1 to n dimension):

- A person's (height, weight, salary), say [\[1.78, 72, 100\]](#): once defined, we have to follow it.

$$\mathbf{x} = \begin{bmatrix} 1 \\ 3 \\ 4 \end{bmatrix} \quad \mathbf{x} = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix}$$



# Basic Math - Vector Operations

## Vector operation rules

1.  $\mathbf{x} + \mathbf{y} \in \mathbb{R}^n$
2.  $\alpha \cdot \mathbf{x} \in \mathbb{R}^n$
3.  $\mathbf{x} + \mathbf{y} = \mathbf{y} + \mathbf{x} \in \mathbb{R}^n$  *(commutativity)*
4.  $\alpha \cdot (\mathbf{x} + \mathbf{y}) = \alpha \cdot \mathbf{x} + \alpha \cdot \mathbf{y}$  *(distributivity)*
5.  $(\alpha + \beta) \cdot \mathbf{x} = \alpha \cdot \mathbf{x} + \beta \cdot \mathbf{x}$  *(distributivity)*
6.  $(\mathbf{x} + \mathbf{y}) + \mathbf{z} = \mathbf{x} + (\mathbf{y} + \mathbf{z})$  *(associativity)*
7.  $(\alpha\beta) \cdot \mathbf{x} = \alpha \cdot (\beta \cdot \mathbf{x})$  *(associativity)*



# Basic Math - Vector Operations

## Vector Operation

### *1.1.2. Vector Addition*

Addition of vectors is defined:

$$\mathbf{x} + \mathbf{y} = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} + \begin{bmatrix} y_1 \\ y_2 \\ \vdots \\ y_n \end{bmatrix} = \begin{bmatrix} x_1 + y_1 \\ x_2 + y_2 \\ \vdots \\ x_n + y_n \end{bmatrix}$$

**Example:**

$$\mathbf{x} + \mathbf{y} = \begin{bmatrix} 2 \\ 6 \\ -5 \end{bmatrix} + \begin{bmatrix} 0 \\ 3 \\ 4 \end{bmatrix} = \begin{bmatrix} 2 \\ 9 \\ -1 \end{bmatrix}$$



# Basic Math - Vector Operations

## Vector Operation

### *1.1.4. Zero Vector*

The **zero** vector **sometimes denoted  $\mathbf{0}$**  is a vector having all elements equal to zero, e.g., the 2-dimensional  **$\mathbf{0}$**  vector:

$$\mathbf{x} = \begin{bmatrix} 0 \\ 0 \end{bmatrix} \quad (\text{A.7})$$





# Basic Math - Vector Operations

## Vector Operation

### 1.1.9. Inner Product

The **inner** or **dot** product of two vectors **x** and **y** of the same dimension is a **scalar** defined by:

$$\mathbf{x}^T \cdot \mathbf{y} = (\mathbf{x}, \mathbf{y}) = x_1y_1 + x_2y_2 + \cdots + x_ny_n = \sum_{i=1}^n x_iy_i \quad (\text{A.11})$$

Note that the inner product of vector **x** and **y** requires that a transposed vector **x** be multiplied by the **y** vector. Sometimes the inner product is denoted simply by juxtaposition of the vectors **x** and **y**, for example, as  $\langle \mathbf{x}, \mathbf{y} \rangle$  or  $(\mathbf{x}, \mathbf{y})$ .

**Example:** The inner product of two vectors  $\mathbf{x} = \begin{bmatrix} 4 \\ 1 \\ 7 \end{bmatrix}$  and  $\mathbf{y} = \begin{bmatrix} 0 \\ 2 \\ -3 \end{bmatrix}$

$$\mathbf{x}^T \mathbf{y} = [4 \ 1 \ 7]^T \begin{bmatrix} 0 \\ 2 \\ -3 \end{bmatrix} = 4 \cdot 0 + 1 \cdot 2 + 7 \cdot (-3) = 19$$



# Basic Math - Vector Operations

## Vector Operation

### 1.1.10. Orthogonal Vectors

Two vectors  $\mathbf{x}$  and  $\mathbf{y}$  are said to be **orthogonal** if their inner product is equal to zero

$$\mathbf{x}^T \mathbf{y} = 0 \quad (\text{A.12})$$

here 0 is a scalar.

**Example:** Two vectors  $\mathbf{x} = \begin{bmatrix} 4 \\ 0 \end{bmatrix}$  and  $\mathbf{y} = \begin{bmatrix} 0 \\ 2 \end{bmatrix}$  and are orthogonal, since their inner product is equal to zero

$$\mathbf{x}^T \cdot \mathbf{y} = \begin{bmatrix} 4 \\ 0 \end{bmatrix}^T = [0 \ 2] = 4 \cdot 0 + 0 \cdot 2 = 0$$



# Basic Math - Vector Operations

## Vector Operation

### 1.1.11. Vector Norm

The magnitude of a vector may be measured in different ways. One method, called the vector **norm**, is a function from  $\mathbb{R}^n$  into  $\mathbb{R}$  for  $\mathbf{x}$  an element of  $\mathbb{R}^n$ . It is denoted  $\|\mathbf{x}\|$  and satisfies the following conditions:

1.  $\|\mathbf{x}\| \geq 0$ , and the equality holds if and only if  $\mathbf{x} = \mathbf{0}$
2.  $\|\alpha\mathbf{x}\| = |\alpha| \cdot \|\mathbf{x}\|$ , where  $|\alpha|$  is the absolute value of scalar  $\alpha$

and is defined as:

$$\|\mathbf{x}\| = \sqrt{\mathbf{x}^T \mathbf{x}} = \sqrt{x_1^2 + x_2^2 + \cdots + x_n^2} \quad (\text{A.13})$$

**Example:** For the vector  $\mathbf{x} = \begin{bmatrix} 4 \\ 3 \end{bmatrix}$  the norm is

$$\|\mathbf{x}\| = \sqrt{\mathbf{x}^T \mathbf{x}} = \sqrt{4^2 + 3^2} = 5$$



**QA**