



CIS 678 - Machine Learning

- Linear to Polynomial Regression
- Model Regularization




Plan

- LR to Polynomial Regression
- Regularization
 - Theory
 - Practical - Notebook presentation



Non linear data/function

- Does this data points seem familiar matching a known function?




Non linear data/function

- Does this data points seem familiar matching a known function?

- A Sinusoidal function


$$y(t) = A \sin(\omega t + \varphi) = A \sin(2\pi f t + \varphi)$$



Non linear data/function

- Does this data points seem familiar matching a known function?
- A Sinusoidal function

$$y(t) = A \sin(\omega t + \varphi) = A \sin(2\pi f t + \varphi)$$





Clearly this is not a linear function; right?

Non linear data/function

- Does this data points seem familiar matching a known function?
- Can we approximate this function using LR?

$$\hat{y} = \beta_0 + \beta_1 x$$



LR will not work; right?



What no-linear functions we are aware of?


- Can you recall any nonlinear function you learned at your high school/colleges?

What no-linear functions we are aware of?

- Can you recall any nonlinear function you learned at your high school/colleges?
- **Quadratic (x^2)**


$$f(x) = x^2, \quad f(x) = 2x^2, \quad f(x) = 5x^2.$$

What is the impact of changing the coefficient of x^2 as we have done in these examples? One way to find out is to sketch the graphs of the functions.



What no-linear functions we are aware of?

- Can you recall any nonlinear function you learned at your high school/colleges?
- **Cubic (x^3)**



What no-linear functions we are aware of?

- Can you recall any nonlinear function you learned at your high school/colleges?
- Quadratic (x^2)
- Cubic (x^3)
-

Linear (x)

$$\hat{y} = \beta_0 + \beta_1 x$$


$$\hat{y} = \beta_0 + \beta_1 x + \beta_2 x^2$$

$$\hat{y} = \beta_0 + \beta_1 x + \beta_2 x^2 + \beta_2 x^3$$

$$\hat{y} = \beta_0 + \beta_1 x + \beta_2 x^2 + \beta_2 x^3 + \dots$$

What no-linear functions we are aware of?

- Can you recall any nonlinear function you learned at your high school/colleges?
- Quadratic (x^2)
- Cubic (x^3)
-



$$\begin{aligned}\hat{y} &= \beta_0 + \beta_1 x \\ \hat{y} &= \beta_0 + \beta_1 x + \beta_2 x^2 \\ \hat{y} &= \beta_0 + \beta_1 x + \beta_2 x^2 + \beta_2 x^3 \\ \hat{y} &= \beta_0 + \beta_1 x + \beta_2 x^2 + \beta_2 x^3 + \dots\end{aligned}$$

What no-linear functions we are aware of?

- Can you recall any nonlinear function you learned at your high school/colleges?
- Quadratic (x^2)
- Cubic (x^3)
-

$$\hat{y} = \beta_0 + \beta_1 x$$

$$\hat{y} = \beta_0 + \beta_1 x + \beta_2 x^2$$

$$\hat{y} = \beta_0 + \beta_1 x + \beta_2 x^2 + \beta_2 x^3$$

$$\hat{y} = \beta_0 + \beta_1 x + \beta_2 x^2 + \beta_2 x^3 + \dots$$



What no-linear functions we are aware of?

- Can you recall any nonlinear function you learned at your high school/colleges?
- Quadratic (x^2)
- Cubic (x^3)
-

$$\hat{y} = \beta_0 + \beta_1 x$$

$$\hat{y} = \beta_0 + \beta_1 x + \beta_2 x^2$$

$$\hat{y} = \beta_0 + \beta_1 x + \beta_2 x^2 + \beta_3 x^3$$

$$\hat{y} = \beta_0 + \beta_1 x + \beta_2 x^2 + \beta_3 x^3 + \dots$$

Linear (x)


Quadratic (x^2)

Cubic (x^3)

LR to Polynomial Regression

- Polynomial function
 - M is the order/degree of polynomial ..

$$M=0$$




$$\hat{y} = \beta_0$$

LR to Polynomial Regression

- Polynomial function
 - M is the order/degree of polynomial ..

$$M=1$$

$$\hat{y} = \beta_0 + \beta_1 x$$



LR to Polynomial Regression

- Polynomial function
 - M is the order/degree of polynomial ..


$$M=3$$

$$\hat{y} = \beta_0 + \beta_1 x$$

$$\hat{y} = \beta_0 + \beta_1 x + \beta_2 x^2$$

$$\hat{y} = \beta_0 + \beta_1 x + \beta_2 x^2 + \beta_3 x^3$$

$$\hat{y} = \beta_0 + \beta_1 x + \beta_2 x^2 + \beta_3 x^3 + \dots$$



LR to Polynomial Regression

- Polynomial function
 - M is the order/degree of polynomial ..


$$M=9$$

$$\hat{y} = \beta_0 + \beta_1 x$$

$$\hat{y} = \beta_0 + \beta_1 x + \beta_2 x^2$$

$$\hat{y} = \beta_0 + \beta_1 x + \beta_2 x^2 + \beta_2 x^3$$

$$\hat{y} = \beta_0 + \beta_1 x + \beta_2 x^2 + \beta_2 x^3 + \dots$$



LR to Polynomial Regression

- Polynomial function
 - M is the order/degree of polynomial ..





$M=?$
Which one is
preferable?

$$\hat{y} = \beta_0 + \beta_1 x$$

$$\hat{y} = \beta_0 + \beta_1 x + \beta_2 x^2$$

$$\hat{y} = \beta_0 + \beta_1 x + \beta_2 x^2 + \beta_3 x^3$$

$$\hat{y} = \beta_0 + \beta_1 x + \beta_2 x^2 + \beta_3 x^3 + \dots$$





Regularization



Regularization

*Let's recall the definition of
our optimization function*

Model

$$\hat{y} = \beta_0 + \beta_1 x$$

$$\Theta = \{\beta_0, \beta_1\}$$

$$\epsilon = |\hat{y} - y|$$

Optimization function

$$E_{\Theta} = \frac{1}{2} \sum_{i=1}^N (\hat{y}_i - y_i)^2$$

$$\Theta^* = \operatorname{argmin}_{\Theta} E\{(x_i, y_i)\}_{i=1, \dots, N}$$

Regularization

$$E(\mathbf{w}) = \frac{1}{2} \sum_{n=1}^N \{y(x_n, \mathbf{w}) - t_n\}^2$$

Same model, two different notations

Model

$$\hat{y} = \beta_0 + \beta_1 x$$

$$\Theta = \{\beta_0, \beta_1\}$$

$$\epsilon = |\hat{y} - y|$$

Optimization function

$$E_{\Theta} = \frac{1}{2} \sum_{i=1}^N (\hat{y}_i - y_i)^2$$

$$\Theta^* = \operatorname{argmin}_{\Theta} E\{(x_i, y_i)\}_{i=1, \dots, N}$$

Regularization

$$E(\mathbf{w}) = \frac{1}{2} \sum_{n=1}^N \{y(x_n, \mathbf{w}) - t_n\}^2$$

x : scalar
 \mathbf{x}, \mathbf{x} : vector
 \mathbf{X} : Matrix

Model

$$\hat{y} = \beta_0 + \beta_1 x$$

$$\Theta = \{\beta_0, \beta_1\}$$

$$\epsilon = |\hat{y} - y|$$

Essentially, the same formulation

Optimization function

Generally **ML** vs **Math** conventions

$$W^* = \operatorname{argmin}_W E\{(x_i, t_i)\}_{i=1, \dots, N}$$

$$E_{\Theta} = \frac{1}{2} \sum_{i=1}^N (\hat{y}_i - y_i)^2$$

$$\Theta^* = \operatorname{argmin}_{\Theta} E\{(x_i, y_i)\}_{i=1, \dots, N}$$

Regularization




Table 1.1 Table of the coefficients w^* for polynomials of various order. Observe how the typical magnitude of the coefficients increases dramatically as the order of the polynomial increases.

| | $M=0$ | $M=1$ | $M=6$ | $M=9$ |
|---------|-------|-------|--------|-------------|
| w_0^* | 0.19 | 0.82 | 0.31 | 0.35 |
| w_1^* | | -1.27 | 7.99 | 232.37 |
| w_2^* | | | -25.43 | -5321.83 |
| w_3^* | | | 17.37 | 48568.31 |
| w_4^* | | | | -231639.30 |
| w_5^* | | | | 640042.26 |
| w_6^* | | | | -1061800.52 |
| w_7^* | | | | 1042400.18 |
| w_8^* | | | | -557682.99 |
| w_9^* | | | | 125201.43 |

Regularization





Table 1.1 Table of the coefficients w^* for polynomials of various order. Observe how the typical magnitude of the coefficients increases dramatically as the order of the polynomial increases.

| | $M=0$ | $M=1$ | $M=6$ | $M=9$ |
|---------|-------|-------|--------|-------------|
| w_0^* | 0.19 | 0.82 | 0.31 | 0.35 |
| w_1^* | | -1.27 | 7.99 | 232.37 |
| w_2^* | | | -25.43 | -5321.83 |
| w_3^* | | | 17.37 | 48568.31 |
| w_4^* | | | | -231639.30 |
| w_5^* | | | | 640042.26 |
| w_6^* | | | | -1067800.52 |
| w_7^* | | | | 1042400.18 |
| w_8^* | | | | -557682.99 |
| w_9^* | | | | 125201.43 |

Absolute values
are increasing

Regularization



$$E(\mathbf{w}) = \frac{1}{2} \sum_{n=1}^N \{y(x_n, \mathbf{w}) - t_n\}^2$$

Regularizer

$$\tilde{E}(\mathbf{w}) = \frac{1}{2} \sum_{n=1}^N \{y(x_n, \mathbf{w}) - t_n\}^2 + \frac{\lambda}{2} \|\mathbf{w}\|^2$$

Regularization

$$E(\mathbf{w}) = \frac{1}{2} \sum_{n=1}^N \{y(x_n, \mathbf{w}) - t_n\}^2$$

Regularizer

$$\tilde{E}(\mathbf{w}) = \frac{1}{2} \sum_{n=1}^N \{y(x_n, \mathbf{w}) - t_n\}^2 + \frac{1}{2} \|\mathbf{w}\|^2$$

Table 1.1 Table of the coefficients \mathbf{w}^* for polynomials of various order. Observe how the typical magnitude of the coefficients increases dramatically as the order of the polynomial increases.


| | $M = 0$ | $M = 1$ | $M = 6$ | $M = 9$ |
|---------|---------|---------|---------|-------------|
| w_0^* | 0.19 | 0.82 | 0.31 | 0.35 |
| w_1^* | | -1.27 | 7.99 | 232.37 |
| w_2^* | | | -25.43 | -5321.83 |
| w_3^* | | | 17.37 | 48568.31 |
| w_4^* | | | | -231639.30 |
| w_5^* | | | | 640042.26 |
| w_6^* | | | | -1067800.52 |
| w_7^* | | | | 1042400.18 |
| w_8^* | | | | -557682.99 |
| w_9^* | | | | 125201.43 |

Absolute values are increasing

Regularization

$$E(\mathbf{w}) = \frac{1}{2} \sum_{n=1}^N \{y(x_n, \mathbf{w}) - t_n\}^2$$

$$\tilde{E}(\mathbf{w}) = \frac{1}{2} \sum_{n=1}^N \{y(x_n, \mathbf{w}) - t_n\}^2 + \frac{1}{2} \|\mathbf{w}\|^2$$



decreasing

Table 1.1 Table of the coefficients \mathbf{w}^* for polynomials of various order. Observe how the typical magnitude of the coefficients increases dramatically as the order of the polynomial increases.

| | $M = 0$ | $M = 1$ | $M = 6$ | $M = 9$ |
|---------|---------|---------|---------|-------------|
| w_0^* | 0.19 | 0.82 | 0.31 | 0.35 |
| w_1^* | | -1.27 | 7.99 | 232.37 |
| w_2^* | | | -25.43 | -5321.83 |
| w_3^* | | | 17.37 | 48568.31 |
| w_4^* | | | | -231639.30 |
| w_5^* | | | | 640042.26 |
| w_6^* | | | | -1067800.52 |
| w_7^* | | | | 1042400.18 |
| w_8^* | | | | -557682.99 |
| w_9^* | | | | 125201.43 |

Absolute values are increasing

Regularization

$$E(\mathbf{w}) = \frac{1}{2} \sum_{n=1}^N \{y(x_n, \mathbf{w}) - t_n\}^2$$

$$\tilde{E}(\mathbf{w}) = \frac{1}{2} \sum_{n=1}^N \{y(x_n, \mathbf{w}) - t_n\}^2 + \frac{1}{2} \|\mathbf{w}\|^2$$





Table 1.1 Table of the coefficients \mathbf{w}^* for polynomials of various order. Observe how the typical magnitude of the coefficients increases dramatically as the order of the polynomial increases.

| | $M = 0$ | $M = 1$ | $M = 6$ | $M = 9$ |
|---------|---------|---------|---------|-------------|
| w_0^* | 0.19 | 0.82 | 0.31 | 0.35 |
| w_1^* | | -1.27 | 7.99 | 232.37 |
| w_2^* | | | -25.43 | -5321.83 |
| w_3^* | | | 17.37 | 48568.31 |
| w_4^* | | | | -231639.30 |
| w_5^* | | | | 640042.26 |
| w_6^* | | | | -1067800.52 |
| w_7^* | | | | 1042400.18 |
| w_8^* | | | | -557682.99 |
| w_9^* | | | | 125201.43 |

Absolute values
are increasing

Regularization

$$E(\mathbf{w}) = \frac{1}{2} \sum_{n=1}^N \{y(x_n, \mathbf{w}) - t_n\}^2$$

$$\tilde{E}(\mathbf{w}) = \frac{1}{2} \sum_{n=1}^N \{y(x_n, \mathbf{w}) - t_n\}^2 + \frac{1}{2} \|\mathbf{w}\|^2$$




Table 1.1 Table of the coefficients \mathbf{w}^* for polynomials of various order. Observe how the typical magnitude of the coefficients increases dramatically as the order of the polynomial increases.

| | $M = 0$ | $M = 1$ | $M = 6$ | $M = 9$ |
|---------|---------|---------|---------|-------------|
| w_0^* | 0.19 | 0.82 | 0.31 | 0.35 |
| w_1^* | | -1.27 | 7.99 | 232.37 |
| w_2^* | | | -25.43 | -5321.83 |
| w_3^* | | | 17.37 | 48568.31 |
| w_4^* | | | | -231639.30 |
| w_5^* | | | | 640042.26 |
| w_6^* | | | | -1067800.52 |
| w_7^* | | | | 1042400.18 |
| w_8^* | | | | -557682.99 |
| w_9^* | | | | 125201.43 |

Absolute values are increasing

How to control this?

Regularization




$$E(\mathbf{w}) = \frac{1}{2} \sum_{n=1}^N \{y(x_n, \mathbf{w}) - t_n\}^2$$

Table 1.1 Table of the coefficients \mathbf{w}^* for polynomials of various order. Observe how the typical magnitude of the coefficients increases dramatically as the order of the polynomial increases.

| | $M = 0$ | $M = 1$ | $M = 6$ | $M = 9$ |
|---------|---------|---------|---------|-------------|
| w_0^* | 0.19 | 0.82 | 0.31 | 0.35 |
| w_1^* | | -1.27 | 7.99 | 232.37 |
| w_2^* | | | -25.43 | -5321.83 |
| w_3^* | | | 17.37 | 48568.31 |
| w_4^* | | | | -231639.30 |
| w_5^* | | | | 640042.26 |
| w_6^* | | | | -1061800.52 |
| w_7^* | | | | 1042400.18 |
| w_8^* | | | | -557682.99 |
| w_9^* | | | | 125201.43 |

$$\tilde{E}(\mathbf{w}) = \frac{1}{2} \sum_{n=1}^N \{y(x_n, \mathbf{w}) - t_n\}^2 + \frac{\lambda}{2} \|\mathbf{w}\|^2$$

Who to control this?



Regularization

$$E(\mathbf{w}) = \frac{1}{2} \sum_{n=1}^N \{y(x_n, \mathbf{w}) - t_n\}^2$$

Table 1.1 Table of the coefficients \mathbf{w}^* for polynomials of various order. Observe how the typical magnitude of the coefficients increases dramatically as the order of the polynomial increases.

| | $M = 0$ | $M = 1$ | $M = 6$ | $M = 9$ |
|---------|---------|---------|---------|-------------|
| w_0^* | 0.19 | 0.82 | 0.31 | 0.35 |
| w_1^* | | -1.27 | 7.99 | 232.37 |
| w_2^* | | | -25.43 | -5321.83 |
| w_3^* | | | 17.37 | 48568.31 |
| w_4^* | | | | -231639.30 |
| w_5^* | | | | 640042.26 |
| w_6^* | | | | -1061800.52 |
| w_7^* | | | | 1042400.18 |
| w_8^* | | | | -557682.99 |
| w_9^* | | | | 125201.43 |

$$\tilde{E}(\mathbf{w}) = \frac{1}{2} \sum_{n=1}^N \{y(x_n, \mathbf{w}) - t_n\}^2 + \frac{\lambda}{2} \|\mathbf{w}\|^2$$


Who to control this?

Lambda is called the **Hyper Parameter** of this model

LR to Polynomial Regression

- Polynomial function
 - M is the order/degree of polynomial ..

$M=0$




$$E(\mathbf{w}) = \frac{1}{2} \sum_{n=1}^N \{y(x_n, \mathbf{w}) - t_n\}^2$$

LR to Polynomial Regression

- Polynomial function
 - M is the order/degree of polynomial ..

$M=1$

$$E(\mathbf{w}) = \frac{1}{2} \sum_{n=1}^N \{y(x_n, \mathbf{w}) - t_n\}^2$$




LR to Polynomial Regression

- Polynomial function
 - M is the order/degree of polynomial ..

$$M=3$$

$$E(\mathbf{w}) = \frac{1}{2} \sum_{n=1}^N \{y(x_n, \mathbf{w}) - t_n\}^2$$




LR to Polynomial Regression

- Polynomial function
 - M is the order/degree of polynomial ..

$M=9$

$$E(\mathbf{w}) = \frac{1}{2} \sum_{n=1}^N \{y(x_n, \mathbf{w}) - t_n\}^2$$

Overfitting!!!




LR to Polynomial Regression

- Polynomial function
 - M is the order/degree of polynomial ..

$M=3$

$$\tilde{E}(\mathbf{w}) = \frac{1}{2} \sum_{n=1}^N \{y(x_n, \mathbf{w}) - t_n\}^2 + \frac{\lambda}{2} \|\mathbf{w}\|^2$$




LR to Polynomial Regression


- Polynomial function
 - M is the order/degree of polynomial ..

$M=3$

$$\tilde{E}(\mathbf{w}) = \frac{1}{2} \sum_{n=1}^N \{y(x_n, \mathbf{w}) - t_n\}^2 + \frac{\lambda}{2} \|\mathbf{w}\|^2$$



Polynomial Regression with Regularization




$$\begin{aligned}\hat{y} &= \beta_0 \\ \hat{y} &= \beta_0 + \beta_1 x \\ \hat{y} &= \beta_0 + \beta_1 x + \beta_2 x^2 \\ \hat{y} &= \beta_0 + \beta_1 x + \beta_2 x^2 + \beta_3 x^3 \\ \hat{y} &= \beta_0 + \beta_1 x + \beta_2 x^2 + \beta_3 x^3 + \dots\end{aligned}$$

$$\tilde{E}(\mathbf{w}) = \frac{1}{2} \sum_{n=1}^N \{y(x_n, \mathbf{w}) - t_n\}^2 + \frac{\lambda}{2} \|\mathbf{w}\|^2$$

Learned function is nonlinear

Polynomial Regression with Regularization



Learned function is **nonlinear**

$$\begin{aligned}\hat{y} &= \beta_0 \\ \hat{y} &= \beta_0 + \beta_1 x \\ \hat{y} &= \beta_0 + \beta_1 x + \beta_2 x^2 \\ \hat{y} &= \beta_0 + \beta_1 x + \beta_2 x^2 + \beta_3 x^3 \\ \hat{y} &= \beta_0 + \beta_1 x + \beta_2 x^2 + \beta_3 x^3 + \dots\end{aligned}$$

$$\tilde{E}(\mathbf{w}) = \frac{1}{2} \sum_{n=1}^N \{y(x_n, \mathbf{w}) - t_n\}^2 + \frac{\lambda}{2} \|\mathbf{w}\|^2$$

Model (still) **linear**



Polynomial Regression with Regularization

$$\tilde{E}(\mathbf{w}) = \frac{1}{2} \sum_{n=1}^N \{y(x_n, \mathbf{w}) - t_n\}^2 + \frac{\lambda}{2} \|\mathbf{w}\|^2$$

*Objective function with the
Regularizer!*



Polynomial Regression with Regularization

$$\tilde{E}(\mathbf{w}) = \frac{1}{2} \sum_{n=1}^N \{y(x_n, \mathbf{w}) - t_n\}^2 + \frac{\lambda}{2} \|\mathbf{w}\|^2$$

$$\tilde{E}(\mathbf{W}) = \frac{1}{2} \sum_{n=1}^N \{y(x_n, \mathbf{w}) - t_n\}^2 + \frac{\lambda}{2} \|\mathbf{w}\|_2^2$$

The same but with slightly different notation!



Polynomial Regression with Regularization

$$\tilde{E}(\mathbf{w}) = \frac{1}{2} \sum_{n=1}^N \{y(x_n, \mathbf{w}) - t_n\}^2 + \frac{\lambda}{2} \|\mathbf{w}\|^2$$

$$\tilde{E}(\mathbf{W}) = \frac{1}{2} \sum_{n=1}^N \{y(x_n, \mathbf{w}) - t_n\}^2 + \frac{\lambda}{2} \|\mathbf{w}\|_2^2$$

L2 Regularizer

The same but with slightly different notation!



Polynomial Regression with Regularization

$$\tilde{E}(\mathbf{w}) = \frac{1}{2} \sum_{n=1}^N \{y(x_n, \mathbf{w}) - t_n\}^2 + \frac{\lambda}{2} \|\mathbf{w}\|^2$$

$$\tilde{E}(\mathbf{W}) = \frac{1}{2} \sum_{n=1}^N \{y(x_n, \mathbf{w}) - t_n\}^2 + \frac{\lambda}{2} \|\mathbf{w}\|_2^2$$

The same but with slightly different notation!

L2 Regularizer

Ridge Regression

Polynomial Regression with Regularization

$$\tilde{E}(\mathbf{w}) = \frac{1}{2} \sum_{n=1}^N \{y(x_n, \mathbf{w}) - t_n\}^2 + \frac{\lambda}{2} \|\mathbf{w}\|^2$$

$$\tilde{E}(\mathbf{W}) = \frac{1}{2} \sum_{n=1}^N \{y(x_n, \mathbf{w}) - t_n\}^2 + \frac{\lambda}{2} \|\mathbf{w}\|_2^2$$

The same but with slightly different notation!

L2 Regularizer

Ridge Regression

$$\tilde{E}(\mathbf{W}) = \frac{1}{2} \sum_{n=1}^N \{y(x_n, \mathbf{w}) - t_n\}^2 + \frac{\lambda}{2} \|\mathbf{w}\|_1$$

L1 Regularizer

Polynomial Regression with Regularization

$$\tilde{E}(\mathbf{w}) = \frac{1}{2} \sum_{n=1}^N \{y(x_n, \mathbf{w}) - t_n\}^2 + \frac{\lambda}{2} \|\mathbf{w}\|^2$$

$$\tilde{E}(\mathbf{W}) = \frac{1}{2} \sum_{n=1}^N \{y(x_n, \mathbf{w}) - t_n\}^2 + \frac{\lambda}{2} \|\mathbf{w}\|_2^2$$

The same but with slightly different notation!

L2 Regularizer

Ridge Regression

$$\tilde{E}(\mathbf{W}) = \frac{1}{2} \sum_{n=1}^N \{y(x_n, \mathbf{w}) - t_n\}^2 + \frac{\lambda}{2} \|\mathbf{w}\|_1$$

L1 Regularizer

Lasso Regression



Polynomial Regression with Regularization

$$\tilde{E}(\mathbf{w}) = \frac{1}{2} \sum_{n=1}^N \{y(x_n, \mathbf{w}) - t_n\}^2 + \frac{\lambda}{2} \|\mathbf{w}\|^2$$

$$\tilde{E}(\mathbf{W}) = \frac{1}{2} \sum_{n=1}^N \{y(x_n, \mathbf{w}) - t_n\}^2 + \frac{\lambda}{2} \|\mathbf{w}\|_2^2$$

L2 Regularizer

Ridge Regression

$$\tilde{E}(\mathbf{W}) = \frac{1}{2} \sum_{n=1}^N \{y(x_n, \mathbf{w}) - t_n\}^2 + \frac{\lambda}{2} \|\mathbf{w}\|_1$$

L1 Regularizer

Lasso Regression

How to choose lambda (λ)? Through CV



QA



Classification

- General Idea (two steps process)
 - LR (Bias Only)
 - LR (general)



Notebook presentation

- Without regularizer
- With regularizer

Predictive modeling: [Regression \(diabetes\)](#)

Predictive modeling: [Classification](#)



Break!



Maximum Likelihood Learning




Regularization

Least Squares

$$E(\mathbf{w}) = \frac{1}{2} \sum_{n=1}^N \{y(x_n, \mathbf{w}) - t_n\}^2$$

MLE EQN

Regularization



$$E(\mathbf{w}) = \frac{1}{2} \sum_{n=1}^N \{y(x_n, \mathbf{w}) - t_n\}^2$$

Regularizer

$$\tilde{E}(\mathbf{w}) = \frac{1}{2} \sum_{n=1}^N \{y(x_n, \mathbf{w}) - t_n\}^2 + \frac{\lambda}{2} \|\mathbf{w}\|^2$$

Our model got a little bigger: 2 params to M param



GPT

I know one of your tricks; get you soon!!



Our model yesterday

Our model got a little bigger: 2 params to M param



GPT

I know one of your tricks; get you soon!!



Our model today

LR to Polynomial Regression

- Polynomial function
 - M is the order/degree of polynomial ..
 - **Where to stop? What is the best M ?**





Good news is our gradient descent (iterative learning) remains the same!

$$\hat{y} = \beta_0 + \beta_1 x$$

$$\hat{y} = \beta_0 + \beta_1 x + \beta_2 x^2$$

$$\hat{y} = \beta_0 + \beta_1 x + \beta_2 x^2 + \beta_3 x^3$$

$$\hat{y} = \beta_0 + \beta_1 x + \beta_2 x^2 + \beta_3 x^3 + \dots$$



LR to Polynomial Regression

- Polynomial function
 - M is the order ..
 - **Where to stop? What is the best M?**


- Good news is our gradient descent (iterative learning) remains the same!
- You only need to change your objective function (from LR to Polynomial LR)

$$\hat{y} = \beta_0 + \beta_1 x$$


$$\hat{y} = \beta_0 + \beta_1 x + \beta_2 x^2$$

$$\hat{y} = \beta_0 + \beta_1 x + \beta_2 x^2 + \beta_3 x^3$$

$$\hat{y} = \beta_0 + \beta_1 x + \beta_2 x^2 + \beta_3 x^3 + \dots$$



LR to Polynomial Regression


$$\begin{aligned}\hat{y} &= \beta_0 + \beta_1 x \\ \hat{y} &= \beta_0 + \beta_1 x + \beta_2 x^2 \\ \hat{y} &= \beta_0 + \beta_1 x + \beta_2 x^2 + \beta_3 x^3 \\ \hat{y} &= \beta_0 + \beta_1 x + \beta_2 x^2 + \beta_3 x^3 + \dots\end{aligned}$$

Model

$$\hat{y} = \beta_0 + \beta_1 x$$

$$\Theta = \{\beta_0, \beta_1\}$$

$$\epsilon = |\hat{y} - y|$$

Optimization function

$$E_{\Theta} = \frac{1}{2} \sum_{i=1}^N (\hat{y}_i - y_i)^2$$

$$\Theta^* = \operatorname{argmin}_{\Theta} E\{(x_i, y_i)\}_{i=1, \dots, N}$$