

# On Professor Smale's legacy for asymptotic stability theory<sup>1</sup>

Matthew D. Kvalheim

Department of Mathematics and Statistics  
University of Maryland, Baltimore County

*kvalheim@umbc.edu*

**Slides are available at my website:** [mdkvalheim.github.io](https://mdkvalheim.github.io)

---

<sup>1</sup>Funding from AFOSR award FA9550-24-1-0299 is gratefully acknowledged.

# Motivation

- ▶ Asymptotic stability is a fundamental dynamical systems concept modeling robust steady-state behavior.
- ▶ Engineers often try to make systems asymptotically stable through feedback control. [▶ Example](#)

# Motivation

- ▶ Asymptotic stability is a fundamental dynamical systems concept modeling robust steady-state behavior.
- ▶ Engineers often try to make systems asymptotically stable through feedback control. ▶ Example

**Goal:** allow stabilizing feedback control to depend on parameters to shape transient response (when perturbed from steady state).

**Question:** when can parametric stabilizing feedback be extended from subset of parameter space to all of parameter space?

**Insight:** given sufficient control authority, the answer depends only on the topology of the space of asymptotically stable vector fields.

# Asymptotic stability

Consider an ordinary differential equation


$$\dot{x}(t) = F(x(t)), \quad (1)$$

where  $F$  is a vector field on  $\mathbb{R}^n$ . Unless stated otherwise,  $F$  and everything else in this talk is smooth ( $C^\infty$ ).

Let  $e \in \mathbb{R}^n$  be an **equilibrium**, meaning  $F(e) = 0$ .

We say that  $e \in \mathbb{R}^n$  is (globally) **asymptotically stable** if

- ▶ every solution of (1) converges to  $e$  as  $t \rightarrow \infty$ .
- ▶ for every open  $U \ni e$  there is a smaller open  $V \ni e$  s.t. every solution of (1) starting in  $V$  at  $t = 0$  stays in  $U$  for all  $t \geq 0$ .



# Lyapunov functions

- ▶ A **Lyapunov function** for a vector field  $F$  with equilibrium  $e$  is a proper function  $L: \mathbb{R}^n \rightarrow [0, \infty)$  such that  $L^{-1}(0) = \{e\}$  and  $dL(x) \cdot F(x) \leq 0$  for all  $x$  with equality iff  $x = e$ .
- ▶ **History:**
  - ▶ Lyapunov (1892) discovered:  
Lyapunov function exists  $\implies e$  is asymptotically stable.
  - ▶ Massera (1956), Kurzweil (1956) proved converse:  
 $e$  is asymptotically stable  $\implies (C^\infty)$  Lyapunov function exists.
  - ▶ Wilson (1967) studied the topology of level sets of such Lyapunov functions. In particular, they are  $C^\infty$  spheres for  $n > 5$  by the h-cobordism theorem of Smale (1962).
- ▶ **In this talk**, we turn Wilson's idea on its head:

We use level sets of Lyapunov functions to study the topology of the space of asymptotically stable vector fields.

# Outline

Main results

Proof step 1: equivalence of the space of Lyapunov functions and the nonlinear Grassmannian of unparametrized discs

Step 2: trivial homotopy groups of the nonlinear Grassmannian

Step 3: equivalence of the spaces of asymptotically stable vector fields and Lyapunov functions

Other applications

# Outline

## Main results

Topology of  $\mathcal{L}(\mathbb{R}^n)$  and  $\mathcal{S}(\mathbb{R}^n)$

Boundary value problems

Proof step 1: equivalence of the space of Lyapunov functions and the nonlinear Grassmannian of unparametrized discs

Step 2: trivial homotopy groups of the nonlinear Grassmannian

Step 3: equivalence of the spaces of asymptotically stable vector fields and Lyapunov functions

Other applications

## Results: topology and boundary value problems

$\mathcal{S}(\mathbb{R}^n) := \{\text{asymptotically stable vector fields on } \mathbb{R}^n\}$

$\mathcal{L}(\mathbb{R}^n) := \{\text{proper functions } \mathbb{R}^n \rightarrow [0, \infty) \text{ w/ unique critical value} = 0\}$

Equip both spaces with the compact-open  $C^\infty$  topology.

**Theorem (K 2025).**  $\mathcal{S}(\mathbb{R}^n)$  and  $\mathcal{L}(\mathbb{R}^n)$  are both path-connected and simply connected if  $n \neq 4, 5$ , and weakly contractible if  $n < 4$ .



## Results: topology and boundary value problems

$\mathcal{S}(\mathbb{R}^n) := \{\text{asymptotically stable vector fields on } \mathbb{R}^n\}$

$\mathcal{L}(\mathbb{R}^n) := \{\text{proper functions } \mathbb{R}^n \rightarrow [0, \infty) \text{ w/ unique critical value} = 0\}$

Equip both spaces with the compact-open  $C^\infty$  topology.

**Theorem (K 2025).**  $\mathcal{S}(\mathbb{R}^n)$  and  $\mathcal{L}(\mathbb{R}^n)$  are both path-connected and simply connected if  $n \neq 4, 5$ , and weakly contractible if  $n < 4$ .

$\Downarrow$

**BVP existence theorem.** For any compact manifold  $P$  and  $C^\infty$


$$F_\partial: \partial P \times \mathbb{R}^n \rightarrow \mathbb{R}^n, \quad L_\partial: \partial P \times \mathbb{R}^n \rightarrow [0, \infty)$$

s.t.  $F_\partial(p, \cdot) \in \mathcal{S}(\mathbb{R}^n)$ ,  $L_\partial(p, \cdot) \in \mathcal{L}(\mathbb{R}^n) \forall p \in \partial P$ , there exist  $C^\infty$

$$F: P \times \mathbb{R}^n \rightarrow \mathbb{R}^n, \quad L: P \times \mathbb{R}^n \rightarrow [0, \infty)$$

extending  $F_\partial, L_\partial$  s.t.  $F(p, \cdot) \in \mathcal{S}(\mathbb{R}^n)$ ,  $L(p, \cdot) \in \mathcal{L}(\mathbb{R}^n) \forall p \in P$  if either (i)  $n < 4$  or (ii)  $n > 5$  and  $\dim P < 3$ .

## Illustration of previous theorem (here $n = 2 = \dim P$ )



Previous theorem: families of asymptotically stable vector fields on  $\mathbb{R}^n$  can always be extended from the boundary  $\partial P$  to the entire parameter space  $P$  if either (i)  $n < 4$  or (ii)  $n > 5$  and  $\dim P < 3$ .

# Outline

## Main results

Proof step 1: equivalence of the space of Lyapunov functions and the nonlinear Grassmannian of unparametrized discs

Relies on **Smale's h-cobordism theorem**

Step 2: trivial homotopy groups of the nonlinear Grassmannian

Step 3: equivalence of the spaces of asymptotically stable vector fields and Lyapunov functions

Other applications

# Topology of Lyapunov function sublevel sets for $n \neq 4, 5$

## Key fact:

For any  $L \in \mathcal{L}(\mathbb{R}^n)$ ,  $L^{-1}([0, 1])$  is diffeomorphic to  $D^n$  if  $n \neq 4, 5$ .

## Proof:

- ▶ The flow of  $\nabla L$  induces deformation retractions of  $L^{-1}([0, 1])$  to  $L^{-1}(0)$  and of  $\mathbb{R}^n \setminus \{L^{-1}(0)\}$  to  $L^{-1}(1)$ .
- ▶ Hence  $L^{-1}([0, 1])$  is a contractible manifold with boundary  $L^{-1}(1)$  a homotopy sphere (Wilson 1967).
- ▶ Hence  $L^{-1}([0, 1])$  is diffeomorphic to  $D^n$  for  $n \neq 4, 5$  by
  - ▶ classification of 1D and 2D manifolds for  $n = 1, 2$ ,
  - ▶ solution to 3D Poincaré conjecture (Perelman 2003) for  $n = 3$ ,
  - ▶ the **h-cobordism theorem (Smale 1962)** for  $n > 5$ . □

## The sublevel set map

Let  $\mathcal{L}_0(\mathbb{R}^n) \subset \mathcal{L}(\mathbb{R}^n)$  be subspace of functions with min at  $0 \in \mathbb{R}^n$ .


Consider the space

$$\mathrm{Gr}(D^n, \mathbb{R}^n) := \mathrm{Emb}(D^n, \mathbb{R}^n) / \mathrm{Diff}(D^n)$$

of submanifolds of  $\mathbb{R}^n$  diffeomorphic to  $D^n$ , known as a **nonlinear Grassmannian**, and its open subspace  $\mathrm{Gr}_0(D^n, \mathbb{R}^n)$  of submanifolds whose interiors contain  $0 \in \mathbb{R}^n$ .

By the previous slide, we have a well-defined **sublevel set map**

$$p: \mathcal{L}_0(\mathbb{R}^n) \rightarrow \mathrm{Gr}_0(D^n, \mathbb{R}^n), \quad p(L) := L^{-1}([0, 1]).$$



## The sublevel set map is a weak homotopy equivalence

**Theorem (K 2025).** The sublevel set map

$$p: \mathcal{L}_0(\mathbb{R}^n) \rightarrow \mathrm{Gr}_0(D^n, \mathbb{R}^n), \quad p(L) := L^{-1}([0, 1])$$

is a fiber bundle w/ weakly contractible fibers (hence also a w.h.e.).

# The sublevel set map is a weak homotopy equivalence

**Theorem (K 2025).** The sublevel set map

$$p: \mathcal{L}_0(\mathbb{R}^n) \rightarrow \text{Gr}_0(D^n, \mathbb{R}^n), \quad p(L) := L^{-1}([0, 1])$$

is a fiber bundle w/ weakly contractible fibers (hence also a w.h.e.).

## Proof sketch:


- ▶  $p$  is continuous by implicit function theorem (Hildebrandt and Graves 1927, Abraham 1967).
- ▶ Disc theorem (Palais 1960, Cerf 1961) implies  $\text{Gr}_0(D^n, \mathbb{R}^n)$  is path-connected and  $p$  is surjective.
- ▶ Each  $M \in \text{Gr}_0(D^n, \mathbb{R}^n)$  has neighborhood  $U \subset \text{Gr}_0(D^n, \mathbb{R}^n)$  and map  $\Psi: U \rightarrow \text{Diff}(\mathbb{R}^n)$  s.t.  $\Psi(M)(N) = N$  for all  $N \in U$ .
- ▶ Define  $f: p^{-1}(U) \rightarrow \mathcal{F} := p^{-1}(M)$  by  $f(L) := L \circ \Psi(p(L))$ .
- ▶ Check:  $(p, f): p^{-1}(U) \rightarrow U \times \mathcal{F}$  is a homeomorphism.
- ▶ To show that  $\mathcal{F}$  is weakly contractible...

## Weak contractibility of $\mathcal{F}$

- ▶ Since  $p: \mathcal{L}_0(\mathbb{R}^n) \rightarrow \text{Gr}_0(D^n, \mathbb{R}^n)$  is a fiber bundle over a path-connected base, it suffices to check that  $\mathcal{F} = p^{-1}(M)$  is weakly contractible for  $M = D^n$ . In this case,

$$\mathcal{F} = \{L \in \mathcal{L}_0(\mathbb{R}^n) : L^{-1}([0, 1]) = D^n\}.$$

- ▶ Any parametric family  $P \rightarrow \mathcal{F}$  is nullhomotopic to  $P \rightarrow \{x \mapsto x^2\}$  by “parting the sea” of level sets of  $L(p)$  away from  $\partial D^n = S^{n-1}$ , replacing with level sets of  $x \mapsto x^2$ .  $\square$





# Outline

Main results

Proof step 1: equivalence of the space of Lyapunov functions and the nonlinear Grassmannian of unparametrized discs

Step 2: trivial homotopy groups of the nonlinear Grassmannian

Relies on **Smale's theorem** that  $\text{Diff}_{\partial}(D^2)$  is contractible and Hatcher's proof of the **Smale conjecture** for  $\text{Diff}_{\partial}(D^3)$ .

Step 3: equivalence of the spaces of asymptotically stable vector fields and Lyapunov functions

Other applications

# Toward homotopy groups of the nonlinear Grassmannian

- ▶ Without too much trouble,

$$\mathcal{L}_0(\mathbb{R}^n) \stackrel{\text{w.h.e.}}{\simeq} \mathcal{L}(\mathbb{R}^n) \text{ and } \text{Gr}_0(D^n, \mathbb{R}^n) \stackrel{\text{w.h.e.}}{\simeq} \text{Gr}(D^n, \mathbb{R}^n),$$

so to prove main theorem for  $\mathcal{L}(\mathbb{R}^n)$ , suffices to show that the appropriate homotopy groups of  $\text{Gr}(D^n, \mathbb{R}^n)$  are trivial.

- ▶ Generalizing a theorem of Binz and Fischer (1981) based on an idea implicit in work of Weinstein (1971), Gay-Balmaz and Vizman (2014) proved that the natural quotient map

$$\text{Emb}^+(D^n, \mathbb{R}^n) \rightarrow \text{Gr}(D^n, \mathbb{R}^n), \quad f \mapsto f(D^n)$$

is a principal  $\text{Diff}^+(D^n)$ -bundle, hence also a Serre fibration, so there is a long exact sequence of homotopy groups:

$$\cdots \pi_k \text{Diff}^+(D^n) \longrightarrow \pi_k \text{Emb}^+(D^n, \mathbb{R}^n) \longrightarrow \pi_k \text{Gr}(D^n, \mathbb{R}^n) \cdots$$

## Analyzing the long exact sequence, part 1

That long exact sequence contains the segment

$$\pi_k \text{Diff}^+(D^n) \xrightarrow{\text{surjective}} \pi_k \text{Emb}^+(D^n, \mathbb{R}^n) \longrightarrow \pi_k \text{Gr}(D^n, \mathbb{R}^n)$$

$$\longrightarrow \pi_{k-1} \text{Diff}^+(D^n) \xrightarrow{\text{surjective}} \pi_{k-1} \text{Emb}^+(D^n, \mathbb{R}^n)$$

in which the indicated arrows are surjective because the diagram

$$\begin{array}{ccc} \text{Diff}^+(D^n) & \hookrightarrow & \text{Emb}^+(D^n, \mathbb{R}^n) \\ & \searrow \text{surjective on } \pi_k & \swarrow \simeq \\ & GL^+(n) & \end{array}$$

commutes: the diagonal arrows are “evaluate derivative at 0”, the right one is a homotopy equivalence by “zooming in”, and the left one is  $\pi_k$ -surjective because the composition

$$\text{Diff}^+(D^n) \longrightarrow GL^+(n) \xrightarrow{\text{Gram-Schmidt}} SO(n)$$

is  $\pi_k$ -surjective and Gram-Schmidt is a homotopy equivalence.

## Analyzing the long exact sequence, part 2: Cerf

$$\begin{array}{c} \pi_k \text{Diff}^+(D^n) \xrightarrow{\text{surjective}} \pi_k \text{Emb}^+(D^n, \mathbb{R}^n) \xrightarrow{0} \pi_k \text{Gr}(D^n, \mathbb{R}^n) \\ \downarrow \text{injective} \\ \pi_{k-1} \text{Diff}^+(D^n) \xrightarrow{\text{surjective}} \pi_{k-1} \text{Emb}^+(D^n, \mathbb{R}^n) \end{array}$$

- ▶ We have established surjectivity of the indicated arrows, so exactness  $\implies$  other arrows are 0, injective.
- ▶  $\pi_0 \text{Gr}(D^n, \mathbb{R}^n) = \{*\}$  by disc theorem (mentioned earlier).
- ▶  $\pi_0 \text{Diff}^+(D^n) = \{*\}$  for  $n > 5$  by Cerf's (1970) pseudoisotopy theorem, so taking  $k = 1$  above yields  $\pi_1 \text{Gr}(D^n, \mathbb{R}^n) = \{*\}$ .
- ▶ Remains only to show that  $\pi_k \text{Gr}(D^n, \mathbb{R}^n) = \{*\}$  for all  $k$  when  $n < 4$ . By exactness, suffices to show that the above surjections are bijections in this case.

## Completing the proof for $\mathcal{L}(\mathbb{R}^n)$ , part 1

- Consider the earlier commutative diagram

$$\begin{array}{ccc} \text{Diff}^+(D^n) & \hookrightarrow & \text{Emb}^+(D^n, \mathbb{R}^n) \\ & \searrow \text{surjective on } \pi_k & \swarrow \simeq \\ & GL^+(n) & \end{array} .$$

The top arrow is  $\pi_k$ -surjective; need to show  $\pi_k$ -injective if  $n < 4$ . Suffices to show the same for the left diagonal arrow.

- “Evaluate derivative at 0” diagram is homotopic to “evaluate derivative at  $e_1 \in \mathbb{R}^n$ ” diagram; left arrow is homotopic to

$$\text{Diff}^+(D^n) \xrightarrow{\rho} \text{Diff}^+(S^{n-1}) \xrightarrow{f} GL^+(n),$$

composition of the restriction  $\rho$  and map  $f$  given by adjoining the value and derivative at  $e_1$ .

## Completing the proof for $\mathcal{L}(\mathbb{R}^n)$ , pt 2: Smale and Hatcher

So we need to show  $\pi_k$ -injectivity of the composition

$$\mathrm{Diff}^+(D^n) \xrightarrow{\rho} \mathrm{Diff}^+(S^{n-1}) \xrightarrow{f} GL^+(n),$$

w/  $\rho$  restriction and  $f$  adjoining the value and derivative at  $e_1$ .

- ▶ First fibration theorem of Cerf (1961)  $\implies \rho$  is fiber bundle; fiber over  $\mathrm{id}_{S^{n-1}}$  is

$$\mathrm{Diff}_{\partial}(D^n) := \{\text{diffeomorphisms of } D^n \text{ that are the identity on } \partial D^n\}.$$

- ▶ This fiber is contractible for:
  - ▶  $n = 1$  by a convexity argument,
  - ▶  $n = 2$  by a **theorem of Smale (1957)**, and
  - ▶  $n = 3$  by Hatcher's (1983) proof of the **Smale conjecture (1961)**.
- ▶  $\implies \rho$  is a w.h.e., so it suffices to show that  $f$  is  $\pi_k$ -injective.

## Completing the proof for $\mathcal{L}(\mathbb{R}^n)$ , pt 3: Smale again

Claim:  $f: \text{Diff}^+(S^{n-1}) \rightarrow GL^+(n)$  is a w.h.e. for  $n < 4$ .

Trivial for  $n = 1$ , so assume  $1 < n < 4$ .

Identifying  $GL^+(n)$  with  $\underbrace{\text{Fr}^+(TS^{n-1})}_{+ \text{ frame bundle}}$ ,  $f$  factors as the composition

$$\text{Diff}^+(S^{n-1}) \rightarrow \text{Emb}^+(D_+^{n-1}, S^{n-1}) \xrightarrow{\simeq} \text{Emb}^+(\text{int}(D_+^{n-1}), S^{n-1})$$

$$\searrow \text{Fr}^+(TS^{n-1}) \quad \quad \quad \simeq$$

in which  $D_+^{n-1}$  is upper hemisphere, first two arrows are restrictions, long arrow given by adjoining value & derivative at  $e_1$ .

- ▶ Indicated arrows are well known to be w.h.e.
- ▶ First arrow is fiber bundle again by Cerf (1961), with fiber

$$\text{Diff}(S^{n-1} \text{ rel } D_+^{n-1}) \simeq \text{Diff}_{\partial}(D^{n-1})$$

weakly contractible again for  $n = 2$  by convexity and  $n = 3$  by **theorem of Smale (1957)**, so first arrow is also a w.h.e.  $\square$

# Outline

Main results

Proof step 1: equivalence of the space of Lyapunov functions and the nonlinear Grassmannian of unparametrized discs

Step 2: trivial homotopy groups of the nonlinear Grassmannian

Step 3: equivalence of the spaces of asymptotically stable vector fields and Lyapunov functions

Other applications



## Proving main theorem for $\mathcal{S}(\mathbb{R}^n)$ from $\mathcal{L}(\mathbb{R}^n)$ case

**Theorem (K 2025).**  $\mathcal{S}(\mathbb{R}^n)$  and  $\mathcal{L}(\mathbb{R}^n)$  are both path-connected and simply connected if  $n \neq 4, 5$ , and weakly contractible if  $n < 4$ .

We already sketched the proof for  $\mathcal{L}(\mathbb{R}^n) \stackrel{\text{w.h.e.}}{\simeq} \mathcal{L}_0(\mathbb{R}^n)$ .

To prove for  $\mathcal{S}(\mathbb{R}^n)$ , suffices to prove  $\mathcal{S}(\mathbb{R}^n) \stackrel{\text{w.h.e.}}{\simeq} \mathcal{L}(\mathbb{R}^n)$ :

- ▶ Consider the **negative gradient embedding**

$$-\nabla: \mathcal{L}(\mathbb{R}^n) \hookrightarrow \mathcal{S}(\mathbb{R}^n), \quad L \mapsto -\nabla L. \quad (2)$$

- ▶ Consider any compact family  $P \rightarrow \mathcal{S}(\mathbb{R}^n)$  mapping  $\partial P$  into  $-\nabla(\mathcal{L}(\mathbb{R}^n))$ .
- ▶ Wilson's (1969) converse Lyapunov theorem  $\implies$  this is homotopic to a family  $P \rightarrow -\nabla(\mathcal{L}(\mathbb{R}^n)) \subset \mathcal{S}(\mathbb{R}^n)$ .
- ▶ Use a cutoff function to make the homotopy stationary on  $\partial P$ .

**This completes proof of theorem for  $\mathcal{S}(\mathbb{R}^n) (\stackrel{\text{w.h.e.}}{\simeq} \mathcal{S}_0(\mathbb{R}^n))$ .**  $\square$

# Outline

## Main results

Proof step 1: equivalence of the space of Lyapunov functions and the nonlinear Grassmannian of unparametrized discs

Step 2: trivial homotopy groups of the nonlinear Grassmannian

Step 3: equivalence of the spaces of asymptotically stable vector fields and Lyapunov functions

## Other applications

- Partial answer to question of Conley

- Parametric Hartman-Grobman theorem for non-hyperbolic asymptotically stable equilibria


- Parametric Morse lemma for degenerate minima of functions

## A question of Conley

- ▶ Conley (1978) defined the **Conley index** & proved that two compact isolated invariant sets  $A, B$  for two flows  $\Phi, \Psi$  have isomorphic Conley indices if they are related by *continuation*.
- ▶ In particular, this is the case if there is a continuous family  $(\Theta_s)_{s \in [0,1]}$  of flows interpolating  $\Phi, \Psi$  such that the  $(\Theta_s)$ -induced flow on  $[0, 1] \times (\text{state space})$  has a compact isolated invariant set  $C$  interpolating  $A, B$ .
- ▶ Conley asked to what extent the converse is true: Given a pair of compact isolated invariant sets with isomorphic Conley indices, when are they related by continuation?


## Partial answers to Conley

- ▶ Reineck (1992): in many interesting cases (using **Smale's (1960) Morse fun./handle manipulation techniques!**).
- ▶ However, Reineck's results do not address a natural case of Conley's question: **Given a pair of asymptotically stable vector fields, when can they be interpolated by asymptotically stable vector fields?**
- ▶ Jongeneel (2024) proved that the associated (semi)flows can always be interpolated through asymptotically stable  $C^0$  semiflows for state space dimension  $n \neq 5$ .
- ▶ This does not quite tell us about homotopy through  $C^\infty$  such flows / vector fields, but the “path-connectedness” portion of our main theorem  $\implies$  this is always possible for  $n \neq 4, 5$ .
- ▶ Cutoff functions, etc  $\implies$  same answer for local version.



# Hartman-Grobman without hyperbolicity


- ▶ **Classical Hartman-Grobman (1960, 1959) theorem:** given a  $C^1$  vector field  $F$  with hyperbolic equilibrium  $e$ , there is a local homeomorphism identifying solutions of  $\dot{x} = F(x)$  with those of  $\dot{y} = Ay$  for some nonunique  $A$  ( $= DF(e)$  works).
- ▶ **Theorem (see K-Sontag 2025).** The hyperbolicity and “ $C^1$ ” assumptions can be dropped if we assume that  $e$  is asymptotically stable. Moreover, the homeomorphism is global (and a  $C^k$  diffeo on complement of  $e$  if  $F \in C^k$ ).




- ▶ Proof: use **Smale (1962)**, Freedman (1982), Perelman (2003) to identify a Lyapunov function level set of one vector field with one of the other, then extend identification via flows.

## Parametric Hartman-Grobman without hyperbolicity

- ▶ **Theorem (see K-Sontag 2025).** The hyperbolicity and “ $C^1$ ” assumptions can be dropped if we assume that  $e$  is asymptotically stable. Moreover, the homeomorphism is global (and a  $C^k$  diffeo on complement of  $e$  if  $F \in C^k$ ).



- ▶ **Theorem (K-2025).** If in previous theorem  $F_p$  depends continuously on parameter  $p \in P$ , there is a continuous family  $h_p : B \rightarrow \mathbb{R}^n$  of linearizing homeomorphisms if either (i)  $n < 4$  or (ii)  $n > 5$  and  $\dim P = 1$ .



- ▶ **Proof:** same, but instead of using Smale, Freedman, Perelman to identify Lyapunov function level sets for a pair of vector fields, use main theorem to parametrically identify Lyapunov function level sets for a pair of families of vector fields.

# Remarks on the 4-dimensional smooth Poincaré conjecture

By the way:

- ▶ The  $C^{k \geq 1}$  statement from the “Hartman-Grobman without hyperbolicity” theorem is true for  $n = 5 \iff$  the conjecture is true (K-Sontag 2025).
- ▶ If the path-connectedness statement from the main theorem is true for  $n = 5$ , then the conjecture is true (K-2025).

Of course, **Smale (1960) solved the topological Poincaré conjecture for smooth manifolds of any dimension  $> 4$ , and was awarded the 1966 Fields medal.**

**Thank you for your attention.**

This talk is based on the preprint arXiv:2503.10828:

**“Differential topology of the spaces of asymptotically stable vector fields and Lyapunov functions”, Kvalheim (2025).**

[▶ Link to preprint](#)

Slides are available at my website: [mdkvalheim.github.io](https://mdkvalheim.github.io)



# On Professor Smale's legacy for asymptotic stability theory

## Main results

- Topology of  $\mathcal{L}(\mathbb{R}^n)$  and  $\mathcal{S}(\mathbb{R}^n)$

- Boundary value problems

Proof step 1: equivalence of the space of Lyapunov functions and the nonlinear Grassmannian of unparametrized discs

- Relies on **Smale's h-cobordism theorem**

Step 2: trivial homotopy groups of the nonlinear Grassmannian

- Relies on **Smale's theorem** that  $\text{Diff}_{\partial}(D^2)$  is contractible and Hatcher's proof of the **Smale conjecture** for  $\text{Diff}_{\partial}(D^3)$ .

Step 3: equivalence of the spaces of asymptotically stable vector fields and Lyapunov functions

## Other applications

- Partial answer to question of Conley

- Parametric Hartman-Grobman theorem for non-hyperbolic asymptotically stable equilibria

- Parametric Morse lemma for degenerate minima of functions