

Existence and uniqueness of Koopman eigenfunctions near stable equilibria and limit cycles¹

Matthew D. Kvalheim (joint with Shai Revzen)

kvalheim@umich.edu

December 13, 2022

¹Funding from ARO W911NF-(14-1-0573, 17-1-0306, 18-1-0327) is gratefully acknowledged.

Relevant papers and slides available

- **Existence and uniqueness of global Koopman eigenfunctions for stable fixed points and periodic orbits.** MDK and Shai Revzen. Physica D (2021), arXiv:1911.11996.
- **Generic properties of Koopman eigenfunctions for stable fixed points and periodic orbits.** MDK, David Hong, and Shai Revzen. IFAC-PapersOnline (2021; MTNS conference cancelled), arXiv:2010.04008.
- **Slides available on my website:** mdkvalheim.github.io/assets/NOLTA2022.pdf

Motivation

$$\dot{x} = f(x), \quad x \in \mathbb{R}^n, \quad f(0) = 0, \quad 0 \text{ hyperbolically stable with basin } B \subset \mathbb{R}^n. \quad (1)$$

- $\psi: B \rightarrow \mathbb{C}$ is a **Koopman eigenfunction** if $\exists \lambda \in \mathbb{C}$ s.t.

$$\psi(x(t)) = e^{\lambda t} \psi(x_0), \quad \text{or } \dot{\psi} = \lambda \psi \text{ if } \psi \in C^1. \quad (2)$$

- Sufficiently many “independent” eigenfunctions determine an invertible change of coordinates through which (1) becomes a linear system, a drastic simplification.
- How to find eigenfunctions? If $\mu \in \mathbb{C}$ and $g: B \rightarrow \mathbb{C}$ is such that the limit²

$$g_\mu^*(x_0) = \lim_{T \rightarrow \infty} \frac{1}{T} \int_0^T g(x(t)) e^{-\mu t} dt \quad (3)$$

exists and is not identically zero, then g_μ^* is an eigenfunction with $\lambda = \mu$ in (2).

- **Questions:** Which one? Can there be more than one possible limit (modulo scalar multiplication)? Can the limit depend sensitively on g ? Other numerical issues?
- If we knew that eigenfunctions were unique, we could resolve these questions. **We will discuss uniqueness and more**, including **new convergence results** for (3).

²**Laplace average:** see §3 of Mauroy, Mezić, Moehlis “Isostables...” 2013. See also Mezić “Analysis...” 2012.

Principal eigenfunctions

- C^1 eigenfunctions determining a linearizing diffeomorphism must be **principal**:

$$d\psi_i(0) \neq 0.$$

- (**Fact**: if ψ_i is principal and $\dot{\psi}_i = \lambda\psi_i$, $d\psi_i(0)$ is left eigenvector of D_0f w/ e.val λ .)
- Thus, we will concentrate on existence & uniqueness of C^k principal eigenfunctions.
- In particular we will see that, under some conditions, principal eigenfunctions are uniquely determined by their derivatives at 0.
- (Later we will classify **all** C^∞ eigenfunctions under generic conditions, not just the principal ones.)

Counterexamples \implies some conditions are needed

- **Ex. 1.** Let $k \geq 2$ be an integer, $(x, y) \in \mathbb{R}^2$,

$$\dot{x} = -x, \quad \dot{y} = -ky.$$

Both

$$\psi_1(x, y) = y \text{ and } \psi_2(x, y) = y + x^k$$

are analytic principal eigenfunctions s.t. $d\psi_1(0) = d\psi_2(0)$,

$$\dot{\psi}_i = \lambda \psi_i \text{ with } \lambda = -k.$$

\implies **nonresonance assumptions needed** (explained later).

- **Ex. 2.** Let $a > 1$ not be an integer, $(x, y) \in \mathbb{R}^2$,

$$\dot{x} = -x, \quad \dot{y} = -ay.$$

Both

$$\psi_1(x, y) = y \text{ and } \psi_2(x, y) = y + |x|^a$$

are $C^{\lfloor a \rfloor}$ principal eigenfunctions ($\lfloor a \rfloor$ is the integer part of a) s.t. $d\psi_1(0) = d\psi_2(0)$,

$$\dot{\psi}_i = \lambda \psi_i \text{ with } \lambda = -a.$$

\implies resonance not an issue here, but **spectral spread assumptions needed** (later).

Towards C^k existence and uniqueness, step 1: reduction to discrete-time

- Henceforth assume vector field $f \in C^k$ is complete with C^k flow $(t, x) \mapsto \Phi^t(x)$.
- Can define eigenfunctions for a diffeomorphism $F: B \rightarrow B$: $\psi(F(x)) = e^\lambda \psi(x)$.
- If eigenfunctions for $F = \Phi^1$ are unique, then they are unique for f .
- If a λ -eigenfunction $\tilde{\psi}$ for $F = \Phi^1$ exists, Sternberg's trick³ \implies

$$\psi = \int_0^1 e^{-\lambda t} \tilde{\psi} \circ \Phi^t dt$$

is a λ -eigenfunction for f and $d\psi(0) = d\tilde{\psi}(0)$.

- \implies **suffices to consider discrete time**, i.e. prove existence & uniqueness for principal eigenfunctions of a diffeomorphism $F: \mathbb{R}^n \rightarrow \mathbb{R}^n$, $F(0) = 0$, 0 hyperbolically stable with basin B .
- Existence & uniqueness for $k < \infty$ plus bootstrapping yields existence & uniqueness for $k = \infty$, hence assume $k < \infty$ for now.

³cf. Lemma 4 of Sternberg, "Local contractions and a theorem of Poincaré" (1957).

Step 2: nonresonance and solving polynomial equations

- If $\mu \in \mathbb{C}$, $k \in \mathbb{N}_{\geq 1} \cup \{+\infty\}$ eigenvalues $(D_0 F) = \lambda_1, \dots, \lambda_n$ repeated with multiplicity, $(e^\lambda, D_0 F)$ is **k -nonresonant** if

$$e^\mu \neq \lambda_1^{m_1} \dots \lambda_n^{m_n}$$

whenever $m_1, \dots, m_n \in \mathbb{N}_{\geq 0}$ satisfy $2 \leq \sum_i m_i < k + 1$.

- **Key fact:** If $F \in C^k$ and $\exists w \in \mathbb{R}^n$ s.t. $w D_0 F = e^\lambda w$, k -nonresonance \implies invertibility of certain linear operators on polynomials $\implies \exists!$ polynomial $P: \mathbb{R}^n \rightarrow \mathbb{C}$ such that $P(0) = 0$ and

$$P \circ F = e^\lambda P + o(\|x\|^k), \quad \text{and } P \text{ is } \mathbb{R}\text{-valued if } e^\lambda \in \mathbb{R}.$$

- In other words, k -nonresonance \implies can Taylor expand and solve eigenfunction equation “order by order” to produce polynomial “eigenfunction up to order k ” P .
- Remains only to **find** $o(\|x\|^k)$ **remainder** $\varphi: \mathbb{R}^n \rightarrow \mathbb{C}$ such that $\psi = P + \varphi$ is an eigenfunction exactly.

Step 3: spectral spread and contraction mapping to eliminate remainder

- **Spectral spread** $\nu(e^\mu, D_0 F) := \min \left\{ r \in \mathbb{R} : |e^\mu| \geq \left(\max_{\lambda \in \text{evals}(D_0 F)} |\lambda| \right)^r \right\}$.




Figure: Illustration of $\nu(e^\mu, D_0 F) < k$.

- **Key fact:** if $\nu(e^\lambda, D_0 F) < k$, \exists adapted norm $\|\cdot\|$ and $\varepsilon > 0$ s.t. with $N := B_\varepsilon(0)$

$$S: \{\varphi|_N \in C^k(N, \mathbb{C}) : \varphi|_N \in o(\|x\|^k)\} \circlearrowright$$

$$S(\varphi|_N) := -P|_N + e^{-\lambda}(P|_N + \varphi|_N) \circ F$$

is a contraction mapping $\implies \exists! \varphi|_N$ s.t. $S(\varphi|_N) = \varphi|_N = \lim_{m \rightarrow \infty} S^m(\tilde{\varphi}|_N)$, i.e.

$$\underbrace{(P|_N + \varphi|_N)}_{\psi|_N} \circ F = e^\lambda \underbrace{(P|_N + \varphi|_N)}_{\psi|_N} \quad \text{and} \quad \psi|_N = \lim_{m \rightarrow \infty} e^{-\lambda m} P \circ F|_N.$$

Step 4: globalization \implies discrete-time existence and uniqueness result

- Can globalize $\psi|_N: N \rightarrow \mathbb{C}$ to $\psi: B \rightarrow \mathbb{C}$ as follows: set $\psi(x) := e^{-m\lambda}\psi|_N \circ F^m(x)$ where m is large enough that $F^m(x) \in N$; can show well-defined independent of m .⁴
- **Theorem:** let $k \geq 1$, $F \in C^k(\mathbb{R}^n, \mathbb{R}^n)$, $F(0) = 0$, 0 hyperbolically stable with basin B , (e^λ, D_0F) k -nonresonant, $\nu(e^\lambda, D_0F) < k$, and $wD_0F = e^\lambda w$. Then there **exists** a **unique** C^k principal eigenfunction ψ satisfying $\psi \circ F = e^\lambda \psi$, and moreover

$$\psi = \lim_{m \rightarrow \infty} e^{-\lambda m} P \circ F \quad C^k\text{-uniformly on compacts if } P \circ F = e^\lambda P + o(\|x\|^k). \quad (4)$$

- **Observation:** (4) \implies Theorem hypotheses \implies **convergence of Laplace average**

$$\psi = \lim_{M \rightarrow \infty} \frac{1}{M} \sum_{m=1}^M e^{-\lambda m} P \circ F.$$

⁴Similar techniques are used in Lan and Mezić (2013); Kvalheim, Eldering, and Revzen (2018).

Continuous-time existence and uniqueness with weaker nonresonance

- If $\text{evals}(D_0 f) = \lambda_1, \dots, \lambda_n$ with multiplicity and $F = \Phi^1$, taking logarithm of $e^\mu \neq e^{m_1 \lambda_1} \dots e^{m_n \lambda_n}$ implies that k -nonresonance of $(e^\mu, D_0 F)$ is equivalent to

$$\mu \neq m_1 \lambda_1 + \dots m_n \lambda_n + i 2\pi \ell \quad (5)$$

for any $\ell \in \mathbb{Z}$ and any $m_1, \dots, m_n \in \mathbb{N}_{\geq 0}$ satisfying $2 \leq \sum_i m_i < k + 1$.

- By replacing $F = \Phi^1$ with $F = \Phi^\tau$ for arbitrary $\tau > 0$, (5) becomes

$$\mu \neq m_1 \lambda_1 + \dots m_n \lambda_n + i \frac{2\pi}{\tau} \ell \quad (6)$$

which can be violated for all τ if and only if it is violated for $\ell = 0$.

- \implies **Theorem**:⁵ let $k \geq 1$, vector field $f \in C^k(\mathbb{R}^n, \mathbb{R}^n)$, $f(0) = 0$, 0 hyperbolically stable with basin B , $\nu(e^\lambda, e^{D_0 f}) < k$, λ not equal to any integer linear combination of eigenvalues of $D_0 f$ with coefficient sum ≥ 2 , and $w D_0 f = \lambda w$. Then there **exists** a **unique** C^k principal eigenfunction ψ satisfying $\psi \circ \Phi^t = e^{\lambda t} \psi$ for all $t \in \mathbb{R}$, and

$$\psi = \lim_{m \rightarrow \infty} e^{-\lambda t} P \circ \Phi^t \quad C^k\text{-uniformly on compacts if } P \circ \Phi^1 = e^\lambda P + o(\|x\|^k). \quad (7)$$

- Observation**: (4) \implies Theorem hypotheses \implies **convergence of Laplace average**

$$\psi = \lim_{T \rightarrow \infty} \frac{1}{T} \int_0^T e^{-\lambda t} P \circ \Phi^t.$$

⁵see Remark 3 of Kvalheim and Revzen (2021) or Proposition 11 of Kvalheim, Hong, and Revzen (2021).

Classification of C^∞ Koopman eigenfunctions

- **Key tool:**⁶ under assumptions of preceding Theorem, if $\varphi \in C^k(B, \mathbb{C})$ satisfies $\varphi \circ \Phi^1 = e^\lambda \varphi$ and $\varphi \in o(\|x\|^k)$, then $\varphi \equiv 0$. In particular, if $\varphi = \psi_1 - \psi_2$, $\psi_1 = \psi_2$.
- Key tool & preceding theorem can be used to prove the following.
- **Classification theorem:** let vector field $f \in C^\infty(\mathbb{R}^n, \mathbb{R}^n)$, $f(0) = 0$, 0 hyperbolically stable with basin B , λ not equal to any integer linear combination of eigenvalues of $D_0 f$ with coefficient sum ≥ 2 , and $D_0 f$ diagonalizable over \mathbb{C} . Then
 - ▶ any Koopman λ -eigenfunction is a finite linear combination of products of n principal eigenfunctions and their complex conjugates.
 - ▶ In particular, λ is a linear combination of eigenvalues of $D_0 f$.

⁶Proposition 1 from Kvalheim and Revzen (2021).

Extension to periodic orbits

- Consider $\dot{x} = f(x)$ having a hyperbolically stable τ -periodic limit cycle with image Γ .
- Apply discrete-time versions of preceding theorems to a Poincaré map with section given by an isochron \implies existence and uniqueness theorems for C^k principal eigenfunctions (those with derivatives nonvanishing on Γ).
- Corresponding classification theorem has a twist involving the unique C^∞ **asymptotic phase** eigenfunction ψ_θ satisfying⁷ $\dot{\psi}_\theta = i\frac{2\pi}{\tau}\psi_\theta$.
- **Classification theorem:** let $f \in C^\infty$ and assume that no Floquet multiplier is an integer linear combination of the others with integer coefficient sum ≥ 2 . Then
 - ▶ **any** Koopman λ -eigenfunction is a finite linear combination of products of $(n-1)$ principal eigenfunctions and ψ_θ and their complex conjugates.
 - ▶ In particular, e^λ is a product of Floquet multipliers.

⁷See Mauroy and Mezić “On the use of Fourier...” (2012), Kvalheim and Revzen (2021).

Remarks on other results from Kvalheim and Revzen (2021)

- Results are given for both continuous-time and discrete-time.
- Main theorem is actually existence/uniqueness of general linearizing **semiconjugacies** (or **factors**): maps $\psi: B \rightarrow \mathbb{C}^m$ s.t. $\psi \circ \Phi^t = e^{At}\psi$ with $A \in \mathbb{C}^{m \times m}$.
- Application in paper: improvements of the Sternberg linearization and Floquet normal form theorems, with uniqueness statement, without assuming diagonalizable linearized dynamics.
- Paper considers $\psi \in C^{k,\alpha}$, i.e. $\psi \in C^k$ such that $D^k\psi$ is locally Hölder continuous with exponent α . With this, results become fairly sharp (as examples in paper show).
- Stronger uniqueness-only statements in paper only require C^1 (not C^k) dynamics, but existence no longer guaranteed for merely C^1 dynamics.
- Paper discusses in detail implications for **isostables** and **isostable coordinates** from literature.
- Also, see Schlosser and Korda “Sparsity structures for Koopman and Perron-Frobenius operators”, SIADS (2022) for an interesting application of the uniqueness results.

Thank you for your time and attention, and thank you to the organizers Milan Korda, Alexandre Mauroy, Igor Mezić, and Yoshihiko Susuki for their kind invitation.