


Approaches to assessing causality in longitudinal data

Mathew Marques
School of Psychology and Public Health
6 November 2026

Methodologies for assessing causality

Experimental design

Longitudinal design ?

Cross-lagged panel model (CLPM) (Campbell, 1963; Kenny, 1975)

Random Intercepts cross-lagged panel model (Hamaker et al., 2015)

Extensions of the RI-CLPM (Mulder & Hamaker, 2021)

Some Logical Assumptions (Mill, 1843)

Time precedence of the cause: The cause has to precede the effect

Covariation: The cause has to be demonstrably related to the effect

There are **no plausible alternative explanations** of the effect

LA TROBE UNIVERSITY
All kinds of clever

Methodological Assumptions

To compare constructs across time, need to establish measurement invariance

Configural invariance (pattern of fixed & free parameters is equivalent across time)

The same questions represent each construct at time points

Weak / metric invariance (respective factor loadings are equal across time)

Each question relates to its construct with the same strength at timepoints

Strong / scalar invariance (respective indicator means are equal across time) *

The item intercepts are equal across time: people with the same underlying level of the construct are expected to give the same average response to each at timepoints

Strict / residual factorial invariance (indicator residuals are exactly equal across time) #

(Osborne & Little , 2024)


LA TROBE UNIVERSITY
All kinds of clever

Cross-lagged panel model

Method to analyse panel data where two or more variables are repeatedly measured at two or more different timepoints

Four variables, there are 6 possible relations

- 2 cross-sectional correlations
- 2 auto-regressive relationships
- 2 cross-lagged relationships



Goals of traditional CLPMS

How stable are two (or more) constructs (autoregressive effects) ?

Does variable X predict changes in variable Y over time (cross-lagged effects)?

Is there a bi-directional relationship between variables X and Y ?

Does variable X have causal predominance over variable Y (Granger causality) ?

(Osborne & Little , 2024)

Social Media Use and Body Dissatisfaction

Body Image 40 (2021) 1–11

Contents lists available at ScienceDirect

Body Image

journal homepage: www.journals.elsevier.com/body-image

A prospective examination of relationships between social media use and body dissatisfaction in a representative sample of adults

Mathew D. Marques[a,*], Susan J. Paxton[a], Siân A. McLean[a], Hannah K. Jarman[a], Chris G. Sibley[b]

[a] School of Psychology and Public Health, La Trobe University, Melbourne, Victoria 3086, Australia
[b] School of Psychology, The University of Auckland, New Zealand

ARTICLE INFO

Article history:
Received 16 February 2021
Received in revised form 20 October 2021
Accepted 25 October 2021
Available online xxxx

Keywords:
Body dissatisfaction
Social media use
Adults
Men and women
Longitudinal

ABSTRACT

This study examined the temporal sequence of the relationship between social media use and body dissatisfaction in adults. A representative sample of adults (19–92 years old; *M* = 52.83, *SD* = 13.43; 62.02% women, 37.98% men) completed measures of social media use, body dissatisfaction, age, gender, BMI, and demographic variables in 2015, 2016, 2017, 2018 and 2019 (*N* = 6258) in the New Zealand Attitudes and Values Study. In the full sample, higher social media use was significantly associated with higher body dissatisfaction one year later, as was higher body dissatisfaction with higher social media use one year later after controlling for body dissatisfaction/social media use (T-1), gender, age, BMI, ethnicity, relationships status, and SES. Effects were small. The prospective pathway from social media use to body dissatisfaction was significant in all age groups but the reverse relationship was only significant in the middle aged and older groups. Both pathway directions were significant in women but only the pathway from body dissatisfaction to social media use was significant in men. The research has limitations and replication is required. However, findings suggest raising awareness about how to use social media positively across the broad community, not merely in adolescents, may be worthwhile.

© 2021 Elsevier Ltd. All rights reserved.

1. Introduction

Social media is a social environment which combines both peer and media aspects. It is a highly engaging, visual, and interactive environment (Perloff, 2014) in which young people spend increasing amounts of time (Auxier & Anderson, 2021). Young adults spend an average of three hours per day on social media (Mingoia, Hutchinson, Gleaves, & Wilson, 2019), although less time is reported with increasing age (Hayes, van Stolk-Cooke, & Muench, 2015). Benefits of social media engagement include opportunities for self-expression, strengthening friendships and social support, and pursuing interests (Weinstein, 2018). However, there is growing cross-sectional (Kelly, Zilanawala, Booker, & Sacker, 2018) and mixed longitudinal (Booker, Kelly, & Sacker, 2018; Coyne, Rogers, Zurcher, Stockdale, & Booth, 2020; Orben, Dienlin, & Przybylski, 2019; Viner et al., 2019) evidence that elevated levels of social media use are associated with poorer mental health in adolescents, although few studies have been conducted in adult samples. One area of well-being that has received particular attention in this regard is body dissatisfaction.


Body dissatisfaction refers to negative thoughts, feelings and perception of one's body (Grogan, 2008) and is frequently assessed as general dissatisfaction with one's appearance or body parts, or weight and shape concerns (Saiphoo & Vahedi, 2019). Body dissatisfaction is not only a negative immediate experience associated with distress and impaired physical and psychosocial quality of life (Griffiths et al., 2016), but also a risk factor for the development of depressive and anxiety symptoms, health risk behaviors (Bornioli, Lewis-Smith, Smith, Slater, & Bray, 2019; Paxton, Neumark-Sztainer, Hannan, & Eisenberg, 2006), and importantly, disordered eating and clinical eating disorders (Stice, Gau, Rohde, & Shaw, 2017). Consequently, understanding risk factors for the development of body dissatisfaction has been a focus of research, one potential risk factor being elevated social media use.

1.1. Social media use as a risk factor for body dissatisfaction

Research into risk factors for body dissatisfaction has been guided by a number of theoretical models, including sociocultural perspectives, of which the dominant one is the tripartite influence model (Thompson, Heinberg, Altabe, & Tantleff-Dunn, 1999). This

* Corresponding author.
E-mail address: m.marques@latrobe.edu.au (M.D. Marques).

https://doi.org/10.1016/j.bodyim.2021.10.008
1740-1445/© 2021 Elsevier Ltd. All rights reserved.



Goals of traditional CLPMS

How stable are two (or more) constructs (autoregressive effects) ?

Does variable X predict changes in variable Y over time (cross-lagged effects)?

Is there a bi-directional relationship between variables X and Y ?

Does variable X have causal predominance over variable Y (Granger causality) ?

(Osborne & Little , 2024)

LA TROBE UNIVERSITY
All kinds of clever

Social Media Use and Body Dissatisfaction

Body Image 40 (2021) 1–11

Contents lists available at ScienceDirect

Body Image

journal homepage: www.journals.elsevier.com/body-image

A prospective examination of relationships between social media use and body dissatisfaction in a representative sample of adults

Mathew D. Marques[a,*], Susan J. Paxton[a], Siân A. McLean[a], Hannah K. Jarman[a], Chris G. Sibley[b]

[a] School of Psychology and Public Health, La Trobe University, Melbourne, Victoria 3086, Australia
[b] School of Psychology, The University of Auckland, New Zealand

ARTICLE INFO

Article history:
Received 16 February 2021
Received in revised form 20 October 2021
Accepted 25 October 2021
Available online xxxx

Keywords:
Body dissatisfaction
Social media use
Adults
Men and women
Longitudinal

ABSTRACT

This study examined the temporal sequence of the relationship between social media use and body dissatisfaction in adults. A representative sample of adults (19–92 years old; *M* = 52.83, *SD* = 13.43; 62.02% women, 37.98% men) completed measures of social media use, body dissatisfaction, age, gender, BMI, and demographic variables in 2015, 2016, 2017, 2018 and 2019 (*N* = 6258) in the New Zealand Attitudes and Values Study. In the full sample, higher social media use was significantly associated with higher body dissatisfaction one year later, as was higher body dissatisfaction with higher social media use one year later after controlling for body dissatisfaction/social media use (T-1), gender, age, BMI, ethnicity, relationships status, and SES. Effects were small. The prospective pathway from social media use to body dissatisfaction was significant in all age groups but the reverse relationship was only significant in the middle aged and older groups. Both pathway directions were significant in women but only the pathway from body dissatisfaction to social media use was significant in men. The research has limitations and replication is required. However, findings suggest raising awareness about how to use social media positively across the broad community, not merely in adolescents, may be worthwhile.

© 2021 Elsevier Ltd. All rights reserved.

1. Introduction

Social media is a social environment which combines both peer and media aspects. It is a highly engaging, visual, and interactive environment (Perloff, 2014) in which young people spend increasing amounts of time (Auxier & Anderson, 2021). Young adults spend an average of three hours per day on social media (Mingoia, Hutchinson, Gleaves, & Wilson, 2019), although less time is reported with increasing age (Hayes, van Stolk-Cooke, & Muench, 2015). Benefits of social media engagement include opportunities for self-expression, strengthening friendships and social support, and pursuing interests (Weinstein, 2018). However, there is growing cross-sectional (Kelly, Zilanawala, Booker, & Sacker, 2018) and mixed longitudinal (Booker, Kelly, & Sacker, 2018; Coyne, Rogers, Zurcher, Stockdale, & Booth, 2020; Orben, Dienlin, & Przybylski, 2019; Viner et al., 2019) evidence that elevated levels of social media use are associated with poorer mental health in adolescents, although few studies have been conducted in adult samples. One area of well-being that has received particular attention in this regard is body dissatisfaction.


Body dissatisfaction refers to negative thoughts, feelings and perception of one's body (Grogan, 2008) and is frequently assessed as general dissatisfaction with one's appearance or body parts, or weight and shape concerns (Saiphoo & Vahedi, 2019). Body dissatisfaction is not only a negative immediate experience associated with distress and impaired physical and psychosocial quality of life (Griffiths et al., 2016), but also a risk factor for the development of depressive and anxiety symptoms, health risk behaviors (Bornioli, Lewis-Smith, Smith, Slater, & Bray, 2019; Paxton, Neumark-Sztainer, Hannan, & Eisenberg, 2006), and importantly, disordered eating and clinical eating disorders (Stice, Gau, Rohde, & Shaw, 2017). Consequently, understanding risk factors for the development of body dissatisfaction has been a focus of research, one potential risk factor being elevated social media use.

1.1. Social media use as a risk factor for body dissatisfaction

Research into risk factors for body dissatisfaction has been guided by a number of theoretical models, including sociocultural perspectives, of which the dominant one is the tripartite influence model (Thompson, Heinberg, Altabe, & Tantleff-Dunn, 1999). This

* Corresponding author.
E-mail address: m.marques@latrobe.edu.au (M.D. Marques).

https://doi.org/10.1016/j.bodyim.2021.10.008
1740-1445/© 2021 Elsevier Ltd. All rights reserved.



Goals of traditional CLPMS

How stable are two (or more) constructs (autoregressive effects) ?

Does variable X predict changes in variable Y over time (cross-lagged effects)?

Is there a bi-directional relationship between variables X and Y ?

Does variable X have causal predominance over variable Y (Granger causality) ?

(Osborne & Little , 2024)

Social Media Use and Body Dissatisfaction

Body Image 40 (2021) 1–11

Contents lists available at ScienceDirect

Body Image

journal homepage: www.journals.elsevier.com/body-image

A prospective examination of relationships between social media use and body dissatisfaction in a representative sample of adults

Mathew D. Marques[a,*], Susan J. Paxton[a], Siân A. McLean[a], Hannah K. Jarman[a], Chris G. Sibley[b]

[a] School of Psychology and Public Health, La Trobe University, Melbourne, Victoria 3086, Australia
[b] School of Psychology, The University of Auckland, New Zealand

ARTICLE INFO

Article history:
Received 16 February 2021
Received in revised form 20 October 2021
Accepted 25 October 2021
Available online xxxx

Keywords:
Body dissatisfaction
Social media use
Adults
Men and women
Longitudinal

ABSTRACT

This study examined the temporal sequence of the relationship between social media use and body dissatisfaction in adults. A representative sample of adults (19–92 years old; *M* = 52.83, *SD* = 13.43; 62.02% women, 37.98% men) completed measures of social media use, body dissatisfaction, age, gender, BMI, and demographic variables in 2015, 2016, 2017, 2018 and 2019 (*N* = 6258) in the New Zealand Attitudes and Values Study. In the full sample, higher social media use was significantly associated with higher body dissatisfaction one year later, as was higher body dissatisfaction with higher social media use one year later after controlling for body dissatisfaction/social media use (T-1), gender, age, BMI, ethnicity, relationships status, and SES. Effects were small. The prospective pathway from social media use to body dissatisfaction was significant in all age groups but the reverse relationship was only significant in the middle aged and older groups. Both pathway directions were significant in women but only the pathway from body dissatisfaction to social media use was significant in men. The research has limitations and replication is required. However, findings suggest raising awareness about how to use social media positively across the broad community, not merely in adolescents, may be worthwhile.

© 2021 Elsevier Ltd. All rights reserved.

1. Introduction

Social media is a social environment which combines both peer and media aspects. It is a highly engaging, visual, and interactive environment (Perloff, 2014) in which young people spend increasing amounts of time (Auxier & Anderson, 2021). Young adults spend an average of three hours per day on social media (Mingoia, Hutchinson, Gleaves, & Wilson, 2019), although less time is reported with increasing age (Hayes, van Stolk-Cooke, & Muench, 2015). Benefits of social media engagement include opportunities for self-expression, strengthening friendships and social support, and pursuing interests (Weinstein, 2018). However, there is growing cross-sectional (Kelly, Zilanawala, Booker, & Sacker, 2018) and mixed longitudinal (Booker, Kelly, & Sacker, 2018; Coyne, Rogers, Zurcher, Stockdale, & Booth, 2020; Orben, Dienlin, & Przybylski, 2019; Viner et al., 2019) evidence that elevated levels of social media use are associated with poorer mental health in adolescents, although few studies have been conducted in adult samples. One area of well-being that has received particular attention in this regard is body dissatisfaction.


Body dissatisfaction refers to negative thoughts, feelings and perception of one's body (Grogan, 2008) and is frequently assessed as general dissatisfaction with one's appearance or body parts, or weight and shape concerns (Saiphoo & Vahedi, 2019). Body dissatisfaction is not only a negative immediate experience associated with distress and impaired physical and psychosocial quality of life (Griffiths et al., 2016), but also a risk factor for the development of depressive and anxiety symptoms, health risk behaviors (Bornioli, Lewis-Smith, Smith, Slater, & Bray, 2019; Paxton, Neumark-Sztainer, Hannan, & Eisenberg, 2006), and importantly, disordered eating and clinical eating disorders (Stice, Gau, Rohde, & Shaw, 2017). Consequently, understanding risk factors for the development of body dissatisfaction has been a focus of research, one potential risk factor being elevated social media use.

1.1. Social media use as a risk factor for body dissatisfaction

Research into risk factors for body dissatisfaction has been guided by a number of theoretical models, including sociocultural perspectives, of which the dominant one is the tripartite influence model (Thompson, Heinberg, Altabe, & Tantleff-Dunn, 1999). This

* Corresponding author.
E-mail address: m.marques@latrobe.edu.au (M.D. Marques).

https://doi.org/10.1016/j.bodyim.2021.10.008
1740-1445/© 2021 Elsevier Ltd. All rights reserved.



Goals of traditional CLPMS

How stable are two (or more) constructs (autoregressive effects) ?

Does variable X predict changes in variable Y over time (cross-lagged effects)?

Is there a bi-directional relationship between variables X and Y ?

Does variable X have causal predominance over variable Y (Granger causality) ?

(Osborne & Little , 2024)

LA TROBE UNIVERSITY
All kinds of clever

Social Media Use and Body Dissatisfaction

Body Image 40 (2021) 1–11

Contents lists available at ScienceDirect

Body Image

journal homepage: www.journals.elsevier.com/body-image

A prospective examination of relationships between social media use and body dissatisfaction in a representative sample of adults

Mathew D. Marques[a,*], Susan J. Paxton[a], Siân A. McLean[a], Hannah K. Jarman[a], Chris G. Sibley[b]

[a] School of Psychology and Public Health, La Trobe University, Melbourne, Victoria 3086, Australia
[b] School of Psychology, The University of Auckland, New Zealand

ARTICLE INFO

Article history:
Received 16 February 2021
Received in revised form 20 October 2021
Accepted 25 October 2021
Available online xxxx

Keywords:
Body dissatisfaction
Social media use
Adults
Men and women
Longitudinal

ABSTRACT

This study examined the temporal sequence of the relationship between social media use and body dissatisfaction in adults. A representative sample of adults (19–92 years old; *M* = 52.83, *SD* = 13.43; 62.02% women, 37.98% men) completed measures of social media use, body dissatisfaction, age, gender, BMI, and demographic variables in 2015, 2016, 2017, 2018 and 2019 (*N* = 6258) in the New Zealand Attitudes and Values Study. In the full sample, higher social media use was significantly associated with higher body dissatisfaction one year later, as was higher body dissatisfaction with higher social media use one year later after controlling for body dissatisfaction/social media use (T-1), gender, age, BMI, ethnicity, relationships status, and SES. Effects were small. The prospective pathway from social media use to body dissatisfaction was significant in all age groups but the reverse relationship was only significant in the middle aged and older groups. Both pathway directions were significant in women but only the pathway from body dissatisfaction to social media use was significant in men. The research has limitations and replication is required. However, findings suggest raising awareness about how to use social media positively across the broad community, not merely in adolescents, may be worthwhile.

© 2021 Elsevier Ltd. All rights reserved.

1. Introduction

Social media is a social environment which combines both peer and media aspects. It is a highly engaging, visual, and interactive environment (Perloff, 2014) in which young people spend increasing amounts of time (Auxier & Anderson, 2021). Young adults spend an average of three hours per day on social media (Mingoia, Hutchinson, Gleaves, & Wilson, 2019), although less time is reported with increasing age (Hayes, van Stolk-Cooke, & Muench, 2015). Benefits of social media engagement include opportunities for self-expression, strengthening friendships and social support, and pursuing interests (Weinstein, 2018). However, there is growing cross-sectional (Kelly, Zilanawala, Booker, & Sacker, 2018) and mixed longitudinal (Booker, Kelly, & Sacker, 2018; Coyne, Rogers, Zurcher, Stockdale, & Booth, 2020; Orben, Dienlin, & Przybylski, 2019; Viner et al., 2019) evidence that elevated levels of social media use are associated with poorer mental health in adolescents, although few studies have been conducted in adult samples. One area of well-being that has received particular attention in this regard is body dissatisfaction.


Body dissatisfaction refers to negative thoughts, feelings and perception of one's body (Grogan, 2008) and is frequently assessed as general dissatisfaction with one's appearance or body parts, or weight and shape concerns (Saiphoo & Vahedi, 2019). Body dissatisfaction is not only a negative immediate experience associated with distress and impaired physical and psychosocial quality of life (Griffiths et al., 2016), but also a risk factor for the development of depressive and anxiety symptoms, health risk behaviors (Bornioli, Lewis-Smith, Smith, Slater, & Bray, 2019; Paxton, Neumark-Sztainer, Hannan, & Eisenberg, 2006), and importantly, disordered eating and clinical eating disorders (Stice, Gau, Rohde, & Shaw, 2017). Consequently, understanding risk factors for the development of body dissatisfaction has been a focus of research, one potential risk factor being elevated social media use.

1.1. Social media use as a risk factor for body dissatisfaction

Research into risk factors for body dissatisfaction has been guided by a number of theoretical models, including sociocultural perspectives, of which the dominant one is the tripartite influence model (Thompson, Heinberg, Altabe, & Tantleff-Dunn, 1999). This

* Corresponding author.
E-mail address: m.marques@latrobe.edu.au (M.D. Marques).

https://doi.org/10.1016/j.bodyim.2021.10.008
1740-1445/© 2021 Elsevier Ltd. All rights reserved.



Cons of the CLPM

Researchers often test models that are just identified

Yields biased cross-lagged effects when constructs have a trait-like component

Confounds between-person stability and within-person change

(Osborne & Little , 2024)

LA TROBE UNIVERSITY
All kinds of clever

“if stability of the constructs is to some extent of a trait-like, time-invariant nature, the inclusion of autoregressive parameters will fail to adequately control for this. As a result, the estimates of the cross-lagged regression coefficients will be biased, which may lead to erroneous conclusions regarding the underlying causal pattern”

“we demonstrate the spurious results that may arise when using the cross-lagged panel model to analyze data that include stable, trait-like individual differences.”

(Hamaker et al., 2015)

Psychological Methods
2015, Vol. 20, No. 1, 102–116

© 2015 American Psychological Association
1082-989X/15/$12.00 http://dx.doi.org/10.1037/a0038889

A Critique of the Cross-Lagged Panel Model

Ellen L. Hamaker and Rebecca M. Kuiper
Utrecht University

Raoul P. P. P. Grasman
University of Amsterdam

The cross-lagged panel model (CLPM) is believed by many to overcome the problems associated with the use of cross-lagged correlations as a way to study causal influences in longitudinal panel data. The current article, however, shows that if stability of constructs is to some extent of a trait-like, time-invariant nature, the autoregressive relationships of the CLPM fail to adequately account for this. As a result, the lagged parameters that are obtained with the CLPM do not represent the actual within-person relationships over time, and this may lead to erroneous conclusions regarding the *presence*, *predominance*, and *sign* of causal influences. In this article we present an alternative model that separates the within-person process from stable between-person differences through the inclusion of random intercepts, and we discuss how this model is related to existing structural equation models that include cross-lagged relationships. We derive the analytical relationship between the cross-lagged parameters from the CLPM and the alternative model, and use simulations to demonstrate the spurious results that may arise when using the CLPM to analyze data that include stable, trait-like individual differences. We also present a modeling strategy to avoid this pitfall and illustrate this using an empirical data set. The implications for both existing and future cross-lagged panel research are discussed.

Keywords: cross-lagged panel, reciprocal effects, longitudinal model, trait–state models, within-person dynamics

In 1980, Rogosa's seminal article *A Critique of the Cross-Lagged Correlation* was published, which successfully conveyed the message that comparing cross-lagged correlations from longitudinal panel data is an inappropriate basis for making causal inferences.[1] One of the key insights stemming from Rogosa's article is that, if two constructs are characterized by different degrees of stability, the comparison of cross-lagged correlations may lead to spurious conclusions regarding the causal mechanism. Since then, most researchers interested in causality in panel data have abandoned cross-lagged correlations and endorsed the *cross-lagged panel model* (CLPM)—also known as the *cross-lagged path model* or the *cross-lagged regression model*—instead. In the CLPM, stability of the constructs is controlled for through the inclusion of autoregressive relationships, and it is therefore often believed that the cross-lagged regression parameters obtained with this model are the most appropriate measures for studying causality in longitudinal correlational data (e.g., Deary, Allerhand, & Der, 2009; Soenens, Luyckx, Vansteekiste, Duriez, & Goossens, 2008; Wood, Maltby, Gillett, Linley, & Joseph, 2008). Specifically, it is common practice to standardize the cross-lagged regression coefficients and compare their relative strength with determine which variable has a stronger causal influence on the other (Bentler & Speckart, 1981).

The current article forms a sequel to the warning given by Rogosa (1980), in that it will be argued that not only should we *account* for stability, but we also need to *account for the right kind* of stability. It will be shown that if stability of the constructs is to some extent of a trait-like, time-invariant nature, the inclusion of autoregressive parameters will fail to adequately control for this. As a result the estimates of the cross-lagged regression coefficients will be biased, which may lead to erroneous conclusions regarding the underlying causal pattern. This message is not novel in itself: In fact, it has been recognized repeatedly that the “omitted variable problem” may affect the estimation of the cross-lagged coefficients (e.g., Allison, 2009; Dwyer, 1983; Finkel, 1995; Heise, 1970), and diverse modeling strategies have been proposed to

Ellen L. Hamaker and Rebecca M. Kuiper, Methodology and Statistics, Faculty of Social and Behavioural Sciences, Utrecht University; Raoul P. P. P. Grasman, Psychological Methodology, University of Amsterdam.

This study was supported by the Netherlands Organization for Scientific Research (NWO; VIDI Grant 452-10-007).

Correspondence concerning this article should be addressed to Ellen L. Hamaker, Methodology and Statistics, Faculty of Social and Behavioural Sciences, Utrecht University, P.O. Box 80140, 3508 TC, Utrecht, The Netherlands. E-mail: e.l.hamaker@uu.nl

[1] While the omitted variable problem implies that we cannot make strong causal statements based on correlational data, it does not prohibit the use of the concept of *Granger causality* (Granger, 1969). However, many researchers using cross-lagged regression refrain from using the term *causal*, and use terms like *reciprocal relationship* (Erickson, Wolfe, King, King, & Sharkansky, 2001; Lindwall, Larsman, & Hagger, 2011), *role* (Ribeiroet al., 2011), *cross-domain effects* (Burt, Obradović, Long, & Masten, 2008), *exposure* (Cole et al., 2006), *impact* (Gault-Sherman, 2012), or *influence* (Green, Furrer, & McAllister, 2011), instead. It may be argued however, that these alternative terms also imply a causal mechanism, and even more so, that an interest in causality is actually the driving force behind these studies. Therefore, we decided to use the terms *causal* and *causality* in the current article, although we acknowledging that strong causal statements can only be based on experimental designs, and we should confine ourselves to the concept of Granger causality.

(Over) simplification of the issue

Which scenario below shows change?

… but *r* = 1.0 for both !

Scenario 1

ID	Time 1	Time 2	Time 3
1	5	5	5
2	3	3	3
3	2	2	2
…	…	…	…
100	7	7	7

Scenario 2

ID	Time 1	Time 2	Time 3
1	4	5	6
2	3	4	5
3	1	2	3
…	…	…	…
100	5	6	7

(Osborne & Little , 2024)

LA TROBE UNIVERSITY
All kinds of clever

Simplification of the solution

Random intercepts account for the trait-like stability of the constructs

Scenario 1

ID	Time 1	Time 2	Time 3	*M*
1	5	5	5	5
2	3	3	3	3
3	2	2	2	2
…	…	…	…	…
100	7	7	7	7

Scenario 2

ID	Time 1	Time 2	Time 3	*M*
1	4	5	6	5
2	3	4	5	4
3	1	2	3	2
…	…	…	…	…
100	5	6	7	6


(Osborne & Little , 2024)

LA TROBE UNIVERSITY
All kinds of clever

Random-Intercepts CLPM

Separates between-person stability (i.e., stable trait-like differences) from within-person (i.e., time variant state like) changes thus overcoming drawbacks of previous modelling techniques

- Between person components indicate on average associations over time
- Auto-regressive terms now indicate changes from a person's trait-level mean in each respective measure at the next timepoint (inertia or carry over effects)
- Cross-lagged terms indicate change from individuals' prior state – difference between observed and expected (grand mean + random intercept) score



Sleep and psychological distress



Sleep Health 7 (2021) 368–374

Contents lists available at ScienceDirect

Sleep Health

Journal of the National Sleep Foundation

journal homepage: sleephealthjournal.org

Increased sleep predicts annual decreases in psychological distress: Results from a 6-year longitudinal panel sample

Mathew D. Marques, PhD[a,*], Bradley J. Wright, PhD[a], Carol H.J. Lee, MSc[b], Chris G. Sibley, PhD[b]

[a] School of Psychology and Public Health, La Trobe University, Melbourne, Victoria, Australia
[b] School of Psychology, The University of Auckland, Auckland, New Zealand

ARTICLE INFO

Keywords:
Sleep duration
Psychological distress
Cross-lagged panel model
New Zealand

ABSTRACT

Objective: To investigate the annual temporal ordering of sleep quantity and psychological distress, separating between-person stability from within-person change.
Design: Random-intercepts cross-lagged panel model using 6 annual waves of longitudinal data from the New Zealand Attitudes Values Study postal questionnaire.
Participants: New Zealand Attitudes Values Study respondents in 2013, 2014, 2015, 2016, 2017, and 2018 (*N*s = 17,880; 15,757; 13,904; 21,849; 17,031; and 47,462).
Measurements: Participants were asked, "During the past month, on average, how many hours of actual sleep did you get per night?" and responded to the K6 psychological distress scale each year. They also reported their demographic characteristics.
Results: Identified longitudinal associations between sleep duration and psychological distress in a traditional cross-lagged panel model were mostly attributable to the stability of the between-person differences in sleep duration and psychological distress. We provide evidence to suggest that increased sleep duration as indicated over a short period of time (ie, 1 month) predicted lower within-person levels of psychological distress the following year. Psychological distress did not predict sleep duration, in contrast.
Conclusions: Our analyses suggest that sleep duration in this sample of New Zealanders precedes psychological distress. This is significant given the propensity for short sleep in this sample and issues of poor mental health and short sleep among low SES indigenous members of this community. The promotion of adequate sleep duration may yield positive gains in psychological well-being.
© 2020 National Sleep Foundation. Published by Elsevier Inc. All rights reserved.


Introduction

Sleep disturbances such as insomnia, sleep apnea, and narcolepsy are related with the incidence of mood disorder diagnoses (eg, major depression, bipolar disorders, and cyclothymic disorder).[1] Emerging evidence from large community prospective investigations reveal relationships between short sleep duration with self-reported psychological distress[2] and psychological distress with short sleep duration.[3] Meta analyses[4] from other prospective studies confirmed associations of both insufficient[5–10] and long sleep[7–10] duration with negative affect (ie, frequency of depressive and anxious emotions). While these large data sets have been useful, differences in study design, measurement of sleep and negative affect, quantification of "long" and "insufficient" sleep, and statistical approach, have hindered conclusions about the direction of association between sleep duration and negative affect. Given the costs to both the individual and society associated with these conditions, an understanding of the temporal order of the sleep duration-negative affect pathway is required to identify individuals "at risk" and to inform public health campaigns and evidence-based intervention.

A systematic review[1] of 9 studies (8 prospective, *N* = 7336) reported a bidirectional relationship between sleep disturbance and mood disorders in 4 studies, a unidirectional relationship between sleep disturbance and depression in 3 studies and no relationship between both variables in one study. The authors concluded there is a bidirectional relationship between insomnia and depression. While there is evidence of an association between sleep disorders and mood disorders, there is a dearth of research that has assessed the directional pathway between sleep duration and nonclinical levels of negative affect in community samples. A measure of nonspecific psychological distress such as the Kessler psychological distress scale (K6) is frequently used in community-based epidemiological research as it is a measure that captures elements of primarily depressive and anxiety-

*Corresponding author: Mathew D. Marques, School of Psychology and Public Health, La Trobe University, Kingsbury Drive Bundoora, Melbourne, Victoria 3086, Australia.
Tel.: +61394792024
E-mail address: m.marques@latrobe.edu.au (M.D. Marques).

https://doi.org/10.1016/j.sleh.2020.12.005
2352-7218/© 2020 National Sleep Foundation. Published by Elsevier Inc. All rights reserved.



Interpretations between CLPM and RI-CLPM

	TRADITIONAL CLPM	RI-CLPM
Autoregressive effect ($X_{T-1} \rightarrow X_T$)	Rank order stability of a construct (after adjusting for the cross-lagged effect) over time. "Those who are relatively high on sleep minutes at one time point tend to be relatively high on sleep minutes at a later timepoint".	Tendency for deviations from a trait-level mean to carry over into a later timepoint (after adjusting for the cross-lagged effect); often referred to as inertia. "A deviation from a person's trait level sleep duration persists as a deviation from their trait level sleep duration at a later timepoint".
Cross-lagged effect ($X_{T-1} \rightarrow Y_T$)	Extent to which one's relative position on variable X at T-1 predicts change in one's relative position on Variable Y at a later timepoint (after adjusting for the stability of Variable Y). "Relative to those lower on sleep minutes, those who are higher on sleep minutes show a larger increase in distress at a later timepoint".	Extent to which a deviation from a person's average score on Variable X predicts a deviation from their average score on Variable Y at a later assessment (after adjusting for inertia in Y). "When a person's sleep duration decreases below their trait level, they display an increase above their trait level in distress at a later point in time".

(Osborne & Little , 2024)

LA TROBE UNIVERSITY
All kinds of clever

Sleep and psychological distress



Sleep Health 7 (2021) 368–374

Contents lists available at ScienceDirect

Sleep Health

Journal of the National Sleep Foundation

journal homepage: sleephealthjournal.org

Increased sleep predicts annual decreases in psychological distress: Results from a 6-year longitudinal panel sample

Mathew D. Marques, PhD[a,*], Bradley J. Wright, PhD[a], Carol H.J. Lee, MSc[b], Chris G. Sibley, PhD[b]

[a] School of Psychology and Public Health, La Trobe University, Melbourne, Victoria, Australia
[b] School of Psychology, The University of Auckland, Auckland, New Zealand

ARTICLE INFO

Keywords:
Sleep duration
Psychological distress
Cross-lagged panel model
New Zealand

ABSTRACT

Objective: To investigate the annual temporal ordering of sleep quantity and psychological distress, separating between-person stability from within-person change.
Design: Random-intercepts cross-lagged panel model using 6 annual waves of longitudinal data from the New Zealand Attitudes Values Study postal questionnaire.
Participants: New Zealand Attitudes Values Study respondents in 2013, 2014, 2015, 2016, 2017, and 2018 (*N*s = 17,880; 15,757; 13,904; 21,849; 17,031; and 47,462).
Measurements: Participants were asked, "During the past month, on average, how many hours of actual sleep did you get per night?" and responded to the K6 psychological distress scale each year. They also reported their demographic characteristics.
Results: Identified longitudinal associations between sleep duration and psychological distress in a traditional cross-lagged panel model were mostly attributable to the stability of the between-person differences in sleep duration and psychological distress. We provide evidence to suggest that increased sleep duration as indicated over a short period of time (ie, 1 month) predicted lower within-person levels of psychological distress the following year. Psychological distress did not predict sleep duration, in contrast.
Conclusions: Our analyses suggest that sleep duration in this sample of New Zealanders precedes psychological distress. This is significant given the propensity for short sleep in this sample and issues of poor mental health and short sleep among low SES indigenous members of this community. The promotion of adequate sleep duration may yield positive gains in psychological well-being.
© 2020 National Sleep Foundation. Published by Elsevier Inc. All rights reserved.


Introduction

Sleep disturbances such as insomnia, sleep apnea, and narcolepsy are related with the incidence of mood disorder diagnoses (eg, major depression, bipolar disorders, and cyclothymic disorder).[1] Emerging evidence from large community prospective investigations reveal relationships between short sleep duration with self-reported psychological distress[2] and psychological distress with short sleep duration.[3] Meta analyses[4] from other prospective studies confirmed associations of both insufficient[5–10] and long sleep[7–10] duration with negative affect (ie, frequency of depressive and anxious emotions). While these large data sets have been useful, differences in study design, measurement of sleep and negative affect, quantification of "long" and "insufficient" sleep, and statistical approach, have hindered conclusions about the direction of association between sleep duration and negative affect. Given the costs to both the individual and society associated with these conditions, an understanding of the temporal order of the sleep duration-negative affect pathway is required to identify individuals "at risk" and to inform public health campaigns and evidence-based intervention.

A systematic review[1] of 9 studies (8 prospective, *N* = 7336) reported a bidirectional relationship between sleep disturbance and mood disorders in 4 studies, a unidirectional relationship between sleep disturbance and depression in 3 studies and no relationship between both variables in one study. The authors concluded there is a bidirectional relationship between insomnia and depression. While there is evidence of an association between sleep disorders and mood disorders, there is a dearth of research that has assessed the directional pathway between sleep duration and nonclinical levels of negative affect in community samples. A measure of nonspecific psychological distress such as the Kessler psychological distress scale (K6) is frequently used in community-based epidemiological research as it is a measure that captures elements of primarily depressive and anxiety-

*Corresponding author: Mathew D. Marques, School of Psychology and Public Health, La Trobe University, Kingsbury Drive Bundoora, Melbourne, Victoria 3086, Australia.
Tel.: +61394792024
E-mail address: m.marques@latrobe.edu.au (M.D. Marques).

https://doi.org/10.1016/j.sleh.2020.12.005
2352-7218/© 2020 National Sleep Foundation. Published by Elsevier Inc. All rights reserved.



Thoughts and considerations



CLPM requires 2T; RI-CLPM >2T

Intervals between T are equal

Can constrain lagged parameters over time

Possible to consider >2 vars @ each T

Multigroup models are possible

How to handle missing data – FIML (Enders, 2001; Enders & Bandalos, 2001)

RI-CLPM Pros and Cons

Requires relatively few waves of data (three or more)
Separates variance into between-person and within-person components
Conceptually similar to CLPM
Fits data better than traditional CLPMs (CLPM is nested under RI-CLPM)

Autoregressive coefficients are counter-intuitive (inertia, not stability)
Effects tend to be small
Ignores between-person change

(Osborne & Little , 2024)

LA TROBE UNIVERSITY
All kinds of clever



Resources

RI-CLPM and extensions in *R* and *Mplus*
https://jeroendmulder.github.io/RI-CLPM/index.html

Osborne, D., & Little, T. D. (2024). The random intercept cross-lagged panel model. In *Longitudinal structural equation modeling* (Second edition, pp. 286–337). The Guilford Press.

Knowledge → action

Let's use *R* and *lavaan* to recreate the RI-CLPM (and CLPM) presented earlier on sleep minutes and distress.

LA TROBE UNIVERSITY
All kinds of clever

CLPM vs. RI-CLPM
When does it matter?

High degree of between-person stability

Table 3
Path coefficients of the associations between sleep duration and psychological distress

		Traditional cross-lagged		Random intercepts cross-lagged	
		Panel model		Panel model	
Outcome	Predictor $_{T-1}$	*B (SE)*	95% CI	*B (SE)*	95% CI
Sleep duration					
	Sleep duration	0.631 (0.002)*	(0.627, 0.636)	0.146 (0.006)*	(0.134, 0.158)
	Psychological distress	−0.062 (0.004)*	(−0.070, −0.053)	−0.005 (0.011)	(−0.027, 0.017)
Psychological distress					
	Psychological distress	0.751 (0.002)*	(0.747, 0.755)	0.127 (0.006)*	(0.115, 0.138)
	Sleep duration	−0.012 (0.001)*	(−0.014, −0.009)	−0.009 (0.003)*	(−0.014, −0.004)
Slope contrast		0.050 (0.005)*	(0.040, 0.059)	0.004 (0.011)	(−0.017, 0.025)
Model fit					
χ^2			16457.234		526.573
df			56		53
P			<.0001		<.0001
CFI			0.841		0.995
SRMR			0.132		0.022
RMSEA			0.071 (0.070, 0.072)		0.012 (0.011, 0.013)

Notes. Slope contrast refers to difference between cross-lagged effects in the given model.
*$P < .001$.

LA TROBE UNIVERSITY
All kinds of clever

Thanks to

Professor Susan Paxton

Dr Siân McLean

Dr Hannah Jarman

Dr Brad Wright

Dr Carol Lee

Professor Chris Sibley

Associate Professor Danny Osborne



References

Campbell, D. T. (1963). Campbell, D. T. From description to experimentation: Interpreting trends as quasi-experiments. In C. W. Harris (Ed.), Problems in measuring change. Madison: University of Wisconsin Press.

Enders, C. K. (2001). A primer on maximum likelihood algorithms available for use with missing data. *Structural Equation Modeling, 8*(1), 128-141.

Enders, C. K., & Bandalos, D. L. (2001). The relative performance of full information maximum likelihood estimation for missing data in structural equation models. *Structural equation modeling, 8*(3), 430-457.

Kenny, D. A. (1975). Cross-lagged panel correlation: A test for spuriousness. *Psychological Bulletin*, *82*(6), 887-903.

Marques, M. D., Wright, B. J., Lee, C. H., & Sibley, C. G. (2021). Increased sleep predicts annual decreases in psychological distress: Results from a 6-year longitudinal panel sample. *Sleep Health, 7*(3), 368-374.

Marques, M. D., Paxton, S. J., McLean, S. A., Jarman, H. K., & Sibley, C. G. (2022). A prospective examination of relationships between social media use and body dissatisfaction in a representative sample of adults. *Body Image, 40*, 1-11.

Mill, J. S. (2012). A system of logic, ratiocinative and inductive: Being a connected view of the principles of evidence, and the methods of scientific investigation (Reprint der Original ausgabe London 1843). Cambridge University Press.

Osborne, D., & Little, T. D. (2024). The random intercept cross-lagged panel model. In *Longitudinal structural equation modeling* (Second edition, pp. 286–337). The Guilford Press.

Thank You

Questions?

m.marques@latrobe.edu.au

