

# Linear regression of the outputs of a device and optimizing the linear model using linear programming

Dristy Parveg  
Mohsin Khan  
Finland

March 20, 2017

It has been observed that the device has two outputs  $y_1, y_2 \in \mathbb{R}$  which are functions of 5 variables  $x_1, x_2, x_3, x_4, x_5 \in \mathbb{R}$ . We are interested to know the values of the variables  $x_1, x_2, x_3, x_4, x_5$  that produce the outputs  $y_1, y_2$  which are close to optimal values.

Let  $f_i : \mathbb{R} \rightarrow \mathbb{R}$  be the function that takes the variable  $x_i$  as input while the other input variables  $x_j$  for  $j \in [5] \setminus i$  are kept at constant values and  $f_i(x_i) = y_1$ . We define the function  $g_i : \mathbb{R} \rightarrow \mathbb{R}$  in the same way so that  $g_i(x_i) = y_2$ . From the result of the experiments, presented in previous section, it is evident that  $f_i$  and  $g_i$  are almost, if not exactly, linear functions.

We are interested to find models  $f : \mathbb{R}^5 \rightarrow \mathbb{R}$ ,  $g : \mathbb{R}^5 \rightarrow \mathbb{R}$  where  $f(x_1, x_2, x_3, x_4, x_5) = y_1$  and  $g(x_1, x_2, x_3, x_4, x_5) = y_2$ . We make an educated guess that both  $f$  and  $g$  are linear. We do not provide an analytical argument to justify the linearity of  $f$  and  $g$ . Instead, we use linear regression on the experimental results. There are 20 experiments. We randomly choose 15 experiments and use them as training data to obtain the linear models

$$\begin{aligned} f &= a_1x_1 + a_2x_2 + a_3x_3 + a_4x_4 + a_5x_5 \\ g &= b_1x_1 + b_2x_2 + b_3x_3 + b_4x_4 + b_5x_5 \end{aligned}$$

If  $x_1, x_2, x_3, x_4, x_5, y_1, y_2$  be an experimental result then  $\frac{|f(x_1, x_2, x_3, x_4, x_5) - y_1|}{y_1} \times 100$  and  $\frac{|f(x_1, x_2, x_3, x_4, x_5) - y_2|}{y_2} \times 100$  are the error percentages of the obtained models  $f$  and  $g$ . We compute the training error of the models using the 15 experimental results and find that the average training error is around ?? percent. We use the rest of the 5 experimental results to compute the test error of the obtained linear models. We find the error percentage to be below ?? percent in both of the cases of  $f$  and  $g$ .

We reckon, this error percentages indicate that the underlying models are linear enough and the obtained approximated linear models can be used to find the approximately optimal values of  $y_1$  and  $y_2$  and also the corresponding inputs  $x_1, \dots, x_5$  that produces outputs  $y_1, y_2$ . As the underlying models are not not exactly linear and our obtained models are only a linear approximation, the optimal values obtained from the model will have some errors. We expect these errors to be similar to the test errors of the models. In this paper we look forward to find the inputs for which  $y_1$  is maximized and  $y_2$  is minimized. In other words we look forward to maximize the expression

$$\begin{aligned}
y_1 - y_2 &= f(x_1, \dots, x_5) - g(x_1, \dots, x_5) \\
&= (a_1x_1 + a_2x_2 + a_3x_3 + a_4x_4 + a_5x_5) - \\
&\quad (b_1x_1 + b_2x_2 + b_3x_3 + b_4x_4 + b_5x_5) \\
&= (a_1 - b_1)x_1 + (a_2 - b_2)x_2 + (a_3 - b_3)x_3 + (a_4 - b_4)x_4 + (a_5 - b_5)x_5 \\
&= \sum_{i \in [5]} (a_i - b_i)x_i
\end{aligned}$$

We model this maximization problem as a linear programming problem as follows:

$$\begin{aligned}
\text{maximize} \quad & : \sum_{i \in [5]} (a_i - b_i)x_i \\
\text{with constraints} \quad & : 0 < x_i \leq c_i \text{ for } i \in [5]
\end{aligned}$$

This is a fairly small instance of a linear programming problem. There exist efficient algorithms, namely, simplex, ellipsoid, etc. for solving linear programming problems. In practice, being oblivious to the algorithm used, this linear programming problem can easily be solved by any standard linear programming solver like `lp_solve`. Let  $x'_1, x'_2, x'_3, x'_4, x'_5$  be the solution of the above linear programming problem.

We run an experiment using these input values and get output  $y'_1, y'_2$ . It can be expected that  $\frac{|f(x_1, \dots, x_5) - y'_1|}{y'_1} \times 100, \frac{|f(x_1, \dots, x_5) - y'_2|}{y'_2} \times 100$  are approximately equal to the test error of  $f$  and  $g$  respectively.