$$\bar{x} = \frac{\sum_{i=1}^{n} x_i}{n} \qquad s^2 = \frac{\sum_{i=1}^{n} (x_i - \bar{x})}{n-1}$$

$$P(A^c) = 1 - P(A) \qquad P(A \cup B) = P(A) + P(B) - P(A \cap B)$$

$$P(A|B) = \frac{P(A \cap B)}{P(B)} = \frac{P(B|A)P(A)}{P(B|A)P(A) + P(B|A^c)P(A^c)}$$

Discrete Random Variables

Let X be a discrete random variable with a pmf of p(x) then

$$E[X] = \sum_{x} x p(x)$$

Let X be a random variable, then

$$Var(X) = E[X^2] - E[X]^2$$

Linear combination of random variables X and Y and fixed numbers a and b:

$$E[aX + bY] = aE[X] + bE[Y]$$
$$Var(aX + bY) = a^{2}Var(X) + b^{2}Var(Y)$$

| Distribution | pmf | E(X) | Var(X) |
|------------------|------------------------------------|-----------------|--------------------------|
| Binomial | $\binom{n}{k}p^k(1-p)^{n-k}$ | $\mid np \mid$ | np(1-p) |
| Geometric | $(1-p)^{n-1}p$ | $\frac{1}{p}$ | $\frac{1-p}{p^2}$ |
| Discrete Uniform | $\frac{1}{n}$ | $\frac{a+b}{2}$ | $\frac{(b-a+1)^2-1}{12}$ |
| Poisson | $\frac{\lambda^k}{k!}e^{-\lambda}$ | λ | λ |

Continuous Random Variables

$$E[X] = \int_{x \in \Omega_x} x f(x) \, dx$$

| Distribution | pmf | E(X) | Var(X) |
|-------------------|-----------------------------------------------------------------|---------------------|-----------------------|
| Continous Uniform | $\frac{1}{b-a}$ | $\frac{a+b}{2}$ | $\frac{(b-a)^2}{12}$ |
| Exponential | $\lambda e^{-\lambda x}$ | $\frac{1}{\lambda}$ | $\frac{1}{\lambda^2}$ |
| Normal | $\frac{1}{\sqrt{2\pi\sigma^2}}e^{-\frac{(x-\mu)^2}{2\sigma^2}}$ | μ | σ^2 |

Central Limit Theorem

Under certain conditions:

$$\bar{x} \sim N(\mu, \frac{\sigma^2}{n})$$

$$(\bar{x}_1 - \bar{x}_2) \sim N(\mu_1 - \mu_2, \frac{\sigma_1^2}{n_1} + \frac{\sigma_2^2}{n_2})$$

$$\hat{p} \sim N(p, \frac{p(1-p)}{n})$$

$$(\hat{p}_1 - \hat{p}_2) \sim N((p_1 - p_2), \frac{p_1(1-p_1)}{n_1} + \frac{p_2(1-p_2)}{n_2})$$

Linear Regression

Maximum Likelihood