

# Native Plant Cuttings Guide

Western Australian native species – Joondalup, WA

Version 1.6 · September 2026 · 40 species

## Species in this edition (alphabetical)

Full index on page 2

|                                    |    |                                     |    |                                 |    |
|------------------------------------|----|-------------------------------------|----|---------------------------------|----|
| ● <i>Anthocercis littorea</i>      | 8  | ● <i>Hibbertia spicata</i> *        | 36 | ● <i>Rhagodia baccata</i> *     | 64 |
| ● <i>Atriplex isatidea</i> *       | 10 | ● <i>Hypocalymma robustum</i>       | 38 | ● <i>Scaevola canescens</i> *   | 66 |
| ● <i>Calothamnus quadrifidus</i> * | 12 | ● <i>Kennedia prostrata</i>         | 40 | ● <i>Scaevola crassifolia</i>   | 68 |
| ● <i>Carpobrotus virescens</i>     | 14 | ● <i>Lechenaultia linarioides</i> * | 42 | ● <i>Scaevola globulifera</i> * | 70 |
| ● <i>Eremophila glabra</i>         | 16 | ● <i>Leucophyta brownii</i> *       | 44 | ● <i>Scaevola nitida</i> *      | 72 |
| ● <i>Grevillea crithmifolia</i> *  | 18 | ● <i>Melaleuca huegelii</i> *       | 46 | ● <i>Scaevola repens</i> *      | 74 |
| ● <i>Grevillea preissii</i>        | 20 | ● <i>Melaleuca systema</i> *        | 48 | ● <i>Scaevola thesioides</i> *  | 76 |
| ● <i>Grevillea vestita</i>         | 22 | ● <i>Myoporum caprarioides</i> *    | 50 | ● <i>Spinifex hirsutus</i> *    | 78 |
| ● <i>Guichenotia ledifolia</i> *   | 24 | ● <i>Myoporum insulare</i>          | 52 | ● <i>Spinifex longifolius</i> * | 80 |
| ● <i>Hardenbergia comptoniana</i>  | 26 | ● <i>Olearia axillaris</i> *        | 54 | ● <i>Templetonia retusa</i> *   | 82 |
| ● <i>Hemiandra pungens</i>         | 28 | ● <i>Phyllanthus calycinus</i> *    | 56 | ● <i>Thomasia triphylla</i> *   | 84 |
| ● <i>Hibbertia cuneiformis</i> *   | 30 | ● <i>Pimelea calcicola</i> *        | 58 | ● <i>Threlkeldia diffusa</i> *  | 86 |
| ● <i>Hibbertia hypericoides</i>    | 32 | ● <i>Pimelea sulphurea</i> *        | 60 |                                 |    |
| ● <i>Hibbertia racemosa</i> *      | 34 | ● <i>Pithocarpa cordata</i> *       | 62 |                                 |    |

● Easy   ● Easy-Mod.   ● Moderate   ● Mod.-Difficult   ● Difficult

\* Provisional rating – based on genus behaviour; confirm with your own nursery records.

Compiled from published sources (see References). Windows and ratings are starting points for local practice.

© 2026 Mitchell Donald. All rights reserved.

# How to use this guide

Each species has its own page set, listed alphabetically by scientific name. Every page follows the same layout: quick facts, a cutting calendar, a plant snapshot, the full propagation method, common problems, and a blank record table so the nursery can log strike rates and refine the advice over time.

Timing windows are written for the Joondalup / Perth coastal climate: hot, dry summers and mild, wet winters. They combine general native-propagation practice with each species' flowering period from FloraBase. Rooting is fastest in warm weather and slows markedly in winter unless bottom heat is used.

## Species index – at a glance

| Species                                                              | Page | Difficulty      | Wood type                | Best months             | Length           | Hormone (IBA)           |
|----------------------------------------------------------------------|------|-----------------|--------------------------|-------------------------|------------------|-------------------------|
| <b><i>Anthocercis littorea</i></b><br>Yellow Tailflower              | 8    | Easy-Moderate   | Semi-hardwood            | Dec – Apr               | 75–100 mm        | 3 g/L IBA               |
| <b><i>Atriplex isatidea</i></b><br>Coast Saltbush                    | 10   | Easy-Moderate * | Semi-hardwood tips       | Sep – Nov               | 100–150 mm       | 1.5–3 g/L IBA           |
| <b><i>Calothamnus quadrifidus</i></b><br>One-sided Bottlebrush       | 12   | Easy-Moderate * | Semi-hardwood            | Jan – Apr               | 75–100 mm        | 3 g/L IBA               |
| <b><i>Carpobrotus virescens</i></b><br>Coastal Pigface               | 14   | Easy            | Succulent stem pieces    | Sep – Nov;<br>Mar – Apr | 10–15 cm (tubes) | Not needed              |
| <b><i>Eremophila glabra</i></b><br>Tar Bush / Common Emu Bush        | 16   | Easy            | Semi-hardwood            | Jan – Apr               | 75–100 mm        | 3 g/L IBA               |
| <b><i>Grevillea crithmifolia</i></b>                                 | 18   | Moderate *      | Semi-hardwood            | Dec – Apr               | 75–100 mm        | 3 g/L IBA               |
| <b><i>Grevillea preissii</i></b><br>Spider-net Grevillea             | 20   | Moderate        | Semi-hardwood            | Nov – Mar               | 75–100 mm        | 3 g/L IBA               |
| <b><i>Grevillea vestita</i></b>                                      | 22   | Easy            | Semi-hardwood            | Dec – Apr               | 80–120 mm        | 3 g/L IBA               |
| <b><i>Guichenotia ledifolia</i></b>                                  | 24   | Easy-Moderate * | Semi-hardwood            | Dec – Apr               | 50–75 mm         | 3 g/L IBA               |
| <b><i>Hardenbergia comptoniana</i></b><br>Native Wisteria            | 26   | Easy-Moderate   | Semi-hardwood            | Dec – Mar               | 50–75 mm         | 3 g/L IBA               |
| <b><i>Hemiandra pungens</i></b><br>Snakebush                         | 28   | Easy            | Softwood – semi-hardwood | Oct – Mar               | 50–75 mm         | 1.5–3 g/L IBA           |
| <b><i>Hibbertia cuneiformis</i></b><br>Cutleaf Hibbertia             | 30   | Easy-Moderate * | Semi-hardwood            | Mar – May               | 75–100 mm        | 3 g/L IBA               |
| <b><i>Hibbertia hypericoides</i></b><br>Yellow Buttercups            | 32   | Difficult       | Semi-hardwood            | Dec – Feb               | 50–75 mm         | 3 g/L IBA (trial range) |
| <b><i>Hibbertia racemosa</i></b><br>Stalked Guinea Flower            | 34   | Moderate *      | Semi-hardwood            | Jan – Apr               | 50–75 mm         | 1.5–3 g/L IBA           |
| <b><i>Hibbertia spicata</i></b>                                      | 36   | Moderate *      | Semi-hardwood            | Jan – Mar               | 50–75 mm         | 1.5–3 g/L IBA           |
| <b><i>Hypocalymma robustum</i></b><br>Swan River Myrtle              | 38   | Easy            | Semi-hardwood            | Dec – Mar               | 75–100 mm        | 3 g/L IBA               |
| <b><i>Kennedia prostrata</i></b><br>Scarlet Runner / Running Postman | 40   | Easy-Moderate   | Semi-hardwood            | Dec – Mar               | 75–100 mm        | 3 g/L IBA               |
| <b><i>Lechenaultia linarioides</i></b><br>Yellow Leschenaultia       | 42   | Easy *          | Softwood – semi-hardwood | Feb – Apr               | 50–75 mm         | 1.5–3 g/L IBA           |
| <b><i>Leucophyta brownii</i></b><br>Cushion Bush                     | 44   | Easy-Moderate * | Semi-hardwood            | Mar – May;<br>Sep – Oct | 50–75 mm         | 1.5–3 g/L IBA           |

| Species                                                              | Page | Difficulty      | Wood type                | Best months             | Length                 | Hormone (IBA)      |
|----------------------------------------------------------------------|------|-----------------|--------------------------|-------------------------|------------------------|--------------------|
| <b><i>Melaleuca huegelii</i></b><br>Chenille Honeymyrtle             | 46   | Easy *          | Semi-hardwood            | Feb – Apr               | 75–100 mm              | 3 g/L IBA          |
| <b><i>Melaleuca systema</i></b><br>Coastal Honeymyrtle               | 48   | Easy *          | Semi-hardwood            | Dec – Jan;<br>Apr – May | 75–100 mm              | 3 g/L IBA          |
| <b><i>Myoporum caprarioides</i></b><br>Slender Myoporum              | 50   | Easy *          | Semi-hardwood            | Nov – Mar               | 75–100 mm              | 3 g/L IBA          |
| <b><i>Myoporum insulare</i></b><br>Blueberry Tree / Common Boobialla | 52   | Easy            | Semi-hardwood            | Feb – Apr               | 75–100 mm              | 3 g/L IBA          |
| <b><i>Olearia axillaris</i></b><br>Coastal Daisybush                 | 54   | Easy *          | Semi-hardwood            | Aug – Oct               | 75–100 mm              | 1.5–3 g/L IBA      |
| <b><i>Phyllanthus calycinus</i></b><br>False Boronia                 | 56   | Moderate *      | Semi-hardwood            | Feb – Apr               | 50–75 mm               | 3 g/L IBA          |
| <b><i>Pimelea calcicola</i></b><br>Coastal Banjine                   | 58   | Moderate *      | Semi-hardwood            | Dec – Mar               | 50–75 mm               | 1.5–3 g/L IBA      |
| <b><i>Pimelea sulphurea</i></b><br>Yellow Banjine                    | 60   | Moderate *      | Semi-hardwood            | Dec – Mar               | 50–75 mm               | 1.5–3 g/L IBA      |
| <b><i>Pithocarpa cordata</i></b><br>Tangle Daisy                     | 62   | Easy–Moderate * | Semi-hardwood tips       | Sep – Nov;<br>May       | 50–75 mm               | 1.5–3 g/L IBA      |
| <b><i>Rhagodia baccata</i></b><br>Berry Saltbush                     | 64   | Moderate *      | Semi-hardwood tips       | Sep – Nov               | 75–100 mm              | 3 g/L IBA          |
| <b><i>Scaevola canescens</i></b><br>Grey Scaevola                    | 66   | Easy–Moderate * | Softwood – semi-hardwood | Nov – Jan               | 50–80 mm               | 1.5–3 g/L IBA      |
| <b><i>Scaevola crassifolia</i></b><br>Thick-leaved Fan-flower        | 68   | Easy            | Semi-hardwood            | Mar – May               | 75–100 mm              | 3 g/L IBA          |
| <b><i>Scaevola globulifera</i></b>                                   | 70   | Easy *          | Softwood – semi-hardwood | Jan – Apr               | 50–80 mm               | 1.5–3 g/L IBA      |
| <b><i>Scaevola nitida</i></b><br>Shining Fanflower                   | 72   | Easy *          | Semi-hardwood            | Jan – Apr               | 75–100 mm              | 3 g/L IBA          |
| <b><i>Scaevola repens</i></b>                                        | 74   | Easy *          | Softwood – semi-hardwood | Mar – May               | 50–80 mm               | 1.5–3 g/L IBA      |
| <b><i>Scaevola thesioides</i></b>                                    | 76   | Easy–Moderate * | Softwood – semi-hardwood | Apr – May               | 50–75 mm               | 1.5–3 g/L IBA      |
| <b><i>Spinifex hirsutus</i></b><br>Hairy Spinifex                    | 78   | Moderate *      | Runner (stolon) sections | Mar – May               | 10–15 cm,<br>1–2 nodes | Optional (1.5 g/L) |
| <b><i>Spinifex longifolius</i></b><br>Beach Spinifex                 | 80   | Moderate *      | Runner (stolon) sections | Mar – May               | 10–15 cm,<br>1–2 nodes | Optional (1.5 g/L) |
| <b><i>Templetonia retusa</i></b><br>Cockies Tongues                  | 82   | Easy–Moderate * | Semi-hardwood            | Dec – Mar               | 50–75 mm               | 3 g/L IBA          |
| <b><i>Thomasia triphylla</i></b>                                     | 84   | Easy–Moderate * | Semi-hardwood            | Dec – Apr               | 50–75 mm               | 3 g/L IBA          |
| <b><i>Threlkeldia diffusa</i></b><br>Coast Bonefruit                 | 86   | Moderate *      | Softwood tips            | Dec – Mar               | 50–75 mm               | 1.5 g/L IBA        |

\* Provisional rating – based on genus behaviour; update once nursery strike rates are recorded.

## Cutting calendar

| Species                         | Jan  | Feb  | Mar  | Apr  | May  | Jun | Jul | Aug  | Sep  | Oct  | Nov  | Dec  |
|---------------------------------|------|------|------|------|------|-----|-----|------|------|------|------|------|
| <i>Anthocercis littorea</i>     | Best | Best | Best | Best | OK   | –   | –   | –    | –    | –    | OK   | Best |
| <i>Atriplex isatidea</i>        | OK   | OK   | –    | –    | –    | –   | –   | OK   | Best | Best | Best | OK   |
| <i>Calothamnus quadrifidus</i>  | Best | Best | Best | Best | OK   | –   | –   | –    | –    | –    | –    | OK   |
| <i>Carpobrotus virescens</i>    | OK   | OK   | Best | Best | OK   | –   | –   | OK   | Best | Best | Best | OK   |
| <i>Eremophila glabra</i>        | Best | Best | Best | Best | OK   | –   | –   | –    | –    | –    | OK   | OK   |
| <i>Grevillea crithmifolia</i>   | Best | Best | Best | Best | OK   | –   | –   | –    | –    | –    | OK   | Best |
| <i>Grevillea preissii</i>       | Best | Best | Best | OK   | –    | –   | –   | –    | –    | OK   | Best | Best |
| <i>Grevillea vestita</i>        | Best | Best | Best | Best | OK   | –   | –   | –    | –    | –    | OK   | Best |
| <i>Guichenotia ledifolia</i>    | Best | Best | Best | Best | OK   | –   | –   | –    | –    | –    | OK   | Best |
| <i>Hardenbergia comptoniana</i> | Best | Best | Best | OK   | –    | –   | –   | –    | –    | –    | OK   | Best |
| <i>Hemiandra pungens</i>        | Best | Best | Best | OK   | –    | –   | –   | –    | OK   | Best | Best | Best |
| <i>Hibbertia cuneiformis</i>    | OK   | OK   | Best | Best | Best | –   | –   | –    | –    | OK   | OK   | OK   |
| <i>Hibbertia hypericoides</i>   | Best | Best | OK   | –    | –    | –   | –   | –    | –    | –    | OK   | Best |
| <i>Hibbertia racemosa</i>       | Best | Best | Best | Best | OK   | –   | –   | –    | –    | –    | –    | OK   |
| <i>Hibbertia spicata</i>        | Best | Best | Best | OK   | –    | –   | –   | –    | –    | –    | –    | OK   |
| <i>Hypocalymma robustum</i>     | Best | Best | Best | OK   | –    | –   | –   | –    | –    | –    | OK   | Best |
| <i>Kennedia prostrata</i>       | Best | Best | Best | OK   | –    | –   | –   | –    | –    | –    | OK   | Best |
| <i>Lechenaultia linarioides</i> | OK   | Best | Best | Best | OK   | –   | –   | –    | –    | OK   | OK   | OK   |
| <i>Leucophyta brownii</i>       | –    | OK   | Best | Best | Best | OK  | –   | OK   | Best | Best | OK   | –    |
| <i>Melaleuca huegelii</i>       | OK   | Best | Best | Best | OK   | –   | –   | –    | –    | –    | –    | OK   |
| <i>Melaleuca systema</i>        | Best | OK   | OK   | Best | Best | OK  | –   | –    | –    | –    | OK   | Best |
| <i>Myoporum caprarioides</i>    | Best | Best | Best | OK   | –    | –   | –   | –    | –    | OK   | Best | Best |
| <i>Myoporum insulare</i>        | OK   | Best | Best | Best | OK   | –   | –   | –    | –    | OK   | OK   | OK   |
| <i>Olearia axillaris</i>        | –    | –    | OK   | OK   | –    | –   | OK  | Best | Best | Best | OK   | –    |
| <i>Phyllanthus calycinus</i>    | OK   | Best | Best | Best | OK   | –   | –   | –    | –    | –    | –    | –    |
| <i>Pimelea calcicola</i>        | Best | Best | Best | OK   | –    | –   | –   | –    | –    | –    | OK   | Best |
| <i>Pimelea sulphurea</i>        | Best | Best | Best | OK   | –    | –   | –   | –    | –    | –    | OK   | Best |
| <i>Pithocarpa cordata</i>       | –    | –    | –    | OK   | Best | OK  | –   | OK   | Best | Best | Best | OK   |
| <i>Rhagodia baccata</i>         | OK   | OK   | OK   | –    | –    | –   | –   | OK   | Best | Best | Best | OK   |
| <i>Scaevola canescens</i>       | Best | OK   | OK   | OK   | OK   | –   | –   | –    | –    | OK   | Best | Best |
| <i>Scaevola crassifolia</i>     | OK   | OK   | Best | Best | Best | –   | –   | –    | –    | OK   | OK   | OK   |
| <i>Scaevola globulifera</i>     | Best | Best | Best | Best | OK   | –   | –   | –    | –    | –    | OK   | OK   |
| <i>Scaevola nitida</i>          | Best | Best | Best | Best | OK   | –   | –   | –    | –    | –    | OK   | OK   |
| <i>Scaevola repens</i>          | OK   | OK   | Best | Best | Best | –   | –   | –    | OK   | OK   | OK   | OK   |

|                             |                                                               |      |        |      |      |                                 |   |   |        |    |      |      |
|-----------------------------|---------------------------------------------------------------|------|--------|------|------|---------------------------------|---|---|--------|----|------|------|
| <i>Scaevola thesioides</i>  | OK                                                            | OK   | OK     | Best | Best | –                               | – | – | OK     | OK | OK   | OK   |
| <i>Spinifex hirsutus</i>    | OK                                                            | OK   | Best   | Best | Best | –                               | – | – | OK     | OK | OK   | OK   |
| <i>Spinifex longifolius</i> | OK                                                            | OK   | Best   | Best | Best | –                               | – | – | OK     | OK | OK   | OK   |
| <i>Templetonia retusa</i>   | Best                                                          | Best | Best   | OK   | –    | –                               | – | – | –      | –  | OK   | Best |
| <i>Thomasia triphylla</i>   | Best                                                          | Best | Best   | Best | OK   | –                               | – | – | –      | –  | OK   | Best |
| <i>Threlkeldia diffusa</i>  | Best                                                          | Best | Best   | OK   | –    | –                               | – | – | –      | –  | OK   | Best |
| Perth season                | Summer                                                        |      | Autumn |      |      | Winter                          |   |   | Spring |    | Sum. |      |
| <div></div> Best window     | <div></div> Workable – see ‘Selecting material’ in the method |      |        |      |      | <div></div> Avoid / low success |   |   |        |    |      |      |

# Key to terms and ratings

## Wood types and the bend test

| Wood type            | What it looks like                                                       | Bend test                         | Use                                                                                       |
|----------------------|--------------------------------------------------------------------------|-----------------------------------|-------------------------------------------------------------------------------------------|
| <b>Softwood</b>      | Soft, flexible, bright new growth from the current flush.                | Flops or stays bent.              | Roots fast but wilts and rots easily. Suits soft-stemmed plants such as <i>Scaevola</i> . |
| <b>Semi-hardwood</b> | Current season's growth that has begun to firm; bark starting to colour. | Bends about 60° and springs back. | The standard for most WA native shrubs. More forgiving than softwood.                     |
| <b>Hardwood</b>      | Fully mature, woody stems from previous seasons.                         | Stiff; snaps rather than bends.   | Slow to root. Rarely needed for the species in this guide.                                |

## Difficulty ratings

| Rating                    | Meaning                                                                                   |
|---------------------------|-------------------------------------------------------------------------------------------|
| <b>Easy</b>               | Strikes reliably with the standard method.                                                |
| <b>Easy-Moderate</b>      | Usually strikes; one factor (e.g. moisture, fragile stems) needs extra care.              |
| <b>Moderate</b>           | Strikes, but results depend on timing, material and moisture control. Set extra cuttings. |
| <b>Moderate-Difficult</b> | Low or variable strike; needs careful timing and good facilities.                         |
| <b>Difficult</b>          | Low or erratic strike; specialist techniques or alternative methods usually needed.       |

\* **Provisional** – rating is based on how the genus generally behaves because little species-specific data is published. Replace it with the nursery's own figure once a few batches have been recorded.

## Confidence

| Level         | Meaning                                                                                                                                        |
|---------------|------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>High</b>   | Species-specific propagation information is published and consistent.                                                                          |
| <b>Medium</b> | Built from genus-level information plus the species' own traits (habit, leaf type, flowering time). Sound starting point – verify with trials. |
| <b>Low</b>    | Little information available; treat as a trial protocol.                                                                                       |

# Standard cutting method

Species pages refer back to these steps and only note where a species differs.

| Step                | What to do                                                                                                                                                                                                                          |
|---------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| 1. Collect          | Take material early in the morning from healthy, pest-free, well-watered, <b>non-flowering</b> stock. Seal in a moist plastic bag and keep in an esky out of the sun. Process the same day.                                         |
| 2. Prepare          | Use a sharp, clean blade (wipe with alcohol between stock plants). Cut just below a node. Strip leaves from the lower half to two-thirds without tearing the bark. Reduce large leaves by about one-third. Remove flowers and buds. |
| 3. Wound (optional) | On firmer stems, remove a thin sliver of bark from the bottom 15–20 mm to expose more rooting area.                                                                                                                                 |
| 4. Hormone          | Decant a small amount into a clean container, dip the base to about 1 cm, and discard what's left. Store hormone in the fridge – it has a limited shelf life.                                                                       |
| 5. Set              | Make a hole with a dibber so the hormone isn't wiped off, insert to the bottom of the hole, and firm the mix. Single tubes make potting on easier than community pots.                                                              |
| 6. Environment      | Water in and place in a humid frame or propagation house in bright shade. Bottom heat so the base is 5–10°C warmer than the air. Keep the mix moist, not wet. Mist sparingly on hairy-leaved species.                               |
| 7. Check            | Most cuttings strike in 4–8 weeks in warm weather. Look for roots at the drainage holes. Remove any rotting cuttings promptly.                                                                                                      |
| 8. Harden off       | About 2 weeks with cover closed by day and open at night, then about 2 weeks in shade, then into the open. Many struck cuttings are lost by skipping this step.                                                                     |
| 9. Pot on           | After 1–3 months, pot on into larger containers. Don't let roots become pot-bound.                                                                                                                                                  |

## Rooting hormone guide

| Product (example) | IBA strength | Use for                                 |
|-------------------|--------------|-----------------------------------------|
| Clonex Green      | 1.5 g/L      | Softwood and soft tip cuttings          |
| Clonex Purple     | 3 g/L        | Semi-hardwood – the general all-rounder |
| Clonex Red        | 8 g/L        | Hardwood cuttings                       |

Equivalent powders or liquids work too – match the strength to the wood type. Callus forming without roots usually means the hormone is too strong for that material.

## Mix, hygiene and collecting

| Topic           | Guidance                                                                                                                                                                                                             |
|-----------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Propagation mix | About 75–80% washed coarse river sand with 20–25% coir or peat. Add extra perlite or coarse sand for rot-prone species.                                                                                              |
| Hygiene         | Fresh mix, clean tubes and benches. Many WA natives (including <i>Hibbertia</i> and <i>Grevillea</i> ) are susceptible to <i>Phytophthora</i> . Discard diseased cuttings; use fungicide only if a problem persists. |
| Collecting      | Collecting from bushland generally requires a DBCA flora licence and landholder permission. Use local-provenance stock and label every batch with its source.                                                        |

# Anthocercis littorea

Yellow Tailflower

Family  
Solanaceae

| DIFFICULTY    | WOOD TYPE     |                                                   |      |      | BEST MONTHS |     |     | CUTTING LENGTH      |     |     | HORMONE (IBA) |      |
|---------------|---------------|---------------------------------------------------|------|------|-------------|-----|-----|---------------------|-----|-----|---------------|------|
| Easy-Moderate | Semi-hardwood |                                                   |      |      | Dec – Apr   |     |     | 75–100 mm           |     |     | 3 g/L IBA     |      |
|               | Jan           | Feb                                               | Mar  | Apr  | May         | Jun | Jul | Aug                 | Sep | Oct | Nov           | Dec  |
| Cuttings      | Best          | Best                                              | Best | Best | OK          | –   | –   | –                   | –   | –   | OK            | Best |
|               | Best window   | Workable – see ‘Selecting material’ in the method |      |      |             |     |     | Avoid / low success |     |     |               |      |

Plant snapshot

|            |                                                                                                                                               |
|------------|-----------------------------------------------------------------------------------------------------------------------------------------------|
| Habit      | Erect, rarely sprawling shrub, 0.6–3 (–4) m.                                                                                                  |
| Flowering  | All year, mainly Jun – Nov (yellow, star-like with long narrow petals).                                                                       |
| Habitat    | Calcareous sand; coastal limestone, sand dunes and sandplains.                                                                                |
| Local note | Recorded in the Wanneroo LGA. <b>Poisonous</b> – all <i>Anthocercis</i> contain tropane alkaloids. Wear gloves and wash hands after handling. |

Propagation options

| Method   | Suitability | Notes                                                     | Details in |
|----------|-------------|-----------------------------------------------------------|------------|
| Cuttings | Main method | Hardened current season's growth strikes reasonably well. | This guide |
| Seed     | Possible    | GA, smoke or karrikinolide raise germination (research).  | –          |
| Division | Not used    | Woody shrub.                                              | –          |

Propagation method

| Item                       | Details                                                                                                                                                                                                                   |
|----------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Best time to take cuttings | Dec – Apr, after the main flowering, once new growth has hardened. Nov and May are workable. Avoid Jun – Oct (main flowering).                                                                                            |
| Wood type                  | Semi-hardwood – hardened current season's growth (ANPSA).                                                                                                                                                                 |
| Cutting size               | 75–100 mm.                                                                                                                                                                                                                |
| Selecting material         | Non-flowering shoots; the plant flowers most of the year, so snip off buds.                                                                                                                                               |
| Pre-treatment              | <ul style="list-style-type: none"> <li>Wear gloves.</li> <li>Strip leaves from the lower half; reduce large leaves by a third.</li> <li>Light wound on firmer stems.</li> <li>Dip the base in rooting hormone.</li> </ul> |
| Rooting hormone            | Semi-hardwood strength, 3 g/L IBA (Clonex Purple).                                                                                                                                                                        |
| Propagation mix            | Standard free-draining cutting mix.                                                                                                                                                                                       |
| Environment                | Humid frame, bright shade, bottom heat.                                                                                                                                                                                   |
| Expected strike time       | Not documented – check from about 4–6 weeks and record the result.                                                                                                                                                        |
| Post-treatment / aftercare | <ul style="list-style-type: none"> <li>Standard hardening off.</li> <li>Grow on in full sun; tip-prune for a bushier plant.</li> </ul>                                                                                    |

## Common problems

| Problem         | Likely cause and fix                                                          |
|-----------------|-------------------------------------------------------------------------------|
| Cuttings flower | Material taken from flowering shoots. Remove buds; pick non-flowering growth. |
| Base rotting    | Mix too wet. Water less; improve drainage.                                    |
| Skin irritation | Handle with gloves – the plant is toxic.                                      |

## Notes

|               |                                                                                                                                                       |
|---------------|-------------------------------------------------------------------------------------------------------------------------------------------------------|
| Difficulty    | <b>Easy-Moderate.</b> Cuttings should strike reasonably well; seed is hard to obtain and slow.                                                        |
| Other methods | Seed – gibberellic acid, smoke water or karrikinolide substantially increased germination in research; untreated seed gave 0% in the only Alcoa test. |
| Confidence    | <b>Medium.</b> ANPSA says cuttings of hardened current season's growth should strike reasonably well; details follow standard practice.               |
| Sources       | See References: 4, 32, 64, 65, 68                                                                                                                     |

## Nursery records

Log each batch here to confirm or update the timing and difficulty above.

| Date set | Stock source / provenance | No. set | No. struck | Strike % | Notes |
|----------|---------------------------|---------|------------|----------|-------|
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |

# Atriplex isatidea

Coast Saltbush

Family  
Amaranthaceae  
(chenopods)

|                 |                    |             |                |               |
|-----------------|--------------------|-------------|----------------|---------------|
| DIFFICULTY      | WOOD TYPE          | BEST MONTHS | CUTTING LENGTH | HORMONE (IBA) |
| Easy-Moderate * | Semi-hardwood tips | Sep – Nov   | 100–150 mm     | 1.5–3 g/L IBA |

|          | Jan         | Feb                                               | Mar | Apr | May | Jun | Jul | Aug                 | Sep  | Oct  | Nov  | Dec |
|----------|-------------|---------------------------------------------------|-----|-----|-----|-----|-----|---------------------|------|------|------|-----|
| Cuttings | OK          | OK                                                | –   | –   | –   | –   | –   | OK                  | Best | Best | Best | OK  |
|          | Best window | Workable – see ‘Selecting material’ in the method |     |     |     |     |     | Avoid / low success |      |      |      |     |

Plant snapshot

|            |                                                                                        |
|------------|----------------------------------------------------------------------------------------|
| Habit      | Grey-leaved, spreading shrub, 0.5–2.5 m. Male and female flowers on the same plant.    |
| Flowering  | Mainly Mar – Jun (small, inconspicuous).                                               |
| Habitat    | White sand on coastal dunes.                                                           |
| Local note | Recorded in the Wanneroo LGA. Very salt- and wind-tolerant – a useful dune stabiliser. |

Propagation options

| Method   | Suitability | Notes                                                                    | Details in                 |
|----------|-------------|--------------------------------------------------------------------------|----------------------------|
| Cuttings | Good        | Saltbush cuttings are generally reliable (provisional for this species). | <a href="#">This guide</a> |
| Seed     | Main method | Soak 4 days in changed water, then smoke water.                          | → Seed Sowing Guide        |
| Division | Not used    | Woody shrub.                                                             | –                          |

Propagation method

| Item                       | Details                                                                                                                                                                                               |
|----------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Best time to take cuttings | Sep – Nov, in active spring growth (the time recommended for the related Old Man Saltbush). Aug and Dec – Feb are workable. Avoid Mar – Jul (flowering, then cool, slow growth).                      |
| Wood type                  | Semi-hardwood terminal (tip) cuttings of current season's growth.                                                                                                                                     |
| Cutting size               | 100–150 mm. Terminal cuttings of about 15 cm are used for <i>A. nummularia</i> .                                                                                                                      |
| Selecting material         | Vigorous, non-flowering tips from healthy plants. Avoid old, woody stems.                                                                                                                             |
| Pre-treatment              | <ul style="list-style-type: none"><li>Strip leaves from the lower half.</li><li>No wounding needed.</li><li>Dip the base in rooting hormone.</li></ul>                                                |
| Rooting hormone            | Softwood to semi-hardwood strength (1.5–3 g/L IBA, <b>Clonex Green</b> or <b>Purple</b> ). Related saltbushes root without hormone, so it mainly evens out the batch.                                 |
| Propagation mix            | Free-draining mix of sand and potting mix.                                                                                                                                                            |
| Environment                | <b>Good ventilation and no mist</b> – misting encourages a leaf-spotting fungus on saltbush cuttings (ANBG, <i>A. nummularia</i> ). Bottom heat helps; keep the mix just moist.                       |
| Expected strike time       | Not documented for this species. In one trial, 20 cm unrooted <i>A. halimus</i> cuttings with no hormone gave about <b>49% survival at 8 weeks</b> . Check from about 4 weeks and record your result. |
| Post-treatment / aftercare | <ul style="list-style-type: none"><li>Harden off, then grow on in full sun.</li><li>Water sparingly once established.</li><li>Tip-prune for a bushier tube.</li></ul>                                 |

## Common problems

| Problem                     | Likely cause and fix                                                |
|-----------------------------|---------------------------------------------------------------------|
| Leaf spots, leaves dropping | Too humid or misted. Stop misting, open the frame, improve airflow. |
| Base rotting                | Mix too wet. Use a sandier mix and water less.                      |
| Leggy, pale growth          | Not enough light. Grow on in full sun.                              |

## Notes

|            |                                                                                                                                                                              |
|------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Difficulty | <b>Easy-Moderate (provisional).</b> Saltbushes generally strike readily from tip cuttings; the main risk is leaf disease under mist. Seed is the more common nursery method. |
| Confidence | <b>Medium.</b> No species-specific cutting data. Method follows published results for related saltbushes ( <i>A. nummularia</i> , <i>A. halimus</i> ).                       |
| Sources    | See References: 4, 36, 37, 38                                                                                                                                                |

## Nursery records

Log each batch here to confirm or update the timing and difficulty above.

| Date set | Stock source / provenance | No. set | No. struck | Strike % | Notes |
|----------|---------------------------|---------|------------|----------|-------|
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |

# Calothamnus quadrifidus

One-sided Bottlebrush

Family  
Myrtaceae

| DIFFICULTY      | WOOD TYPE     | BEST MONTHS | CUTTING LENGTH | HORMONE (IBA) |
|-----------------|---------------|-------------|----------------|---------------|
| Easy-Moderate * | Semi-hardwood | Jan – Apr   | 75–100 mm      | 3 g/L IBA     |

|          | Jan  | Feb  | Mar  | Apr  | May | Jun | Jul | Aug | Sep | Oct | Nov | Dec |
|----------|------|------|------|------|-----|-----|-----|-----|-----|-----|-----|-----|
| Cuttings | Best | Best | Best | Best | OK  | –   | –   | –   | –   | –   | –   | OK  |

Best window
Workable – see 'Selecting material' in the method
Avoid / low success

## Plant snapshot

|            |                                                                                                    |
|------------|----------------------------------------------------------------------------------------------------|
| Habit      | Erect, compact or spreading shrub, usually 0.9–2 m.                                                |
| Flowering  | Jun – Dec (red, one-sided brushes).                                                                |
| Habitat    | Wide variety of soils and habitats.                                                                |
| Local note | Recorded in the Wanneroo LGA. Several subspecies exist – label stock by subspecies and provenance. |

## Propagation options

| Method   | Suitability | Notes                                                     | Details in          |
|----------|-------------|-----------------------------------------------------------|---------------------|
| Cuttings | Good        | Presumed to strike like related melaleucas (provisional). | This guide          |
| Seed     | Main method | Easy from seed held in old capsules.                      | → Seed Sowing Guide |
| Division | Not used    | Woody shrub.                                              | –                   |

## Propagation method

| Item                       | Details                                                                                                                                                                                           |
|----------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Best time to take cuttings | Jan – Apr, after flowering, once the new growth has firmed. Dec and May are workable. Avoid Jun – Nov (flowering).                                                                                |
| Wood type                  | Semi-hardwood – firm current season's growth.                                                                                                                                                     |
| Cutting size               | 75–100 mm, as for <i>Melaleuca</i> .                                                                                                                                                              |
| Selecting material         | Non-flowering side shoots. Keep subspecies and forms separate.                                                                                                                                    |
| Pre-treatment              | <ul style="list-style-type: none"> <li>Strip the needle-like leaves from the lower half to two-thirds.</li> <li>Light wound on firmer stems.</li> <li>Dip the base in rooting hormone.</li> </ul> |
| Rooting hormone            | Semi-hardwood strength, 3 g/L IBA ( <b>Clonex Purple</b> ).                                                                                                                                       |
| Propagation mix            | Standard free-draining cutting mix.                                                                                                                                                               |
| Environment                | Humid frame, bright shade, bottom heat.                                                                                                                                                           |
| Expected strike time       | Not documented – check from about 6 weeks and record the result.                                                                                                                                  |
| Post-treatment / aftercare | <ul style="list-style-type: none"> <li>Standard hardening off.</li> <li>Pot on into a free-draining native mix.</li> </ul>                                                                        |

## Common problems

| Problem            | Likely cause and fix                                                        |
|--------------------|-----------------------------------------------------------------------------|
| Cuttings brown off | Too dry or too soft. Use firmer material and keep humidity up until rooted. |
| Callus, no roots   | Trim and re-set; try a lower hormone strength.                              |
| Slow in winter     | Use bottom heat, or wait for the Jan – Apr window.                          |

## Notes

|            |                                                                                                                                         |
|------------|-----------------------------------------------------------------------------------------------------------------------------------------|
| Difficulty | <b>Easy-Moderate (provisional).</b> Expected to behave like related melaleucas, but untested here. Seed is easy and is the main method. |
| Confidence | <b>Low.</b> ANPSA notes the genus is 'presumably' grown from cuttings. Method follows related melaleucas – trial protocol.              |
| Sources    | See References: 1, 4, 13, 41                                                                                                            |

## Nursery records

Log each batch here to confirm or update the timing and difficulty above.

| Date set | Stock source / provenance | No. set | No. struck | Strike % | Notes |
|----------|---------------------------|---------|------------|----------|-------|
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |

# Carpobrotus virescens

Coastal Pigface

Family  
Aizoaceae

| DIFFICULTY  | WOOD TYPE                                         |     |      |      | BEST MONTHS          |     | CUTTING LENGTH   |     | HORMONE (IBA) |      |      |     |
|-------------|---------------------------------------------------|-----|------|------|----------------------|-----|------------------|-----|---------------|------|------|-----|
| Easy        | Succulent stem pieces                             |     |      |      | Sep - Nov; Mar - Apr |     | 10-15 cm (tubes) |     | Not needed    |      |      |     |
|             | Jan                                               | Feb | Mar  | Apr  | May                  | Jun | Jul              | Aug | Sep           | Oct  | Nov  | Dec |
| Cuttings    | OK                                                | OK  | Best | Best | OK                   | -   | -                | OK  | Best          | Best | Best | OK  |
| Best window | Workable - see 'Selecting material' in the method |     |      |      | Avoid / low success  |     |                  |     |               |      |      |     |

## Plant snapshot

|            |                                                                                                                                                                   |
|------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Habit      | Prostrate succulent groundcover, 0.1–0.3 m high and 0.5–3 m wide, rooting along the stems.                                                                        |
| Flowering  | Jun – Dec or Jan (purple-pink, sometimes white), followed by fleshy fruit.                                                                                        |
| Habitat    | Coastal limestone and sand dunes.                                                                                                                                 |
| Local note | Recorded in the Wanneroo LGA. Collect only from plants you are sure are native – the introduced Hottentot Fig ( <i>C. edulis</i> ) is naturalised on some coasts. |

## Propagation options

| Method   | Suitability | Notes                                                       | Details in                          |
|----------|-------------|-------------------------------------------------------------|-------------------------------------|
| Cuttings | Main method | Stem pieces root very easily; rooted runners can be lifted. | <a href="#">This guide</a>          |
| Seed     | Good        | Ferment fruit, sow fine seed on the surface; 6–8 days.      | → <a href="#">Seed Sowing Guide</a> |
| Division | Possible    | Lift stems that have rooted – treat as cuttings.            | –                                   |

## Propagation method

| Item                       | Details                                                                                                                                                                                                                        |
|----------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Best time to take cuttings | Sep – Nov and Mar – Apr, when plants are actively growing. Aug, Dec – Feb and May are workable. Avoid Jun – Jul (slow growth; cuttings rot in cold, wet mix).                                                                  |
| Wood type                  | Succulent stem (runner) pieces – not woody cuttings. Use firm stems with plump leaves.                                                                                                                                         |
| Cutting size               | Tubes: 10–15 cm pieces with 2–3 leaf pairs.<br>Direct planting / layering: about 30 cm, with at least 5 cm left above the sand (ANBG, <i>C. glaucescens</i> ).                                                                 |
| Selecting material         | Healthy runners. Pieces that have already rooted at the nodes are ideal. Avoid shrivelled or damaged stems.                                                                                                                    |
| Pre-treatment              | <ul style="list-style-type: none"> <li>Remove the leaves from the lowest 1–2 nodes.</li> <li>Leave the cut ends to dry in the shade for a day or two before setting – this reduces rot.</li> <li>No hormone needed.</li> </ul> |
| Rooting hormone            | Not needed – stems root readily at the nodes.                                                                                                                                                                                  |
| Propagation mix            | Coarse washed sand or a very sandy, free-draining mix.                                                                                                                                                                         |
| Environment                | Open bench in bright light or light shade. <b>No mist and no humid frame</b> – succulent leaves rot. Water sparingly until rooted.                                                                                             |
| Expected strike time       | Not documented, but usually quick – check from about 3 weeks.                                                                                                                                                                  |

| Item                       | Details                                                                                                                                                                                                          |
|----------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Post-treatment / aftercare | <ul style="list-style-type: none"> <li>• Grow on in full sun.</li> <li>• Pot on or plant out once roots hold the mix together.</li> <li>• Water sparingly – overwatering is the main cause of losses.</li> </ul> |

### Common problems

| Problem                | Likely cause and fix                                                               |
|------------------------|------------------------------------------------------------------------------------|
| Stems go soft and rot  | Too wet or humid. Dry the cut ends first, water less, move out of the humid frame. |
| Leaves shrivel         | Too dry for too long. Water lightly once set.                                      |
| Pale, stretched growth | Not enough light. Move to full sun.                                                |

### Notes

|               |                                                                                                                                       |
|---------------|---------------------------------------------------------------------------------------------------------------------------------------|
| Difficulty    | <b>Easy.</b> Pigface roots readily from stem pieces, and rooted runners can simply be cut off and potted up.                          |
| Other methods | Layering – pin a runner onto a pot of sand beside the parent and cut it free once rooted.                                             |
| Confidence    | <b>Medium.</b> Method from published advice for the closely related <i>C. glaucescens</i> ; pigface is widely grown from stem pieces. |
| Sources       | See References: 4, 12, 33                                                                                                             |

### Nursery records

Log each batch here to confirm or update the timing and difficulty above.

| Date set | Stock source / provenance | No. set | No. struck | Strike % | Notes |
|----------|---------------------------|---------|------------|----------|-------|
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |

# Eremophila glabra

Tar Bush / Common Emu Bush

Family  
Scrophulariaceae

| DIFFICULTY | WOOD TYPE     | BEST MONTHS | CUTTING LENGTH | HORMONE (IBA) |
|------------|---------------|-------------|----------------|---------------|
| Easy       | Semi-hardwood | Jan – Apr   | 75–100 mm      | 3 g/L IBA     |

|          | Jan  | Feb  | Mar  | Apr  | May | Jun | Jul | Aug | Sep | Oct | Nov | Dec |
|----------|------|------|------|------|-----|-----|-----|-----|-----|-----|-----|-----|
| Cuttings | Best | Best | Best | Best | OK  | –   | –   | –   | –   | –   | OK  | OK  |

Best window
Workable – see 'Selecting material' in the method
Avoid / low success

## Plant snapshot

|            |                                                                                                                                                                           |
|------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Habit      | Very variable, from prostrate to 3 m. Perth coastal plants are mostly subsp. <i>albicans</i> : low, spreading, to about 0.4 m high and 2 m wide, with grey, felty leaves. |
| Flowering  | Species: Mar – Dec. Subsp. <i>albicans</i> : May – Jan. Tubular flowers, yellow, orange, red or green.                                                                    |
| Habitat    | Sand and limestone; dunes, sandplains and winter-wet flats, sometimes saline.                                                                                             |
| Local note | Recorded in the Wanneroo LGA and along the Perth coast. Several subspecies and many forms – label stock by subspecies and form.                                           |

## Propagation options

| Method   | Suitability | Notes                                                  | Details in                 |
|----------|-------------|--------------------------------------------------------|----------------------------|
| Cuttings | Main method | Most forms strike readily from semi-hardwood cuttings. | <a href="#">This guide</a> |
| Seed     | Difficult   | Unreliable – hard fruits; slow, erratic germination.   | –                          |
| Division | Not used    | Woody shrub.                                           | –                          |

## Propagation method

| Item                       | Details                                                                                                                                                                                                            |
|----------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Best time to take cuttings | Jan – Apr, using hardened current season's growth while the weather is warm and plants are mostly out of flower. Nov – Dec and May are workable. Avoid Jun – Oct (cool weather, flowering).                        |
| Wood type                  | Semi-hardwood – hardened current season's growth.                                                                                                                                                                  |
| Cutting size               | 75–100 mm.                                                                                                                                                                                                         |
| Selecting material         | Non-flowering tips or side shoots. On grey-leaved forms pick clean foliage with no sign of mould.                                                                                                                  |
| Pre-treatment              | <ul style="list-style-type: none"> <li>Carefully remove leaves from the lower two-thirds.</li> <li>Wound: remove a thin sliver of bark from the base.</li> <li>Dip the base in rooting hormone.</li> </ul>         |
| Rooting hormone            | Semi-hardwood strength, e.g. <b>Clonex Purple</b> (3 g/L IBA).                                                                                                                                                     |
| Propagation mix            | Very free-draining, sand-based mix.                                                                                                                                                                                |
| Environment                | <b>Grey, felty-leaved forms rot under heavy mist or high humidity</b> – keep foliage as dry as possible, water the mix, give good airflow. Green-leaved forms tolerate normal misting. Bottom heat speeds rooting. |
| Expected strike time       | About <b>4–8 weeks</b> in warm weather.                                                                                                                                                                            |
| Post-treatment / aftercare | <ul style="list-style-type: none"> <li>Standard hardening off.</li> <li>Pot on into a free-draining mix and grow on in full sun.</li> <li>Keep water off grey foliage; water in the morning.</li> </ul>            |

## Common problems

| Problem                            | Likely cause and fix                                                 |
|------------------------------------|----------------------------------------------------------------------|
| Leaves blacken or grow grey mould  | Too humid. Cut back misting, open the frame, remove affected leaves. |
| Base rotting                       | Mix too wet. Water less often and use a sandier mix.                 |
| Cuttings flower instead of rooting | Use non-flowering shoots and pinch off buds.                         |

## Notes

|               |                                                                                                                                                                   |
|---------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Difficulty    | Easy. Most forms of <i>E. glabra</i> strike readily from cuttings.                                                                                                |
| Other methods | Seed – unreliable in <i>Eremophila</i> (hard fruits, slow and erratic germination). Grafting is used for difficult eremophilas but isn't needed for this species. |
| Confidence    | High. Published sources agree most forms strike readily from cuttings.                                                                                            |
| Sources       | See References: 1, 4, 12, 13, 20, 21, 22, 30                                                                                                                      |

## Nursery records

Log each batch here to confirm or update the timing and difficulty above.

| Date set | Stock source / provenance | No. set | No. struck | Strike % | Notes |
|----------|---------------------------|---------|------------|----------|-------|
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |

# Grevillea crithmifolia

No widely used common name

Family  
Proteaceae

|                          |                            |                          |                             |                            |
|--------------------------|----------------------------|--------------------------|-----------------------------|----------------------------|
| DIFFICULTY<br>Moderate * | WOOD TYPE<br>Semi-hardwood | BEST MONTHS<br>Dec – Apr | CUTTING LENGTH<br>75–100 mm | HORMONE (IBA)<br>3 g/L IBA |
|--------------------------|----------------------------|--------------------------|-----------------------------|----------------------------|

|          |             |      |                                                   |      |     |     |     |     |     |     |                     |      |
|----------|-------------|------|---------------------------------------------------|------|-----|-----|-----|-----|-----|-----|---------------------|------|
|          | Jan         | Feb  | Mar                                               | Apr  | May | Jun | Jul | Aug | Sep | Oct | Nov                 | Dec  |
| Cuttings | Best        | Best | Best                                              | Best | OK  | –   | –   | –   | –   | –   | OK                  | Best |
|          | Best window |      | Workable – see ‘Selecting material’ in the method |      |     |     |     |     |     |     | Avoid / low success |      |

Plant snapshot

|            |                                                                                                        |
|------------|--------------------------------------------------------------------------------------------------------|
| Habit      | Dense, much-branched shrub, (0.3–) 0.6–2 m. Compact shrub and low groundcover forms exist.             |
| Flowering  | Jun or Aug – Nov (white to white-pink). Some people dislike the scent.                                 |
| Habitat    | White or yellow sand over limestone; dunes, hillslopes and sandplains.                                 |
| Local note | Recorded in the City of Joondalup. Dieback susceptible. Keep the shrub and groundcover forms separate. |

Propagation options

| Method   | Suitability | Notes                                            | Details in |
|----------|-------------|--------------------------------------------------|------------|
| Cuttings | Main method | The usual method, but may be slow to root.       | This guide |
| Seed     | Difficult   | About 9% in one nursery test; smoke recommended. | –          |
| Division | Not used    | –                                                | –          |

Propagation method

| Item                       | Details                                                                                                                                                                                                                             |
|----------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Best time to take cuttings | Dec – Apr, after flowering, once new growth has firmed. Nov and May are workable. Avoid Jun – Oct (flowering).                                                                                                                      |
| Wood type                  | Semi-hardwood – firm current season's growth.                                                                                                                                                                                       |
| Cutting size               | 75–100 mm (ANPSA Grevillea guidance).                                                                                                                                                                                               |
| Selecting material         | Non-flowering shoots from a plant of the form you want (shrub or groundcover).                                                                                                                                                      |
| Pre-treatment              | <ul style="list-style-type: none"><li>Strip leaves from the lower half to two-thirds (cut, don't tear).</li><li>Wound the base – a thin sliver of bark from the bottom 15–20 mm.</li><li>Dip the base in rooting hormone.</li></ul> |
| Rooting hormone            | Semi-hardwood strength, 3 g/L IBA (Clonex Purple). Wounding and hormone improve strike in grevilleas.                                                                                                                               |
| Propagation mix            | Free-draining, clean cutting mix. Strict hygiene – the species is dieback susceptible.                                                                                                                                              |
| Environment                | Humid frame, bright shade, bottom heat. Be patient – roots can be slow.                                                                                                                                                             |
| Expected strike time       | Not documented, but reported slow – allow at least 8–12 weeks before discarding. Record your results.                                                                                                                               |
| Post-treatment / aftercare | <ul style="list-style-type: none"><li>Standard hardening off.</li><li>Pot on into a low-phosphorus native mix; feed only with low-P fertiliser.</li></ul>                                                                           |

## Common problems

| Problem                            | Likely cause and fix                                                        |
|------------------------------------|-----------------------------------------------------------------------------|
| No roots after 8 weeks             | Normal for this species – keep them going. Next time wound and use hormone. |
| Heavy callus, no roots             | Trim the callus and re-set; try a lower hormone strength.                   |
| Plants yellow or die after potting | Phosphorus or dieback. Low-P mix, clean pots off the ground.                |

## Notes

|               |                                                                                                                                       |
|---------------|---------------------------------------------------------------------------------------------------------------------------------------|
| Difficulty    | <b>Moderate (provisional).</b> Cuttings are the usual method but may be slow to produce roots.                                        |
| Other methods | Seed – poor: about 9% in a single Alcoa nursery test; a seed supplier recommends smoke treatment.                                     |
| Confidence    | <b>Medium.</b> Grown commercially from cuttings but reported slow to root; size and treatment follow ANPSA <i>Grevillea</i> guidance. |
| Sources       | See References: 4, 32, 56, 57, 59, 61                                                                                                 |

## Nursery records

Log each batch here to confirm or update the timing and difficulty above.

| Date set | Stock source / provenance | No. set | No. struck | Strike % | Notes |
|----------|---------------------------|---------|------------|----------|-------|
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |

# Grevillea preissii

Spider-net Grevillea

Family  
ProteaceaeDIFFICULTY  
**Moderate**WOOD TYPE  
**Semi-hardwood**BEST MONTHS  
**Nov – Mar**CUTTING LENGTH  
**75–100 mm**HORMONE (IBA)  
**3 g/L IBA**

|          | Jan  | Feb  | Mar  | Apr | May | Jun | Jul | Aug | Sep | Oct | Nov  | Dec  |
|----------|------|------|------|-----|-----|-----|-----|-----|-----|-----|------|------|
| Cuttings | Best | Best | Best | OK  | –   | –   | –   | –   | –   | OK  | Best | Best |

Best window

Workable – see 'Selecting material' in the method

Avoid / low success

## Plant snapshot

|            |                                                                                                                                                                                         |
|------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Habit      | Shrub, 0.15–1.7 m.                                                                                                                                                                      |
| Flowering  | Jun – Sep (red spider flowers; attract honeyeaters).                                                                                                                                    |
| Habitat    | White, grey or yellow sand, sandy clay and limestone; ridges, outcrops and seasonally wet areas.                                                                                        |
| Local note | Recorded in Joondalup and Wanneroo. The former subsp. <i>glabrilimba</i> (including 'Magic Lantern') is now a separate species, <i>G. glabrilimba</i> – keep it apart from local stock. |

## Propagation options

| Method   | Suitability | Notes                                       | Details in |
|----------|-------------|---------------------------------------------|------------|
| Cuttings | Main method | Firm growth; slow, hormone improves strike. | This guide |
| Seed     | Possible    | Nick the seed coat; smoke water.            | –          |
| Division | Not used    | –                                           | –          |

## Propagation method

| Item                       | Details                                                                                                                                                                                                                        |
|----------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Best time to take cuttings | <b>Nov – Mar</b> , after flowering, once the current season's growth has firmed. Oct and Apr are workable. Avoid May – Sep (flowering).                                                                                        |
| Wood type                  | <b>Semi-hardwood</b> – firm current season's growth (ANPSA).                                                                                                                                                                   |
| Cutting size               | 75–100 mm (ANPSA <i>Grevillea</i> guidance).                                                                                                                                                                                   |
| Selecting material         | Non-flowering shoots from healthy, local-provenance plants.                                                                                                                                                                    |
| Pre-treatment              | <ul style="list-style-type: none"> <li>Strip leaves from the lower half to two-thirds.</li> <li>Wound the base on firmer stems.</li> <li>Dip the base in rooting hormone – ANPSA notes it clearly improves success.</li> </ul> |
| Rooting hormone            | Semi-hardwood strength, 3 g/L IBA ( <b>Clonex Purple</b> ).                                                                                                                                                                    |
| Propagation mix            | Free-draining, clean cutting mix.                                                                                                                                                                                              |
| Environment                | Humid frame, bright shade, bottom heat.                                                                                                                                                                                        |
| Expected strike time       | Not documented, but reported slow – allow at least <b>8–12 weeks</b> before discarding. Record your results.                                                                                                                   |
| Post-treatment / aftercare | <ul style="list-style-type: none"> <li>Standard hardening off.</li> <li>Pot on into a <b>low-phosphorus</b> native mix.</li> </ul>                                                                                             |

## Common problems

| Problem                    | Likely cause and fix                                 |
|----------------------------|------------------------------------------------------|
| Slow to strike             | Normal. Use hormone and bottom heat; be patient.     |
| Heavy callus, no roots     | Trim and re-set; try a lower hormone strength.       |
| Yellowing after potting on | Phosphorus in mix or fertiliser. Use low-P products. |

## Notes

|               |                                                                                                         |
|---------------|---------------------------------------------------------------------------------------------------------|
| Difficulty    | <b>Moderate.</b> Cuttings of firm growth can be slow to strike; hormone improves success.               |
| Other methods | Seed – nick the seed coat before sowing (ANPSA); a seed supplier recommends smoke treatment.            |
| Confidence    | <b>High.</b> ANPSA gives species-specific cutting advice; size follows ANPSA <i>Grevillea</i> guidance. |
| Sources       | See References: 4, 58, 59, 60, 61                                                                       |

## Nursery records

Log each batch here to confirm or update the timing and difficulty above.

| Date set | Stock source / provenance | No. set | No. struck | Strike % | Notes |
|----------|---------------------------|---------|------------|----------|-------|
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |

# Grevillea vestita

No widely accepted common name

Family  
Proteaceae

| DIFFICULTY | WOOD TYPE     | BEST MONTHS | CUTTING LENGTH | HORMONE (IBA) |
|------------|---------------|-------------|----------------|---------------|
| Easy       | Semi-hardwood | Dec – Apr   | 80–120 mm      | 3 g/L IBA     |

|          | Jan  | Feb  | Mar  | Apr  | May | Jun | Jul | Aug | Sep | Oct | Nov | Dec  |
|----------|------|------|------|------|-----|-----|-----|-----|-----|-----|-----|------|
| Cuttings | Best | Best | Best | Best | OK  | –   | –   | –   | –   | –   | OK  | Best |

Best window
Workable – see 'Selecting material' in the method
Avoid / low success

## Plant snapshot

|            |                                                                                                                               |
|------------|-------------------------------------------------------------------------------------------------------------------------------|
| Habit      | Erect, spreading, prickly shrub, usually 0.5–3 m. Hairy branchlets; lobed leaves with sharp-pointed tips.                     |
| Flowering  | Mainly Jun – Nov (white to cream, often pink-tinged, scented). Odd flowers Jan – Apr.                                         |
| Habitat    | Sandy soils over limestone or granite. Occurs on the Swan Coastal Plain, including the Perth area.                            |
| Local note | Two subspecies exist (subsp. <i>vestita</i> and subsp. <i>isopogoides</i> ). Label stock plants by subspecies and provenance. |

## Propagation options

| Method   | Suitability | Notes                                                              | Details in                          |
|----------|-------------|--------------------------------------------------------------------|-------------------------------------|
| Cuttings | Main method | Firm current season's growth strikes reliably.                     | <a href="#">This guide</a>          |
| Seed     | Good        | Grows reasonably easily from seed; seedlings vary from the parent. | → <a href="#">Seed Sowing Guide</a> |
| Division | Possible    | Can sucker – lift rooted suckers.                                  | –                                   |

## Propagation method

| Item                       | Details                                                                                                                                                                                                                                                                                                                                                                                                                   |
|----------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Best time to take cuttings | <b>Dec – Apr</b> (late spring to autumn), once the spring flush has firmed and flowering has finished. Nov and May are workable. Avoid Jun – Oct: plants are flowering, new growth is soft and cool weather slows rooting.                                                                                                                                                                                                |
| Wood type                  | <b>Semi-hardwood</b> – firm, current season's growth. It should bend about 60° and spring back.                                                                                                                                                                                                                                                                                                                           |
| Cutting size               | 80–120 mm long with 3–5 nodes. Tip or sub-tip pieces; two cuttings can often be taken from one shoot.                                                                                                                                                                                                                                                                                                                     |
| Selecting material         | Healthy, vigorous, <b>non-flowering</b> shoots. Skip shoots carrying buds, old woody stems and soft new tips.                                                                                                                                                                                                                                                                                                             |
| Pre-treatment              | <ul style="list-style-type: none"> <li>Wear gloves: leaves are prickly and grevillea sap can irritate skin.</li> <li>Remove any flowers or buds.</li> <li>Strip leaves from the lower half to two-thirds (cut, don't tear the bark).</li> <li>Trim remaining lobed leaves by about one-third.</li> <li>Wound: remove a thin sliver of bark from the bottom 15–20 mm.</li> <li>Dip the base in rooting hormone.</li> </ul> |
| Rooting hormone            | Semi-hardwood strength, e.g. <b>Clonex Purple (3 g/L IBA)</b> . If cuttings form a heavy callus but no roots, trim the callus back to clean wood and re-set; try a lower strength on the next batch.                                                                                                                                                                                                                      |
| Propagation mix            | Free-draining, low nutrient: about 75–80% washed coarse river sand + 20–25% coir or peat (or a perlite/coir mix).                                                                                                                                                                                                                                                                                                         |
| Environment                | Humid propagation house or frame in bright shade. Bottom heat with the base of the cutting 5–10°C warmer than the air. Light misting only – the hairy foliage can rot if kept constantly wet.                                                                                                                                                                                                                             |

| Item                       | Details                                                                                                                                                                                                                                                                                                                                                                                                                                             |
|----------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Expected strike time       | About <b>6–8 weeks</b> in warm weather. Slower in the cooler months.                                                                                                                                                                                                                                                                                                                                                                                |
| Post-treatment / aftercare | <ul style="list-style-type: none"> <li>• Harden off: ~2 weeks with cover closed by day and open at night, then ~2 weeks in shade before full sun.</li> <li>• Pot on when roots show at the drainage holes, into a <b>low-phosphorus</b> native mix.</li> <li>• Feed only with a low-phosphorus native fertiliser (Proteaceae are phosphorus-sensitive).</li> <li>• Responds well to pruning – tip-prune young plants for a bushier tube.</li> </ul> |

### Common problems

| Problem                                   | Likely cause and fix                                                    |
|-------------------------------------------|-------------------------------------------------------------------------|
| Heavy callus, no roots                    | Trim callus back and re-set. Lower the hormone strength next time.      |
| Base rotting                              | Mix too wet or material too soft. Use firmer wood and water less often. |
| Leaf yellowing or scorch after potting on | Check phosphorus in the potting mix and fertiliser.                     |

### Notes

|               |                                                                                                                                                                               |
|---------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Difficulty    | <b>Easy.</b> Cuttings from firm current season's growth usually strike reliably.                                                                                              |
| Other methods | Seed – grows reasonably easily from seed, but seedlings are not identical to the parent. Grafting onto <i>G. robusta</i> is used on the humid east coast; not needed locally. |
| Confidence    | <b>High.</b> Species-specific propagation information is published and consistent across sources.                                                                             |
| Sources       | See References: 1, 2, 3, 4, 13, 15, 17                                                                                                                                        |

### Nursery records

Log each batch here to confirm or update the timing and difficulty above.

| Date set | Stock source / provenance | No. set | No. struck | Strike % | Notes |
|----------|---------------------------|---------|------------|----------|-------|
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |

# Guichenotia ledifolia

No widely used common name

Family  
Malvaceae

|                 |               |             |                |               |
|-----------------|---------------|-------------|----------------|---------------|
| DIFFICULTY      | WOOD TYPE     | BEST MONTHS | CUTTING LENGTH | HORMONE (IBA) |
| Easy-Moderate * | Semi-hardwood | Dec – Apr   | 50–75 mm       | 3 g/L IBA     |

|          | Jan         | Feb                                               | Mar  | Apr  | May | Jun | Jul | Aug | Sep | Oct | Nov                 | Dec  |
|----------|-------------|---------------------------------------------------|------|------|-----|-----|-----|-----|-----|-----|---------------------|------|
| Cuttings | Best        | Best                                              | Best | Best | OK  | –   | –   | –   | –   | –   | OK                  | Best |
|          | Best window | Workable – see ‘Selecting material’ in the method |      |      |     |     |     |     |     |     | Avoid / low success |      |

Plant snapshot

|            |                                                                                       |
|------------|---------------------------------------------------------------------------------------|
| Habit      | Spreading shrub, 0.4–1.5 (–2) m.                                                      |
| Flowering  | Jul – Oct (blue-purple or pink).                                                      |
| Habitat    | Sandy or rocky clay; coastal limestone, sandplains and granite rocks.                 |
| Local note | Recorded in Joondalup and Wanneroo. Tolerates drought and alkaline (limestone) soils. |

Propagation options

| Method   | Suitability | Notes                                         | Details in          |
|----------|-------------|-----------------------------------------------|---------------------|
| Cuttings | Main method | Often grown from cuttings; semi-hardwood.     | This guide          |
| Seed     | Good        | Hard coat: boiling water over seed; variable. | → Seed Sowing Guide |
| Division | Not used    | Shrub – not divided.                          | –                   |

Propagation method

| Item                       | Details                                                                                                                                                                    |
|----------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Best time to take cuttings | Dec – Apr, after flowering, once new growth has firmed. Nov and May are workable. Avoid Jun – Oct (cool, and plants are flowering).                                        |
| Wood type                  | Semi-hardwood – firm current season's growth.                                                                                                                              |
| Cutting size               | 50–75 mm.                                                                                                                                                                  |
| Selecting material         | Healthy, non-flowering side shoots.                                                                                                                                        |
| Pre-treatment              | <ul style="list-style-type: none"><li>Strip leaves from the lower half; the leaves are small, so no trimming is needed.</li><li>Dip the base in rooting hormone.</li></ul> |
| Rooting hormone            | 3 g/L IBA (Clonex Purple).                                                                                                                                                 |
| Propagation mix            | Very free-draining cutting mix.                                                                                                                                            |
| Environment                | Humid frame, bright shade, bottom heat. Moist, not wet; good airflow.                                                                                                      |
| Expected strike time       | Not documented – check from about 4–6 weeks and record the result.                                                                                                         |
| Post-treatment / aftercare | <ul style="list-style-type: none"><li>Standard hardening off.</li><li>Grow on in full sun in a free-draining mix.</li><li>Tip-prune for a denser plant.</li></ul>          |

## Common problems

| Problem          | Likely cause and fix                                        |
|------------------|-------------------------------------------------------------|
| Leaves go mouldy | Too humid or crowded. Ventilate the frame; don't over-mist. |
| Base rotting     | Too wet. Coarser mix; water less.                           |
| Slow to root     | Use bottom heat and take cuttings in the warm months.       |

## Notes

|               |                                                                                                                                                                           |
|---------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Difficulty    | <b>Easy-Moderate (provisional).</b> Firm growth of the related <i>G. macrantha</i> usually strikes reliably; untested here.                                               |
| Other methods | Seed – hard-coated. Pour boiling water over sown seed or soak it 24 h in water just off the boil; germination is variable and takes 30+ days (see the Seed Sowing Guide). |
| Confidence    | <b>Medium.</b> WA seed notes say the genus is often grown from cuttings; semi-hardwood cuttings are listed for this species. No published strike rates.                   |
| Sources       | See References: 4, 12, 13, 74, 75, 76, 77                                                                                                                                 |

## Nursery records

Log each batch here to confirm or update the timing and difficulty above.

| Date set | Stock source / provenance | No. set | No. struck | Strike % | Notes |
|----------|---------------------------|---------|------------|----------|-------|
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |

# Hardenbergia comptoniana

Native Wisteria

Family  
Fabaceae

|               |               |             |                |               |
|---------------|---------------|-------------|----------------|---------------|
| DIFFICULTY    | WOOD TYPE     | BEST MONTHS | CUTTING LENGTH | HORMONE (IBA) |
| Easy-Moderate | Semi-hardwood | Dec - Mar   | 50-75 mm       | 3 g/L IBA     |

|          | Jan         | Feb                                               | Mar  | Apr | May | Jun | Jul | Aug | Sep | Oct | Nov                 | Dec  |
|----------|-------------|---------------------------------------------------|------|-----|-----|-----|-----|-----|-----|-----|---------------------|------|
| Cuttings | Best        | Best                                              | Best | OK  | -   | -   | -   | -   | -   | -   | OK                  | Best |
|          | Best window | Workable - see 'Selecting material' in the method |      |     |     |     |     |     |     |     | Avoid / low success |      |

Plant snapshot

|            |                                                                                                       |
|------------|-------------------------------------------------------------------------------------------------------|
| Habit      | Vigorous twining climber or scrambling shrub.                                                         |
| Flowering  | Jul - Oct (blue-purple, occasionally white).                                                          |
| Habitat    | Sandy soils; coastal limestone, sandplains and dunes.                                                 |
| Local note | Recorded in Joondalup and Wanneroo. More vigorous than <i>H. violacea</i> - can smother small shrubs. |

Propagation options

| Method   | Suitability | Notes                                      | Details in          |
|----------|-------------|--------------------------------------------|---------------------|
| Cuttings | Good        | Firm current season's growth strikes well. | This guide          |
| Seed     | Main method | Hot water + smoke water; about 10 days.    | → Seed Sowing Guide |
| Division | Not used    | -                                          | -                   |

Propagation method

| Item                       | Details                                                                                                                                                           |
|----------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Best time to take cuttings | Dec - Mar, once the current season's growth has firmed after spring flowering. Nov and Apr are workable. Avoid May - Oct (cool, then flowering).                  |
| Wood type                  | Semi-hardwood - firm current season's growth (ANPSA).                                                                                                             |
| Cutting size               | 50-75 mm, the length ANPSA suggests for pea-family cuttings. One runner gives several pieces.                                                                     |
| Selecting material         | Healthy runners without flowers or pods.                                                                                                                          |
| Pre-treatment              | <ul style="list-style-type: none"><li>Remove the leaves from the lower two-thirds.</li><li>No wounding needed.</li><li>Dip the base in rooting hormone.</li></ul> |
| Rooting hormone            | Semi-hardwood strength, 3 g/L IBA (Clonex Purple).                                                                                                                |
| Propagation mix            | Standard free-draining cutting mix.                                                                                                                               |
| Environment                | Humid frame, bright shade, bottom heat.                                                                                                                           |
| Expected strike time       | Not documented - check from about 4 weeks and record the result.                                                                                                  |
| Post-treatment / aftercare | <ul style="list-style-type: none"><li>Standard hardening off.</li><li>Stake or trim runners so tubes don't tangle; pot on promptly.</li></ul>                     |

## Common problems

| Problem                      | Likely cause and fix                                                 |
|------------------------------|----------------------------------------------------------------------|
| Cuttings wilt                | Material too soft. Use firmer growth; keep humidity up until rooted. |
| Runners tangle between tubes | Space tubes; trim or stake runners.                                  |
| Base rotting                 | Mix too wet. Water less.                                             |

## Notes

|            |                                                                                                                            |
|------------|----------------------------------------------------------------------------------------------------------------------------|
| Difficulty | <b>Easy-Moderate.</b> Cuttings strike well. Seed is also easy after hot-water treatment.                                   |
| Confidence | <b>Medium.</b> ANPSA reports cuttings of firm current season's growth strike well; size follows ANPSA pea-family guidance. |
| Sources    | See References: 4, 13, 43, 44                                                                                              |

## Nursery records

Log each batch here to confirm or update the timing and difficulty above.

| Date set | Stock source / provenance | No. set | No. struck | Strike % | Notes |
|----------|---------------------------|---------|------------|----------|-------|
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |

Family  
**Lamiaceae**

## Lamiaceae

## Plant snapshot

## Propagation options

## Propagation method

Version 1.6 • September 2026

## Common problems

| Problem         | Likely cause and fix                                                      |
|-----------------|---------------------------------------------------------------------------|
| Cuttings rot    | Too humid or wet. Mist less and improve airflow.                          |
| Cuttings flower | Taken from flowering tips. Snip off buds and choose non-flowering shoots. |
| Slow in winter  | Use bottom heat, or wait for the warm window.                             |

## Notes

|               |                                                                                                                                                                     |
|---------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Difficulty    | <b>Easy.</b> ANPSA describes it as relatively easy to propagate from cuttings.                                                                                      |
| Other methods | Seed – rarely available and germinates poorly (about 4% untreated in Alcoa nursery tests). Grafting onto <i>Westringia fruticosa</i> is used where plants struggle. |
| Confidence    | <b>Medium.</b> ANPSA says it is relatively easy from cuttings; timing and size follow standard practice.                                                            |
| Sources       | See References: 4, 32, 45, 46, 47                                                                                                                                   |

## Nursery records

Log each batch here to confirm or update the timing and difficulty above.

| Date set | Stock source / provenance | No. set | No. struck | Strike % | Notes |
|----------|---------------------------|---------|------------|----------|-------|
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |

# Hibbertia cuneiformis

Cutleaf Hibbertia

Family  
Dilleniaceae

| DIFFICULTY      | WOOD TYPE     | BEST MONTHS | CUTTING LENGTH | HORMONE (IBA) |
|-----------------|---------------|-------------|----------------|---------------|
| Easy-Moderate * | Semi-hardwood | Mar – May   | 75–100 mm      | 3 g/L IBA     |

|          | Jan | Feb | Mar  | Apr  | May  | Jun | Jul | Aug | Sep | Oct | Nov | Dec |
|----------|-----|-----|------|------|------|-----|-----|-----|-----|-----|-----|-----|
| Cuttings | OK  | OK  | Best | Best | Best | –   | –   | –   | –   | OK  | OK  | OK  |

|             |                                                   |                     |
|-------------|---------------------------------------------------|---------------------|
| Best window | Workable – see 'Selecting material' in the method | Avoid / low success |
|-------------|---------------------------------------------------|---------------------|

## Plant snapshot

|            |                                                                                                |
|------------|------------------------------------------------------------------------------------------------|
| Habit      | Erect or sprawling shrub, 0.5–3 m (usually 1–2 m). Wedge-shaped leaves, toothed near the tip.  |
| Flowering  | Jan – Mar or Jun – Nov (yellow).                                                               |
| Habitat    | Coastal sand dunes and swampy places near the coast.                                           |
| Local note | Recorded in the City of Joondalup. One of the more vigorous hibbertias – useful for screening. |

## Propagation options

| Method   | Suitability | Notes                                            | Details in |
|----------|-------------|--------------------------------------------------|------------|
| Cuttings | Main method | Commonly propagated by cuttings.                 | This guide |
| Seed     | Difficult   | 0% in a nursery test; Hibbertia seed is dormant. | –          |
| Division | Not used    | –                                                | –          |

## Propagation method

| Item                       | Details                                                                                                                                                                                                              |
|----------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Best time to take cuttings | Mar – May, between the summer and winter flowering periods, once new growth has firmed. Oct – Feb is workable using non-flowering shoots. Avoid Jun – Sep (cool, and plants are flowering).                          |
| Wood type                  | Semi-hardwood – current season's growth that has begun to firm.                                                                                                                                                      |
| Cutting size               | 75–100 mm.                                                                                                                                                                                                           |
| Selecting material         | Non-flowering shoots from healthy plants. Snip off any buds.                                                                                                                                                         |
| Pre-treatment              | <ul style="list-style-type: none"> <li>Strip leaves from the lower half.</li> <li>Reduce large leaves by about one-third.</li> <li>Light wound on firmer stems.</li> <li>Dip the base in rooting hormone.</li> </ul> |
| Rooting hormone            | Semi-hardwood strength, 3 g/L IBA (Clonex Purple).                                                                                                                                                                   |
| Propagation mix            | Very free-draining, sand-based mix. Use fresh mix and clean tubes – hibbertias are prone to <i>Phytophthora</i> .                                                                                                    |
| Environment                | Humid frame, bright shade, bottom heat. Moist, never wet; good airflow.                                                                                                                                              |
| Expected strike time       | Roughly 4–8 weeks in warm weather (typical for the genus). Record your results.                                                                                                                                      |
| Post-treatment / aftercare | <ul style="list-style-type: none"> <li>Standard hardening off.</li> <li>Pot on into a free-draining native mix; avoid overwatering.</li> <li>Tip-prune for a bushier plant.</li> </ul>                               |

## Common problems

| Problem                            | Likely cause and fix                                                             |
|------------------------------------|----------------------------------------------------------------------------------|
| Cuttings collapse at the base      | Too wet, or root rot. Improve drainage, use clean mix, remove affected cuttings. |
| Cuttings flower instead of rooting | Material taken from flowering shoots. Use non-flowering shoots.                  |
| Slow to strike                     | Cool weather. Use bottom heat or wait for the Mar – May window.                  |

## Notes

|               |                                                                                                                                                                                                                    |
|---------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Difficulty    | <b>Easy-Moderate (provisional).</b> Commonly grown from cuttings. Hibbertias usually strike well, but <i>H. hypericoides</i> shows not every WA species does – set extra cuttings until your strike rate is known. |
| Other methods | Seed – 0% germination in the one Alcoa nursery test; <i>Hibbertia</i> seed has complex dormancy.<br>Layering – check low, sprawling stems for roots.                                                               |
| Confidence    | <b>Medium.</b> Commonly propagated from cuttings in the nursery trade; little published detail on timing or strike rate.                                                                                           |
| Sources       | See References: 1, 4, 9, 10, 32, 39, 40                                                                                                                                                                            |

## Nursery records

Log each batch here to confirm or update the timing and difficulty above.

| Date set | Stock source / provenance | No. set | No. struck | Strike % | Notes |
|----------|---------------------------|---------|------------|----------|-------|
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |

# Hibbertia hypericoides

Yellow Buttercups

Family  
Dilleniaceae

| DIFFICULTY       | WOOD TYPE            | BEST MONTHS      | CUTTING LENGTH  | HORMONE (IBA)                  |
|------------------|----------------------|------------------|-----------------|--------------------------------|
| <b>Difficult</b> | <b>Semi-hardwood</b> | <b>Dec – Feb</b> | <b>50–75 mm</b> | <b>3 g/L IBA (trial range)</b> |

|          | Jan  | Feb  | Mar | Apr | May | Jun | Jul | Aug | Sep | Oct | Nov | Dec  |
|----------|------|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|------|
| Cuttings | Best | Best | OK  | –   | –   | –   | –   | –   | –   | –   | OK  | Best |

Best window
Workable – see 'Selecting material' in the method
Avoid / low success

## Plant snapshot

|                   |                                                                                                                                                             |
|-------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Habit</b>      | Spreading shrub, 0.3–0.8 m. Densely hairy branches; small leaves with rolled edges and white-hairy undersides.                                              |
| <b>Flowering</b>  | Apr – Nov (bright yellow, notched petals).                                                                                                                  |
| <b>Habitat</b>    | Sandy soils in Banksia woodland and shrubland; one of the most common understorey plants between Perth and Albany. Resprouts from its rootstock after fire. |
| <b>Local note</b> | Recorded in the City of Joondalup.                                                                                                                          |

## Propagation options

| Method   | Suitability      | Notes                                                       | Details in                 |
|----------|------------------|-------------------------------------------------------------|----------------------------|
| Cuttings | <b>Difficult</b> | Known to be difficult; trial protocol on the cuttings page. | <a href="#">This guide</a> |
| Seed     | <b>Difficult</b> | Complex seed dormancy; rarely practical.                    | –                          |
| Division | <b>Not used</b>  | Resprouts from a rootstock; not practical to divide.        | –                          |

## Propagation method

| Item                              | Details                                                                                                                                                                           |
|-----------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Best time to take cuttings</b> | <b>Dec – Feb</b> , after flowering finishes and once the new flush has started to firm. Nov and Mar are workable. Avoid Apr – Oct (flowering period).                             |
| <b>Wood type</b>                  | <b>Semi-hardwood</b> – new growth that is just firming. Soft tips wilt quickly; old wood rarely roots.                                                                            |
| <b>Cutting size</b>               | 50–75 mm.                                                                                                                                                                         |
| <b>Selecting material</b>         | Vigorous <b>non-flowering</b> shoots. Well-watered, actively growing stock plants (e.g. nursery-held stock) usually give better material than stressed bushland plants.           |
| <b>Pre-treatment</b>              | <ul style="list-style-type: none"> <li>Strip lower leaves by cutting, not pulling.</li> <li>Light wound on the firmer stems.</li> <li>Dip the base in rooting hormone.</li> </ul> |
| <b>Rooting hormone</b>            | Start with 3 g/L IBA ( <b>Clonex Purple</b> ). Because results are unreliable, split trial batches across 1.5, 3 and 8 g/L (Green, Purple, Red) and record which works best.      |
| <b>Propagation mix</b>            | Very free-draining, fresh, clean mix. Strict hygiene – hibbertias are prone to <i>Phytophthora</i> root rot.                                                                      |
| <b>Environment</b>                | Humid frame, bottom heat, bright shade. Mist lightly – the hairy leaf undersides hold water. Good airflow.                                                                        |
| <b>Expected strike time</b>       | Slow and variable – allow <b>8–16 weeks</b> . Leave cuttings in as long as they stay green and healthy.                                                                           |

| Item                       | Details                                                                                                                                                                                                                          |
|----------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Post-treatment / aftercare | <ul style="list-style-type: none"> <li>• Harden off slowly (extend each stage if cuttings wilt).</li> <li>• Pot on gently, disturbing roots as little as possible.</li> <li>• Avoid overwatering; keep benches clean.</li> </ul> |

### Common problems

| Problem                              | Likely cause and fix                                                                |
|--------------------------------------|-------------------------------------------------------------------------------------|
| Cuttings stay green but don't root   | Leave longer; check for heavy callus and trim it; try a different hormone strength. |
| Cuttings collapse or rot             | Too wet or root rot. Improve drainage and hygiene; remove affected cuttings.        |
| Rooted cuttings die after potting on | Roots damaged or hardening too fast. Use single tubes and harden off more slowly.   |

### Notes

|               |                                                                                                                                                                                                                   |
|---------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Difficulty    | <b>Difficult.</b> DBCA notes it is difficult to grow from seed or cuttings, and specialist Perth nurseries describe it as a species other producers often fail with. Set large batches and keep detailed records. |
| Other methods | Seed – also difficult: WA <i>Hibbertia</i> seed has complex dormancy, and treatments are still the subject of research. If numbers are needed quickly, consider buying in tubestock from a specialist grower.     |
| Confidence    | <b>Low.</b> Its difficulty is well documented (DBCA; specialist WA nurseries), but no detailed cutting protocol is published. Treat the method below as a trial protocol.                                         |
| Sources       | See References: 1, 4, 10, 11, 27, 30, 31                                                                                                                                                                          |

### Nursery records

Log each batch here to confirm or update the timing and difficulty above.

| Date set | Stock source / provenance | No. set | No. struck | Strike % | Notes |
|----------|---------------------------|---------|------------|----------|-------|
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |

# Hibbertia racemosa

Stalked Guinea Flower

Family  
Dilleniaceae

|                                                   |               |      |      |                     |             |     |                |     |     |               |     |     |
|---------------------------------------------------|---------------|------|------|---------------------|-------------|-----|----------------|-----|-----|---------------|-----|-----|
| DIFFICULTY                                        | WOOD TYPE     |      |      |                     | BEST MONTHS |     | CUTTING LENGTH |     |     | HORMONE (IBA) |     |     |
| Moderate *                                        | Semi-hardwood |      |      |                     | Jan - Apr   |     | 50-75 mm       |     |     | 1.5-3 g/L IBA |     |     |
|                                                   | Jan           | Feb  | Mar  | Apr                 | May         | Jun | Jul            | Aug | Sep | Oct           | Nov | Dec |
| Cuttings                                          | Best          | Best | Best | Best                | OK          | -   | -              | -   | -   | -             | -   | OK  |
| Best window                                       |               |      |      |                     |             |     |                |     |     |               |     |     |
| Workable - see 'Selecting material' in the method |               |      |      |                     |             |     |                |     |     |               |     |     |
|                                                   |               |      |      | Avoid / low success |             |     |                |     |     |               |     |     |

Plant snapshot

|            |                                                                                                                              |
|------------|------------------------------------------------------------------------------------------------------------------------------|
| Habit      | Small erect or ascending, spreading shrub, 0.1-0.75 m.                                                                       |
| Flowering  | Jul - Dec (yellow, on long stalks).                                                                                          |
| Habitat    | Grey, white or yellow sand in coastal areas: dunes, plains and limestone.                                                    |
| Local note | Recorded in the City of Joondalup. Grey-leaved and prostrate forms are sold in the trade - keep forms separate and labelled. |

Propagation options

| Method   | Suitability | Notes                                                    | Details in |
|----------|-------------|----------------------------------------------------------|------------|
| Cuttings | Main method | Semi-hardwood cuttings (provisional rating).             | This guide |
| Seed     | Difficult   | Hibbertia seed is dormant and unreliable.                | -          |
| Division | Possible    | Low stems may root where they touch the soil (layering). | -          |

Propagation method

| Item                       | Details                                                                                                                                                                                                                                                       |
|----------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Best time to take cuttings | Jan - Apr, after flowering finishes and new growth has begun to firm. Dec and May are workable. Avoid Jun - Nov (cool weather, then flowering).                                                                                                               |
| Wood type                  | Semi-hardwood - new growth that has started to harden. Firm tip cuttings also work.                                                                                                                                                                           |
| Cutting size               | 50-75 mm long. Stems are fine and leaves small, so keep cuttings short.                                                                                                                                                                                       |
| Selecting material         | Non-flowering side shoots from healthy plants. Snip off any flower stalks or buds.                                                                                                                                                                            |
| Pre-treatment              | <ul style="list-style-type: none"><li>Strip leaves from the lower half (cut rather than pull on fine stems).</li><li>Wounding not needed on fine stems; a light wound is optional on firmer material.</li><li>Dip the base in rooting hormone.</li></ul>      |
| Rooting hormone            | Semi-hardwood strength (3 g/L IBA, e.g. Clonex Purple). For softer tips use softwood strength (1.5 g/L IBA, e.g. Clonex Green).                                                                                                                               |
| Propagation mix            | Very free-draining, sand-based mix. Use fresh, clean mix and clean tubes - hibbertias are prone to Phytophthora root rot.                                                                                                                                     |
| Environment                | Humid frame, bright shade, bottom heat. Keep the mix moist but never wet; good airflow.                                                                                                                                                                       |
| Expected strike time       | Roughly 4-8 weeks in warm weather; may be slower. Record your results.                                                                                                                                                                                        |
| Post-treatment / aftercare | <ul style="list-style-type: none"><li>Standard hardening off (2 weeks closed by day/open at night, then 2 weeks shade).</li><li>Pot on into a free-draining native mix; avoid overwatering.</li><li>Keep good hygiene on benches to limit root rot.</li></ul> |

## Common problems

| Problem                       | Likely cause and fix                                                             |
|-------------------------------|----------------------------------------------------------------------------------|
| Cuttings collapse at the base | Too wet, or root rot. Improve drainage, use clean mix, remove affected cuttings. |
| Few roots, cuttings flower    | Material was taken from flowering shoots. Use non-flowering side shoots.         |
| Slow to strike in winter      | Use bottom heat, or wait for the Jan – Apr window.                               |

## Notes

|               |                                                                                                                                                                                                                   |
|---------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Difficulty    | <b>Moderate (provisional).</b> Hibbertias are generally easy from cuttings, but some WA species (e.g. <i>H. hypericoides</i> ) are known to be difficult. Set larger batches until your own strike rate is known. |
| Other methods | Seed – generally unreliable in <i>Hibbertia</i> (complex dormancy). Layering – several hibbertias root where stems touch the ground; check low stems on prostrate forms and pot up any rooted pieces.             |
| Confidence    | <b>Medium.</b> Little species-specific data published. Based on how the genus behaves and this species' growth cycle.                                                                                             |
| Sources       | See References: 1, 4, 8, 9, 10, 11                                                                                                                                                                                |

## Nursery records

Log each batch here to confirm or update the timing and difficulty above.

| Date set | Stock source / provenance | No. set | No. struck | Strike % | Notes |
|----------|---------------------------|---------|------------|----------|-------|
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |

# Hibbertia spicata

No widely used common name

Perth coastal plants are now *Hibbertia leptotheca* (formerly *H. spicata* subsp. *leptotheca*)

Family  
Dilleniaceae

| DIFFICULTY | WOOD TYPE     | BEST MONTHS | CUTTING LENGTH | HORMONE (IBA) |
|------------|---------------|-------------|----------------|---------------|
| Moderate * | Semi-hardwood | Jan – Mar   | 50–75 mm       | 1.5–3 g/L IBA |

|          | Jan  | Feb  | Mar  | Apr | May | Jun | Jul | Aug | Sep | Oct | Nov | Dec |
|----------|------|------|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| Cuttings | Best | Best | Best | OK  | –   | –   | –   | –   | –   | –   | –   | OK  |

Best window

Workable – see ‘Selecting material’ in the method

Avoid / low success

Plant snapshot

|            |                                                                                                                                                                                                                                                                                                                                                                                                      |
|------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Habit      | Small erect or spreading shrub, 0.2–0.7 m ( <i>H. leptotheca</i> 0.2–0.5 m).                                                                                                                                                                                                                                                                                                                         |
| Flowering  | Jul – Nov (yellow).                                                                                                                                                                                                                                                                                                                                                                                  |
| Habitat    | Sand, limestone and laterite; near-coastal limestone ridges, outcrops and cliffs.                                                                                                                                                                                                                                                                                                                    |
| Local note | A recent revision split <i>H. spicata</i> (in the broad sense) into six species. True <i>H. spicata</i> is recorded around Geraldton – Shark Bay and Augusta – Margaret River. The Perth coastal plant is <i>H. leptotheca</i> (recorded in Wanneroo). It is a <b>Priority Three</b> species, so take cuttings only from nursery stock or under the right DBCA licence, and check provenance labels. |

Propagation options

| Method   | Suitability | Notes                                                  | Details in                 |
|----------|-------------|--------------------------------------------------------|----------------------------|
| Cuttings | Main method | Most hibbertias are grown from cuttings (provisional). | <a href="#">This guide</a> |
| Seed     | Difficult   | Dormant; 9% in the one Alcoa test; GA may help.        | –                          |
| Division | Not used    | Shrub – not divided.                                   | –                          |

Propagation method

| Item                       | Details                                                                                                                                                                                          |
|----------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Best time to take cuttings | Jan – Mar, after flowering, once new growth has firmed. <i>H. empetrifolia</i> cuttings strike readily in Jan – Feb. Dec and Apr are workable. Avoid May – Nov (cool, and plants are flowering). |
| Wood type                  | Semi-hardwood – current season's growth that has begun to firm.                                                                                                                                  |
| Cutting size               | 50–75 mm.                                                                                                                                                                                        |
| Selecting material         | Non-flowering shoots from healthy plants. Snip off any buds.                                                                                                                                     |
| Pre-treatment              | <ul style="list-style-type: none"><li>Strip leaves from the lower half.</li><li>Dip the base in rooting hormone.</li></ul>                                                                       |
| Rooting hormone            | 1.5 g/L IBA ( <b>Clonex Green</b> ) for soft tips; 3 g/L ( <b>Clonex Purple</b> ) for firmer stems.                                                                                              |
| Propagation mix            | Very free-draining, sand-based mix. Use fresh mix and clean tubes – hibbertias are prone to root rot.                                                                                            |
| Environment                | Humid frame, bright shade, bottom heat. Moist, never wet; good airflow.                                                                                                                          |
| Expected strike time       | Not documented – check from about 6 weeks and record the result.                                                                                                                                 |
| Post-treatment / aftercare | <ul style="list-style-type: none"><li>Standard hardening off.</li><li>Pot on into a free-draining native mix; avoid overwatering.</li></ul>                                                      |

## Common problems

| Problem                            | Likely cause and fix                                                               |
|------------------------------------|------------------------------------------------------------------------------------|
| Cuttings collapse at the base      | Too wet, or root rot. Improve drainage; clean mix; remove affected cuttings.       |
| Cuttings flower instead of rooting | Material from flowering shoots. Use non-flowering shoots.                          |
| Very low strike                    | Some WA hibbertias are hard. Try softer tips, bottom heat, and record which works. |

## Notes

|               |                                                                                                                                                                      |
|---------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Difficulty    | <b>Moderate (provisional).</b> Untested for this species. Set extra cuttings until your strike rate is known.                                                        |
| Other methods | Seed – dormant: 9% in the one Alcoa nursery test (about 30 seedlings per gram). Removing the seed coat or using gibberellic acid (25 mg/L) can help (WA seed notes). |
| Confidence    | <b>Low.</b> No species data. WA seed notes say most hibbertias are grown from cuttings, but <i>H. hypericoides</i> shows some WA species are hard.                   |
| Sources       | See References: 1, 4, 9, 13, 32, 69, 70                                                                                                                              |

## Nursery records

Log each batch here to confirm or update the timing and difficulty above.

| Date set | Stock source / provenance | No. set | No. struck | Strike % | Notes |
|----------|---------------------------|---------|------------|----------|-------|
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |

# Hypocalymma robustum

Swan River Myrtle

Family  
Myrtaceae

| DIFFICULTY  | WOOD TYPE                                         |      |      |     | BEST MONTHS |     |                     | CUTTING LENGTH |     |     | HORMONE (IBA) |      |  |
|-------------|---------------------------------------------------|------|------|-----|-------------|-----|---------------------|----------------|-----|-----|---------------|------|--|
| Easy        | Semi-hardwood                                     |      |      |     | Dec - Mar   |     |                     | 75-100 mm      |     |     | 3 g/L IBA     |      |  |
|             | Jan                                               | Feb  | Mar  | Apr | May         | Jun | Jul                 | Aug            | Sep | Oct | Nov           | Dec  |  |
| Cuttings    | Best                                              | Best | Best | OK  | -           | -   | -                   | -              | -   | -   | OK            | Best |  |
| Best window | Workable - see 'Selecting material' in the method |      |      |     |             |     | Avoid / low success |                |     |     |               |      |  |

## Plant snapshot

|            |                                                                            |
|------------|----------------------------------------------------------------------------|
| Habit      | Erect shrub, 0.4–1 (–1.5) m.                                               |
| Flowering  | Jun – Nov (pink to pink-red, along the stems).                             |
| Habitat    | Gravelly lateritic soils and sandy soils; undulating ground and ridges.    |
| Local note | Recorded in Joondalup and Wanneroo. Prune yearly for a dense, bushy plant. |

## Propagation options

| Method   | Suitability | Notes                                                  | Details in |
|----------|-------------|--------------------------------------------------------|------------|
| Cuttings | Main method | Strikes readily from hardened current season's growth. | This guide |
| Seed     | Possible    | Smoke water likely helps; about 35% in one test.       | –          |
| Division | Not used    | –                                                      | –          |

## Propagation method

| Item                       | Details                                                                                                                                                                        |
|----------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Best time to take cuttings | Dec – Mar (January is suggested for the Geraldton wax group), once the current season's growth has hardened. Nov and Apr are workable. Avoid May – Oct (cool, then flowering). |
| Wood type                  | Semi-hardwood – hardened current season's growth (ANPSA).                                                                                                                      |
| Cutting size               | 75–100 mm.                                                                                                                                                                     |
| Selecting material         | Non-flowering tips from healthy plants.                                                                                                                                        |
| Pre-treatment              | <ul style="list-style-type: none"> <li>Carefully remove the leaves from the lower two-thirds.</li> <li>Wound the base.</li> <li>Dip the base in rooting hormone.</li> </ul>    |
| Rooting hormone            | Semi-hardwood strength, 3 g/L IBA ( <b>Clonex Purple</b> ) – wounding and hormone both improve results.                                                                        |
| Propagation mix            | Standard free-draining cutting mix. Clean mix and tubes – the genus includes dieback-affected species.                                                                         |
| Environment                | Humid frame, bright shade, bottom heat.                                                                                                                                        |
| Expected strike time       | Not documented – usually quick; check from about 4 weeks.                                                                                                                      |
| Post-treatment / aftercare | <ul style="list-style-type: none"> <li>Standard hardening off.</li> <li>Pot on into a free-draining native mix; tip-prune for bushiness.</li> </ul>                            |

## Common problems

| Problem            | Likely cause and fix                                        |
|--------------------|-------------------------------------------------------------|
| Cuttings brown off | Too soft or too dry. Use hardened growth; keep humidity up. |
| Base rotting       | Mix too wet. Water less, improve drainage.                  |
| Leggy plants       | Not pruned. Tip-prune young plants.                         |

## Notes

|               |                                                                                                                                            |
|---------------|--------------------------------------------------------------------------------------------------------------------------------------------|
| Difficulty    | <b>Easy.</b> Cuttings are the preferred method and strike readily (ANPSA).                                                                 |
| Other methods | Seed – possible; smoke water is likely to help (about 35% in a single Alcoa test with smoke).                                              |
| Confidence    | <b>Medium.</b> ANPSA says cuttings are the preferred method and strike readily; size and treatment follow the related Geraldton wax group. |
| Sources       | See References: 4, 32, 53, 54, 55                                                                                                          |

## Nursery records

Log each batch here to confirm or update the timing and difficulty above.

| Date set | Stock source / provenance | No. set | No. struck | Strike % | Notes |
|----------|---------------------------|---------|------------|----------|-------|
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |

# Kennedia prostrata

Scarlet Runner / Running Postman

Family  
Fabaceae

|               |               |             |                |               |
|---------------|---------------|-------------|----------------|---------------|
| DIFFICULTY    | WOOD TYPE     | BEST MONTHS | CUTTING LENGTH | HORMONE (IBA) |
| Easy-Moderate | Semi-hardwood | Dec - Mar   | 75-100 mm      | 3 g/L IBA     |

|          | Jan         | Feb  | Mar                                               | Apr | May | Jun | Jul | Aug | Sep | Oct | Nov | Dec                 |
|----------|-------------|------|---------------------------------------------------|-----|-----|-----|-----|-----|-----|-----|-----|---------------------|
| Cuttings | Best        | Best | Best                                              | OK  | -   | -   | -   | -   | -   | -   | OK  | Best                |
|          | Best window |      | Workable - see 'Selecting material' in the method |     |     |     |     |     |     |     |     | Avoid / low success |

Plant snapshot

|            |                                                             |
|------------|-------------------------------------------------------------|
| Habit      | Prostrate or twining groundcover, spreading to about 2.5 m. |
| Flowering  | Apr - Nov (red pea flowers).                                |
| Habitat    | Usually sandy, gravelly soils.                              |
| Local note | Recorded in Joondalup and Wanneroo.                         |

Propagation options

| Method   | Suitability | Notes                                      | Details in          |
|----------|-------------|--------------------------------------------|---------------------|
| Cuttings | Good        | Firm current season's growth strikes well. | This guide          |
| Seed     | Main method | Hot water + smoke water; about 8 days.     | → Seed Sowing Guide |
| Division | Not used    | -                                          | -                   |

Propagation method

| Item                       | Details                                                                                                                                                                                                 |
|----------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Best time to take cuttings | Dec - Mar, when the current season's runners have firmed. Nov and Apr are workable. Avoid May - Oct (flowering and cool weather).                                                                       |
| Wood type                  | Semi-hardwood - firm current season's growth (ANPSA).                                                                                                                                                   |
| Cutting size               | 75-100 mm with 2-3 nodes. One runner can be cut into several pieces.                                                                                                                                    |
| Selecting material         | Healthy runners without flowers or pods.                                                                                                                                                                |
| Pre-treatment              | <ul style="list-style-type: none"><li>Remove the leaves from the lower nodes; cut remaining leaflets back by about half.</li><li>No wounding needed.</li><li>Dip the base in rooting hormone.</li></ul> |
| Rooting hormone            | Semi-hardwood strength, 3 g/L IBA (Clonex Purple).                                                                                                                                                      |
| Propagation mix            | Standard free-draining cutting mix.                                                                                                                                                                     |
| Environment                | Humid frame, bright shade, bottom heat.                                                                                                                                                                 |
| Expected strike time       | Not documented - check from about 4 weeks and record the result.                                                                                                                                        |
| Post-treatment / aftercare | <ul style="list-style-type: none"><li>Standard hardening off.</li><li>Pot on promptly - runners tangle across the bench. Trim runners to keep tubes tidy.</li></ul>                                     |

## Common problems

| Problem                       | Likely cause and fix                                        |
|-------------------------------|-------------------------------------------------------------|
| Cuttings wilt and collapse    | Material too soft. Use firmer runners and keep humidity up. |
| Runners tangled between tubes | Space tubes out and trim runners regularly.                 |
| Base rotting                  | Mix too wet. Water less, improve drainage.                  |

## Notes

|            |                                                                                                                              |
|------------|------------------------------------------------------------------------------------------------------------------------------|
| Difficulty | <b>Easy-Moderate.</b> Cuttings of firm growth strike well. Seed is quick and is the usual nursery method.                    |
| Confidence | <b>Medium.</b> ANPSA reports cuttings of firm current season's growth strike well; timing and size follow standard practice. |
| Sources    | See References: 4, 13, 42                                                                                                    |

## Nursery records

Log each batch here to confirm or update the timing and difficulty above.

| Date set | Stock source / provenance | No. set | No. struck | Strike % | Notes |
|----------|---------------------------|---------|------------|----------|-------|
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |

# Lechenaultia linarioides

Yellow Leschenaultia

Family  
Goodeniaceae

| DIFFICULTY | WOOD TYPE                | BEST MONTHS | CUTTING LENGTH | HORMONE (IBA) |
|------------|--------------------------|-------------|----------------|---------------|
| Easy *     | Softwood - semi-hardwood | Feb - Apr   | 50-75 mm       | 1.5-3 g/L IBA |

|          | Jan | Feb  | Mar  | Apr  | May | Jun | Jul | Aug | Sep | Oct | Nov | Dec |
|----------|-----|------|------|------|-----|-----|-----|-----|-----|-----|-----|-----|
| Cuttings | OK  | Best | Best | Best | OK  | -   | -   | -   | -   | OK  | OK  | OK  |

|             |                                                   |                     |
|-------------|---------------------------------------------------|---------------------|
| Best window | Workable - see 'Selecting material' in the method | Avoid / low success |
|-------------|---------------------------------------------------|---------------------|

Plant snapshot

|            |                                                                                                 |
|------------|-------------------------------------------------------------------------------------------------|
| Habit      | Sprawling, open shrub to 1.5 m, with narrow, linear leaves.                                     |
| Flowering  | Can flower in any month. Flowers yellow with red or pink tones, sometimes cream.                |
| Habitat    | Sand (often over limestone) and sandy clay; sandplains, low dunes, ridges and winter-wet flats. |
| Local note | Recorded in the Wanneroo LGA.                                                                   |

Propagation options

| Method   | Suitability | Notes                                               | Details in                 |
|----------|-------------|-----------------------------------------------------|----------------------------|
| Cuttings | Main method | Cuttings strike easily in this genus.               | <a href="#">This guide</a> |
| Seed     | Difficult   | Slow; trays may need a second season.               | -                          |
| Division | Possible    | Some leschenaultias sucker - pot up rooted suckers. | -                          |

Propagation method

| Item                       | Details                                                                                                                                                                                      |
|----------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Best time to take cuttings | <b>Feb - Apr</b> (late summer to autumn) while it is warm. One grower reports roots in 4-6 weeks from late-February leschenaultia cuttings. Oct - Jan and May are workable. Avoid Jun - Sep. |
| Wood type                  | <b>Softwood to semi-hardwood</b> tip cuttings.                                                                                                                                               |
| Cutting size               | 50-75 mm.                                                                                                                                                                                    |
| Selecting material         | <b>Non-flowering</b> tips. This species can flower in any month, so check each cutting and remove buds.                                                                                      |
| Pre-treatment              | <ul style="list-style-type: none"><li>Strip leaves from the lower half (narrow leaves strip easily).</li><li>No wounding needed.</li><li>Dip the base in rooting hormone.</li></ul>          |
| Rooting hormone            | 1.5 g/L IBA ( <b>Clonex Green</b> ) for soft tips; 3 g/L ( <b>Clonex Purple</b> ) for firmer material.                                                                                       |
| Propagation mix            | Very free-draining - leschenaultias dislike wet roots.                                                                                                                                       |
| Environment                | Humid frame, bottom heat, bright shade. Keep the mix moist, never wet.                                                                                                                       |
| Expected strike time       | About <b>4-6 weeks</b> in warm weather.                                                                                                                                                      |
| Post-treatment / aftercare | <ul style="list-style-type: none"><li>Standard hardening off.</li><li>Pot on into a free-draining mix; water sparingly.</li><li>Tip-prune to counter the sprawling habit.</li></ul>          |

## Common problems

| Problem              | Likely cause and fix                                                                    |
|----------------------|-----------------------------------------------------------------------------------------|
| Base rotting         | Overwatering. Water less often; add coarse sand or perlite to the mix.                  |
| Wilting in the frame | Material too soft or humidity too low. Use slightly firmer tips; keep the frame closed. |
| Cuttings flowering   | Remove buds; pick vegetative shoots.                                                    |

## Notes

|               |                                                                                                                                                                                                                                 |
|---------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Difficulty    | <b>Easy (provisional).</b> <i>Lechenaultia</i> cuttings generally strike very easily; seed is difficult.                                                                                                                        |
| Other methods | Seed – slow and unreliable. The Geraldton community nursery soaked seed in smoke water, saw germination at about 46 days, and kept trays for a second year. Division – some <i>Lechenaultias</i> sucker; pot up rooted suckers. |
| Confidence    | <b>Medium.</b> No species-specific cutting data found. Based on the genus, whose cuttings strike very easily.                                                                                                                   |
| Sources       | See References: 1, 4, 12, 13, 24, 25                                                                                                                                                                                            |

## Nursery records

Log each batch here to confirm or update the timing and difficulty above.

| Date set | Stock source / provenance | No. set | No. struck | Strike % | Notes |
|----------|---------------------------|---------|------------|----------|-------|
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |

# Leucophyta brownii

Cushion Bush

Family  
Asteraceae

| DIFFICULTY      | WOOD TYPE     | BEST MONTHS          | CUTTING LENGTH | HORMONE (IBA) |
|-----------------|---------------|----------------------|----------------|---------------|
| Easy-Moderate * | Semi-hardwood | Mar – May; Sep – Oct | 50–75 mm       | 1.5–3 g/L IBA |

|          | Jan         | Feb                                               | Mar  | Apr  | May  | Jun | Jul | Aug | Sep  | Oct  | Nov | Dec                 |
|----------|-------------|---------------------------------------------------|------|------|------|-----|-----|-----|------|------|-----|---------------------|
| Cuttings | –           | OK                                                | Best | Best | Best | OK  | –   | OK  | Best | Best | OK  | –                   |
|          | Best window | Workable – see 'Selecting material' in the method |      |      |      |     |     |     |      |      |     | Avoid / low success |

## Plant snapshot

|            |                                                                                     |
|------------|-------------------------------------------------------------------------------------|
| Habit      | Compact, rounded shrub to about 1 m, densely covered in silvery-white woolly hairs. |
| Flowering  | Dec or Jan – Feb (small yellow button heads).                                       |
| Habitat    | Coastal foredunes and cliffs.                                                       |
| Local note | Recorded in the Wanneroo LGA. Very salt-tolerant; grows poorly in high humidity.    |

## Propagation options

| Method   | Suitability | Notes                                            | Details in                 |
|----------|-------------|--------------------------------------------------|----------------------------|
| Cuttings | Main method | Semi-hardened cuttings are the preferred method. | <a href="#">This guide</a> |
| Seed     | Good        | No pre-treatment; 10–30 days.                    | → Seed Sowing Guide        |
| Division | Not used    | –                                                | –                          |

## Propagation method

| Item                       | Details                                                                                                                                                                                                                             |
|----------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Best time to take cuttings | Mar – May, after flowering, and Sep – Oct as spring growth firms. Feb, Jun, Aug and Nov are workable. Avoid Dec – Jan (flowering and heat) and Jul (cold).                                                                          |
| Wood type                  | Semi-hardwood – ANBG notes cuttings should come from semi-hardened branches.                                                                                                                                                        |
| Cutting size               | 50–75 mm tip cuttings.                                                                                                                                                                                                              |
| Selecting material         | Non-flowering tips from compact, healthy plants. Avoid soft, sappy growth.                                                                                                                                                          |
| Pre-treatment              | <ul style="list-style-type: none"> <li>• Rub the small leaves off the lower half.</li> <li>• No wounding.</li> <li>• Dip the base in rooting hormone.</li> </ul>                                                                    |
| Rooting hormone            | 1.5 g/L IBA ( <b>Clonex Green</b> ) for firm tips; 3 g/L for firmer material.                                                                                                                                                       |
| Propagation mix            | Very free-draining – add extra coarse sand or perlite.                                                                                                                                                                              |
| Environment                | <b>Keep it airy.</b> The woolly foliage rots in humidity: use a well-ventilated frame in bright light with little or no mist. Bottom heat helps. Keep the mix only just moist.                                                      |
| Expected strike time       | Not documented – check from about 4 weeks and record the result.                                                                                                                                                                    |
| Post-treatment / aftercare | <ul style="list-style-type: none"> <li>• Harden off promptly and grow on in full sun.</li> <li>• Free-draining mix; never let pots sit wet – plants die easily if waterlogged.</li> <li>• Tip-prune for a dense cushion.</li> </ul> |

Common problems

| Problem                            | Likely cause and fix                                            |
|------------------------------------|-----------------------------------------------------------------|
| Cuttings turn grey-brown and mushy | Too humid or wet. Reduce mist, open the frame, improve airflow. |
| Slow to root in winter             | Use bottom heat, or wait for the autumn or spring window.       |
| Loses its dense shape              | Tip-prune young plants.                                         |

Notes

|            |                                                                                                                                          |
|------------|------------------------------------------------------------------------------------------------------------------------------------------|
| Difficulty | <b>Easy-Moderate (provisional).</b> Cuttings are the preferred method; the main risk is rot from too much moisture or humidity.          |
| Confidence | <b>Medium.</b> ANBG growing notes say semi-hardened cuttings are the preferred method; timing and details follow general daisy practice. |
| Sources    | See References: 4, 34, 35                                                                                                                |

Nursery records

Log each batch here to confirm or update the timing and difficulty above.

| Date set | Stock source / provenance | No. set | No. struck | Strike % | Notes |
|----------|---------------------------|---------|------------|----------|-------|
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |

# Melaleuca huegelii

Chenille Honeymyrtle

Family  
Myrtaceae

|                      |                            |                          |                             |                            |
|----------------------|----------------------------|--------------------------|-----------------------------|----------------------------|
| DIFFICULTY<br>Easy * | WOOD TYPE<br>Semi-hardwood | BEST MONTHS<br>Feb – Apr | CUTTING LENGTH<br>75–100 mm | HORMONE (IBA)<br>3 g/L IBA |
|----------------------|----------------------------|--------------------------|-----------------------------|----------------------------|

|          | Jan         | Feb                                               | Mar  | Apr  | May | Jun | Jul | Aug | Sep                 | Oct | Nov | Dec |
|----------|-------------|---------------------------------------------------|------|------|-----|-----|-----|-----|---------------------|-----|-----|-----|
| Cuttings | OK          | Best                                              | Best | Best | OK  | –   | –   | –   | –                   | –   | –   | OK  |
|          | Best window | Workable – see ‘Selecting material’ in the method |      |      |     |     |     |     | Avoid / low success |     |     |     |

Plant snapshot

|            |                                                                           |
|------------|---------------------------------------------------------------------------|
| Habit      | Shrub or small tree, 0.5–5 m.                                             |
| Flowering  | Sep – Dec or Jan (pink, white or pink-purple).                            |
| Habitat    | Sand; limestone cliffs, coastal plains and dunes.                         |
| Local note | Recorded in Joondalup and Wanneroo. Cuttings keep a chosen flower colour. |

Propagation options

| Method   | Suitability | Notes                                        | Details in          |
|----------|-------------|----------------------------------------------|---------------------|
| Cuttings | Good        | Reliable in this genus; keeps a chosen form. | This guide          |
| Seed     | Main method | Surface-sow fine seed; no pre-treatment.     | → Seed Sowing Guide |
| Division | Not used    | Woody shrub.                                 | –                   |

Propagation method

| Item                       | Details                                                                                                                                                                          |
|----------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Best time to take cuttings | Feb – Apr, after flowering, once new growth has firmed. Jan, May and Dec are workable. Avoid Jun – Nov.                                                                          |
| Wood type                  | Semi-hardwood – firm current season's growth.                                                                                                                                    |
| Cutting size               | 75–100 mm (ANPSA).                                                                                                                                                               |
| Selecting material         | Non-flowering tips. Take from a plant with the flower colour you want.                                                                                                           |
| Pre-treatment              | <ul style="list-style-type: none"><li>Strip leaves from the lower half to two-thirds.</li><li>Wound the base on firmer stems.</li><li>Dip the base in rooting hormone.</li></ul> |
| Rooting hormone            | Semi-hardwood strength, 3 g/L IBA (Clonex Purple).                                                                                                                               |
| Propagation mix            | Standard free-draining cutting mix.                                                                                                                                              |
| Environment                | Humid frame, bright shade, bottom heat.                                                                                                                                          |
| Expected strike time       | Typically 4–8 weeks in warm weather. Record your results.                                                                                                                        |
| Post-treatment / aftercare | <ul style="list-style-type: none"><li>Standard hardening off.</li><li>Pot on into a free-draining native mix; tip-prune for bushiness.</li></ul>                                 |

## Common problems

| Problem                  | Likely cause and fix                                     |
|--------------------------|----------------------------------------------------------|
| Callus, no roots         | Trim and re-set; lower the hormone strength next batch.  |
| Leaves drop in the frame | Too dry or too hot. Shade more and keep humidity steady. |
| Base rotting             | Mix too wet. Water less.                                 |

## Notes

|            |                                                                                                     |
|------------|-----------------------------------------------------------------------------------------------------|
| Difficulty | <b>Easy (provisional).</b> Melaleucas are generally reliable from cuttings.                         |
| Confidence | <b>Medium.</b> Based on ANPSA <i>Melaleuca</i> genus guidance – cuttings are reliable in the genus. |
| Sources    | See References: 4, 13, 41                                                                           |

## Nursery records

Log each batch here to confirm or update the timing and difficulty above.

| Date set | Stock source / provenance | No. set | No. struck | Strike % | Notes |
|----------|---------------------------|---------|------------|----------|-------|
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |

# Melaleuca systema

Coastal Honeymyrtle

Family  
Myrtaceae

| DIFFICULTY | WOOD TYPE     | BEST MONTHS          | CUTTING LENGTH | HORMONE (IBA) |
|------------|---------------|----------------------|----------------|---------------|
| Easy *     | Semi-hardwood | Dec – Jan; Apr – May | 75–100 mm      | 3 g/L IBA     |

|          | Jan  | Feb | Mar | Apr  | May  | Jun | Jul | Aug | Sep | Oct | Nov | Dec  |
|----------|------|-----|-----|------|------|-----|-----|-----|-----|-----|-----|------|
| Cuttings | Best | OK  | OK  | Best | Best | OK  | –   | –   | –   | –   | OK  | Best |

Best window

Workable – see ‘Selecting material’ in the method

Avoid / low success

Plant snapshot

|            |                                                                                                                        |
|------------|------------------------------------------------------------------------------------------------------------------------|
| Habit      | Erect to spreading shrub, usually 0.5–2 m.                                                                             |
| Flowering  | Feb – Mar or Aug – Dec (yellow-cream).                                                                                 |
| Habitat    | Sand over laterite or limestone; stabilised coastal dunes and rocky limestone.                                         |
| Local note | Recorded in Joondalup and Wanneroo. Seed germination is low (about 14% in nursery tests), so cuttings are worth using. |

Propagation options

| Method   | Suitability | Notes                                                             | Details in                 |
|----------|-------------|-------------------------------------------------------------------|----------------------------|
| Cuttings | Good        | Reliable in this genus; a good option given low seed germination. | <a href="#">This guide</a> |
| Seed     | Main method | Surface-sow fine seed; no pre-treatment.                          | → Seed Sowing Guide        |
| Division | Not used    | Woody shrub.                                                      | –                          |

Propagation method

| Item                       | Details                                                                                                                                                                          |
|----------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Best time to take cuttings | Dec – Jan and Apr – May – the gaps between its two flowering periods. Feb – Mar, Jun and Nov are workable with non-flowering shoots. Avoid Jul – Oct.                            |
| Wood type                  | Semi-hardwood – firm current season's growth.                                                                                                                                    |
| Cutting size               | 75–100 mm (ANPSA).                                                                                                                                                               |
| Selecting material         | Non-flowering tips from healthy plants.                                                                                                                                          |
| Pre-treatment              | <ul style="list-style-type: none"><li>Strip leaves from the lower half to two-thirds.</li><li>Wound the base on firmer stems.</li><li>Dip the base in rooting hormone.</li></ul> |
| Rooting hormone            | Semi-hardwood strength, 3 g/L IBA (Clonex Purple).                                                                                                                               |
| Propagation mix            | Standard free-draining cutting mix.                                                                                                                                              |
| Environment                | Humid frame, bright shade, bottom heat. Autumn batches root more slowly.                                                                                                         |
| Expected strike time       | Typically 4–8 weeks in warm weather; longer in May. Record your results.                                                                                                         |
| Post-treatment / aftercare | <ul style="list-style-type: none"><li>Standard hardening off.</li><li>Pot on into a free-draining native mix.</li></ul>                                                          |

## Common problems

| Problem          | Likely cause and fix                                                            |
|------------------|---------------------------------------------------------------------------------|
| Callus, no roots | Trim and re-set; lower the hormone strength next batch.                         |
| Cuttings flower  | Taken from flowering shoots. Use non-flowering tips, between flowering periods. |
| Base rotting     | Mix too wet. Water less.                                                        |

## Notes

|            |                                                                                                     |
|------------|-----------------------------------------------------------------------------------------------------|
| Difficulty | <b>Easy (provisional).</b> Melaleucas are generally reliable from cuttings.                         |
| Confidence | <b>Medium.</b> Based on ANPSA <i>Melaleuca</i> genus guidance – cuttings are reliable in the genus. |
| Sources    | See References: 4, 13, 41                                                                           |

## Nursery records

Log each batch here to confirm or update the timing and difficulty above.

| Date set | Stock source / provenance | No. set | No. struck | Strike % | Notes |
|----------|---------------------------|---------|------------|----------|-------|
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |

# Myoporum caprarioides

Slender Myoporum

Family  
Scrophulariaceae

| DIFFICULTY | WOOD TYPE     | BEST MONTHS | CUTTING LENGTH | HORMONE (IBA) |
|------------|---------------|-------------|----------------|---------------|
| Easy *     | Semi-hardwood | Nov – Mar   | 75–100 mm      | 3 g/L IBA     |

|          | Jan  | Feb  | Mar  | Apr | May | Jun | Jul | Aug | Sep | Oct | Nov  | Dec  |
|----------|------|------|------|-----|-----|-----|-----|-----|-----|-----|------|------|
| Cuttings | Best | Best | Best | OK  | –   | –   | –   | –   | –   | OK  | Best | Best |

Best window
Workable – see 'Selecting material' in the method
Avoid / low success

## Plant snapshot

|            |                                                                                                               |
|------------|---------------------------------------------------------------------------------------------------------------|
| Habit      | Erect, slender, compact or sprawling shrub, (0.3–) 0.5–3 m.                                                   |
| Flowering  | Variable – from Feb, Apr or Jun to Dec (white, sometimes blue-pink tinged).                                   |
| Habitat    | White-grey to black sand, peaty sand and clay loam; seasonally wet flats, swamps, dunes and limestone ridges. |
| Local note | Recorded in Joondalup and Wanneroo. Suits sites that stay damp in winter. Small berries attract birds.        |

## Propagation options

| Method   | Suitability | Notes                                                                | Details in                 |
|----------|-------------|----------------------------------------------------------------------|----------------------------|
| Cuttings | Main method | Hardened current season's growth strikes readily in the genus.       | <a href="#">This guide</a> |
| Seed     | Possible    | Not documented; the related <i>M. insulare</i> is fermented + smoke. | –                          |
| Division | Not used    | Woody shrub.                                                         | –                          |

## Propagation method

| Item                       | Details                                                                                                                                                             |
|----------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Best time to take cuttings | Nov – Mar, in the warm months once current growth has hardened. Oct and Apr are workable. Avoid May – Sep (slow rooting).                                           |
| Wood type                  | Semi-hardwood – hardened current season's growth.                                                                                                                   |
| Cutting size               | 75–100 mm.                                                                                                                                                          |
| Selecting material         | Non-flowering tips from a plant with the form you want (upright or sprawling).                                                                                      |
| Pre-treatment              | <ul style="list-style-type: none"> <li>Strip leaves from the lower half.</li> <li>Light wound on firmer stems.</li> <li>Dip the base in rooting hormone.</li> </ul> |
| Rooting hormone            | Semi-hardwood strength, 3 g/L IBA ( <b>Clonex Purple</b> ).                                                                                                         |
| Propagation mix            | Standard free-draining cutting mix.                                                                                                                                 |
| Environment                | Humid frame, bright shade, bottom heat.                                                                                                                             |
| Expected strike time       | Not documented – usually quick in the genus; check from about 4 weeks.                                                                                              |
| Post-treatment / aftercare | <ul style="list-style-type: none"> <li>Standard hardening off.</li> <li>Pot on into a native mix that holds some moisture.</li> </ul>                               |

## Common problems

| Problem       | Likely cause and fix                                    |
|---------------|---------------------------------------------------------|
| Base rotting  | Mix too wet in the frame. Water less; improve drainage. |
| Cuttings wilt | Material too soft. Use hardened growth.                 |
| Leggy plants  | Tip-prune young plants.                                 |

## Notes

|               |                                                                                                                                         |
|---------------|-----------------------------------------------------------------------------------------------------------------------------------------|
| Difficulty    | <b>Easy (provisional).</b> Most myoporums strike readily; this species is untested here.                                                |
| Other methods | Seed – not documented for this species. The related <i>M. insulare</i> is fermented, then soaked in smoke water (Geraldton).            |
| Confidence    | <b>Medium.</b> No species data. ANPSA: most <i>Myoporum</i> strike readily from hardened current season's growth, the preferred method. |
| Sources       | See References: 4, 12, 62, 63                                                                                                           |

## Nursery records

Log each batch here to confirm or update the timing and difficulty above.

| Date set | Stock source / provenance | No. set | No. struck | Strike % | Notes |
|----------|---------------------------|---------|------------|----------|-------|
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |

# Myoporum insulare

Blueberry Tree / Common Boobialla

Family  
**Scrophulariaceae**

| DIFFICULTY              |  | WOOD TYPE                                                     |      |      |      | BEST MONTHS |     | CUTTING LENGTH                  |     |     | HORMONE (IBA) |     |     |
|-------------------------|--|---------------------------------------------------------------|------|------|------|-------------|-----|---------------------------------|-----|-----|---------------|-----|-----|
| Easy                    |  | Semi-hardwood                                                 |      |      |      | Feb – Apr   |     | 75–100 mm                       |     |     | 3 g/L IBA     |     |     |
|                         |  |                                                               |      |      |      |             |     |                                 |     |     |               |     |     |
|                         |  | Jan                                                           | Feb  | Mar  | Apr  | May         | Jun | Jul                             | Aug | Sep | Oct           | Nov | Dec |
| Cuttings                |  | OK                                                            | Best | Best | Best | OK          | –   | –                               | –   | –   | OK            | OK  | OK  |
| <div></div> Best window |  | <div></div> Workable – see ‘Selecting material’ in the method |      |      |      |             |     | <div></div> Avoid / low success |     |     |               |     |     |

Plant snapshot

|            |                                                                                                                                  |
|------------|----------------------------------------------------------------------------------------------------------------------------------|
| Habit      | Dense, low spreading to erect shrub or small tree, 0.25–5 m, with thick, fleshy leaves.                                          |
| Flowering  | Jul – Dec, also Jan – Feb (white, purple-spotted), followed by purple berries.                                                   |
| Habitat    | White or grey sand on coastal dunes and limestone.                                                                               |
| Local note | Recorded in Joondalup and Wanneroo. Forms range from prostrate to tree-like – take cuttings from a plant with the form you want. |

Propagation options

| Method   | Suitability | Notes                                                           | Details in          |
|----------|-------------|-----------------------------------------------------------------|---------------------|
| Cuttings | Main method | Firm young growth roots readily.                                | This guide          |
| Seed     | Good        | Ferment berries, smoke water; 15–30 days. Seedling form varies. | → Seed Sowing Guide |
| Division | Not used    | Woody shrub.                                                    | –                   |

Propagation method

| Item                       | Details                                                                                                                                                                                  |
|----------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Best time to take cuttings | Feb – Apr – warm, and after the main flowering. Oct – Jan and May are workable. Avoid Jun – Sep.                                                                                         |
| Wood type                  | Semi-hardwood – firm young growth.                                                                                                                                                       |
| Cutting size               | 75–100 mm.                                                                                                                                                                               |
| Selecting material         | Firm, <b>non-flowering</b> young shoots from a stock plant with the growth form you need (prostrate for groundcover, erect for screening).                                               |
| Pre-treatment              | <ul style="list-style-type: none"><li>Strip leaves from the lower half.</li><li>Reduce the remaining thick leaves by about one-third.</li><li>Dip the base in rooting hormone.</li></ul> |
| Rooting hormone            | 3 g/L IBA ( <b>Clonex Purple</b> ). Strikes readily without it, but hormone gives faster, more even rooting.                                                                             |
| Propagation mix            | Standard sand/coir cutting mix.                                                                                                                                                          |
| Environment                | Standard humid frame with bottom heat. Hairless leaves tolerate normal misting.                                                                                                          |
| Expected strike time       | About <b>4–8 weeks</b> in warm weather.                                                                                                                                                  |
| Post-treatment / aftercare | <ul style="list-style-type: none"><li>Standard hardening off.</li><li>Fast-growing – pot on promptly before roots circle.</li><li>Tip-prune for a bushier plant.</li></ul>               |

## Common problems

| Problem                       | Likely cause and fix                                                |
|-------------------------------|---------------------------------------------------------------------|
| Leaf drop in the frame        | Too dark or too wet. Give brighter shade and water less.            |
| Roots circling in tubes       | Pot on sooner – it grows quickly.                                   |
| Plants not the expected shape | Seed-grown or mixed stock. Propagate from a known-form stock plant. |

## Notes

|               |                                                                                                                                                                                                |
|---------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Difficulty    | <b>Easy.</b> Firm young growth roots readily; even branchlets placed straight into soil can strike.                                                                                            |
| Other methods | Seed – the Geraldton community nursery ferments ripe (darkest purple) berries for 4–5 days, washes off the pulp and soaks in smoke water; germination took 15–30 days. Seedlings vary in form. |
| Confidence    | <b>High.</b> Published sources agree it strikes readily from firm young growth.                                                                                                                |
| Sources       | See References: 1, 4, 12, 13, 23                                                                                                                                                               |

## Nursery records

Log each batch here to confirm or update the timing and difficulty above.

| Date set | Stock source / provenance | No. set | No. struck | Strike % | Notes |
|----------|---------------------------|---------|------------|----------|-------|
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |

# Olearia axillaris

Coastal Daisybush

Family  
Asteraceae

|                      |                            |                          |                             |                                |
|----------------------|----------------------------|--------------------------|-----------------------------|--------------------------------|
| DIFFICULTY<br>Easy * | WOOD TYPE<br>Semi-hardwood | BEST MONTHS<br>Aug – Oct | CUTTING LENGTH<br>75–100 mm | HORMONE (IBA)<br>1.5–3 g/L IBA |
|----------------------|----------------------------|--------------------------|-----------------------------|--------------------------------|

|                                                              | Jan | Feb | Mar | Apr | May | Jun | Jul | Aug  | Sep  | Oct  | Nov | Dec |
|--------------------------------------------------------------|-----|-----|-----|-----|-----|-----|-----|------|------|------|-----|-----|
| Cuttings                                                     | –   | –   | OK  | OK  | –   | –   | OK  | Best | Best | Best | OK  | –   |
| <div>Best window</div>                                       |     |     |     |     |     |     |     |      |      |      |     |     |
| <div>Workable – see ‘Selecting material’ in the method</div> |     |     |     |     |     |     |     |      |      |      |     |     |
| <div>Avoid / low success</div>                               |     |     |     |     |     |     |     |      |      |      |     |     |

Plant snapshot

|            |                                                                                      |
|------------|--------------------------------------------------------------------------------------|
| Habit      | Erect, much-branched grey shrub, 0.2–3 m.                                            |
| Flowering  | Jan – Jul or Nov – Dec (small white to yellow daisies).                              |
| Habitat    | White, grey or red sand and loam; coastal limestone, sand dunes and rocky hillsides. |
| Local note | Recorded in Joondalup and Wanneroo.                                                  |

Propagation options

| Method   | Suitability | Notes                                                  | Details in                          |
|----------|-------------|--------------------------------------------------------|-------------------------------------|
| Cuttings | Good        | Perennial daisies strike quickly from vigorous growth. | <a href="#">This guide</a>          |
| Seed     | Main method | Fresh seed, aired 3 months; surface-sow.               | → <a href="#">Seed Sowing Guide</a> |
| Division | Not used    | Woody shrub.                                           | –                                   |

Propagation method

| Item                       | Details                                                                                                                                                               |
|----------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Best time to take cuttings | <b>Aug – Oct</b> , on vigorous spring growth between flowering periods. Mar – Apr, Jul and Nov are workable with non-flowering shoots. Avoid Dec – Feb and May – Jun. |
| Wood type                  | <b>Semi-hardwood</b> – vigorous current growth that has begun to firm.                                                                                                |
| Cutting size               | 75–100 mm.                                                                                                                                                            |
| Selecting material         | Vigorous, healthy, non-flowering shoots (ANPSA).                                                                                                                      |
| Pre-treatment              | <ul style="list-style-type: none"><li>Strip leaves from the lower half.</li><li>No wounding needed.</li><li>Dip the base in rooting hormone.</li></ul>                |
| Rooting hormone            | 1.5 g/L IBA ( <b>Clonex Green</b> ) for softer tips; 3 g/L for firmer material.                                                                                       |
| Propagation mix            | Free-draining: about 3 parts sand to 1 part peat or coir (ANPSA daisy mix).                                                                                           |
| Environment                | Humid frame, bright shade, bottom heat. The grey, hairy leaves rot if kept wet – mist lightly.                                                                        |
| Expected strike time       | Usually quick – check from about 3–4 weeks.                                                                                                                           |
| Post-treatment / aftercare | <ul style="list-style-type: none"><li>Standard hardening off.</li><li>Grow on in full sun; tip-prune for a compact plant.</li></ul>                                   |

## Common problems

| Problem                       | Likely cause and fix                                            |
|-------------------------------|-----------------------------------------------------------------|
| Leaves go grey-brown and soft | Too wet or humid. Mist less, improve airflow.                   |
| Cuttings flower               | Taken from flowering shoots. Use vigorous non-flowering growth. |
| Leggy plants                  | Too much shade. Move to full sun after hardening off.           |

## Notes

|            |                                                                                                               |
|------------|---------------------------------------------------------------------------------------------------------------|
| Difficulty | <b>Easy (provisional).</b> Perennial daisies usually strike quickly and easily.                               |
| Confidence | <b>Medium.</b> ANPSA: stem cuttings of most perennial Australian daisies strike quickly from vigorous growth. |
| Sources    | See References: 4, 13, 35                                                                                     |

## Nursery records

Log each batch here to confirm or update the timing and difficulty above.

| Date set | Stock source / provenance | No. set | No. struck | Strike % | Notes |
|----------|---------------------------|---------|------------|----------|-------|
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |

# Phyllanthus calycinus

False Boronia

Current WA Herbarium name: *Lysiandra calycina*

Family  
Phyllanthaceae

| DIFFICULTY | WOOD TYPE     | BEST MONTHS | CUTTING LENGTH | HORMONE (IBA) |
|------------|---------------|-------------|----------------|---------------|
| Moderate * | Semi-hardwood | Feb - Apr   | 50-75 mm       | 3 g/L IBA     |

|          | Jan | Feb  | Mar  | Apr  | May | Jun | Jul | Aug | Sep | Oct | Nov | Dec |
|----------|-----|------|------|------|-----|-----|-----|-----|-----|-----|-----|-----|
| Cuttings | OK  | Best | Best | Best | OK  | -   | -   | -   | -   | -   | -   | -   |

Best window

Workable - see 'Selecting material' in the method

Avoid / low success

Plant snapshot

|            |                                                                                                                                                                                           |
|------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Habit      | Small erect shrub, 0.2-1.2 m, with smooth branchlets.                                                                                                                                     |
| Flowering  | Jun - Dec or Jan (white-cream to pink). Sepals enlarge to enclose the fruit.                                                                                                              |
| Habitat    | Usually sandy soils, including coastal alkaline sands.                                                                                                                                    |
| Name note  | FloraBase now lists this plant as <i>Lysiandra calycina</i> ; many nurseries still sell it as <i>Phyllanthus calycinus</i> . Kept under <i>Phyllanthus</i> here to match your plant list. |
| Local note | Recorded in Joondalup and Wanneroo.                                                                                                                                                       |

Propagation options

| Method   | Suitability | Notes                                               | Details in |
|----------|-------------|-----------------------------------------------------|------------|
| Cuttings | Trial       | No published cutting data; trial protocol.          | This guide |
| Seed     | Possible    | Fresh seed is recommended; little published detail. | -          |
| Division | Not used    | Woody shrub.                                        | -          |

Propagation method

| Item                       | Details                                                                                                                                                         |
|----------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Best time to take cuttings | Feb - Apr - warm, and after the long flowering season ends. Jan and May are workable. Avoid Jun - Dec (flowering, and cool in winter).                          |
| Wood type                  | Semi-hardwood.                                                                                                                                                  |
| Cutting size               | 50-75 mm.                                                                                                                                                       |
| Selecting material         | Non-flowering shoots from healthy plants.                                                                                                                       |
| Pre-treatment              | <ul style="list-style-type: none"><li>Strip leaves from the lower half.</li><li>Light wound on firmer stems.</li><li>Dip the base in rooting hormone.</li></ul> |
| Rooting hormone            | 3 g/L IBA (Clonex Purple); trial 1.5 g/L (Clonex Green) on softer tips.                                                                                         |
| Propagation mix            | Free-draining sand-based mix.                                                                                                                                   |
| Environment                | Standard humid frame, bottom heat, bright shade.                                                                                                                |
| Expected strike time       | Not documented - start checking from about 6 weeks and record the result.                                                                                       |
| Post-treatment / aftercare | <ul style="list-style-type: none"><li>Standard hardening off.</li><li>Pot on into a free-draining native mix.</li></ul>                                         |

## Common problems

| Problem                  | Likely cause and fix                                                                 |
|--------------------------|--------------------------------------------------------------------------------------|
| No roots after 10+ weeks | Try a different hormone strength or timing on the next batch; record what you tried. |
| Rotting                  | Mix too wet. Water less often; improve drainage.                                     |
| Cuttings flowering       | Use non-flowering shoots only.                                                       |

## Notes

|               |                                                                                                                 |
|---------------|-----------------------------------------------------------------------------------------------------------------|
| Difficulty    | <b>Moderate (provisional).</b> No cutting data found for this species; the rating is a cautious starting point. |
| Other methods | Seed – the method most guides mention; use fresh seed. Few published details on pre-treatment.                  |
| Confidence    | <b>Low.</b> No published cutting data found. Treat as a trial protocol and record results.                      |
| Sources       | See References: 1, 4, 29, 30                                                                                    |

## Nursery records

Log each batch here to confirm or update the timing and difficulty above.

| Date set | Stock source / provenance | No. set | No. struck | Strike % | Notes |
|----------|---------------------------|---------|------------|----------|-------|
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |

# Pimelea calcicola

Coastal Banjine

Family  
Thymelaeaceae

|                                 |                                   |                                 |                                   |                                       |
|---------------------------------|-----------------------------------|---------------------------------|-----------------------------------|---------------------------------------|
| DIFFICULTY<br><b>Moderate</b> * | WOOD TYPE<br><b>Semi-hardwood</b> | BEST MONTHS<br><b>Dec - Mar</b> | CUTTING LENGTH<br><b>50-75 mm</b> | HORMONE (IBA)<br><b>1.5-3 g/L IBA</b> |
|---------------------------------|-----------------------------------|---------------------------------|-----------------------------------|---------------------------------------|

|          | Jan         | Feb                                               | Mar  | Apr | May | Jun | Jul | Aug | Sep | Oct | Nov | Dec                 |
|----------|-------------|---------------------------------------------------|------|-----|-----|-----|-----|-----|-----|-----|-----|---------------------|
| Cuttings | Best        | Best                                              | Best | OK  | -   | -   | -   | -   | -   | -   | OK  | Best                |
|          | Best window | Workable - see 'Selecting material' in the method |      |     |     |     |     |     |     |     |     | Avoid / low success |

Plant snapshot

|            |                                                                                                                                                         |
|------------|---------------------------------------------------------------------------------------------------------------------------------------------------------|
| Habit      | Erect to spreading shrub, 0.2-1 m.                                                                                                                      |
| Flowering  | Sep - Nov (pink heads).                                                                                                                                 |
| Habitat    | Sand on coastal limestone ridges.                                                                                                                       |
| Local note | Recorded in Joondalup and Wanneroo. <b>Priority Three</b> conservation species - take cuttings only from nursery stock or under the right DBCA licence. |

Propagation options

| Method   | Suitability | Notes                                                                    | Details in |
|----------|-------------|--------------------------------------------------------------------------|------------|
| Cuttings | Main method | Firm growth strikes; snip leaves so the bark doesn't tear (provisional). | This guide |
| Seed     | Difficult   | Pimeleas are hard from seed; smoke helps some.                           | -          |
| Division | Not used    | -                                                                        | -          |

Propagation method

| Item                       | Details                                                                                                                                                                                           |
|----------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Best time to take cuttings | <b>Dec - Mar</b> , after flowering, once new growth has firmed. Nov and Apr are workable. Avoid May - Oct.                                                                                        |
| Wood type                  | <b>Semi-hardwood</b> - firm current season's growth.                                                                                                                                              |
| Cutting size               | 50-75 mm.                                                                                                                                                                                         |
| Selecting material         | Healthy, non-flowering side shoots.                                                                                                                                                               |
| Pre-treatment              | <ul style="list-style-type: none"><li>• <b>Snip leaves off with fine scissors</b> - pulling them tears the thin bark.</li><li>• No wounding.</li><li>• Dip the base in rooting hormone.</li></ul> |
| Rooting hormone            | 1.5 g/L IBA ( <b>Clonex Green</b> ) for softer tips; 3 g/L for firmer stems.                                                                                                                      |
| Propagation mix            | Very free-draining cutting mix.                                                                                                                                                                   |
| Environment                | Humid frame, bright shade, bottom heat. Moist, not wet.                                                                                                                                           |
| Expected strike time       | Not documented - check from about 4-6 weeks and record the result.                                                                                                                                |
| Post-treatment / aftercare | <ul style="list-style-type: none"><li>• Standard hardening off.</li><li>• Grow on in full sun in a free-draining mix.</li></ul>                                                                   |

## Common problems

| Problem               | Likely cause and fix                               |
|-----------------------|----------------------------------------------------|
| Bark torn at the base | Leaves were pulled off. Snip them with scissors.   |
| Base rotting          | Too wet. Coarser mix; water less.                  |
| Slow to root          | Use bottom heat; take cuttings in the warm months. |

## Notes

|               |                                                                                                                                           |
|---------------|-------------------------------------------------------------------------------------------------------------------------------------------|
| Difficulty    | <b>Moderate (provisional).</b> Pimeleas strike from cuttings, but the bark damages easily; untested for this species.                     |
| Other methods | Seed – pimeleas are difficult from seed; smoke water helps some species. The only Alcoa test of this species gave 0%.                     |
| Confidence    | <b>Medium.</b> No species data. ANPSA: <i>Pimelea</i> cuttings are successful and firm growth usually strikes, but the bark tears easily. |
| Sources       | See References: 4, 32, 66, 67                                                                                                             |

## Nursery records

Log each batch here to confirm or update the timing and difficulty above.

| Date set | Stock source / provenance | No. set | No. struck | Strike % | Notes |
|----------|---------------------------|---------|------------|----------|-------|
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |

# Pimelea sulphurea

Yellow Banjine

Family  
Thymelaeaceae

|                          |                            |                          |                            |                                |
|--------------------------|----------------------------|--------------------------|----------------------------|--------------------------------|
| DIFFICULTY<br>Moderate * | WOOD TYPE<br>Semi-hardwood | BEST MONTHS<br>Dec - Mar | CUTTING LENGTH<br>50-75 mm | HORMONE (IBA)<br>1.5-3 g/L IBA |
|--------------------------|----------------------------|--------------------------|----------------------------|--------------------------------|

|          | Jan         | Feb                                               | Mar  | Apr | May | Jun | Jul | Aug | Sep | Oct | Nov | Dec                 |
|----------|-------------|---------------------------------------------------|------|-----|-----|-----|-----|-----|-----|-----|-----|---------------------|
| Cuttings | Best        | Best                                              | Best | OK  | -   | -   | -   | -   | -   | -   | OK  | Best                |
|          | Best window | Workable - see 'Selecting material' in the method |      |     |     |     |     |     |     |     |     | Avoid / low success |

Plant snapshot

|            |                                                                                       |
|------------|---------------------------------------------------------------------------------------|
| Habit      | Erect, spindly or open shrub, 0.15-0.7 m.                                             |
| Flowering  | Jul - Nov (yellow heads).                                                             |
| Habitat    | White, grey or yellow sand, sometimes with gravel or laterite; sandplains and ridges. |
| Local note | Recorded in Joondalup and Wanneroo.                                                   |

Propagation options

| Method   | Suitability | Notes                                                | Details in |
|----------|-------------|------------------------------------------------------|------------|
| Cuttings | Main method | Firm growth; snip leaves - bark tears (provisional). | This guide |
| Seed     | Difficult   | Winter sowing with smoke water, and patience.        | -          |
| Division | Not used    | Shrub - not divided.                                 | -          |

Propagation method

| Item                       | Details                                                                                                                                                                              |
|----------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Best time to take cuttings | Dec - Mar, after flowering, once new growth has firmed. Nov and Apr are workable. Avoid May - Oct.                                                                                   |
| Wood type                  | Semi-hardwood - firm current season's growth.                                                                                                                                        |
| Cutting size               | 50-75 mm. Stems are thin - take extra.                                                                                                                                               |
| Selecting material         | Healthy, non-flowering side shoots.                                                                                                                                                  |
| Pre-treatment              | <ul style="list-style-type: none"><li>Snip leaves off with fine scissors - pulling them tears the thin bark.</li><li>No wounding.</li><li>Dip the base in rooting hormone.</li></ul> |
| Rooting hormone            | 1.5 g/L IBA (Clonex Green) for softer tips; 3 g/L for firmer stems.                                                                                                                  |
| Propagation mix            | Very free-draining cutting mix.                                                                                                                                                      |
| Environment                | Humid frame, bright shade, bottom heat. Moist, not wet.                                                                                                                              |
| Expected strike time       | Not documented - check from about 4-6 weeks and record the result.                                                                                                                   |
| Post-treatment / aftercare | <ul style="list-style-type: none"><li>Standard hardening off.</li><li>Grow on in full sun in a free-draining mix.</li></ul>                                                          |

## Common problems

| Problem               | Likely cause and fix                               |
|-----------------------|----------------------------------------------------|
| Bark torn at the base | Leaves were pulled off. Snip them with scissors.   |
| Base rotting          | Too wet. Coarser mix; water less.                  |
| Slow to root          | Use bottom heat; take cuttings in the warm months. |

## Notes

|               |                                                                                                                                              |
|---------------|----------------------------------------------------------------------------------------------------------------------------------------------|
| Difficulty    | <b>Moderate (provisional).</b> Pimeleas strike from cuttings, but the bark damages easily; untested for this species.                        |
| Other methods | Seed – pimeleas are difficult from seed. Sowing in winter with smoke water, and patience, is the best bet. Not in the Alcoa nursery records. |
| Confidence    | <b>Low.</b> No species data. Other WA pimeleas strike from firm growth, but the bark tears easily.                                           |
| Sources       | See References: 4, 13, 66, 67, 73                                                                                                            |

## Nursery records

Log each batch here to confirm or update the timing and difficulty above.

| Date set | Stock source / provenance | No. set | No. struck | Strike % | Notes |
|----------|---------------------------|---------|------------|----------|-------|
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |

# Pithocarpa cordata

Tangle Daisy

Previously *Ozothamnus cordatus* / *Helichrysum cordatum*Family  
Asteraceae

| DIFFICULTY      | WOOD TYPE          | BEST MONTHS    | CUTTING LENGTH | HORMONE (IBA) |
|-----------------|--------------------|----------------|----------------|---------------|
| Easy-Moderate * | Semi-hardwood tips | Sep – Nov; May | 50–75 mm       | 1.5–3 g/L IBA |

|          | Jan         | Feb                                               | Mar | Apr | May  | Jun | Jul | Aug | Sep  | Oct  | Nov  | Dec |
|----------|-------------|---------------------------------------------------|-----|-----|------|-----|-----|-----|------|------|------|-----|
| Cuttings | –           | –                                                 | –   | OK  | Best | OK  | –   | OK  | Best | Best | Best | OK  |
|          | Best window | Workable – see 'Selecting material' in the method |     |     |      |     |     |     |      |      |      |     |

## Plant snapshot

|            |                                                                                                                  |
|------------|------------------------------------------------------------------------------------------------------------------|
| Habit      | Scrambling, tangled shrub, 1–2 m, with small heart-shaped leaves.                                                |
| Flowering  | Jan – Apr (white to yellow).                                                                                     |
| Habitat    | Coastal limestone and dunes.                                                                                     |
| Local note | Recorded in the Wanneroo LGA. Older books list it as <i>Ozothamnus cordatus</i> or <i>Helichrysum cordatum</i> . |

## Propagation options

| Method   | Suitability | Notes                                                    | Details in                 |
|----------|-------------|----------------------------------------------------------|----------------------------|
| Cuttings | Good        | Perennial daisy – expected to strike well (provisional). | <a href="#">This guide</a> |
| Seed     | Possible    | Not documented; daisy seed often needs light.            | –                          |
| Division | Not used    | –                                                        | –                          |

## Propagation method

| Item                       | Details                                                                                                                                                              |
|----------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Best time to take cuttings | <b>Sep – Nov</b> on vigorous spring growth, and <b>May</b> after flowering. Apr, Jun, Aug and Dec are workable. Avoid Jan – Mar (flowering and heat) and Jul (cold). |
| Wood type                  | <b>Semi-hardwood tips</b> – vigorous growth that has started to firm.                                                                                                |
| Cutting size               | 50–75 mm.                                                                                                                                                            |
| Selecting material         | Vigorous, healthy, non-flowering shoots.                                                                                                                             |
| Pre-treatment              | <ul style="list-style-type: none"> <li>Strip leaves from the lower half.</li> <li>No wounding.</li> <li>Dip the base in rooting hormone.</li> </ul>                  |
| Rooting hormone            | 1.5 g/L IBA ( <b>Clonex Green</b> ); 3 g/L for firmer stems.                                                                                                         |
| Propagation mix            | Free-draining: about 3 parts sand to 1 part peat or coir.                                                                                                            |
| Environment                | Humid frame, bright shade, bottom heat. Moist, not wet.                                                                                                              |
| Expected strike time       | Not documented – check from about 3–4 weeks and record the result.                                                                                                   |
| Post-treatment / aftercare | <ul style="list-style-type: none"> <li>Standard hardening off.</li> <li>Grow on in full sun; tip-prune to control the scrambling habit.</li> </ul>                   |

## Common problems

| Problem                  | Likely cause and fix                                                               |
|--------------------------|------------------------------------------------------------------------------------|
| Cuttings rot             | Too wet or humid. Mist less, improve airflow.                                      |
| No roots after 8 weeks   | Try firmer tips, a stronger hormone, or a different season, and record the result. |
| Tangled, sprawling tubes | Tip-prune and space tubes.                                                         |

## Notes

|               |                                                                                                                                                         |
|---------------|---------------------------------------------------------------------------------------------------------------------------------------------------------|
| Difficulty    | <b>Easy-Moderate (provisional).</b> Most perennial daisies strike quickly from vigorous growth, but this species is untested. Start with a trial batch. |
| Other methods | Seed – not documented. Daisy-family seed often needs light, so surface-sow if trialling.                                                                |
| Confidence    | <b>Low.</b> No published propagation data. Based on perennial Australian daisies, which usually strike quickly – trial protocol.                        |
| Sources       | See References: 4, 35                                                                                                                                   |

## Nursery records

Log each batch here to confirm or update the timing and difficulty above.

| Date set | Stock source / provenance | No. set | No. struck | Strike % | Notes |
|----------|---------------------------|---------|------------|----------|-------|
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |

# Rhagodia baccata

Berry Saltbush

Also listed as *Chenopodium baccatum*

Family  
**Amaranthaceae**  
(chenopods)

| DIFFICULTY | WOOD TYPE          | BEST MONTHS | CUTTING LENGTH | HORMONE (IBA) |
|------------|--------------------|-------------|----------------|---------------|
| Moderate * | Semi-hardwood tips | Sep – Nov   | 75–100 mm      | 3 g/L IBA     |

|          | Jan         | Feb                                               | Mar | Apr | May | Jun | Jul | Aug | Sep  | Oct  | Nov  | Dec |
|----------|-------------|---------------------------------------------------|-----|-----|-----|-----|-----|-----|------|------|------|-----|
| Cuttings | OK          | OK                                                | OK  | –   | –   | –   | –   | OK  | Best | Best | Best | OK  |
|          | Best window | Workable – see ‘Selecting material’ in the method |     |     |     |     |     |     |      |      |      |     |

Plant snapshot

|            |                                                                                                      |
|------------|------------------------------------------------------------------------------------------------------|
| Habit      | Spreading shrub, 0.3–2 m, with grey-green leaves.                                                    |
| Flowering  | Variable – Feb, Apr – May or Oct – Dec (tiny), followed by red berries eaten by birds and bobtails.  |
| Habitat    | Sand dunes and coastal rock.                                                                         |
| Local note | Recorded in Joondalup and Wanneroo. Salt-tolerant; a breeding plant for the saltbush blue butterfly. |

Propagation options

| Method   | Suitability | Notes                                                | Details in                 |
|----------|-------------|------------------------------------------------------|----------------------------|
| Cuttings | Good        | Chenopod shrubs generally strike well (provisional). | <a href="#">This guide</a> |
| Seed     | Main method | Ferment berries, then smoke water.                   | → Seed Sowing Guide        |
| Division | Not used    | Woody shrub.                                         | –                          |

Propagation method

| Item                       | Details                                                                                                                                                             |
|----------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Best time to take cuttings | Sep – Nov, in active spring growth. Aug and Dec – Mar are workable with non-flowering shoots. Avoid Apr – Jul.                                                      |
| Wood type                  | Semi-hardwood tips of current season's growth.                                                                                                                      |
| Cutting size               | 75–100 mm.                                                                                                                                                          |
| Selecting material         | Vigorous tips without flowers or berries.                                                                                                                           |
| Pre-treatment              | <ul style="list-style-type: none"> <li>Strip leaves from the lower half.</li> <li>Light wound on firmer stems.</li> <li>Dip the base in rooting hormone.</li> </ul> |
| Rooting hormone            | Semi-hardwood strength, 3 g/L IBA ( <b>Clonex Purple</b> ).                                                                                                         |
| Propagation mix            | Free-draining mix of sand and potting mix.                                                                                                                          |
| Environment                | Bright shade with <b>good ventilation and little or no mist</b> (misting encourages leaf disease on saltbush cuttings). Bottom heat helps.                          |
| Expected strike time       | Not documented – check from about 4–6 weeks and record the result.                                                                                                  |
| Post-treatment / aftercare | <ul style="list-style-type: none"> <li>Harden off, then grow on in full sun.</li> <li>Water sparingly once established.</li> </ul>                                  |

## Common problems

| Problem                     | Likely cause and fix                                           |
|-----------------------------|----------------------------------------------------------------|
| Leaf spots, leaves dropping | Too humid. Stop misting and improve airflow.                   |
| Base rotting                | Mix too wet. Use a sandier mix, water less.                    |
| No roots                    | Try firmer material or the seed method, and record the result. |

## Notes

|            |                                                                                                                           |
|------------|---------------------------------------------------------------------------------------------------------------------------|
| Difficulty | <b>Moderate (provisional).</b> Untested here; related saltbushes strike from tip cuttings. Seed is the documented method. |
| Confidence | <b>Low.</b> No published cutting data found. Method follows related saltbushes – trial protocol.                          |
| Sources    | See References: 4, 20, 36, 37                                                                                             |

## Nursery records

Log each batch here to confirm or update the timing and difficulty above.

| Date set | Stock source / provenance | No. set | No. struck | Strike % | Notes |
|----------|---------------------------|---------|------------|----------|-------|
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |

# Scaevola canescens

Grey Scaevola

Family  
Goodeniaceae

| DIFFICULTY      | WOOD TYPE                | BEST MONTHS | CUTTING LENGTH | HORMONE (IBA) |
|-----------------|--------------------------|-------------|----------------|---------------|
| Easy-Moderate * | Softwood – semi-hardwood | Nov – Jan   | 50–80 mm       | 1.5–3 g/L IBA |

|          | Jan  | Feb | Mar | Apr | May | Jun | Jul | Aug | Sep | Oct | Nov  | Dec  |
|----------|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|------|------|
| Cuttings | Best | OK  | OK  | OK  | OK  | –   | –   | –   | –   | OK  | Best | Best |

|             |                                                   |                     |
|-------------|---------------------------------------------------|---------------------|
| Best window | Workable – see 'Selecting material' in the method | Avoid / low success |
|-------------|---------------------------------------------------|---------------------|

## Plant snapshot

|            |                                                                                        |
|------------|----------------------------------------------------------------------------------------|
| Habit      | Prostrate to ascending shrub, 0.1–0.5 m. Densely hairy grey leaves; stems are brittle. |
| Flowering  | Mar – Oct (white to white-purple), in spikes in the leaf axils.                        |
| Habitat    | Sand, often with lateritic gravel or limestone.                                        |
| Local note | Recorded in the northern Perth area (Wanneroo and Stirling LGAs).                      |

## Propagation options

| Method   | Suitability | Notes                                               | Details in                 |
|----------|-------------|-----------------------------------------------------|----------------------------|
| Cuttings | Main method | Soft to semi-hardwood tips; keep hairy foliage dry. | <a href="#">This guide</a> |
| Seed     | Difficult   | Scaevola seed is generally difficult.               | –                          |
| Division | Possible    | Only if stems have rooted or suckered.              | –                          |

## Propagation method

| Item                       | Details                                                                                                                                                                                                           |
|----------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Best time to take cuttings | Nov – Jan, on new growth after flowering ends, while it is warm. Oct and Feb – May are workable if you choose shoots without flower spikes. Avoid Jun – Sep (cool, peak flowering).                               |
| Wood type                  | Softwood to semi-hardwood tip cuttings.                                                                                                                                                                           |
| Cutting size               | 50–80 mm tips.                                                                                                                                                                                                    |
| Selecting material         | Vigorous <b>non-flowering</b> tips. Flowers form in the leaf axils along the stem, so check the whole cutting and pinch off any spikes.                                                                           |
| Pre-treatment              | <ul style="list-style-type: none"> <li>Handle gently – stems snap easily. Carry in a rigid tray or box.</li> <li>Strip lower leaves by cutting, not tearing.</li> <li>Dip the base in rooting hormone.</li> </ul> |
| Rooting hormone            | Softwood strength (1.5 g/L IBA, e.g. <b>Clonex Green</b> ) for soft tips; semi-hardwood strength (3 g/L, <b>Clonex Purple</b> ) for firmer material.                                                              |
| Propagation mix            | Extra free-draining: standard sand/coir mix with extra coarse sand or perlite.                                                                                                                                    |
| Environment                | <b>Keep foliage as dry as possible.</b> Dense hairy leaves rot under heavy mist. Water the mix rather than the leaves, give good airflow, and use bottom heat (root zone about 21–24°C).                          |
| Expected strike time       | About <b>4–8 weeks</b> in warm weather.                                                                                                                                                                           |
| Post-treatment / aftercare | <ul style="list-style-type: none"> <li>Standard hardening off.</li> <li>Pot on into a free-draining mix; keep water off the foliage.</li> <li>Remove any leaves showing grey mould promptly.</li> </ul>           |

## Common problems

| Problem                         | Likely cause and fix                                                         |
|---------------------------------|------------------------------------------------------------------------------|
| Grey fuzz or blackened leaves   | Too humid/wet. Cut back misting, remove affected material, improve airflow.  |
| Snapped stems                   | Brittle stems – transport in rigid containers and use a dibber when setting. |
| Cuttings flowering, not rooting | Take shoots without axillary flower spikes.                                  |

## Notes

|               |                                                                                                                                        |
|---------------|----------------------------------------------------------------------------------------------------------------------------------------|
| Difficulty    | <b>Easy-Moderate (provisional).</b> <i>Scaevola</i> cuttings generally strike very readily; the hairy foliage makes rot the main risk. |
| Other methods | Seed – <i>Scaevola</i> seed is generally difficult. Division – possible if a plant has suckered or rooted along its stems.             |
| Confidence    | <b>Medium.</b> Little species-specific data published. Based on the genus (strikes readily) plus this species' hairy foliage.          |
| Sources       | See References: 1, 4, 5, 6, 7, 12, 13, 14, 16                                                                                          |

## Nursery records

Log each batch here to confirm or update the timing and difficulty above.

| Date set | Stock source / provenance | No. set | No. struck | Strike % | Notes |
|----------|---------------------------|---------|------------|----------|-------|
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |

# Scaevola crassifolia

Thick-leaved Fan-flower

Family  
Goodeniaceae

| DIFFICULTY              |  | WOOD TYPE                                                     |     |      | BEST MONTHS |      | CUTTING LENGTH                  |     |     | HORMONE (IBA) |     |     |     |
|-------------------------|--|---------------------------------------------------------------|-----|------|-------------|------|---------------------------------|-----|-----|---------------|-----|-----|-----|
| Easy                    |  | Semi-hardwood                                                 |     |      | Mar – May   |      | 75–100 mm                       |     |     | 3 g/L IBA     |     |     |     |
|                         |  |                                                               |     |      |             |      |                                 |     |     |               |     |     |     |
|                         |  | Jan                                                           | Feb | Mar  | Apr         | May  | Jun                             | Jul | Aug | Sep           | Oct | Nov | Dec |
| Cuttings                |  | OK                                                            | OK  | Best | Best        | Best | –                               | –   | –   | –             | OK  | OK  | OK  |
| <div></div> Best window |  | <div></div> Workable – see ‘Selecting material’ in the method |     |      |             |      | <div></div> Avoid / low success |     |     |               |     |     |     |

Plant snapshot

|            |                                                                                                                             |
|------------|-----------------------------------------------------------------------------------------------------------------------------|
| Habit      | Erect or spreading shrub, 0.1–1.5 m and up to 3 m wide. Young growth is sticky; thick, toothed, egg-shaped to round leaves. |
| Flowering  | Jul – Dec, also Jan – Feb (blue, mauve or white fans).                                                                      |
| Habitat    | Sand on coastal dunes, plains and limestone cliffs.                                                                         |
| Local note | Recorded in Joondalup and Wanneroo. Flower colour and habit vary – cuttings keep the parent's form.                         |

Propagation options

| Method   | Suitability | Notes                                          | Details in |
|----------|-------------|------------------------------------------------|------------|
| Cuttings | Main method | Standard method; keeps flower colour and form. | This guide |
| Seed     | Possible    | Large seeds sometimes germinate naturally.     | –          |
| Division | Not used    | –                                              | –          |

Propagation method

| Item                       | Details                                                                                                                                                                                                                           |
|----------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Best time to take cuttings | Mar – May – warm, and outside the main flowering period. Oct – Feb is workable with non-flowering shoots (shade well in summer). Avoid Jun – Sep.                                                                                 |
| Wood type                  | Semi-hardwood.                                                                                                                                                                                                                    |
| Cutting size               | 75–100 mm.                                                                                                                                                                                                                        |
| Selecting material         | Non-flowering tips. Flower spikes carry leafy bracts – check closely and pinch them off.                                                                                                                                          |
| Pre-treatment              | <ul style="list-style-type: none"><li>• New growth is sticky – wipe the blade often.</li><li>• Strip lower leaves; cut the large remaining leaves back by a third to a half.</li><li>• Dip the base in rooting hormone.</li></ul> |
| Rooting hormone            | 3 g/L IBA (Clonex Purple).                                                                                                                                                                                                        |
| Propagation mix            | Free-draining sand/coir mix.                                                                                                                                                                                                      |
| Environment                | Standard humid frame with bottom heat, bright shade.                                                                                                                                                                              |
| Expected strike time       | About 4–8 weeks in warm weather.                                                                                                                                                                                                  |
| Post-treatment / aftercare | <ul style="list-style-type: none"><li>• Standard hardening off.</li><li>• Pot on into a free-draining mix.</li><li>• Tip-prune young plants for dense growth.</li></ul>                                                           |

## Common problems

| Problem                   | Likely cause and fix                                           |
|---------------------------|----------------------------------------------------------------|
| Wilting in the first week | Too much leaf area. Trim leaves harder; keep the frame closed. |
| Rotting                   | Mix too wet. Water less often; improve drainage.               |
| Cuttings flowering        | Use non-flowering tips or wait for the Mar – May window.       |

## Notes

|               |                                                                                                                            |
|---------------|----------------------------------------------------------------------------------------------------------------------------|
| Difficulty    | <b>Easy.</b> Semi-hardwood cuttings in the warmer months are the standard method.                                          |
| Other methods | Seed – large seeds are produced and sometimes germinate naturally; cuttings are preferred to keep flower colour and habit. |
| Confidence    | <b>High.</b> Cuttings are the standard method in WA nurseries (e.g. Geraldton community nursery).                          |
| Sources       | See References: 1, 4, 5, 12, 13, 26, 30                                                                                    |

## Nursery records

Log each batch here to confirm or update the timing and difficulty above.

| Date set | Stock source / provenance | No. set | No. struck | Strike % | Notes |
|----------|---------------------------|---------|------------|----------|-------|
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |

# Scaevola globulifera

No widely used common name

Family  
Goodeniaceae

|            |                          |             |                |               |
|------------|--------------------------|-------------|----------------|---------------|
| DIFFICULTY | WOOD TYPE                | BEST MONTHS | CUTTING LENGTH | HORMONE (IBA) |
| Easy *     | Softwood - semi-hardwood | Jan - Apr   | 50-80 mm       | 1.5-3 g/L IBA |

|          |      |      |      |      |     |     |     |     |     |     |     |     |
|----------|------|------|------|------|-----|-----|-----|-----|-----|-----|-----|-----|
|          | Jan  | Feb  | Mar  | Apr  | May | Jun | Jul | Aug | Sep | Oct | Nov | Dec |
| Cuttings | Best | Best | Best | Best | OK  | -   | -   | -   | -   | -   | OK  | OK  |

|             |                                                   |                     |
|-------------|---------------------------------------------------|---------------------|
| Best window | Workable - see 'Selecting material' in the method | Avoid / low success |
|-------------|---------------------------------------------------|---------------------|

Plant snapshot

|            |                                                               |
|------------|---------------------------------------------------------------|
| Habit      | Spreading, low-lying to ascending shrub, 0.2-1.2 m. Hairless. |
| Flowering  | Aug - Dec (blue to white).                                    |
| Habitat    | Sand, sandhills and limestone ridges.                         |
| Local note | Recorded in Joondalup and Wanneroo.                           |

Propagation options

| Method   | Suitability | Notes                                            | Details in                 |
|----------|-------------|--------------------------------------------------|----------------------------|
| Cuttings | Main method | Soft to semi-hardwood tips (provisional rating). | <a href="#">This guide</a> |
| Seed     | Difficult   | Scaevola seed is generally difficult.            | -                          |
| Division | Possible    | Check stock plants for rooted stems.             | -                          |

Propagation method

| Item                       | Details                                                                                                                                                            |
|----------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Best time to take cuttings | Jan - Apr - warm, and after flowering finishes in Dec. Nov - Dec and May are workable. Avoid Jun - Oct.                                                            |
| Wood type                  | Softwood to semi-hardwood tip cuttings.                                                                                                                            |
| Cutting size               | 50-80 mm.                                                                                                                                                          |
| Selecting material         | Non-flowering tips; pinch off any buds.                                                                                                                            |
| Pre-treatment              | <ul style="list-style-type: none"><li>Strip lower leaves.</li><li>Trim larger leaves by about one-third.</li><li>Dip the base in rooting hormone.</li></ul>        |
| Rooting hormone            | 1.5 g/L IBA (Clonex Green) for soft tips; 3 g/L (Clonex Purple) for firmer material.                                                                               |
| Propagation mix            | Free-draining sand/coir mix.                                                                                                                                       |
| Environment                | Humid frame, bottom heat, bright shade. Hairless leaves tolerate normal misting.                                                                                   |
| Expected strike time       | About 4-8 weeks in warm weather.                                                                                                                                   |
| Post-treatment / aftercare | <ul style="list-style-type: none"><li>Standard hardening off.</li><li>Pot on into a free-draining native mix.</li><li>Pinch tips to encourage branching.</li></ul> |

## Common problems

| Problem            | Likely cause and fix                                               |
|--------------------|--------------------------------------------------------------------|
| Wilting            | Low humidity or too much leaf. Trim leaves; keep the frame closed. |
| Rotting            | Mix too wet. Water less often.                                     |
| Cuttings flowering | Use non-flowering tips, or wait for the Jan – Apr window.          |

## Notes

|               |                                                                                                         |
|---------------|---------------------------------------------------------------------------------------------------------|
| Difficulty    | <b>Easy (provisional).</b> <i>Scaevola</i> cuttings generally strike very readily.                      |
| Other methods | Seed – <i>Scaevola</i> seed is generally difficult. Division – check stock plants for rooted stems.     |
| Confidence    | <b>Medium.</b> No species-specific data found. Based on the genus, which strikes readily from cuttings. |
| Sources       | See References: 1, 4, 5, 6, 7, 13, 14                                                                   |

## Nursery records

Log each batch here to confirm or update the timing and difficulty above.

| Date set | Stock source / provenance | No. set | No. struck | Strike % | Notes |
|----------|---------------------------|---------|------------|----------|-------|
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |

# Scaevola nitida

Shining Fanflower

Family  
Goodeniaceae

|                      |                            |                          |                             |                            |
|----------------------|----------------------------|--------------------------|-----------------------------|----------------------------|
| DIFFICULTY<br>Easy * | WOOD TYPE<br>Semi-hardwood | BEST MONTHS<br>Jan – Apr | CUTTING LENGTH<br>75–100 mm | HORMONE (IBA)<br>3 g/L IBA |
|----------------------|----------------------------|--------------------------|-----------------------------|----------------------------|

|          | Jan         | Feb                                               | Mar  | Apr  | May | Jun | Jul | Aug | Sep | Oct | Nov                 | Dec |
|----------|-------------|---------------------------------------------------|------|------|-----|-----|-----|-----|-----|-----|---------------------|-----|
| Cuttings | Best        | Best                                              | Best | Best | OK  | –   | –   | –   | –   | –   | OK                  | OK  |
|          | Best window | Workable – see ‘Selecting material’ in the method |      |      |     |     |     |     |     |     | Avoid / low success |     |

Plant snapshot

|            |                                                                                                 |
|------------|-------------------------------------------------------------------------------------------------|
| Habit      | Erect shrub, 0.3–3 m.                                                                           |
| Flowering  | Aug – Dec (blue-purple fans).                                                                   |
| Habitat    | White or grey sand and clay; coastal limestone cliffs and dunes.                                |
| Local note | Recorded in Joondalup and Wanneroo. Tolerates full coastal exposure, sand blast and salt spray. |

Propagation options

| Method   | Suitability | Notes                                     | Details in |
|----------|-------------|-------------------------------------------|------------|
| Cuttings | Main method | Scaevola cuttings strike very readily.    | This guide |
| Seed     | Possible    | Smoke; about 21–42 days (seed suppliers). | –          |
| Division | Not used    | Shrub – not divided.                      | –          |

Propagation method

| Item                       | Details                                                                                                                                                            |
|----------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Best time to take cuttings | Jan – Apr, after the main flowering, while it's warm. Nov – Dec and May are workable with non-flowering shoots. Avoid Jun – Oct.                                   |
| Wood type                  | Semi-hardwood – firm current season's growth.                                                                                                                      |
| Cutting size               | 75–100 mm.                                                                                                                                                         |
| Selecting material         | Non-flowering tips. Pinch off any buds.                                                                                                                            |
| Pre-treatment              | <ul style="list-style-type: none"><li>Strip lower leaves; cut large remaining leaves back by a third to a half.</li><li>Dip the base in rooting hormone.</li></ul> |
| Rooting hormone            | 3 g/L IBA (Clonex Purple).                                                                                                                                         |
| Propagation mix            | Free-draining sand/coir mix.                                                                                                                                       |
| Environment                | Standard humid frame with bottom heat, bright shade.                                                                                                               |
| Expected strike time       | About 4–8 weeks in warm weather (typical for the genus). Record your results.                                                                                      |
| Post-treatment / aftercare | <ul style="list-style-type: none"><li>Standard hardening off.</li><li>Pot on into a free-draining mix.</li><li>Tip-prune young plants for dense growth.</li></ul>  |

## Common problems

| Problem                   | Likely cause and fix                                           |
|---------------------------|----------------------------------------------------------------|
| Wilting in the first week | Too much leaf area. Trim leaves harder; keep the frame closed. |
| Rotting                   | Mix too wet. Water less often; improve drainage.               |
| Cuttings flowering        | Use non-flowering tips, or wait for the Jan – Apr window.      |

## Notes

|               |                                                                                                                                           |
|---------------|-------------------------------------------------------------------------------------------------------------------------------------------|
| Difficulty    | <b>Easy (provisional).</b> Scaevolas strike very readily; untested for this species.                                                      |
| Other methods | Seed – smoke treatment is recommended by seed suppliers; germination in about 21–42 days. Not in the Alcoa records.                       |
| Confidence    | <b>Medium.</b> No species-specific cutting data. Based on the genus – ANPSA says cuttings of <i>Scaevola</i> species strike very readily. |
| Sources       | See References: 1, 4, 5, 13, 51, 71, 72, 78                                                                                               |

## Nursery records

Log each batch here to confirm or update the timing and difficulty above.

| Date set | Stock source / provenance | No. set | No. struck | Strike % | Notes |
|----------|---------------------------|---------|------------|----------|-------|
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |

# Scaevola repens

No widely used common name ('repens' = creeping)

Family  
Goodeniaceae

| DIFFICULTY | WOOD TYPE                | BEST MONTHS | CUTTING LENGTH | HORMONE (IBA) |
|------------|--------------------------|-------------|----------------|---------------|
| Easy *     | Softwood – semi-hardwood | Mar – May   | 50–80 mm       | 1.5–3 g/L IBA |

|                                                   | Jan | Feb | Mar  | Apr  | May  | Jun | Jul | Aug | Sep | Oct | Nov | Dec |
|---------------------------------------------------|-----|-----|------|------|------|-----|-----|-----|-----|-----|-----|-----|
| Cuttings                                          | OK  | OK  | Best | Best | Best | –   | –   | –   | OK  | OK  | OK  | OK  |
| Best window                                       |     |     |      |      |      |     |     |     |     |     |     |     |
| Workable – see 'Selecting material' in the method |     |     |      |      |      |     |     |     |     |     |     |     |
| Avoid / low success                               |     |     |      |      |      |     |     |     |     |     |     |     |

## Plant snapshot

|            |                                                                                                                             |
|------------|-----------------------------------------------------------------------------------------------------------------------------|
| Habit      | Prostrate to ascending or erect shrub, 0.05–1 m; trailing branches to about 50 cm.                                          |
| Flowering  | Jun – Dec, sometimes Jan – Feb (white or cream, occasionally blue-purple; golden hairs on the outside of the flower).       |
| Habitat    | Sand dunes and plains on sand, gravel, laterite or limestone.                                                               |
| Local note | Recorded in the City of Joondalup. A variable species with informal subspecies – use local provenance and keep it labelled. |

## Propagation options

| Method   | Suitability | Notes                                            | Details in                 |
|----------|-------------|--------------------------------------------------|----------------------------|
| Cuttings | Main method | Soft to semi-hardwood tips (provisional rating). | <a href="#">This guide</a> |
| Seed     | Difficult   | Scaevola seed is generally difficult.            | –                          |
| Division | Possible    | Some scaevolae sucker; check for rooted pieces.  | –                          |

## Propagation method

| Item                       | Details                                                                                                                                                                                                                 |
|----------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Best time to take cuttings | Mar – May – warm autumn weather, mostly outside the main flowering period. Sep – Feb is workable using non-flowering shoots (shade well in summer). Avoid Jun – Aug.                                                    |
| Wood type                  | Softwood to semi-hardwood tip cuttings.                                                                                                                                                                                 |
| Cutting size               | 50–80 mm tips from the trailing stems.                                                                                                                                                                                  |
| Selecting material         | Non-flowering tips. Flower spikes sit in the leaf axils – pinch off any you find.                                                                                                                                       |
| Pre-treatment              | <ul style="list-style-type: none"> <li>Strip lower leaves.</li> <li>Leaves can be long (up to 90 mm) – trim remaining leaves by about one-third to cut water loss.</li> <li>Dip the base in rooting hormone.</li> </ul> |
| Rooting hormone            | Softwood strength (1.5 g/L IBA, <b>Clonex Green</b> ) for soft tips; 3 g/L ( <b>Clonex Purple</b> ) for firmer material.                                                                                                |
| Propagation mix            | Free-draining sand/coir mix.                                                                                                                                                                                            |
| Environment                | Humid frame with bottom heat, bright shade. Keep the mix moist but on the drier side rather than wet.                                                                                                                   |
| Expected strike time       | About 4–8 weeks in warm weather.                                                                                                                                                                                        |
| Post-treatment / aftercare | <ul style="list-style-type: none"> <li>Standard hardening off.</li> <li>Pot on into a free-draining native mix.</li> <li>Pinch tips once growing to encourage branching.</li> </ul>                                     |

## Common problems

| Problem                         | Likely cause and fix                                                           |
|---------------------------------|--------------------------------------------------------------------------------|
| Wilting in the first week       | Too much leaf area or low humidity. Trim leaves harder; keep the frame closed. |
| Rotting                         | Mix too wet. Water less often; improve drainage.                               |
| Cuttings flowering, not rooting | Take non-flowering tips, or wait for the Mar – May window.                     |

## Notes

|               |                                                                                                                                    |
|---------------|------------------------------------------------------------------------------------------------------------------------------------|
| Difficulty    | <b>Easy (provisional).</b> <i>Scaevola</i> cuttings generally strike very readily.                                                 |
| Other methods | Seed – <i>Scaevola</i> seed is generally difficult. Division – some <i>scaevolas</i> sucker; check stock plants for rooted pieces. |
| Confidence    | <b>Medium.</b> Little species-specific data published. Based on the genus, which strikes readily from cuttings.                    |
| Sources       | See References: 1, 4, 5, 6, 7, 12, 13, 14, 16                                                                                      |

## Nursery records

Log each batch here to confirm or update the timing and difficulty above.

| Date set | Stock source / provenance | No. set | No. struck | Strike % | Notes |
|----------|---------------------------|---------|------------|----------|-------|
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |

# Scaevola thesioides

No widely used common name

Family  
Goodeniaceae

|                 |                          |             |                |               |
|-----------------|--------------------------|-------------|----------------|---------------|
| DIFFICULTY      | WOOD TYPE                | BEST MONTHS | CUTTING LENGTH | HORMONE (IBA) |
| Easy-Moderate * | Softwood - semi-hardwood | Apr - May   | 50-75 mm       | 1.5-3 g/L IBA |

|          | Jan         | Feb                                               | Mar | Apr  | May  | Jun | Jul | Aug | Sep                 | Oct | Nov | Dec |
|----------|-------------|---------------------------------------------------|-----|------|------|-----|-----|-----|---------------------|-----|-----|-----|
| Cuttings | OK          | OK                                                | OK  | Best | Best | -   | -   | -   | OK                  | OK  | OK  | OK  |
|          | Best window | Workable - see 'Selecting material' in the method |     |      |      |     |     |     | Avoid / low success |     |     |     |

### Plant snapshot

|            |                                                                                                              |
|------------|--------------------------------------------------------------------------------------------------------------|
| Habit      | Spreading subshrub or perennial herb, 0.15-1 m. Mostly hairless; thread-like to narrow leaves; ribbed stems. |
| Flowering  | Aug - Dec and Jan - Mar (pale blue to white), in spikes at the ends of branches.                             |
| Habitat    | Coastal sand and limestone.                                                                                  |
| Local note | The coastal Perth plant is subsp. <i>thesioides</i> ; recorded in the Wanneroo LGA.                          |

### Propagation options

| Method   | Suitability | Notes                                           | Details in                 |
|----------|-------------|-------------------------------------------------|----------------------------|
| Cuttings | Main method | Non-flowering side shoots (provisional rating). | <a href="#">This guide</a> |
| Seed     | Difficult   | Scaevola seed is generally difficult.           | -                          |
| Division | Possible    | Only if stock plants have suckered.             | -                          |

### Propagation method

| Item                       | Details                                                                                                                                                                                                                              |
|----------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Best time to take cuttings | <b>Apr - May</b> - warm autumn weather between flowering periods. Sep - Mar is workable using non-flowering side shoots. Avoid Jun - Aug.                                                                                            |
| Wood type                  | <b>Softwood to semi-hardwood</b> - firm but still green tip or side-shoot cuttings.                                                                                                                                                  |
| Cutting size               | 50-75 mm.                                                                                                                                                                                                                            |
| Selecting material         | <b>Non-flowering side shoots.</b> Flowers form at the branch tips, so main tips often carry buds.                                                                                                                                    |
| Pre-treatment              | <ul style="list-style-type: none"> <li>Strip leaves from the lower half; leaves are narrow so little trimming is needed.</li> <li>Stems are slender - always use a dibber hole.</li> <li>Dip the base in rooting hormone.</li> </ul> |
| Rooting hormone            | Softwood strength (1.5 g/L IBA, <b>Clonex Green</b> ); 3 g/L ( <b>Clonex Purple</b> ) for firmer material.                                                                                                                           |
| Propagation mix            | Free-draining sand/coir mix.                                                                                                                                                                                                         |
| Environment                | Hairless leaves tolerate misting better than <i>S. canescens</i> , but fine stems collapse quickly if the mix is waterlogged or dries out. Keep evenly moist, with bottom heat and bright shade.                                     |
| Expected strike time       | About <b>4-8 weeks</b> in warm weather.                                                                                                                                                                                              |
| Post-treatment / aftercare | <ul style="list-style-type: none"> <li>Standard hardening off.</li> <li>Pot on gently - fine roots break easily. Single cuttings per tube make this easier.</li> <li>Pinch tips once growing to encourage branching.</li> </ul>      |

## Common problems

| Problem                            | Likely cause and fix                                           |
|------------------------------------|----------------------------------------------------------------|
| Fine stems wilt or collapse        | Mix drying out or waterlogged. Keep evenly moist; check daily. |
| Cuttings flower instead of rooting | Pinch off terminal flower buds; use side shoots.               |
| Root damage when potting on        | Strike one cutting per tube instead of community pots.         |

## Notes

|               |                                                                                                                      |
|---------------|----------------------------------------------------------------------------------------------------------------------|
| Difficulty    | <b>Easy-Moderate (provisional).</b> The genus strikes readily; the fine stems are less forgiving of watering errors. |
| Other methods | Seed – <i>Scaevola</i> seed is generally difficult. Division – possible if stock plants have suckered.               |
| Confidence    | <b>Medium.</b> Little species-specific data published. Based on the genus plus this species' fine, slender growth.   |
| Sources       | See References: 1, 4, 5, 6, 7, 12, 13, 14, 16                                                                        |

## Nursery records

Log each batch here to confirm or update the timing and difficulty above.

| Date set | Stock source / provenance | No. set | No. struck | Strike % | Notes |
|----------|---------------------------|---------|------------|----------|-------|
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |

# Spinifex hirsutus

## Hairy Spinifex

Family  
**Poaceae**

|            |                          |             |                     |                    |
|------------|--------------------------|-------------|---------------------|--------------------|
| DIFFICULTY | WOOD TYPE                | BEST MONTHS | CUTTING LENGTH      | HORMONE (IBA)      |
| Moderate * | Runner (stolon) sections | Mar – May   | 10–15 cm, 1-2 nodes | Optional (1.5 g/L) |

|          | Jan | Feb | Mar  | Apr  | May  | Jun | Jul | Aug | Sep | Oct | Nov | Dec |
|----------|-----|-----|------|------|------|-----|-----|-----|-----|-----|-----|-----|
| Cuttings | OK  | OK  | Best | Best | Best | –   | –   | –   | OK  | OK  | OK  | OK  |

### Best window

Workable – see ‘Selecting material’ in the method

Avoid / low success

## Plant snapshot

|                   |                                                                                                                                        |
|-------------------|----------------------------------------------------------------------------------------------------------------------------------------|
| <b>Habit</b>      | Rhizomatous, tussock-forming grass, 0.3–0.9 m, sending long runners across the sand. Broad, flat, silvery leaves, hairy on both sides. |
| <b>Flowering</b>  | Oct – Jan. Male and female flowers are on separate plants; female heads are large spiky balls that roll in the wind.                   |
| <b>Habitat</b>    | White sand on coastal dunes; an important stabiliser at the toe of unstable dunes.                                                     |
| <b>Local note</b> | Recorded in the City of Joondalup. Less common than <i>S. longifolius</i> on the Cottesloe foreshore.                                  |

## Propagation options

| Method   | Suitability | Notes                                                              | Details in                          |
|----------|-------------|--------------------------------------------------------------------|-------------------------------------|
| Cuttings | Good        | Runner (stolon) sections with 1–2 nodes.                           | <a href="#">This guide</a>          |
| Seed     | Main method | The usual nursery method; cover seed (needs dark).                 | → <a href="#">Seed Sowing Guide</a> |
| Division | Possible    | Tussock divisions work but survived less well than rooted runners. | → <a href="#">Division Guide</a>    |

## Propagation method

| Item                       | Details                                                                                                                                                                                                                                                    |
|----------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Best time to take cuttings | <b>Mar – May</b> , when runners are young, green and actively growing, giving plants ready for winter planting. Sep – Feb is workable. Avoid Jun – Aug (slow growth).                                                                                      |
| Wood type                  | <b>Runner (stolon) sections</b> – not woody cuttings. Use young, green runners, not old brown ones.                                                                                                                                                        |
| Cutting size               | <b>Nursery tubes:</b> 10–15 cm sections with at least one (ideally two) nodes.<br><b>Direct planting on site:</b> 40–50 cm runners, buried about halfway in damp sand.                                                                                     |
| Selecting material         | Actively growing runners from the dune edge. Plants are male or female, so collect from several plants if you want seed-producing plantings later.                                                                                                         |
| Pre-treatment              | <ul style="list-style-type: none"> <li>• Keep runners moist and shaded after collecting – they dry out fast.</li> <li>• Cut just below a node; trim leaves back by about half.</li> <li>• Hormone optional – roots form naturally at the nodes.</li> </ul> |
| Rooting hormone            | Optional. If used, softwood strength (1.5 g/L IBA, <b>Clonex Green</b> ) on the node.                                                                                                                                                                      |
| Propagation mix            | Coarse washed sand or a very sandy mix. Deep tubes suit the root system.                                                                                                                                                                                   |
| Environment                | Bottom heat and misting gave good rooting at the nodes in <i>S. sericeus</i> trials. The hairy leaves of this species can hold water, so mist lightly.                                                                                                     |
| Expected strike time       | Nodes root readily, but plants are <b>slow to reach planting size</b> – in one trial about 80% of cuttings were ready to plant within 18 months.                                                                                                           |

| Item                       | Details                                                                                                                                                                                                          |
|----------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Post-treatment / aftercare | <ul style="list-style-type: none"> <li>• Harden off, then grow on in full sun.</li> <li>• Pot on into deep tubes or larger pots.</li> <li>• Plant out in late autumn or winter when rain is reliable.</li> </ul> |

### Common problems

| Problem                        | Likely cause and fix                                              |
|--------------------------------|-------------------------------------------------------------------|
| Runners dry out before setting | Carry in damp bags in an esky; set the same day.                  |
| Nodes rot                      | Mix too wet or too much mist. Use coarser sand; mist less.        |
| Slow growth after rooting      | Normal for spinifex – allow time; feed lightly and give full sun. |

### Notes

|               |                                                                                                                                                                                                                                                                                                                                                |
|---------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Difficulty    | <b>Moderate (provisional).</b> Runner sections root well under bottom heat and mist, but it's labour-intensive and slow. Seed is the more common nursery method.                                                                                                                                                                               |
| Other methods | Seed – the usual nursery method. The Geraldton community nursery advises collecting from female plants growing near males. <i>S. sericeus</i> research found seed germinates far better in darkness (56%) than light (14%), so cover seed. Division of tussocks also works but survived less well than rooted cuttings in trials (31% vs 40%). |
| Confidence    | <b>Medium.</b> Method based on published trials with the closely related <i>S. sericeus</i> and general spinifex practice.                                                                                                                                                                                                                     |
| Sources       | See References: 4, 12, 18, 19, 20                                                                                                                                                                                                                                                                                                              |

### Nursery records

Log each batch here to confirm or update the timing and difficulty above.

| Date set | Stock source / provenance | No. set | No. struck | Strike % | Notes |
|----------|---------------------------|---------|------------|----------|-------|
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |

# Spinifex longifolius

Beach Spinifex

Family  
Poaceae

| DIFFICULTY | WOOD TYPE                | BEST MONTHS | CUTTING LENGTH      | HORMONE (IBA)      |
|------------|--------------------------|-------------|---------------------|--------------------|
| Moderate * | Runner (stolon) sections | Mar – May   | 10–15 cm, 1–2 nodes | Optional (1.5 g/L) |

|          | Jan | Feb | Mar  | Apr  | May  | Jun | Jul | Aug | Sep | Oct | Nov | Dec |
|----------|-----|-----|------|------|------|-----|-----|-----|-----|-----|-----|-----|
| Cuttings | OK  | OK  | Best | Best | Best | –   | –   | –   | OK  | OK  | OK  | OK  |

Best window
Workable – see 'Selecting material' in the method
Avoid / low success

## Plant snapshot

|            |                                                                                                                         |
|------------|-------------------------------------------------------------------------------------------------------------------------|
| Habit      | Rhizomatous, tussock-forming grass, 0.3–1 m high and up to 2 m wide, with long runners through the sand.                |
| Flowering  | Apr – Dec or Jan (green-brown). Male and female plants are separate; female heads up to 250 mm across roll in the wind. |
| Habitat    | White sand on coastal dunes; a key foredune stabiliser.                                                                 |
| Local note | Recorded in the Wanneroo LGA; the more common of the two spinifex species on the Cottesloe foreshore.                   |

## Propagation options

| Method   | Suitability | Notes                                                              | Details in                 |
|----------|-------------|--------------------------------------------------------------------|----------------------------|
| Cuttings | Good        | Runner (stolon) sections with 1–2 nodes.                           | <a href="#">This guide</a> |
| Seed     | Main method | The usual nursery method; smoke water, about 10 days.              | → Seed Sowing Guide        |
| Division | Possible    | Tussock divisions work but survived less well than rooted runners. | → Division Guide           |

## Propagation method

| Item                       | Details                                                                                                                                                                                                                                              |
|----------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Best time to take cuttings | Mar – May, when runners are young, green and actively growing, giving plants ready for winter planting. Sep – Feb is workable. Avoid Jun – Aug (slow growth).                                                                                        |
| Wood type                  | Runner (stolon) sections – not woody cuttings. Use young, green runners, not old brown ones.                                                                                                                                                         |
| Cutting size               | Nursery tubes: 10–15 cm sections with at least one (ideally two) nodes.<br>Direct planting on site: 40–50 cm runners, buried about halfway in damp sand.                                                                                             |
| Selecting material         | Actively growing runners from the dune edge. Plants are male or female, so collect from several plants if you want seed-producing plantings later.                                                                                                   |
| Pre-treatment              | <ul style="list-style-type: none"> <li>Keep runners moist and shaded after collecting – they dry out fast.</li> <li>Cut just below a node; trim leaves back by about half.</li> <li>Hormone optional – roots form naturally at the nodes.</li> </ul> |
| Rooting hormone            | Optional. If used, softwood strength (1.5 g/L IBA, <b>Clonex Green</b> ) on the node.                                                                                                                                                                |
| Propagation mix            | Coarse washed sand or a very sandy mix. Deep tubes suit the root system.                                                                                                                                                                             |
| Environment                | Bottom heat and misting gave good rooting at the nodes in <i>S. sericeus</i> trials. Keep the sand moist, not wet.                                                                                                                                   |
| Expected strike time       | Nodes root readily, but plants are <b>slow to reach planting size</b> – in one trial about 80% of cuttings were ready to plant within 18 months.                                                                                                     |

| Item                       | Details                                                                                                                                                                                                          |
|----------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Post-treatment / aftercare | <ul style="list-style-type: none"> <li>• Harden off, then grow on in full sun.</li> <li>• Pot on into deep tubes or larger pots.</li> <li>• Plant out in late autumn or winter when rain is reliable.</li> </ul> |

### Common problems

| Problem                        | Likely cause and fix                                              |
|--------------------------------|-------------------------------------------------------------------|
| Runners dry out before setting | Carry in damp bags in an esky; set the same day.                  |
| Nodes rot                      | Mix too wet or too much mist. Use coarser sand; mist less.        |
| Slow growth after rooting      | Normal for spinifex – allow time; feed lightly and give full sun. |

### Notes

|               |                                                                                                                                                                                                                                                                                                                                                       |
|---------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Difficulty    | <b>Moderate (provisional).</b> Runner sections root well under bottom heat and mist, but it's labour-intensive and slow. Seed is the more common nursery method.                                                                                                                                                                                      |
| Other methods | Seed – the usual nursery method. The Geraldton community nursery collects from female plants growing close to males, separates the seed, cuts off the awns and sows en masse with smoke water (about 10 days to germinate). <i>S. sericeus</i> research found seed germinates far better in darkness, so cover seed. Division of tussocks also works. |
| Confidence    | <b>Medium.</b> Method based on published trials with the closely related <i>S. sericeus</i> and general spinifex practice.                                                                                                                                                                                                                            |
| Sources       | See References: 4, 12, 18, 19, 20                                                                                                                                                                                                                                                                                                                     |

### Nursery records

Log each batch here to confirm or update the timing and difficulty above.

| Date set | Stock source / provenance | No. set | No. struck | Strike % | Notes |
|----------|---------------------------|---------|------------|----------|-------|
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |

# Templetonia retusa

Cookies Tongues

Family  
Fabaceae

| DIFFICULTY      | WOOD TYPE     | BEST MONTHS | CUTTING LENGTH | HORMONE (IBA) |
|-----------------|---------------|-------------|----------------|---------------|
| Easy-Moderate * | Semi-hardwood | Dec - Mar   | 50-75 mm       | 3 g/L IBA     |

|          | Jan  | Feb  | Mar  | Apr | May | Jun | Jul | Aug | Sep | Oct | Nov | Dec  |
|----------|------|------|------|-----|-----|-----|-----|-----|-----|-----|-----|------|
| Cuttings | Best | Best | Best | OK  | -   | -   | -   | -   | -   | -   | OK  | Best |

Best window
Workable - see 'Selecting material' in the method
Avoid / low success

## Plant snapshot

|            |                                                                                     |
|------------|-------------------------------------------------------------------------------------|
| Habit      | Much-branched shrub, 0.3-4 m.                                                       |
| Flowering  | Apr - Nov (red, sometimes white-yellow).                                            |
| Habitat    | White sand, red clay, gravel and limestone; coastal cliffs, hillsides and plateaus. |
| Local note | Recorded in the Wanneroo LGA. Cuttings keep a chosen flower colour.                 |

## Propagation options

| Method   | Suitability | Notes                                       | Details in          |
|----------|-------------|---------------------------------------------|---------------------|
| Cuttings | Good        | Firm current season's growth is successful. | This guide          |
| Seed     | Main method | Hot water + smoke water; 8-12 days.         | → Seed Sowing Guide |
| Division | Not used    | Woody shrub.                                | -                   |

## Propagation method

| Item                       | Details                                                                                                                                                                        |
|----------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Best time to take cuttings | Dec - Mar, after flowering, once the new growth has firmed. Nov and Apr are workable. Avoid May - Oct (flowering).                                                             |
| Wood type                  | Semi-hardwood - firm current season's growth (ANPSA).                                                                                                                          |
| Cutting size               | 50-75 mm (ANPSA pea-family guidance).                                                                                                                                          |
| Selecting material         | Non-flowering side shoots from healthy plants.                                                                                                                                 |
| Pre-treatment              | <ul style="list-style-type: none"> <li>Remove the leaves from the lower two-thirds.</li> <li>Light wound on firmer stems.</li> <li>Dip the base in rooting hormone.</li> </ul> |
| Rooting hormone            | Semi-hardwood strength, 3 g/L IBA (Clonex Purple).                                                                                                                             |
| Propagation mix            | Standard free-draining cutting mix.                                                                                                                                            |
| Environment                | Humid frame, bright shade, bottom heat.                                                                                                                                        |
| Expected strike time       | Not documented - check from about 6 weeks and record the result.                                                                                                               |
| Post-treatment / aftercare | <ul style="list-style-type: none"> <li>Standard hardening off.</li> <li>Pot on into a free-draining mix.</li> </ul>                                                            |

## Common problems

| Problem                   | Likely cause and fix                                    |
|---------------------------|---------------------------------------------------------|
| <b>Callus, no roots</b>   | Trim and re-set; try a lower hormone strength.          |
| <b>Cuttings brown off</b> | Too dry or too soft. Firmer material; keep humidity up. |
| <b>Base rotting</b>       | Mix too wet. Water less.                                |

## Notes

|                   |                                                                                                                                |
|-------------------|--------------------------------------------------------------------------------------------------------------------------------|
| <b>Difficulty</b> | <b>Easy-Moderate (provisional).</b> Cuttings are reported successful; no strike rates published. Seed is easy after hot water. |
| <b>Confidence</b> | <b>Medium.</b> ANPSA reports cuttings of firm current season's growth are successful; size follows ANPSA pea-family guidance.  |
| <b>Sources</b>    | See References: 4, 13, 43, 48                                                                                                  |

## Nursery records

Log each batch here to confirm or update the timing and difficulty above.

| Date set | Stock source / provenance | No. set | No. struck | Strike % | Notes |
|----------|---------------------------|---------|------------|----------|-------|
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |

# Thomasia triphylla

No widely used common name

Family  
Malvaceae

|                                                               |                                                            |                                                          |                                                            |                                                            |
|---------------------------------------------------------------|------------------------------------------------------------|----------------------------------------------------------|------------------------------------------------------------|------------------------------------------------------------|
| <div> <div>DIFFICULTY</div> <div>Easy-Moderate *</div> </div> | <div> <div>WOOD TYPE</div> <div>Semi-hardwood</div> </div> | <div> <div>BEST MONTHS</div> <div>Dec – Apr</div> </div> | <div> <div>CUTTING LENGTH</div> <div>50–75 mm</div> </div> | <div> <div>HORMONE (IBA)</div> <div>3 g/L IBA</div> </div> |
|---------------------------------------------------------------|------------------------------------------------------------|----------------------------------------------------------|------------------------------------------------------------|------------------------------------------------------------|

|          | Jan  | Feb  | Mar  | Apr  | May | Jun | Jul | Aug | Sep | Oct | Nov | Dec  |
|----------|------|------|------|------|-----|-----|-----|-----|-----|-----|-----|------|
| Cuttings | Best | Best | Best | Best | OK  | –   | –   | –   | –   | –   | OK  | Best |

Best window

Workable – see ‘Selecting material’ in the method

Avoid / low success

Plant snapshot

|            |                                                                                    |
|------------|------------------------------------------------------------------------------------|
| Habit      | Multi-stemmed shrub, 0.3–1.2 m.                                                    |
| Flowering  | Jul – Nov (pink-purple or white).                                                  |
| Habitat    | Calcareous sand; coastal limestone and dunes.                                      |
| Local note | Recorded in the Wanneroo LGA. Suits alkaline limestone sand and dry coastal sites. |

Propagation options

| Method   | Suitability | Notes                                                        | Details in |
|----------|-------------|--------------------------------------------------------------|------------|
| Cuttings | Main method | Cuttings usually strike reliably in the genus (provisional). | This guide |
| Seed     | Difficult   | Unreliable in the genus; a supplier recommends smoke.        | –          |
| Division | Not used    | –                                                            | –          |

Propagation method

| Item                       | Details                                                                                                                                                                                             |
|----------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Best time to take cuttings | Dec – Apr, after flowering, once new growth has firmed. Nov and May are workable. Avoid Jun – Oct (cool, then flowering).                                                                           |
| Wood type                  | Semi-hardwood – firm current season's growth.                                                                                                                                                       |
| Cutting size               | 50–75 mm.                                                                                                                                                                                           |
| Selecting material         | Non-flowering side shoots from healthy plants.                                                                                                                                                      |
| Pre-treatment              | <ul style="list-style-type: none"> <li>Strip leaves from the lower half; reduce large leaves by a third.</li> <li>Light wound on firmer stems.</li> <li>Dip the base in rooting hormone.</li> </ul> |
| Rooting hormone            | Semi-hardwood strength, 3 g/L IBA (Clonex Purple).                                                                                                                                                  |
| Propagation mix            | Standard free-draining cutting mix.                                                                                                                                                                 |
| Environment                | Humid frame, bright shade, bottom heat. Keep the mix moist, not wet, and mist lightly.                                                                                                              |
| Expected strike time       | Not documented – check from about 4–6 weeks and record the result.                                                                                                                                  |
| Post-treatment / aftercare | <ul style="list-style-type: none"> <li>Standard hardening off.</li> <li>Grow on in full sun; tip-prune for a bushier plant.</li> </ul>                                                              |

## Common problems

| Problem                  | Likely cause and fix                            |
|--------------------------|-------------------------------------------------|
| Leaves go brown and soft | Too wet or humid. Mist less, improve airflow.   |
| Callus, no roots         | Trim and re-set; try a lower hormone strength.  |
| Slow in winter           | Use bottom heat, or wait for the summer window. |

## Notes

|               |                                                                                                                                                           |
|---------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------|
| Difficulty    | <b>Easy-Moderate (provisional).</b> <i>Thomasia</i> cuttings usually strike reliably, but this species is untested – start with a trial batch.            |
| Other methods | Seed – unreliable in <i>Thomasia</i> ; a seed supplier recommends smoke treatment.                                                                        |
| Confidence    | <b>Medium.</b> No species data. ANPSA says cuttings of firm growth usually strike reliably in <i>Thomasia</i> , though at least one species is difficult. |
| Sources       | See References: 4, 49, 50, 51, 52                                                                                                                         |

## Nursery records

Log each batch here to confirm or update the timing and difficulty above.

| Date set | Stock source / provenance | No. set | No. struck | Strike % | Notes |
|----------|---------------------------|---------|------------|----------|-------|
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |

# Threlkeldia diffusa

Coast Bonefruit

Family  
Amaranthaceae  
(chenopods)

|                                 |                                   |                                 |                                   |                                     |
|---------------------------------|-----------------------------------|---------------------------------|-----------------------------------|-------------------------------------|
| DIFFICULTY<br><b>Moderate</b> * | WOOD TYPE<br><b>Softwood tips</b> | BEST MONTHS<br><b>Dec – Mar</b> | CUTTING LENGTH<br><b>50–75 mm</b> | HORMONE (IBA)<br><b>1.5 g/L IBA</b> |
|---------------------------------|-----------------------------------|---------------------------------|-----------------------------------|-------------------------------------|

|          | Jan         | Feb                                               | Mar  | Apr | May | Jun | Jul | Aug | Sep | Oct | Nov                 | Dec  |
|----------|-------------|---------------------------------------------------|------|-----|-----|-----|-----|-----|-----|-----|---------------------|------|
| Cuttings | Best        | Best                                              | Best | OK  | –   | –   | –   | –   | –   | –   | OK                  | Best |
|          | Best window | Workable – see ‘Selecting material’ in the method |      |     |     |     |     |     |     |     | Avoid / low success |      |

Plant snapshot

|            |                                                                                                         |
|------------|---------------------------------------------------------------------------------------------------------|
| Habit      | Much-branched, prostrate to erect perennial, 0.1–0.4 m, forming a spreading mat. Semi-succulent leaves. |
| Flowering  | Oct – Nov (tiny, green), followed by small red to black fruits.                                         |
| Habitat    | White or grey sand over limestone, coastal areas and saline flats. Salt- and drought-tolerant.          |
| Local note | Recorded in the City of Joondalup.                                                                      |

Propagation options

| Method   | Suitability | Notes                                         | Details in          |
|----------|-------------|-----------------------------------------------|---------------------|
| Cuttings | Trial       | No published cutting data; trial protocol.    | This guide          |
| Seed     | Main method | Smoke-water soak; 10–15 days in cool weather. | → Seed Sowing Guide |
| Division | Not used    | Not documented.                               | –                   |

Propagation method

| Item                       | Details                                                                                                                           |
|----------------------------|-----------------------------------------------------------------------------------------------------------------------------------|
| Best time to take cuttings | Dec – Mar – warm, with plants actively growing after flowering. Nov and Apr are workable. Avoid May – Oct.                        |
| Wood type                  | Softwood tip cuttings of the semi-succulent stems.                                                                                |
| Cutting size               | 50–75 mm.                                                                                                                         |
| Selecting material         | Healthy tips without developing fruit.                                                                                            |
| Pre-treatment              | <ul style="list-style-type: none"><li>Strip lower leaves.</li><li>No wounding.</li><li>Dip the base in rooting hormone.</li></ul> |
| Rooting hormone            | Softwood strength, 1.5 g/L IBA (Clonex Green).                                                                                    |
| Propagation mix            | Very free-draining sandy mix.                                                                                                     |
| Environment                | Semi-succulent leaves rot if kept wet – mist very little, keep the mix on the dry side, give good airflow and bright light.       |
| Expected strike time       | Not documented – start checking from about 4 weeks and record the result.                                                         |
| Post-treatment / aftercare | <ul style="list-style-type: none"><li>Standard hardening off.</li><li>Grow on in full sun; water sparingly.</li></ul>             |

## Common problems

| Problem                    | Likely cause and fix                                      |
|----------------------------|-----------------------------------------------------------|
| Cuttings turn soft and rot | Too wet or humid. Water less, mist less, improve airflow. |
| Weak, stretched growth     | Not enough light. Move to a brighter position.            |
| No roots                   | Try firmer tips or the seed method.                       |

## Notes

|               |                                                                                                                                                                                           |
|---------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Difficulty    | <b>Moderate (provisional).</b> No cutting data found; seed is the proven method (see Other methods).                                                                                      |
| Other methods | Seed – the documented method. The Geraldton community nursery soaks seed in smoke water (1:9) for 24 hours; germination takes 10–15 days and happens in cooler weather, so sow in autumn. |
| Confidence    | <b>Low.</b> No published cutting data found. Seed is the documented method; cuttings are a trial.                                                                                         |
| Sources       | See References: 1, 4, 12, 28                                                                                                                                                              |

## Nursery records

Log each batch here to confirm or update the timing and difficulty above.

| Date set | Stock source / provenance | No. set | No. struck | Strike % | Notes |
|----------|---------------------------|---------|------------|----------|-------|
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |
|          |                           |         |            |          |       |

# References

Numbers match the source list on each species page. Web addresses shortened; accessed September 2026.

| #  | Source                                                                                                                                                                                                                                                                                               | Address                                                                                                                                   |
|----|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------|
| 1  | Australian Native Plants Society (ANPSA) – Plant Propagation: cuttings, hormones, mixes, hardening off                                                                                                                                                                                               | <a href="https://anpsa.org.au/plant-propagation/">anpsa.org.au/plant-propagation/</a>                                                     |
| 2  | ANPSA – Grevillea vestita plant profile                                                                                                                                                                                                                                                              | <a href="https://anpsa.org.au/plant_profiles/grevillea-vestita/">anpsa.org.au/plant_profiles/grevillea-vestita/</a>                       |
| 3  | Australian Plants Society NSW – Grevillea vestita                                                                                                                                                                                                                                                    | <a href="https://resources.austplants.com.au/plant/grevillea-vestita/">resources.austplants.com.au/plant/grevillea-vestita/</a>           |
| 4  | FloraBase, WA Herbarium (DBCA) – species profiles for every species in this guide (e.g. 2463, 2798, 5117, 5162, 7603, 11341, 12585, 12824, 13152, 16449, 42281, 3961, 4256, 5105, 6839, 1982, 5825, 8839, 5237, 6949, 7289, 5011, 5171, 5268, 7626, 49637; others as cited in the Seed Sowing Guide) | <a href="https://florabase.dbca.wa.gov.au">florabase.dbca.wa.gov.au</a>                                                                   |
| 5  | ANPSA – Scaevola ramosissima profile (genus note: cuttings strike very readily, seed difficult)                                                                                                                                                                                                      | <a href="https://anpsa.org.au/plant_profiles/scaevola-ramosissima/">anpsa.org.au/plant_profiles/scaevola-ramosissima/</a>                 |
| 6  | Sustainable Gardening Australia – Fan Flower: take cuttings during the growing season                                                                                                                                                                                                                | <a href="https://sgaonline.org.au/fan-flower-fabulous-native-ground-cover/">sgaonline.org.au/fan-flower-fabulous-native-ground-cover/</a> |
| 7  | Greenhouse Product News – Scaevola propagation (IBA rates, flower-free cuttings, bottom heat)                                                                                                                                                                                                        | <a href="https://gpnmag.com/article/scaevola-outback-fan-flower/">gpnmag.com/article/scaevola-outback-fan-flower/</a>                     |
| 8  | ANPSA – Hibbertia vestita profile (genus note: easy from cuttings, seed unreliable)                                                                                                                                                                                                                  | <a href="https://anpsa.org.au/plant_profiles/hibbertia-vestita/">anpsa.org.au/plant_profiles/hibbertia-vestita/</a>                       |
| 9  | Australian National Botanic Gardens – Hibbertia pedunculata and H. empetrifolia (cutting timing, root rot)                                                                                                                                                                                           | <a href="https://anbg.gov.au/gnp/gnp11/hibbertia-pedunculata.html">anbg.gov.au/gnp/gnp11/hibbertia-pedunculata.html</a>                   |
| 10 | Trees and Shrubs Online – Hibbertia (nodal cuttings while in growth; seed dormancy)                                                                                                                                                                                                                  | <a href="https://treesandshrubsonline.org/articles/hibbertia/">treesandshrubsonline.org/articles/hibbertia/</a>                           |
| 11 | DBCA Library – wildflower booklet: H. hypericoides difficult from seed or cuttings                                                                                                                                                                                                                   | <a href="https://library.dbca.wa.gov.au/FullTextFiles/072542.pdf">library.dbca.wa.gov.au/FullTextFiles/072542.pdf</a>                     |
| 12 | City of Greater Geraldton Community Nursery – Native Plant Propagation Guide (2019)                                                                                                                                                                                                                  | <a href="https://cgg.wa.gov.au">cgg.wa.gov.au</a>                                                                                         |
| 13 | Growth Technology – Clonex rooting gel strengths (Green 1.5, Purple 3, Red 8 g/L IBA)                                                                                                                                                                                                                | <a href="https://growthtechnology.com.au">growthtechnology.com.au</a>                                                                     |
| 14 | EverydayWits – Propagating Scaevola (small-nursery practice, seasonal success rates)                                                                                                                                                                                                                 | <a href="https://everydaywits.com/propagate-scaevola/">everydaywits.com/propagate-scaevola/</a>                                           |
| 15 | Minimalist Gardener – Propagating Australian natives from cuttings (grevillea strike times, sap irritant)                                                                                                                                                                                            | <a href="https://minimalistgardener.com.au">minimalistgardener.com.au</a>                                                                 |
| 16 | Wikipedia – Scaevola canescens, S. repens, S. thesioides (descriptions)                                                                                                                                                                                                                              | <a href="https://en.wikipedia.org">en.wikipedia.org</a>                                                                                   |
| 17 | CompleteGrow – How to propagate Grevillea (low-phosphorus feeding)                                                                                                                                                                                                                                   | <a href="https://completegrow.com.au">completegrow.com.au</a>                                                                             |
| 18 | NZ Department of Conservation – Rehabilitation of coastal foredunes using indigenous sand-binding species (Spinifex sericeus stolon cuttings, seed)                                                                                                                                                  | <a href="https://doc.govt.nz/Documents/science-and-technical/sfc122.pdf">doc.govt.nz/Documents/science-and-technical/sfc122.pdf</a>       |
| 19 | ABC Gardening Australia – Spinifex (40–50 cm runner cuttings)                                                                                                                                                                                                                                        | <a href="https://abc.net.au/gardening/how-to/spinifex/9432152">abc.net.au/gardening/how-to/spinifex/9432152</a>                           |
| 20 | Cottesloe Coastcare Association – plant pages: Spinifex longifolius, S. hirsutus, Eremophila glabra, Rhagodia baccata                                                                                                                                                                                | <a href="https://cottesloecoastcare.org">cottesloecoastcare.org</a>                                                                       |
| 21 | ANPSA – Eremophila glabra plant profile                                                                                                                                                                                                                                                              | <a href="https://anpsa.org.au/plant_profiles/eremophila-glabra/">anpsa.org.au/plant_profiles/eremophila-glabra/</a>                       |
| 22 | Australian Bushfoods – The Eremophila Page (cutting length, wounding, hormone)                                                                                                                                                                                                                       | <a href="https://ausbushfoods.com/bushfoodsonline/reports/Plants/EREMO.TXT">ausbushfoods.com/bushfoodsonline/reports/Plants/EREMO.TXT</a> |
| 23 | Australian National Botanic Gardens – Myoporum insulare                                                                                                                                                                                                                                              | <a href="https://anbg.gov.au/gnp/interns-2010/myoporum-insulare.html">anbg.gov.au/gnp/interns-2010/myoporum-insulare.html</a>             |
| 24 | ANPSA – Lechenaultia biloba profile (cuttings strike very easily; seed difficult)                                                                                                                                                                                                                    | <a href="https://anpsa.org.au/plant_profiles/lechenaultia-biloba/">anpsa.org.au/plant_profiles/lechenaultia-biloba/</a>                   |
| 25 | Better Homes and Gardens – fastest and easiest native plants to propagate (leschenaultia timing)                                                                                                                                                                                                     | <a href="https://bhg.com.au/how-to-propagate-natives">bhg.com.au/how-to-propagate-natives</a>                                             |

| #  | Source                                                                                                                                                   | Address                                                                                                                                                                                              |
|----|----------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| 26 | Nurseries Online – <i>Scaevola crassifolia</i> (semi-hardwood cuttings in warmer months)                                                                 | <a href="http://nurseriesonline.com.au/plant-index/australian-native-plants/scaevola-crassifolia/">nurseriesonline.com.au/plant-index/australian-native-plants/scaevola-crassifolia/</a>             |
| 27 | Natural Area Nursery – WA native plant propagation ( <i>H. hypericoides</i> difficult to propagate)                                                      | <a href="http://naturalarea.com.au/revegetation-plant-production/wa-native-plant-propagation/">naturalarea.com.au/revegetation-plant-production/wa-native-plant-propagation/</a>                     |
| 28 | APACE WA – <i>Threlkeldia diffusa</i>                                                                                                                    | <a href="http://apacewa.org.au/plant/threlkeldia-diffusa/">apacewa.org.au/plant/threlkeldia-diffusa/</a>                                                                                             |
| 29 | Geographe Community Landcare Nursery – <i>Phyllanthus calycinus</i> ( <i>Lysiandra calycina</i> )                                                        | <a href="http://gcln.org.au/product/phyllanthus-calycinus/">gcln.org.au/product/phyllanthus-calycinus/</a>                                                                                           |
| 30 | Wikipedia – <i>Hibbertia hypericoides</i> , <i>Eremophila glabra</i> subsp. <i>albicans</i> , <i>Phyllanthus calycinus</i> , <i>Scaevola crassifolia</i> | <a href="http://en.wikipedia.org">en.wikipedia.org</a>                                                                                                                                               |
| 31 | PMC – Sympatric species of <i>Hibbertia</i> vary in dormancy break and germination requirements                                                          | <a href="http://pmc.ncbi.nlm.nih.gov/articles/PMC3336950">pmc.ncbi.nlm.nih.gov/articles/PMC3336950</a>                                                                                               |
| 32 | Cromer E.L. (2007) Seed Germination and Research Records from Alcoa's Marrinup Nursery (seed test figures quoted under Other methods)                    | <a href="http://library.dbca.wa.gov.au/static/FullTextFiles/924754.pdf">library.dbca.wa.gov.au/static/FullTextFiles/924754.pdf</a>                                                                   |
| 33 | Australian National Botanic Gardens – <i>Carpobrotus glaucescens</i> (layering, 30 cm stem pieces, division)                                             | <a href="http://anbg.gov.au/gnp/interns-2005/carpobrotus-glaucescens.html">anbg.gov.au/gnp/interns-2005/carpobrotus-glaucescens.html</a>                                                             |
| 34 | Australian National Botanic Gardens – <i>Leucophyta brownii</i> (semi-hardened cuttings; dislikes humidity)                                              | <a href="http://anbg.gov.au/gnp/interns-2014/leucophyta-brownii.html">anbg.gov.au/gnp/interns-2014/leucophyta-brownii.html</a>                                                                       |
| 35 | ANPSA – Australian Daisies (perennial stem cuttings strike quickly; 3:1 sand:peat mix)                                                                   | <a href="http://anpsa.org.au/genera/daisies/">anpsa.org.au/genera/daisies/</a>                                                                                                                       |
| 36 | Australian National Botanic Gardens – <i>Atriplex nummularia</i> (ventilate, avoid mist on cuttings)                                                     | <a href="http://anbg.gov.au/gnp/interns-2007/atriples-nummularia.html">anbg.gov.au/gnp/interns-2007/atriples-nummularia.html</a>                                                                     |
| 37 | Nurseries Online – <i>Atriplex nummularia</i> (15 cm terminal cuttings in spring)                                                                        | <a href="http://nurseriesonline.com.au/plant-index/australian-native-plants/atriples-nummularia/">nurseriesonline.com.au/plant-index/australian-native-plants/atriples-nummularia/</a>               |
| 38 | Applied Sciences (MDPI, 2025) – Propagation of <i>Atriplex halimus</i> by unrooted cuttings                                                              | <a href="http://mdpi.com/2076-3417/15/13/7027">mdpi.com/2076-3417/15/13/7027</a>                                                                                                                     |
| 39 | Peppy Plants – <i>Hibbertia cuneiformis</i> (propagated by cuttings)                                                                                     | <a href="http://peppyplants.com.au/hibbertia-cuneiformis-unveiling-the-charms-of-the-cutleaf-hibbertia/">peppyplants.com.au/hibbertia-cuneiformis-unveiling-the-charms-of-the-cutleaf-hibbertia/</a> |
| 40 | Wikipedia – <i>Hibbertia cuneiformis</i> (description, flowering, habitat)                                                                               | <a href="http://en.wikipedia.org/wiki/Hibbertia_cuneiformis">en.wikipedia.org/wiki/Hibbertia_cuneiformis</a>                                                                                         |
| 41 | ANPSA – <i>Melaleuca</i> genus page (75–100 mm cuttings, strip, wound, hormone)                                                                          | <a href="http://anpsa.org.au/genera/melaleuca/">anpsa.org.au/genera/melaleuca/</a>                                                                                                                   |
| 42 | ANPSA – <i>Kennedia prostrata</i> profile (cuttings of firm current season's growth)                                                                     | <a href="http://anpsa.org.au/plant_profiles/kennedia-prostrata/">anpsa.org.au/plant_profiles/kennedia-prostrata/</a>                                                                                 |
| 43 | ANPSA – Pea family (Fabaceae) genus page (50–75 mm cuttings, leaves off the lower two-thirds)                                                            | <a href="http://anpsa.org.au/genera/pea-family/">anpsa.org.au/genera/pea-family/</a>                                                                                                                 |
| 44 | ANPSA – <i>Hardenbergia comptoniana</i> profile (cuttings of firm current season's growth strike well)                                                   | <a href="http://anpsa.org.au/plant_profiles/hardenbergia-comptoniana/">anpsa.org.au/plant_profiles/hardenbergia-comptoniana/</a>                                                                     |
| 45 | ANPSA – <i>Hemiandra pungens</i> profile (relatively easy from cuttings; unreliable in humid conditions; grafting)                                       | <a href="http://anpsa.org.au/plant_profiles/hemiandra-pungens/">anpsa.org.au/plant_profiles/hemiandra-pungens/</a>                                                                                   |
| 46 | Gardening With Angus – <i>Hemiandra pungens</i> (softwood and semi-hardwood cuttings)                                                                    | <a href="http://gardeningwithangus.com.au/hemiandra-pungens-snake-plant/">gardeningwithangus.com.au/hemiandra-pungens-snake-plant/</a>                                                               |
| 47 | APACE WA – <i>Hemiandra pungens</i> (dieback resistant)                                                                                                  | <a href="http://apacewa.org.au/plant/hemiandra-pungens/">apacewa.org.au/plant/hemiandra-pungens/</a>                                                                                                 |
| 48 | ANPSA – <i>Templetonia retusa</i> profile (cuttings of firm current season's growth are successful)                                                      | <a href="http://anpsa.org.au/plant_profiles/templetonia-retusa/">anpsa.org.au/plant_profiles/templetonia-retusa/</a>                                                                                 |
| 49 | ANPSA – <i>Thomasia grandiflora</i> profile (genus: cuttings of firm growth usually strike reliably; seed unreliable)                                    | <a href="http://anpsa.org.au/plant_profiles/thomasia-grandiflora/">anpsa.org.au/plant_profiles/thomasia-grandiflora/</a>                                                                             |
| 50 | Geographe Community Landcare Nursery – <i>Thomasia triphylla</i> (coastal, drought tolerant, alkaline sand)                                              | <a href="http://gcln.org.au/product/thomasia-triphylla/">gcln.org.au/product/thomasia-triphylla/</a>                                                                                                 |
| 51 | Nindethana Australian Seed – <i>Thomasia triphylla</i> and <i>Scaevola nitida</i> seed (smoke treatment recommended)                                     | <a href="http://nindethana.net.au">nindethana.net.au</a>                                                                                                                                             |
| 52 | ANPSA – Australian Plants Online (2007): <i>Thomasia pygmaea</i> , cuttings difficult to strike                                                          | <a href="http://anpsa.org.au/APOL2007/jul07-2.html">anpsa.org.au/APOL2007/jul07-2.html</a>                                                                                                           |

| #  | Source                                                                                                                                          | Address                                                                                                                                                   |
|----|-------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------|
| 53 | ANPSA – Hypocalymma robustum profile (cuttings preferred, strike readily from hardened growth)                                                  | <a href="https://anpsa.org.au/plant_profiles/hypocalymma-robustum/">anpsa.org.au/plant_profiles/hypocalymma-robustum/</a>                                 |
| 54 | ANPSA – Geraldton wax and relatives (75–100 mm cuttings in January, strip lower two-thirds, wound, hormone)                                     | <a href="https://anpsa.org.au/genera/geraldton-wax-and-relatives/">anpsa.org.au/genera/geraldton-wax-and-relatives/</a>                                   |
| 55 | Gardening With Angus – Hypocalymma robustum (softwood and semi-hardwood cuttings)                                                               | <a href="https://gardeningwithangus.com.au/hypocalymma-robustum-swan-river-myrtle/">gardeningwithangus.com.au/hypocalymma-robustum-swan-river-myrtle/</a> |
| 56 | Australian Plants Society NSW – Grevillea crithmifolia (cuttings may be slow to root; shrub and groundcover forms)                              | <a href="https://resources.austplants.com.au/plant/grevillea-crithmifolia/">resources.austplants.com.au/plant/grevillea-crithmifolia/</a>                 |
| 57 | APACE WA – Grevillea crithmifolia (dieback susceptible)                                                                                         | <a href="https://apacewa.org.au/plant/grevillea-crithmifolia/">apacewa.org.au/plant/grevillea-crithmifolia/</a>                                           |
| 58 | ANPSA – Grevillea preissii profile (cuttings slow; hormone improves success; nick seed)                                                         | <a href="https://anpsa.org.au/plant_profiles/grevillea-preissii/">anpsa.org.au/plant_profiles/grevillea-preissii/</a>                                     |
| 59 | ANPSA – Grevillea genus page (75–100 mm cuttings, strip lower half to two-thirds, wounding, hormone; phosphorus)                                | <a href="https://anpsa.org.au/genera/grevillea/">anpsa.org.au/genera/grevillea/</a>                                                                       |
| 60 | Olde P.M. (2022) Telopea – Grevillea preissii subsp. glabrilimba raised to species rank as G. glabrilimba                                       | <a href="https://openjournals.library.sydney.edu.au/TEL/article/view/15872">openjournals.library.sydney.edu.au/TEL/article/view/15872</a>                 |
| 61 | Nindethana Australian Seed – Grevillea crithmifolia and G. preissii seed (smoke treatment recommended)                                          | <a href="https://nindethana.net.au">nindethana.net.au</a>                                                                                                 |
| 62 | ANPSA – Eremophila and relatives (Myoporum: cuttings of hardened current season's growth strike readily)                                        | <a href="https://anpsa.org.au/genera/eremophila-and-relatives/">anpsa.org.au/genera/eremophila-and-relatives/</a>                                         |
| 63 | APACE WA – Myoporum caprarioides (winter-wet sites; berries attract birds)                                                                      | <a href="https://apacewa.org.au/plant/myoporum-caprarioides/">apacewa.org.au/plant/myoporum-caprarioides/</a>                                             |
| 64 | ANPSA – Anthocercis littorea profile (cuttings of hardened growth strike reasonably well)                                                       | <a href="https://anpsa.org.au/plant_profiles/anthocercis-littorea/">anpsa.org.au/plant_profiles/anthocercis-littorea/</a>                                 |
| 65 | Wikipedia – Anthocercis (tropane alkaloids; poisoning)                                                                                          | <a href="https://en.wikipedia.org/wiki/Anthocercis">en.wikipedia.org/wiki/Anthocercis</a>                                                                 |
| 66 | ANPSA – Pimelea physodes profile (genus: cuttings successful but bark tears; seed difficult, smoke helps)                                       | <a href="https://anpsa.org.au/plant_profiles/pimelea-physodes/">anpsa.org.au/plant_profiles/pimelea-physodes/</a>                                         |
| 67 | ANPSA – Pimelea ferruginea profile (firm current season's growth usually strikes reliably)                                                      | <a href="https://anpsa.org.au/plant_profiles/pimelea-ferruginea/">anpsa.org.au/plant_profiles/pimelea-ferruginea/</a>                                     |
| 68 | Commander L.E. et al. (2009) Journal of Arid Environments – GA, smoke water and karrikinolide raise germination (incl. Anthocercis littorea)    | <a href="https://sciencedirect.com/science/article/abs/pii/S0140196309000287">sciencedirect.com/science/article/abs/pii/S0140196309000287</a>             |
| 69 | Cochrane A. (2007) Seed Notes 17 – Hibbertia (most plants grown from cuttings; dormant seed; GA 25 mg/L). Wildflower Society of WA / DEC        | <a href="https://library.dbca.wa.gov.au/FullTextFiles/PAM02004.pdf">library.dbca.wa.gov.au/FullTextFiles/PAM02004.pdf</a>                                 |
| 70 | Thiele K.R. – Nuytsia 30: the Hibbertia spicata group split into six species, incl. H. leptotheca                                               | <a href="https://florabase.dbca.wa.gov.au/nuytsia/article/924">florabase.dbca.wa.gov.au/nuytsia/article/924</a>                                           |
| 71 | ANPSA – Scaevola aemula profile (cuttings strike very easily from firm current season's growth)                                                 | <a href="https://anpsa.org.au/plant_profiles/scaevola-aemula/">anpsa.org.au/plant_profiles/scaevola-aemula/</a>                                           |
| 72 | Australian Seed – Scaevola nitida (smoke pre-treatment; 21–42 days; tolerates full coastal exposure)                                            | <a href="https://australianseed.com/shop/item/scaevola-nitida">australianseed.com/shop/item/scaevola-nitida</a>                                           |
| 73 | ANGAIR – Propagating the impossibles (Pimelea: winter sowing with smoke water)                                                                  | <a href="https://angair.org.au/propagating-the-impossibles/">angair.org.au/propagating-the-impossibles/</a>                                               |
| 74 | Cochrane A. (2007) Seed Notes 20 – Lasioptalam, Thomasia and Guichenotia (often grown from cuttings; hard seed). Wildflower Society of WA / DEC | <a href="https://library.dbca.wa.gov.au/FullTextFiles/PAM02007.pdf">library.dbca.wa.gov.au/FullTextFiles/PAM02007.pdf</a>                                 |
| 75 | Gardening With Angus – Guichenotia ledifolia (semi-hardwood cuttings)                                                                           | <a href="https://gardeningwithangus.com.au/guichenotia-ledifolia/">gardeningwithangus.com.au/guichenotia-ledifolia/</a>                                   |
| 76 | ANPSA – Guichenotia macrantha profile (cuttings of firm current season's growth usually strike reliably)                                        | <a href="https://anpsa.org.au/plant_profiles/guichenotia-macrantha/">anpsa.org.au/plant_profiles/guichenotia-macrantha/</a>                               |
| 77 | Botanic Gardens of South Australia Plant Selector – Guichenotia ledifolia (drought and lime tolerant)                                           | <a href="https://plantselector.botanicgardens.sa.gov.au">plantselector.botanicgardens.sa.gov.au</a>                                                       |
| 78 | ANPSA – Scaevola calendulacea profile (cuttings of this and other Scaevola species strike very readily)                                         | <a href="https://anpsa.org.au/plant_profiles/scaevola-calendulacea/">anpsa.org.au/plant_profiles/scaevola-calendulacea/</a>                               |

## Revision log

| Version | Date     | Changes                                                                                                                                                                                                                                                                                                                                                                                                                    |
|---------|----------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| 1.0     | Sep 2026 | First edition: <i>Grevillea vestita</i> , <i>Hibbertia racemosa</i> , <i>Scaevola canescens</i> , <i>S. repens</i> , <i>S. thesioides</i> .                                                                                                                                                                                                                                                                                |
| 1.1     | Sep 2026 | Added 10 species: <i>Eremophila glabra</i> , <i>Hibbertia hypericoides</i> , <i>Lechenaultia linarioides</i> , <i>Myoporum insulare</i> , <i>Phyllanthus calycinus</i> , <i>Scaevola crassifolia</i> , <i>S. globulifera</i> , <i>Spinifex hirsutus</i> , <i>S. longifolius</i> , <i>Threlkeldia diffusa</i> . Added page numbers to the species index.                                                                    |
| 1.2     | Sep 2026 | Added 11 species: <i>Atriplex isatidea</i> , <i>Calothamnus quadrifidus</i> , <i>Carpobrotus virescens</i> , <i>Hibbertia cuneiformis</i> , <i>Kennedia prostrata</i> , <i>Leucophyta brownii</i> , <i>Melaleuca huegelii</i> , <i>M. systema</i> , <i>Olearia axillaris</i> , <i>Pithocarpa cordata</i> , <i>Rhagodia baccata</i> . Added a 'Propagation options' table (cuttings, seed, division) to every species page. |
| 1.3     | Sep 2026 | Added 4 species: <i>Hardenbergia comptoniana</i> , <i>Hemiandra pungens</i> , <i>Templetonia retusa</i> , <i>Thomasia triphylla</i> .                                                                                                                                                                                                                                                                                      |
| 1.4     | Sep 2026 | Added 3 species: <i>Grevillea crithmifolia</i> , <i>G. preissii</i> , <i>Hypocalymma robustum</i> .                                                                                                                                                                                                                                                                                                                        |
| 1.5     | Sep 2026 | Added 3 species: <i>Anthocercis littorea</i> , <i>Myoporum caprarioides</i> , <i>Pimelea calcicola</i> .                                                                                                                                                                                                                                                                                                                   |
| 1.6     | Sep 2026 | Added 4 species: <i>Guichenotia ledifolia</i> , <i>Hibbertia spicata</i> (incl. <i>H. leptotheca</i> ), <i>Pimelea sulphurea</i> , <i>Scaevola nitida</i> .                                                                                                                                                                                                                                                                |
|         |          |                                                                                                                                                                                                                                                                                                                                                                                                                            |
|         |          |                                                                                                                                                                                                                                                                                                                                                                                                                            |
|         |          |                                                                                                                                                                                                                                                                                                                                                                                                                            |
|         |          |                                                                                                                                                                                                                                                                                                                                                                                                                            |