

# Native Plant Division Guide

Western Australian native species – Joondalup, WA

Version 1.4 · October 2026 · 12 species

## Species in this edition (alphabetical)

Full index on page 2

|                                                             |       |                      |                                                    |       |                      |
|-------------------------------------------------------------|-------|----------------------|----------------------------------------------------|-------|----------------------|
| <b>Anigozanthos humilis</b><br>Catspaw                      | p. 6  | Moderate-Difficult * | <b>Ficinia nodosa</b><br>Knotted Club Rush         | p. 18 | Easy                 |
| <b>Anigozanthos manglesii</b><br>Red and Green Kangaroo Paw | p. 8  | Moderate             | <b>Lepidosperma gladiatum</b><br>Coast Sword-sedge | p. 20 | Moderate *           |
| <b>Austrostipa elegantissima</b><br>Feather Speargrass      | p. 10 | Easy-Moderate *      | <b>Lomandra maritima</b><br>Maritime Mat Rush      | p. 22 | Easy-Moderate *      |
| <b>Austrostipa flavescens</b><br>Coast Speargrass           | p. 12 | Moderate *           | <b>Poa poiformis</b><br>Coastal Poa                | p. 24 | Easy-Moderate *      |
| <b>Conostylis aculeata</b><br>Prickly Conostylis            | p. 14 | Moderate             | <b>Spinifex hirsutus</b><br>Hairy Spinifex         | p. 26 | Moderate-Difficult * |
| <b>Dianella revoluta</b><br>Blueberry Lily / Flax Lily      | p. 16 | Easy                 | <b>Spinifex longifolius</b><br>Beach Spinifex      | p. 28 | Moderate-Difficult * |

\* Provisional rating – based on related plants; confirm with your own nursery records.

Compiled from published sources (see References). Windows and ratings are starting points for local practice.

© 2026 Mitchell Donald. All rights reserved.

# How to use this guide

Division means splitting one plant into several pieces that each already have roots. It is the quickest way to multiply clump-forming plants such as kangaroo paws, flax lilies and tufted grasses, and every new plant is identical to its parent. Each species has its own page set, listed alphabetically: quick facts, a division calendar, a plant snapshot, the method, common problems and a record table.

Divide nursery stock plants, not plants from bushland. Timing is written for the Joondalup / Perth coastal climate: most clumps are best divided in autumn so the pieces root before winter.

## Species index – at a glance

| Species                                                     | Page | Difficulty           | Division type                   | Best months | Division size                     |
|-------------------------------------------------------------|------|----------------------|---------------------------------|-------------|-----------------------------------|
| <i>Anigozanthos humilis</i><br>Catspaw                      | 6    | Moderate-Difficult * | Rhizome clump division          | Oct – Nov   | 3+ shoots with roots              |
| <i>Anigozanthos manglesii</i><br>Red and Green Kangaroo Paw | 8    | Moderate             | Rhizome clump division          | Mar – May   | 3+ shoots with roots              |
| <i>Austrostipa elegantissima</i><br>Feather Speargrass      | 10   | Easy-Moderate *      | Rhizome / tussock division      | Aug – Sep   | Clumps of 3–5 tillers             |
| <i>Austrostipa flavescens</i><br>Coast Speargrass           | 12   | Moderate *           | Tussock division                | Apr – Jun   | Small clumps, 3–5 tillers         |
| <i>Conostylis aculeata</i><br>Prickly Conostylis            | 14   | Moderate             | Clump division / rooted runners | Mar – May   | Larger clumps, 5+ shoots          |
| <i>Dianella revoluta</i><br>Blueberry Lily / Flax Lily      | 16   | Easy                 | Rhizome clump division          | Mar – May   | 3–4 shoots with rhizome           |
| <i>Ficinia nodosa</i><br>Knotted Club Rush                  | 18   | Easy                 | Rhizome clump division          | Mar – May   | 6–10 stems with rhizome           |
| <i>Lepidosperma gladiatum</i><br>Coast Sword-sedge          | 20   | Moderate *           | Crown (clump) division          | Mar – May   | Large pieces with rhizome & roots |
| <i>Lomandra maritima</i><br>Maritime Mat Rush               | 22   | Easy-Moderate *      | Clump division                  | Aug – Sep   | 3–4 shoots with roots             |
| <i>Poa poiformis</i><br>Coastal Poa                         | 24   | Easy-Moderate *      | Tussock division                | Aug – Sep   | Clumps of 3–5 tillers             |
| <i>Spinifex hirsutus</i><br>Hairy Spinifex                  | 26   | Moderate-Difficult * | Tussock division                | Mar – May   | Rooted clumps with runners        |
| <i>Spinifex longifolius</i><br>Beach Spinifex               | 28   | Moderate-Difficult * | Tussock division                | Mar – May   | Rooted clumps with runners        |

\* Provisional rating – based on related plants; update once your own results are recorded.

## Division calendar

| Species                          | Jan | Feb | Mar  | Apr  | May  | Jun  | Jul | Aug  | Sep  | Oct  | Nov  | Dec |
|----------------------------------|-----|-----|------|------|------|------|-----|------|------|------|------|-----|
| <i>Anigozanthos humilis</i>      | –   | –   | OK   | OK   | –    | –    | –   | –    | –    | Best | Best | –   |
| <i>Anigozanthos manglesii</i>    | –   | OK  | Best | Best | Best | OK   | –   | –    | OK   | OK   | –    | –   |
| <i>Austrostipa elegantissima</i> | –   | –   | OK   | OK   | OK   | OK   | OK  | Best | Best | OK   | –    | –   |
| <i>Austrostipa flavescens</i>    | –   | –   | OK   | Best | Best | Best | OK  | –    | –    | –    | –    | –   |
| <i>Conostylis aculeata</i>       | –   | OK  | Best | Best | Best | OK   | –   | –    | –    | OK   | OK   | OK  |
| <i>Dianella revoluta</i>         | –   | OK  | Best | Best | Best | OK   | –   | OK   | OK   | OK   | –    | –   |
| <i>Ficinia nodosa</i>            | –   | OK  | Best | Best | Best | OK   | OK  | OK   | OK   | OK   | –    | –   |
| <i>Lepidosperma gladiatum</i>    | –   | OK  | Best | Best | Best | OK   | OK  | –    | –    | –    | –    | –   |

|                                                                                               |                                                                                                                                        |    |        |      |                                                                                                         |    |        |      |      |    |    |    |
|-----------------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------|----|--------|------|---------------------------------------------------------------------------------------------------------|----|--------|------|------|----|----|----|
| <i>Lomandra maritima</i>                                                                      | –                                                                                                                                      | –  | OK     | OK   | OK                                                                                                      | OK | OK     | Best | Best | OK | –  | –  |
| <i>Poa poiformis</i>                                                                          | –                                                                                                                                      | –  | OK     | OK   | OK                                                                                                      | –  | OK     | Best | Best | OK | –  | –  |
| <i>Spinifex hirsutus</i>                                                                      | OK                                                                                                                                     | OK | Best   | Best | Best                                                                                                    | –  | –      | –    | OK   | OK | OK | OK |
| <i>Spinifex longifolius</i>                                                                   | OK                                                                                                                                     | OK | Best   | Best | Best                                                                                                    | –  | –      | –    | OK   | OK | OK | OK |
| Perth season                                                                                  | Summer                                                                                                                                 |    | Autumn |      | Winter                                                                                                  |    | Spring |      | Sum. |    |    |    |
|  Best window |  Workable – slower to re-establish or needs more care |    |        |      |  Avoid / low success |    |        |      |      |    |    |    |

## Other species that can be divided or layered

These have no page in this guide because another method works better, but they can be split, or rooted pieces lifted, when stock plants allow.

| Species                         | How                                                                  | Main method |
|---------------------------------|----------------------------------------------------------------------|-------------|
| <i>Acanthocarpus preissii</i>   | Rhizomatous clumps; division not documented.                         | Seed        |
| <i>Atriplex cinerea</i>         | Low stems layer themselves – lift rooted pieces.                     | Cuttings    |
| <i>Carpobrotus virescens</i>    | Lift stems that have rooted – treat as cuttings.                     | Cuttings    |
| <i>Conostylis candicans</i>     | Less easy than kangaroo paws – large pieces (3+ big shoots), autumn. | Seed        |
| <i>Conostylis setigera</i>      | Large pieces in autumn, as for other <i>Conostylis</i> .             | Seed        |
| <i>Grevillea vestita</i>        | Can sucker – lift rooted suckers.                                    | Cuttings    |
| <i>Hibbertia racemosa</i>       | Low stems may root where they touch the soil (layering).             | Cuttings    |
| <i>Lechenaultia linarioides</i> | Some <i>leschenaultias</i> sucker – pot up rooted suckers.           | Cuttings    |
| <i>Opercularia vaginata</i>     | May sucker; untested.                                                | –           |
| <i>Orthrosanthus laxus</i>      | Clumps can be divided in spring (RHS); untested here.                | Seed        |
| <i>Patersonia occidentalis</i>  | Large pieces in autumn; ANBG calls division unreliable.              | Seed        |
| <i>Scaevola canescens</i>       | Only if stems have rooted or suckered.                               | Cuttings    |
| <i>Scaevola globulifera</i>     | Check stock plants for rooted stems.                                 | Cuttings    |
| <i>Scaevola repens</i>          | Some <i>scaevolae</i> sucker; check for rooted pieces.               | Cuttings    |
| <i>Scaevola thesioides</i>      | Only if stock plants have suckered.                                  | Cuttings    |
| <i>Tricoryne elatior</i>        | Usually divided in eastern Australia – small rhizome clumps.         | –           |

# Types of division

| Type            | What it is                                                                                                                | Examples                                                                    |
|-----------------|---------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------|
| Clump / rhizome | Grass-like clumps with a creeping underground stem. Cut into pieces of rhizome, each with shoots and roots.               | <i>Anigozanthos</i> , <i>Dianella</i> , <i>Lomandra</i> , <i>Patersonia</i> |
| Tussock         | True grasses. Split a clump into smaller clumps of shoots (tillers) with roots.                                           | <i>Austrostipa</i> , <i>Spinifex</i>                                        |
| Suckers         | New shoots arising from roots a short way from the parent. Dig up with roots attached and pot individually.               | Some <i>Lechenaultia</i> , <i>Scaevola</i> , <i>Grevillea</i>               |
| Layering        | Stems that root where they touch the soil. Cut into rooted pieces and pot up, or pin a stem into a pot beside the parent. | <i>Hibbertia</i> , <i>Carpobrotus</i> , spinifex runners                    |

Based on ANPSA's propagation notes (source 1).

## Difficulty ratings

| Rating             | Meaning                                                                     |
|--------------------|-----------------------------------------------------------------------------|
| Easy               | Divisions re-establish reliably with the standard method.                   |
| Easy-Moderate      | Usually re-establish; one factor (e.g. moisture, size of piece) needs care. |
| Moderate           | Re-establish, but results depend on timing, piece size and aftercare.       |
| Moderate-Difficult | Many divisions fail; another method is usually better.                      |
| Difficult          | Rarely successful; use another method.                                      |

## Confidence

| Level  | Meaning                                                                                      |
|--------|----------------------------------------------------------------------------------------------|
| High   | A published division method exists for the species or genus.                                 |
| Medium | Based on closely related plants or general practice for this plant type. Verify with trials. |
| Low    | Little information available; treat as a trial protocol.                                     |

# Standard division method

Species pages note where a species differs.

| Step              | What to do                                                                                                                                                                             |
|-------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| 1. Choose stock   | Healthy, vigorous, pest- and disease-free nursery plants with plenty of shoots. Label each stock plant's provenance.                                                                   |
| 2. Time it        | Warmer months give the best results; late-summer to autumn divisions root well before winter (ANPSA). Summer divisions need extra shelter.                                             |
| 3. Water first    | Water the stock plant the day before so roots and shoots are full of water.                                                                                                            |
| 4. Lift and clean | Lift the clump or knock it out of its pot. Shake or wash off the mix so you can see where to cut.                                                                                      |
| 5. Divide         | Cut with a clean, sharp knife or spade. Aim for pieces with <b>three or four leaf shoots</b> , a good section of rhizome and healthy roots (ANPSA). Smaller pieces need a humid house. |
| 6. Trim           | Cut off damaged roots. Cut leaves back by about <b>one-third</b> to reduce water loss.                                                                                                 |
| 7. Pot            | Pot into a free-draining mix with shoots at the <b>same level as before</b> . Firm in and water well.                                                                                  |
| 8. Shelter        | Keep in a sheltered, shaded spot (or humid house for small pieces) until new leaves grow and roots show at the drainage holes. Small divisions typically need 4–6 weeks.               |
| 9. Harden off     | About a week with nights exposed, then a week in a sheltered spot, then into the open (ANPSA).                                                                                         |
| 10. Pot on        | Pot on or plant out once roots fill the pot.                                                                                                                                           |

## Hygiene

| Topic      | Guidance                                                                                                                                                                                             |
|------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Tools      | Clean, sharp knives and spades. Wipe with alcohol between stock plants.                                                                                                                              |
| Disease    | Don't divide plants showing disease. Kangaroo paws are prone to ink disease (black leaf spots); many WA natives are susceptible to <i>Phytophthora</i> – keep pots off the ground and use fresh mix. |
| Collecting | Digging plants from bushland needs a DBCA licence and landholder permission. Divide nursery stock instead, and label every batch with its source.                                                    |

# Anigozanthos humilis

Catpaw

Family  
Haemodoraceae

| DIFFICULTY           | DIVISION TYPE          | BEST MONTHS | DIVISION SIZE        | TIME TO ESTABLISH   |
|----------------------|------------------------|-------------|----------------------|---------------------|
| Moderate-Difficult * | Rhizome clump division | Oct – Nov   | 3+ shoots with roots | 4–6 weeks sheltered |

|          | Jan         | Feb | Mar                                                  | Apr | May | Jun | Jul | Aug | Sep | Oct  | Nov  | Dec                 |
|----------|-------------|-----|------------------------------------------------------|-----|-----|-----|-----|-----|-----|------|------|---------------------|
| Division | –           | –   | OK                                                   | OK  | –   | –   | –   | –   | –   | Best | Best | –                   |
|          | Best window |     | Workable – slower to re-establish or needs more care |     |     |     |     |     |     |      |      | Avoid / low success |

Plant snapshot

|            |                                                                                                                      |
|------------|----------------------------------------------------------------------------------------------------------------------|
| Habit      | Low rhizomatous herb, usually 0.1–0.3 m locally. Often short-lived – about five years at most in humid-summer areas. |
| Flowering  | Jul – Oct (yellow, orange and red furry paws).                                                                       |
| Habitat    | Sand, sandy loam, clay, laterite and limestone; well-drained ground to winter-wet swamps.                            |
| Local note | Recorded in Joondalup and Wanneroo (subsp. <i>humilis</i> ). Prone to rust.                                          |

Propagation options

| Method   | Suitability | Notes                                             | Details in          |
|----------|-------------|---------------------------------------------------|---------------------|
| Cuttings | Not used    | Kangaroo paws are not grown from cuttings.        | –                   |
| Seed     | Main method | No pre-treatment needed; smoke lifts germination. | → Seed Sowing Guide |
| Division | Possible    | After flowering, vigorous plants only.            | This guide          |

Division method

| Item                | Details                                                                                                                                                                                                          |
|---------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Best time to divide | Oct – Nov, just after flowering (ANPSA). Mar – Apr is workable, as for other kangaroo paws. Avoid Dec – Feb (heat) and May – Sep (plants are growing and flowering).                                             |
| Division type       | Rhizome clump division.                                                                                                                                                                                          |
| Division size       | At least 3 shoots with a piece of rhizome and roots. Plants are small, so you may only get two or three pieces.                                                                                                  |
| Selecting stock     | Only vigorous plants – weak or older plants rarely recover. Choose plants free of rust.                                                                                                                          |
| Preparation         | <ul style="list-style-type: none"><li>Water the stock plant the day before.</li><li>Lift the clump and cut the rhizome apart with a clean, sharp knife.</li><li>Trim the leaves back by about a third.</li></ul> |
| Potting             | Pot into a very free-draining mix at the original depth.                                                                                                                                                         |
| Environment         | Sheltered, shaded spot until new growth starts. Keep the leaves dry – water the mix, not the plant.                                                                                                              |
| Time to establish   | About 4–6 weeks (record yours).                                                                                                                                                                                  |
| Aftercare           | <ul style="list-style-type: none"><li>Harden off gradually.</li><li>Grow on in full sun with good airflow.</li></ul>                                                                                             |

## Common problems

| Problem                          | Likely cause and fix                                                          |
|----------------------------------|-------------------------------------------------------------------------------|
| <b>Divisions fail</b>            | Stock plant too weak. Divide only vigorous plants.                            |
| <b>Orange pustules on leaves</b> | Rust. Remove affected leaves; keep foliage dry; don't divide infected plants. |
| <b>Base rots</b>                 | Too wet. Very free-draining mix; water less.                                  |

## Notes

|                   |                                                                                                                                       |
|-------------------|---------------------------------------------------------------------------------------------------------------------------------------|
| <b>Difficulty</b> | <b>Moderate–Difficult (provisional).</b> Works on vigorous plants, but the species is small and short-lived. Seed is the main method. |
| <b>Confidence</b> | <b>Medium.</b> ANPSA: divide after flowering and only if the plant is vigorous. Timing and size follow the kangaroo paw genus method. |
| <b>Sources</b>    | See References: 2, 3, 15                                                                                                              |

## Nursery records

Log each batch of divisions here to confirm or update the timing and difficulty above.

| Date divided | Stock plant / provenance | No. pieces | No. established | Success % | Notes |
|--------------|--------------------------|------------|-----------------|-----------|-------|
|              |                          |            |                 |           |       |
|              |                          |            |                 |           |       |
|              |                          |            |                 |           |       |
|              |                          |            |                 |           |       |
|              |                          |            |                 |           |       |

# Anigozanthos manglesii

Red and Green Kangaroo Paw

Family  
Haemodoraceae

| DIFFICULTY | DIVISION TYPE          | BEST MONTHS | DIVISION SIZE        | TIME TO ESTABLISH   |
|------------|------------------------|-------------|----------------------|---------------------|
| Moderate   | Rhizome clump division | Mar – May   | 3+ shoots with roots | 4–6 weeks sheltered |

|          | Jan         | Feb                                                  | Mar  | Apr  | May  | Jun | Jul | Aug                 | Sep | Oct | Nov | Dec |
|----------|-------------|------------------------------------------------------|------|------|------|-----|-----|---------------------|-----|-----|-----|-----|
| Division | –           | OK                                                   | Best | Best | Best | OK  | –   | –                   | OK  | OK  | –   | –   |
|          | Best window | Workable – slower to re-establish or needs more care |      |      |      |     |     | Avoid / low success |     |     |     |     |

Plant snapshot

|            |                                                                                                             |
|------------|-------------------------------------------------------------------------------------------------------------|
| Habit      | Rhizomatous perennial herb, 0.2–1.1 m (flower stems). Often short-lived – best treated as a 2–3 year plant. |
| Flowering  | Aug – Nov (green and red). WA's floral emblem.                                                              |
| Habitat    | White, yellow or grey sand and sandy loam.                                                                  |
| Local note | Recorded in the Wanneroo LGA. Less vigorous and shorter-lived than most garden hybrids.                     |

Propagation options

| Method   | Suitability | Notes                                                            | Details in          |
|----------|-------------|------------------------------------------------------------------|---------------------|
| Cuttings | Not used    | Clumping herb – no stems to cut.                                 | –                   |
| Seed     | Main method | Dry heat or aged seed; autumn sowing.                            | → Seed Sowing Guide |
| Division | Good        | Split vigorous clumps into rhizome pieces with shoots and roots. | This guide          |

Division method

| Item                | Details                                                                                                                                                                                                                                                                                                                                                                 |
|---------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Best time to divide | Mar – May – ANPSA says clumps are best divided in autumn, though other times also work. Feb, Jun and Sep – Oct are workable; summer divisions need protection until re-established. Avoid Jul – Aug (cold, wet) and Nov – Jan (heat).                                                                                                                                   |
| Division type       | Rhizome clump division – the clump is cut into pieces of rhizome, each with leaf shoots and roots.                                                                                                                                                                                                                                                                      |
| Division size       | At least three large shoots per piece gives the best chance of success (ANPSA). Single-shoot pieces with some roots can work but need a humid glasshouse until new roots form.                                                                                                                                                                                          |
| Selecting stock     | Vigorous, healthy clumps, ideally in their second year – this species is often short-lived. Don't divide plants with black leaf spotting (ink disease).                                                                                                                                                                                                                 |
| Preparation         | <ul style="list-style-type: none"><li>Water the stock plant the day before.</li><li>Lift the clump (or knock it out of its pot) and shake or wash off the mix so you can see the rhizome.</li><li>Cut through the rhizome with a clean, sharp knife.</li><li>Trim damaged roots.</li><li>Cut all leaves back by about one-third to reduce water loss (ANPSA).</li></ul> |
| Potting             | Pot into a well-drained potting mix with the shoots at the same level as before (ANPSA). Don't bury the crown. Water in.                                                                                                                                                                                                                                                |
| Environment         | Sheltered, shaded spot until new leaf growth is well advanced and roots show at the drainage holes. Small divisions go in a humid house.                                                                                                                                                                                                                                |
| Time to establish   | Small divisions need about 4–6 weeks in humid conditions (ANPSA); larger pieces re-establish in the shelter over a similar time.                                                                                                                                                                                                                                        |

| Item      | Details                                                                                                                                                                                                                                                                         |
|-----------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Aftercare | <ul style="list-style-type: none"> <li>• Harden off gradually – about a week with nights exposed, then a week in a sheltered spot, then into the open.</li> <li>• Keep foliage dry and airflow good to limit ink disease.</li> <li>• Pot on once roots fill the pot.</li> </ul> |

### Common problems

| Problem                         | Likely cause and fix                                                                             |
|---------------------------------|--------------------------------------------------------------------------------------------------|
| Division rots at the base       | Too wet, too cold, or planted too deep. Pot at the original level; water less; divide in autumn. |
| Black spots spreading on leaves | Ink disease. Discard affected pieces; keep foliage dry; improve airflow.                         |
| Division wilts and dies         | Too few roots or leaves not trimmed. Use larger pieces, trim leaves by a third, keep sheltered.  |

### Notes

|            |                                                                                                                                                            |
|------------|------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Difficulty | <b>Moderate.</b> Kangaroo paws divide well, but this species is shorter-lived than many hybrids, so divide young, healthy clumps. Seed is the main method. |
| Confidence | <b>High.</b> ANPSA publishes a detailed division method for kangaroo paws.                                                                                 |
| Sources    | See References: 1, 2, 3, 5                                                                                                                                 |

### Nursery records

Log each batch of divisions here to confirm or update the timing and difficulty above.

| Date divided | Stock plant / provenance | No. pieces | No. established | Success % | Notes |
|--------------|--------------------------|------------|-----------------|-----------|-------|
|              |                          |            |                 |           |       |
|              |                          |            |                 |           |       |
|              |                          |            |                 |           |       |
|              |                          |            |                 |           |       |
|              |                          |            |                 |           |       |

# Austrostipa elegantissima

Feather Speargrass

Previously *Stipa elegantissima*

Family  
Poaceae

| DIFFICULTY      | DIVISION TYPE              | BEST MONTHS | DIVISION SIZE         | TIME TO ESTABLISH |
|-----------------|----------------------------|-------------|-----------------------|-------------------|
| Easy-Moderate * | Rhizome / tussock division | Aug – Sep   | Clumps of 3-5 tillers | 4-6 weeks         |

|          | Jan         | Feb                                                  | Mar | Apr | May | Jun | Jul | Aug  | Sep  | Oct                 | Nov | Dec |
|----------|-------------|------------------------------------------------------|-----|-----|-----|-----|-----|------|------|---------------------|-----|-----|
| Division | –           | –                                                    | OK  | OK  | OK  | OK  | OK  | Best | Best | OK                  | –   | –   |
|          | Best window | Workable – slower to re-establish or needs more care |     |     |     |     |     |      |      | Avoid / low success |     |     |

## Plant snapshot

|            |                                                                                                           |
|------------|-----------------------------------------------------------------------------------------------------------|
| Habit      | Rhizomatous, tufted, often sprawling perennial grass, 0.35–2 (–3) m.                                      |
| Flowering  | Aug – Dec or Jan (fluffy, feathery green heads).                                                          |
| Habitat    | Sand, loam, clay and granite, occasionally saline.                                                        |
| Local note | Recorded around Perth (Cottesloe, Cambridge, Cockburn); FloraBase has no Joondalup or Wanneroo specimens. |

## Propagation options

| Method   | Suitability | Notes                                | Details in          |
|----------|-------------|--------------------------------------|---------------------|
| Cuttings | Not used    | Grasses are not grown from cuttings. | –                   |
| Seed     | Main method | 1–2 year old seed; smoke-water soak. | → Seed Sowing Guide |
| Division | Good        | Late winter – early spring (genus).  | This guide          |

## Division method

| Item                | Details                                                                                                                                                                                                                                        |
|---------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Best time to divide | <b>Aug – Sep</b> – late winter to early spring gives high success for <i>Austrostipa</i> . Mar – Jul and Oct are workable. Avoid Nov – Feb (heat).                                                                                             |
| Division type       | <b>Rhizome / tussock division</b> – split the clump into pieces of shoots (tillers), each with roots.                                                                                                                                          |
| Division size       | Clumps of <b>3–5 tillers</b> with a good root mass. Very small pieces establish poorly.                                                                                                                                                        |
| Selecting stock     | Healthy, established pot-grown clumps that are actively growing.                                                                                                                                                                               |
| Preparation         | <ul style="list-style-type: none"> <li>Water the stock plant the day before.</li> <li>Knock the clump out and cut it apart with a sharp knife or spade.</li> <li>Cut the stems and leaves back to about 15 cm to reduce water loss.</li> </ul> |
| Potting             | Pot into a free-draining mix at the original depth. Firm in and water well.                                                                                                                                                                    |
| Environment         | Shade house until new leaves appear. Keep the mix moist, not wet.                                                                                                                                                                              |
| Time to establish   | Roughly <b>4–6 weeks</b> . Record your results.                                                                                                                                                                                                |
| Aftercare           | <ul style="list-style-type: none"> <li>Move gradually to full sun once new growth shows.</li> <li>Give tubes room – plants sprawl.</li> </ul>                                                                                                  |

## Common problems

| Problem                    | Likely cause and fix                                                          |
|----------------------------|-------------------------------------------------------------------------------|
| <b>Divisions brown off</b> | Too small, or roots dried out. Keep 3–5 tillers per piece; work in the shade. |
| <b>Base rots</b>           | Too wet or planted too deep.                                                  |
| <b>No new growth</b>       | Divided in summer. Divide in late winter – early spring.                      |

## Notes

|                   |                                                                                                                                          |
|-------------------|------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Difficulty</b> | <b>Easy–Moderate (provisional).</b> Division works well in the genus; untested for this species.                                         |
| <b>Confidence</b> | <b>Medium.</b> No species method published. ANBG reports high success from division of <i>Austrostipa</i> in late winter – early spring. |
| <b>Sources</b>    | See References: 1, 3, 16                                                                                                                 |

## Nursery records

Log each batch of divisions here to confirm or update the timing and difficulty above.

| Date divided | Stock plant / provenance | No. pieces | No. established | Success % | Notes |
|--------------|--------------------------|------------|-----------------|-----------|-------|
|              |                          |            |                 |           |       |
|              |                          |            |                 |           |       |
|              |                          |            |                 |           |       |
|              |                          |            |                 |           |       |
|              |                          |            |                 |           |       |

# Austrostipa flavescens

Coast Speargrass

Family  
Poaceae

| DIFFICULTY                                                                                                                     | DIVISION TYPE    |     |     | BEST MONTHS |      |      | DIVISION SIZE             |     |     | TIME TO ESTABLISH |     |     |
|--------------------------------------------------------------------------------------------------------------------------------|------------------|-----|-----|-------------|------|------|---------------------------|-----|-----|-------------------|-----|-----|
| Moderate *                                                                                                                     | Tussock division |     |     | Apr – Jun   |      |      | Small clumps, 3-5 tillers |     |     | 4-6 weeks         |     |     |
|                                                                                                                                | Jan              | Feb | Mar | Apr         | May  | Jun  | Jul                       | Aug | Sep | Oct               | Nov | Dec |
| Division                                                                                                                       | –                | –   | OK  | Best        | Best | Best | OK                        | –   | –   | –                 | –   | –   |
| <div><div>Best window</div><div>Workable – slower to re-establish or needs more care</div><div>Avoid / low success</div></div> |                  |     |     |             |      |      |                           |     |     |                   |     |     |

Plant snapshot

|            |                                                                              |
|------------|------------------------------------------------------------------------------|
| Habit      | Tufted perennial grass, 0.2–1.2 m.                                           |
| Flowering  | Sep – Oct.                                                                   |
| Habitat    | Grey, red or calcareous sand, sandy loam and limestone; sand dunes.          |
| Local note | Coastal species. Use nursery stock plants rather than digging from bushland. |

Propagation options

| Method   | Suitability | Notes                                          | Details in          |
|----------|-------------|------------------------------------------------|---------------------|
| Cuttings | Not used    | Tufted grass.                                  | –                   |
| Seed     | Main method | Aged seed and smoke water; sow shallow.        | → Seed Sowing Guide |
| Division | Good        | Divide established tussocks into small clumps. | This guide          |

Division method

| Item                | Details                                                                                                                                                                                                                                                                  |
|---------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Best time to divide | Apr – Jun, as autumn rain restarts growth. Mar and Jul are workable. Avoid Aug – Feb (flowering, seed set and summer heat).                                                                                                                                              |
| Division type       | Tussock division – a clump is split into smaller clumps of shoots (tillers), each with roots.                                                                                                                                                                            |
| Division size       | Small clumps of 3-5 tillers with a good root mass. Very small pieces establish poorly.                                                                                                                                                                                   |
| Selecting stock     | Healthy, established pot-grown clumps that are actively growing.                                                                                                                                                                                                         |
| Preparation         | <ul style="list-style-type: none"><li>Water the stock plant the day before.</li><li>Knock the clump out and tease or cut it apart with a sharp knife or spade.</li><li>Trim damaged roots.</li><li>Cut the leaves back to about 10-15 cm to reduce water loss.</li></ul> |
| Potting             | Pot into a free-draining mix at the original depth. Firm in and water well.                                                                                                                                                                                              |
| Environment         | Shade house until new leaves appear. Keep the mix moist, not wet.                                                                                                                                                                                                        |
| Time to establish   | Roughly 4-6 weeks in autumn. Record your results.                                                                                                                                                                                                                        |
| Aftercare           | <ul style="list-style-type: none"><li>Move gradually to full sun once new growth shows.</li><li>Pot on or plant out in winter.</li></ul>                                                                                                                                 |

## Common problems

| Problem                    | Likely cause and fix                                                                  |
|----------------------------|---------------------------------------------------------------------------------------|
| <b>Divisions brown off</b> | Too small, or roots dried out. Keep 3–5 tillers per piece; work quickly in the shade. |
| <b>Base rots</b>           | Too wet or planted too deep. Pot at the original level; water less.                   |
| <b>No new growth</b>       | Divided in the wrong season. Divide as autumn growth starts.                          |

## Notes

|                   |                                                                                                                                                                        |
|-------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Difficulty</b> | <b>Moderate (provisional).</b> Grass clumps usually divide readily, but this species hasn't been tested here. Seed is the main method.                                 |
| <b>Confidence</b> | <b>Low.</b> Division of established clumps works for this grass, but no species-specific method is published. Follows general tussock-grass practice – trial protocol. |
| <b>Sources</b>    | See References: 1, 3                                                                                                                                                   |

## Nursery records

Log each batch of divisions here to confirm or update the timing and difficulty above.

| Date divided | Stock plant / provenance | No. pieces | No. established | Success % | Notes |
|--------------|--------------------------|------------|-----------------|-----------|-------|
|              |                          |            |                 |           |       |
|              |                          |            |                 |           |       |
|              |                          |            |                 |           |       |
|              |                          |            |                 |           |       |
|              |                          |            |                 |           |       |

# Conostylis aculeata

Prickly Conostylis

Family  
Haemodoraceae

| DIFFICULTY  | DIVISION TYPE                                        |     |      |      | BEST MONTHS |     | DIVISION SIZE            |                     |     | TIME TO ESTABLISH   |     |     |
|-------------|------------------------------------------------------|-----|------|------|-------------|-----|--------------------------|---------------------|-----|---------------------|-----|-----|
| Moderate    | Clump division / rooted runners                      |     |      |      | Mar – May   |     | Larger clumps, 5+ shoots |                     |     | 4-8 weeks sheltered |     |     |
|             | Jan                                                  | Feb | Mar  | Apr  | May         | Jun | Jul                      | Aug                 | Sep | Oct                 | Nov | Dec |
| Division    | –                                                    | OK  | Best | Best | Best        | OK  | –                        | –                   | –   | OK                  | OK  | OK  |
| Best window | Workable – slower to re-establish or needs more care |     |      |      |             |     |                          | Avoid / low success |     |                     |     |     |

## Plant snapshot

|            |                                                                                                                                |
|------------|--------------------------------------------------------------------------------------------------------------------------------|
| Habit      | Rhizomatous, tufted perennial, 0.06–0.5 m, with prickly-edged leaves.                                                          |
| Flowering  | Aug – Nov (yellow).                                                                                                            |
| Habitat    | Sand, loam, clay, gravel, limestone or laterite; winter-wet areas and sand dunes.                                              |
| Local note | Recorded in Joondalup and Wanneroo (subsp. <i>cygnorum</i> on the coastal plain). Dieback resistant. Keep subspecies separate. |

## Propagation options

| Method   | Suitability | Notes                                                | Details in          |
|----------|-------------|------------------------------------------------------|---------------------|
| Cuttings | Not used    | Tufted herb.                                         | –                   |
| Seed     | Main method | Smoke water; about 19% in nursery tests.             | → Seed Sowing Guide |
| Division | Good        | Larger divisions, or rooted runners after flowering. | This guide          |

## Division method

| Item                | Details                                                                                                                                                                                                                                                                                                                                   |
|---------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Best time to divide | Mar – May for clump division. Rooted runners are ready <b>just after flowering</b> (about Oct – Dec). Feb and Jun workable. Avoid Jul – Sep for clumps (flowering, cold).                                                                                                                                                                 |
| Division type       | <b>Clump division</b> of the rhizome, or <b>rooted runners</b> : after flowering the runners form aerial roots at the leaf bases.                                                                                                                                                                                                         |
| Division size       | Keep divisions <b>larger than for kangaroo paws</b> – ANPSA notes <i>Conostylis</i> divide less readily and larger pieces do best. Aim for 5 or more shoots with a good piece of rhizome.                                                                                                                                                 |
| Selecting stock     | Vigorous, established nursery clumps. Watch runners after flowering for new roots.                                                                                                                                                                                                                                                        |
| Preparation         | <ul style="list-style-type: none"> <li>• Water the stock plant the day before.</li> <li>• Lift and cut the clump with a sharp knife, keeping pieces large.</li> <li>• <b>Rooted runners</b>: there is only a 2–3 week window after the roots appear – cut and pot them promptly.</li> <li>• Trim leaves back by about a third.</li> </ul> |
| Potting             | Pot into a free-draining mix at the original depth.                                                                                                                                                                                                                                                                                       |
| Environment         | Sheltered, shaded spot (humid house for runners and small pieces) until new growth starts.                                                                                                                                                                                                                                                |
| Time to establish   | About <b>4–8 weeks</b> (record yours).                                                                                                                                                                                                                                                                                                    |
| Aftercare           | <ul style="list-style-type: none"> <li>• Harden off gradually.</li> <li>• Grow on in full sun; don't overwater.</li> </ul>                                                                                                                                                                                                                |

## Common problems

| Problem                       | Likely cause and fix                                                             |
|-------------------------------|----------------------------------------------------------------------------------|
| Small divisions die           | Pieces too small. Keep divisions larger.                                         |
| Runners don't root in the pot | Cut too late or too early. Take them in the 2–3 weeks after aerial roots appear. |
| Base rots                     | Too wet. Free-draining mix; water less.                                          |

## Notes

|            |                                                                                                                                                        |
|------------|--------------------------------------------------------------------------------------------------------------------------------------------------------|
| Difficulty | <b>Moderate.</b> Divides less readily than kangaroo paws; use larger pieces or rooted runners. Seed with smoke is the other option.                    |
| Confidence | <b>Medium.</b> ANPSA notes <i>Conostylis</i> divide less readily than kangaroo paws (larger divisions best); APS NSW describes potting rooted runners. |
| Sources    | See References: 2, 3, 12, 13, 14                                                                                                                       |

## Nursery records

Log each batch of divisions here to confirm or update the timing and difficulty above.

| Date divided | Stock plant / provenance | No. pieces | No. established | Success % | Notes |
|--------------|--------------------------|------------|-----------------|-----------|-------|
|              |                          |            |                 |           |       |
|              |                          |            |                 |           |       |
|              |                          |            |                 |           |       |
|              |                          |            |                 |           |       |
|              |                          |            |                 |           |       |

# Dianella revoluta

Blueberry Lily / Flax Lily

Family  
Asphodelaceae  
(Hemerocallidoideae)

| DIFFICULTY  |  | DIVISION TYPE                                        |     |      |      | BEST MONTHS |     | DIVISION SIZE           |     | TIME TO ESTABLISH |     |     |     |
|-------------|--|------------------------------------------------------|-----|------|------|-------------|-----|-------------------------|-----|-------------------|-----|-----|-----|
| Easy        |  | Rhizome clump division                               |     |      |      | Mar – May   |     | 3-4 shoots with rhizome |     | 4-6 weeks         |     |     |     |
|             |  | Jan                                                  | Feb | Mar  | Apr  | May         | Jun | Jul                     | Aug | Sep               | Oct | Nov | Dec |
| Division    |  | -                                                    | OK  | Best | Best | Best        | OK  | -                       | OK  | OK                | OK  | -   | -   |
| Best window |  | Workable – slower to re-establish or needs more care |     |      |      |             |     | Avoid / low success     |     |                   |     |     |     |

Plant snapshot

|            |                                                                                                    |
|------------|----------------------------------------------------------------------------------------------------|
| Habit      | Rhizomatous, tufted perennial herb, 0.3–1.5 m.                                                     |
| Flowering  | Sep – Dec (blue-purple), followed by blue berries.                                                 |
| Habitat    | White or yellow sand, sandy clay, laterite, loam and clay.                                         |
| Local note | Recorded in Joondalup and Wanneroo. Division is much quicker than seed, which is slow and erratic. |

Propagation options

| Method   | Suitability | Notes                               | Details in          |
|----------|-------------|-------------------------------------|---------------------|
| Cuttings | Not used    | Clumping herb.                      | –                   |
| Seed     | Difficult   | GA + smoke water; 8–11 weeks.       | → Seed Sowing Guide |
| Division | Main method | The quickest way to produce plants. | This guide          |

Division method

| Item                | Details                                                                                                                                                                                                                                                                                   |
|---------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Best time to divide | Mar – May, so divisions root before winter. Feb, Jun and Aug – Oct are workable. Avoid Nov – Jan (heat) and Jul (cold).                                                                                                                                                                   |
| Division type       | Rhizome clump division.                                                                                                                                                                                                                                                                   |
| Division size       | Three or four leaf shoots with a good section of rhizome and healthy roots (ANPSA). Single shoots need a humid house.                                                                                                                                                                     |
| Selecting stock     | Vigorous clumps with plenty of shoots. Keep forms and provenances separate.                                                                                                                                                                                                               |
| Preparation         | <ul style="list-style-type: none"><li>Water the stock plant the day before.</li><li>Lift the clump and shake or wash off the mix.</li><li>Cut the rhizome into pieces with a clean, sharp knife.</li><li>Trim damaged roots and cut leaves back by about one-third to one-half.</li></ul> |
| Potting             | Pot into a free-draining native mix at the original depth. Water in.                                                                                                                                                                                                                      |
| Environment         | Sheltered, shaded spot (humid house for small pieces) until new growth starts.                                                                                                                                                                                                            |
| Time to establish   | About 4–6 weeks in autumn.                                                                                                                                                                                                                                                                |
| Aftercare           | <ul style="list-style-type: none"><li>Harden off gradually – about a week with nights exposed, then a sheltered week.</li><li>Pot on once roots fill the pot.</li></ul>                                                                                                                   |

## Common problems

| Problem                 | Likely cause and fix                                                 |
|-------------------------|----------------------------------------------------------------------|
| <b>Divisions wilt</b>   | Leaves not trimmed or too few roots. Trim harder; use bigger pieces. |
| <b>Rhizome rots</b>     | Too wet or planted too deep. Water less; pot at the original level.  |
| <b>Slow to re-shoot</b> | Divided in cold weather. Divide in autumn or spring.                 |

## Notes

|                   |                                                                                                                               |
|-------------------|-------------------------------------------------------------------------------------------------------------------------------|
| <b>Difficulty</b> | <b>Easy.</b> Clump-forming lilies divide readily, and this is the quickest way to produce plants.                             |
| <b>Confidence</b> | <b>Medium.</b> ANPSA lists <i>Dianella</i> among the grass-like clumps suited to division; details follow the general method. |
| <b>Sources</b>    | See References: 1, 3                                                                                                          |

## Nursery records

Log each batch of divisions here to confirm or update the timing and difficulty above.

| Date divided | Stock plant / provenance | No. pieces | No. established | Success % | Notes |
|--------------|--------------------------|------------|-----------------|-----------|-------|
|              |                          |            |                 |           |       |
|              |                          |            |                 |           |       |
|              |                          |            |                 |           |       |
|              |                          |            |                 |           |       |
|              |                          |            |                 |           |       |

# Ficinia nodosa

Knotted Club Rush

Family  
Cyperaceae (sedges)

| DIFFICULTY  |  | DIVISION TYPE                                        |     |      |      | BEST MONTHS |     | DIVISION SIZE           |     | TIME TO ESTABLISH         |     |     |     |
|-------------|--|------------------------------------------------------|-----|------|------|-------------|-----|-------------------------|-----|---------------------------|-----|-----|-----|
| Easy        |  | Rhizome clump division                               |     |      |      | Mar – May   |     | 6–10 stems with rhizome |     | Plantable in a few months |     |     |     |
|             |  | Jan                                                  | Feb | Mar  | Apr  | May         | Jun | Jul                     | Aug | Sep                       | Oct | Nov | Dec |
| Division    |  | –                                                    | OK  | Best | Best | Best        | OK  | OK                      | OK  | OK                        | OK  | –   | –   |
| Best window |  | Workable – slower to re-establish or needs more care |     |      |      |             |     | Avoid / low success     |     |                           |     |     |     |

Plant snapshot

|            |                                                                                           |
|------------|-------------------------------------------------------------------------------------------|
| Habit      | Erect, clumping, rhizomatous sedge to 1 m high and 0.8 m wide.                            |
| Flowering  | Oct – Dec or Jan (brown knot-like heads).                                                 |
| Habitat    | Calcareous sand and limestone; coastal dunes, seasonally wet swamps and salt-lake shores. |
| Local note | Recorded in Joondalup and Wanneroo. Salt tolerant, fast growing, a strong sand binder.    |

Propagation options

| Method   | Suitability | Notes                                       | Details in          |
|----------|-------------|---------------------------------------------|---------------------|
| Cuttings | Not used    | Sedge.                                      | –                   |
| Seed     | Main method | No pre-treatment; about 55% germination.    | → Seed Sowing Guide |
| Division | Main method | Divides readily; plantable in a few months. | This guide          |

Division method

| Item                | Details                                                                                                                                                                                                                      |
|---------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Best time to divide | Mar – May, so divisions root before winter. Feb and Jun – Oct are workable. Avoid Nov – Jan (heat).                                                                                                                          |
| Division type       | Rhizome clump division.                                                                                                                                                                                                      |
| Division size       | Anything from a short piece of rhizome with <b>6–10 stems</b> up to a 10 × 10 cm clod (NZ restoration guidance).                                                                                                             |
| Selecting stock     | Vigorous nursery clumps or tray-grown stock.                                                                                                                                                                                 |
| Preparation         | <ul style="list-style-type: none"><li>Water the stock plant the day before.</li><li>Lift the clump and cut it into pieces with a sharp knife or spade.</li><li>Trim stems back by about a third to cut water loss.</li></ul> |
| Potting             | Pot into tubes or pots of free-draining mix at the original depth. Water well.                                                                                                                                               |
| Environment         | Shade house until new stems appear; keep the mix moist – it tolerates wetter mix than most.                                                                                                                                  |
| Time to establish   | Divisions are ready for planting <b>within a few months</b> .                                                                                                                                                                |
| Aftercare           | <ul style="list-style-type: none"><li>Harden off, then grow on in full sun.</li><li>Plant out in winter.</li></ul>                                                                                                           |

## Common problems

| Problem                    | Likely cause and fix                                   |
|----------------------------|--------------------------------------------------------|
| <b>Divisions brown off</b> | Pieces dried out. Keep moist and shaded while working. |
| <b>Slow to re-shoot</b>    | Divided in cold weather. Divide in autumn.             |
| <b>Base rots</b>           | Waterlogged in heavy mix. Use a free-draining mix.     |

## Notes

|                   |                                                                                                                    |
|-------------------|--------------------------------------------------------------------------------------------------------------------|
| <b>Difficulty</b> | <b>Easy.</b> Divides readily – transplanted divisions had 100% survival in one NZ planting trial.                  |
| <b>Confidence</b> | <b>Medium.</b> Coastal restoration guidance for this species (NZ); timing follows ANPSA's general division advice. |
| <b>Sources</b>    | See References: 1, 3, 9, 10                                                                                        |

## Nursery records

Log each batch of divisions here to confirm or update the timing and difficulty above.

| Date divided | Stock plant / provenance | No. pieces | No. established | Success % | Notes |
|--------------|--------------------------|------------|-----------------|-----------|-------|
|              |                          |            |                 |           |       |
|              |                          |            |                 |           |       |
|              |                          |            |                 |           |       |
|              |                          |            |                 |           |       |
|              |                          |            |                 |           |       |

# Lepidosperma gladiatum

Coast Sword-sedge

Family  
Cyperaceae (sedges)

| DIFFICULTY  |  | DIVISION TYPE                                        |     |      |      | BEST MONTHS |     | DIVISION SIZE                     |     | TIME TO ESTABLISH    |     |     |     |
|-------------|--|------------------------------------------------------|-----|------|------|-------------|-----|-----------------------------------|-----|----------------------|-----|-----|-----|
| Moderate *  |  | Crown (clump) division                               |     |      |      | Mar – May   |     | Large pieces with rhizome & roots |     | Slow – several weeks |     |     |     |
|             |  | Jan                                                  | Feb | Mar  | Apr  | May         | Jun | Jul                               | Aug | Sep                  | Oct | Nov | Dec |
| Division    |  | –                                                    | OK  | Best | Best | Best        | OK  | OK                                | –   | –                    | –   | –   | –   |
| Best window |  | Workable – slower to re-establish or needs more care |     |      |      |             |     | Avoid / low success               |     |                      |     |     |     |

Plant snapshot

|            |                                                                                                         |
|------------|---------------------------------------------------------------------------------------------------------|
| Habit      | Robust, rhizomatous, tufted sedge, 0.5–1.5 (–3) m, clumps to 1.5 m wide. <b>Knife-sharp leaf edges.</b> |
| Flowering  | Nov – Dec or Jan – May (brown).                                                                         |
| Habitat    | White, grey or calcareous sand, limestone and loam; dunes and creeklines.                               |
| Local note | Recorded in the Wanneroo LGA. Wear heavy gloves and long sleeves when dividing.                         |

Propagation options

| Method   | Suitability | Notes                                             | Details in          |
|----------|-------------|---------------------------------------------------|---------------------|
| Cuttings | Not used    | Sedge.                                            | –                   |
| Seed     | Difficult   | Variable; 41% in one test, 53–72 days.            | → Seed Sowing Guide |
| Division | Possible    | Crown division is used commercially (trial here). | <b>This guide</b>   |

Division method

| Item                | Details                                                                                                                                                                                                                  |
|---------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Best time to divide | Mar – May (trial). Feb, Jun – Jul workable. Avoid Aug – Jan.                                                                                                                                                             |
| Division type       | Crown (clump) division of the rhizome.                                                                                                                                                                                   |
| Division size       | Keep pieces large, with a good section of rhizome and plenty of roots (trial).                                                                                                                                           |
| Selecting stock     | Vigorous nursery clumps. Mark any selected forms.                                                                                                                                                                        |
| Preparation         | <ul style="list-style-type: none"><li>Water the day before.</li><li>Lift the clump; cut through the tough rhizome with a sharp spade or pruning saw.</li><li>Trim leaves back by about half to cut water loss.</li></ul> |
| Potting             | Pot into deep pots of free-draining mix at the original depth. Water well.                                                                                                                                               |
| Environment         | Shade house until new leaves appear.                                                                                                                                                                                     |
| Time to establish   | Slow – allow several weeks; record the result.                                                                                                                                                                           |
| Aftercare           | <ul style="list-style-type: none"><li>Harden off, then grow on in full sun.</li><li>Plant out in late autumn or winter.</li></ul>                                                                                        |

## Common problems

| Problem            | Likely cause and fix                             |
|--------------------|--------------------------------------------------|
| Divisions die back | Too few roots or too small. Keep pieces large.   |
| Hands cut          | Leaf edges are sharp – heavy gloves and sleeves. |
| Slow establishment | Normal. Be patient; full sun once rooted.        |

## Notes

|            |                                                                                                                                                                         |
|------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Difficulty | <b>Moderate (provisional).</b> Used commercially, but untested here. Tissue-cultured stock is also available.                                                           |
| Confidence | <b>Low.</b> Crown division is used commercially (a Joondalup-selected form was first propagated this way) but no method or survival data are published. Trial protocol. |
| Sources    | See References: 3, 11                                                                                                                                                   |

## Nursery records

Log each batch of divisions here to confirm or update the timing and difficulty above.

| Date divided | Stock plant / provenance | No. pieces | No. established | Success % | Notes |
|--------------|--------------------------|------------|-----------------|-----------|-------|
|              |                          |            |                 |           |       |
|              |                          |            |                 |           |       |
|              |                          |            |                 |           |       |
|              |                          |            |                 |           |       |
|              |                          |            |                 |           |       |

# Lomandra maritima

Maritime Mat Rush

Family  
Asparagaceae  
(Lomandroideae)

|                 |                |             |                       |                   |
|-----------------|----------------|-------------|-----------------------|-------------------|
| DIFFICULTY      | DIVISION TYPE  | BEST MONTHS | DIVISION SIZE         | TIME TO ESTABLISH |
| Easy-Moderate * | Clump division | Aug – Sep   | 3–4 shoots with roots | 4–6 weeks         |

|                                                      | Jan | Feb | Mar | Apr | May | Jun | Jul | Aug  | Sep  | Oct | Nov | Dec |
|------------------------------------------------------|-----|-----|-----|-----|-----|-----|-----|------|------|-----|-----|-----|
| Division                                             | –   | –   | OK  | OK  | OK  | OK  | OK  | Best | Best | OK  | –   | –   |
| Best window                                          |     |     |     |     |     |     |     |      |      |     |     |     |
| Workable – slower to re-establish or needs more care |     |     |     |     |     |     |     |      |      |     |     |     |
| Avoid / low success                                  |     |     |     |     |     |     |     |      |      |     |     |     |

Plant snapshot

|            |                                                                                                                  |
|------------|------------------------------------------------------------------------------------------------------------------|
| Habit      | Rhizomatous, tufted perennial, 0.2–0.6 m, clumps to 0.2 m wide. Male and female plants are separate (dioecious). |
| Flowering  | Aug – Nov (purple and yellow).                                                                                   |
| Habitat    | White or grey sand and limestone; coastal dunes and hills.                                                       |
| Local note | Recorded in Joondalup and Wanneroo. A breeding plant for the threatened graceful sun-moth.                       |

Propagation options

| Method   | Suitability | Notes                                                               | Details in |
|----------|-------------|---------------------------------------------------------------------|------------|
| Cuttings | Not used    | Tufted herb.                                                        | –          |
| Seed     | Difficult   | Lomandra seed is strongly dormant; not documented for this species. | –          |
| Division | Main method | Clump division, late winter to early spring (provisional).          | This guide |

Division method

| Item                | Details                                                                                                                                                                                          |
|---------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Best time to divide | Aug – Sep – late winter to early spring is usually best for lomandras. Mar – Jul and Oct are workable. Avoid Nov – Feb.                                                                          |
| Division type       | Clump division.                                                                                                                                                                                  |
| Division size       | Three or four shoots with a piece of rhizome and healthy roots (ANPSA general guidance).                                                                                                         |
| Selecting stock     | Vigorous nursery clumps. Plants are male or female – label stock plants when they flower so plantings include both.                                                                              |
| Preparation         | <ul style="list-style-type: none"><li>Water the day before.</li><li>Lift the clump and cut it apart with a sharp knife.</li><li>Trim the leaves back to about a third of their length.</li></ul> |
| Potting             | Pot into a free-draining mix at the original depth.                                                                                                                                              |
| Environment         | Sheltered, shaded spot until new growth starts.                                                                                                                                                  |
| Time to establish   | About 4–6 weeks (record yours).                                                                                                                                                                  |
| Aftercare           | <ul style="list-style-type: none"><li>Harden off gradually.</li><li>Pot on once roots fill the pot.</li></ul>                                                                                    |

## Common problems

| Problem            | Likely cause and fix                                                                 |
|--------------------|--------------------------------------------------------------------------------------|
| Divisions wilt     | Leaves not trimmed or too few roots. Trim harder; use bigger pieces.                 |
| Base rots          | Too wet or planted too deep.                                                         |
| All plants one sex | Stock came from a single plant. Divide from several labelled male and female plants. |

## Notes

|            |                                                                                                                |
|------------|----------------------------------------------------------------------------------------------------------------|
| Difficulty | <b>Easy-Moderate (provisional).</b> Lomandras generally divide well; untested for this species.                |
| Confidence | <b>Medium.</b> No species method published. Follows general <i>Lomandra</i> and ANPSA clump-division guidance. |
| Sources    | See References: 1, 3, 6, 7, 8                                                                                  |

## Nursery records

Log each batch of divisions here to confirm or update the timing and difficulty above.

| Date divided | Stock plant / provenance | No. pieces | No. established | Success % | Notes |
|--------------|--------------------------|------------|-----------------|-----------|-------|
|              |                          |            |                 |           |       |
|              |                          |            |                 |           |       |
|              |                          |            |                 |           |       |
|              |                          |            |                 |           |       |
|              |                          |            |                 |           |       |

# Poa poiformis

Coastal Poa

Family  
Poaceae

| DIFFICULTY      | DIVISION TYPE    | BEST MONTHS | DIVISION SIZE         | TIME TO ESTABLISH |
|-----------------|------------------|-------------|-----------------------|-------------------|
| Easy-Moderate * | Tussock division | Aug – Sep   | Clumps of 3-5 tillers | 4-6 weeks         |

|                                                      | Jan | Feb | Mar | Apr | May | Jun | Jul | Aug  | Sep  | Oct | Nov | Dec |
|------------------------------------------------------|-----|-----|-----|-----|-----|-----|-----|------|------|-----|-----|-----|
| Division                                             | –   | –   | OK  | OK  | OK  | –   | OK  | Best | Best | OK  | –   | –   |
| Best window                                          |     |     |     |     |     |     |     |      |      |     |     |     |
| Workable – slower to re-establish or needs more care |     |     |     |     |     |     |     |      |      |     |     |     |
| Avoid / low success                                  |     |     |     |     |     |     |     |      |      |     |     |     |

## Plant snapshot

|            |                                                                                                                                                             |
|------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Habit      | Densely tufted perennial tussock grass, 0.15–0.9 m, with fine blue-green leaves.                                                                            |
| Flowering  | Oct – Nov (green to yellow).                                                                                                                                |
| Habitat    | Sand, granite and sandy loam; coastal areas, estuaries and dunes.                                                                                           |
| Local note | Recorded in the Perth area (Cottesloe, Stirling); FloraBase has no Joondalup or Wanneroo specimens. Tolerates salt spray and saline soils once established. |

## Propagation options

| Method   | Suitability | Notes                                 | Details in          |
|----------|-------------|---------------------------------------|---------------------|
| Cuttings | Not used    | Grasses are not grown from cuttings.  | –                   |
| Seed     | Main method | Chill 1-2 weeks; sow at 15°C or more. | → Seed Sowing Guide |
| Division | Good        | Divide mature clumps in early spring. | This guide          |

## Division method

| Item                | Details                                                                                                                                                                                        |
|---------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Best time to divide | Aug – Sep (early spring). Mar – May, Jul and Oct are workable. Avoid Nov – Feb (heat).                                                                                                         |
| Division type       | Tussock division.                                                                                                                                                                              |
| Division size       | Clumps of 3-5 tillers with roots. Very small pieces establish poorly.                                                                                                                          |
| Selecting stock     | Mature, crowded nursery clumps – these benefit from dividing.                                                                                                                                  |
| Preparation         | <ul style="list-style-type: none"> <li>Water the stock plant the day before.</li> <li>Tease or cut the clump apart with a sharp knife.</li> <li>Trim the leaves back by about half.</li> </ul> |
| Potting             | Pot into a free-draining mix at the original depth.                                                                                                                                            |
| Environment         | Shade house until new leaves appear.                                                                                                                                                           |
| Time to establish   | Roughly 4-6 weeks (record yours).                                                                                                                                                              |
| Aftercare           | <ul style="list-style-type: none"> <li>Harden off, then grow on in full sun.</li> <li>Plant out in winter.</li> </ul>                                                                          |

## Common problems

| Problem                    | Likely cause and fix                                             |
|----------------------------|------------------------------------------------------------------|
| <b>Divisions brown off</b> | Pieces too small or dried out. Bigger pieces; work in the shade. |
| <b>Base rots</b>           | Too wet or planted too deep.                                     |
| <b>Slow to re-shoot</b>    | Divided in the heat. Divide in early spring.                     |

## Notes

|                   |                                                                                                                                   |
|-------------------|-----------------------------------------------------------------------------------------------------------------------------------|
| <b>Difficulty</b> | <b>Easy-Moderate (provisional).</b> Tussock grasses usually divide readily; untested for this species.                            |
| <b>Confidence</b> | <b>Low.</b> Only a commercial grower's note: divide mature clumps in early spring. Method follows general tussock-grass practice. |
| <b>Sources</b>    | See References: 1, 3, 17, 18                                                                                                      |

## Nursery records

Log each batch of divisions here to confirm or update the timing and difficulty above.

| Date divided | Stock plant / provenance | No. pieces | No. established | Success % | Notes |
|--------------|--------------------------|------------|-----------------|-----------|-------|
|              |                          |            |                 |           |       |
|              |                          |            |                 |           |       |
|              |                          |            |                 |           |       |
|              |                          |            |                 |           |       |
|              |                          |            |                 |           |       |

# Spinifex hirsutus

Hairy Spinifex

Family  
Poaceae

| DIFFICULTY           | DIVISION TYPE    | BEST MONTHS | DIVISION SIZE              | TIME TO ESTABLISH    |
|----------------------|------------------|-------------|----------------------------|----------------------|
| Moderate-Difficult * | Tussock division | Mar – May   | Rooted clumps with runners | Slow – several weeks |

|          | Jan         | Feb                                                  | Mar  | Apr  | May  | Jun | Jul | Aug                 | Sep | Oct | Nov | Dec |
|----------|-------------|------------------------------------------------------|------|------|------|-----|-----|---------------------|-----|-----|-----|-----|
| Division | OK          | OK                                                   | Best | Best | Best | –   | –   | –                   | OK  | OK  | OK  | OK  |
|          | Best window | Workable – slower to re-establish or needs more care |      |      |      |     |     | Avoid / low success |     |     |     |     |

Plant snapshot

|            |                                                                                                       |
|------------|-------------------------------------------------------------------------------------------------------|
| Habit      | Rhizomatous, tussock-forming grass, 0.3–0.9 m, with long runners and silvery, hairy leaves.           |
| Flowering  | Oct – Jan. Male and female flowers on separate plants.                                                |
| Habitat    | White sand on coastal dunes.                                                                          |
| Local note | Recorded in the City of Joondalup. Dune plants are protected – use nursery stock or approved salvage. |

Propagation options

| Method   | Suitability | Notes                                                              | Details in          |
|----------|-------------|--------------------------------------------------------------------|---------------------|
| Cuttings | Good        | Runner (stolon) sections with 1–2 nodes.                           | → Cuttings Guide    |
| Seed     | Main method | The usual nursery method; cover seed (needs dark).                 | → Seed Sowing Guide |
| Division | Possible    | Tussock divisions work but survived less well than rooted runners. | This guide          |

Division method

| Item                | Details                                                                                                                                                                                         |
|---------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Best time to divide | Mar – May, while growing actively, giving plants ready for winter planting. Sep – Feb workable. Avoid Jun – Aug.                                                                                |
| Division type       | Tussock division – small clumps with rooted runners.                                                                                                                                            |
| Division size       | Small clumps with as much root and rooted runner as possible.                                                                                                                                   |
| Selecting stock     | Vigorous nursery stock plants. Collect male and female plants if you want seed-producing plantings later.                                                                                       |
| Preparation         | <ul style="list-style-type: none"><li>Keep pieces moist and shaded – they dry out fast.</li><li>Cut clumps apart with a sharp spade or knife.</li><li>Trim leaves back by about half.</li></ul> |
| Potting             | Pot into deep tubes of coarse sand or a very sandy mix. Firm in and water well.                                                                                                                 |
| Environment         | Shade house or light shade until new growth shows; keep the sand moist, not wet.                                                                                                                |
| Time to establish   | Slow – allow several weeks for new roots and shoots. Plants take many months to reach planting size.                                                                                            |
| Aftercare           | <ul style="list-style-type: none"><li>Harden off, then grow on in full sun.</li><li>Plant out in late autumn or winter when rain is reliable.</li></ul>                                         |

## Common problems

| Problem                   | Likely cause and fix                                                 |
|---------------------------|----------------------------------------------------------------------|
| Divisions dry out and die | Roots exposed too long. Work in the shade; keep pieces in damp bags. |
| Base rots                 | Mix too wet. Use coarse sand, water less.                            |
| Slow establishment        | Normal for spinifex. Full sun once hardened; deep tubes.             |

## Notes

|            |                                                                                                                                                                  |
|------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Difficulty | <b>Moderate–Difficult (provisional).</b> Divisions survived less well than rooted runner cuttings in trials. Runner cuttings or seed are usually better choices. |
| Confidence | <b>Medium.</b> Based on <i>S. sericeus</i> trials, where divided tussocks survived less well (31%) than rooted runner cuttings (40%).                            |
| Sources    | See References: 3, 4                                                                                                                                             |

## Nursery records

Log each batch of divisions here to confirm or update the timing and difficulty above.

| Date divided | Stock plant / provenance | No. pieces | No. established | Success % | Notes |
|--------------|--------------------------|------------|-----------------|-----------|-------|
|              |                          |            |                 |           |       |
|              |                          |            |                 |           |       |
|              |                          |            |                 |           |       |
|              |                          |            |                 |           |       |
|              |                          |            |                 |           |       |

# Spinifex longifolius

Beach Spinifex

Family  
Poaceae

|                                       |                                   |                          |                                             |                                           |
|---------------------------------------|-----------------------------------|--------------------------|---------------------------------------------|-------------------------------------------|
| DIFFICULTY<br>Moderate-Difficult<br>* | DIVISION TYPE<br>Tussock division | BEST MONTHS<br>Mar – May | DIVISION SIZE<br>Rooted clumps with runners | TIME TO ESTABLISH<br>Slow – several weeks |
|---------------------------------------|-----------------------------------|--------------------------|---------------------------------------------|-------------------------------------------|

|          | Jan         | Feb                                                  | Mar  | Apr  | May  | Jun | Jul | Aug                 | Sep | Oct | Nov | Dec |
|----------|-------------|------------------------------------------------------|------|------|------|-----|-----|---------------------|-----|-----|-----|-----|
| Division | OK          | OK                                                   | Best | Best | Best | –   | –   | –                   | OK  | OK  | OK  | OK  |
|          | Best window | Workable – slower to re-establish or needs more care |      |      |      |     |     | Avoid / low success |     |     |     |     |

Plant snapshot

|            |                                                                                                  |
|------------|--------------------------------------------------------------------------------------------------|
| Habit      | Rhizomatous, tussock-forming grass, 0.3–1 m high and up to 2 m wide, with long runners.          |
| Flowering  | Apr – Dec or Jan. Male and female plants are separate.                                           |
| Habitat    | White sand on coastal dunes; a key foredune stabiliser.                                          |
| Local note | Recorded in the Wanneroo LGA. Dune plants are protected – use nursery stock or approved salvage. |

Propagation options

| Method   | Suitability | Notes                                                              | Details in          |
|----------|-------------|--------------------------------------------------------------------|---------------------|
| Cuttings | Good        | Runner (stolon) sections with 1–2 nodes.                           | → Cuttings Guide    |
| Seed     | Main method | The usual nursery method; smoke water, about 10 days.              | → Seed Sowing Guide |
| Division | Possible    | Tussock divisions work but survived less well than rooted runners. | This guide          |

Division method

| Item                | Details                                                                                                                                                                                         |
|---------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Best time to divide | Mar – May, while growing actively, giving plants ready for winter planting. Sep – Feb workable. Avoid Jun – Aug.                                                                                |
| Division type       | Tussock division – small clumps with rooted runners.                                                                                                                                            |
| Division size       | Small clumps with as much root and rooted runner as possible.                                                                                                                                   |
| Selecting stock     | Vigorous nursery stock plants.                                                                                                                                                                  |
| Preparation         | <ul style="list-style-type: none"><li>Keep pieces moist and shaded – they dry out fast.</li><li>Cut clumps apart with a sharp spade or knife.</li><li>Trim leaves back by about half.</li></ul> |
| Potting             | Pot into deep tubes of coarse sand or a very sandy mix. Firm in and water well.                                                                                                                 |
| Environment         | Shade house or light shade until new growth shows; keep the sand moist, not wet.                                                                                                                |
| Time to establish   | Slow – allow several weeks for new roots and shoots.                                                                                                                                            |
| Aftercare           | <ul style="list-style-type: none"><li>Harden off, then grow on in full sun.</li><li>Plant out in late autumn or winter when rain is reliable.</li></ul>                                         |

## Common problems

| Problem                   | Likely cause and fix                                                 |
|---------------------------|----------------------------------------------------------------------|
| Divisions dry out and die | Roots exposed too long. Work in the shade; keep pieces in damp bags. |
| Base rots                 | Mix too wet. Use coarse sand, water less.                            |
| Slow establishment        | Normal for spinifex. Full sun once hardened; deep tubes.             |

## Notes

|            |                                                                                                                                                                  |
|------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Difficulty | <b>Moderate-Difficult (provisional).</b> Divisions survived less well than rooted runner cuttings in trials. Runner cuttings or seed are usually better choices. |
| Confidence | <b>Medium.</b> Based on <i>S. sericeus</i> trials, where divided tussocks survived less well (31%) than rooted runner cuttings (40%).                            |
| Sources    | See References: 3, 4                                                                                                                                             |

## Nursery records

Log each batch of divisions here to confirm or update the timing and difficulty above.

| Date divided | Stock plant / provenance | No. pieces | No. established | Success % | Notes |
|--------------|--------------------------|------------|-----------------|-----------|-------|
|              |                          |            |                 |           |       |
|              |                          |            |                 |           |       |
|              |                          |            |                 |           |       |
|              |                          |            |                 |           |       |
|              |                          |            |                 |           |       |

# References

Numbers match the source list on each species page. Web addresses shortened; accessed September–October 2026.

| #  | Source                                                                                                                                                                       | Address                                                                                                                                                                                                 |
|----|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| 1  | Australian Native Plants Society (ANPSA) – Plant Propagation: division and layering (grass-like clumps, suckers, self-layering plants; division size, timing, hardening off) | <a href="https://anpsa.org.au/plant-propagation/">anpsa.org.au/plant-propagation/</a>                                                                                                                   |
| 2  | ANPSA – Kangaroo Paws & Relatives: propagation by division (autumn, 3+ shoots, trim leaves by a third)                                                                       | <a href="https://anpsa.org.au/genera/kangaroo-paw-family/">anpsa.org.au/genera/kangaroo-paw-family/</a>                                                                                                 |
| 3  | FloraBase, WA Herbarium (DBCA) – species profiles (descriptions, flowering, habitat, local records)                                                                          | <a href="https://florabase.dbca.wa.gov.au">florabase.dbca.wa.gov.au</a>                                                                                                                                 |
| 4  | NZ Department of Conservation – Rehabilitation of coastal foredunes using indigenous sand-binding species ( <i>Spinifex sericeus</i> division vs stolon cuttings)            | <a href="https://doc.govt.nz/Documents/science-and-technical/sfc122.pdf">doc.govt.nz/Documents/science-and-technical/sfc122.pdf</a>                                                                     |
| 5  | Wikipedia – <i>Anigozanthos manglesii</i> (short-lived habit, ink disease)                                                                                                   | <a href="https://en.wikipedia.org/wiki/Anigozanthos_manglesii">en.wikipedia.org/wiki/Anigozanthos_manglesii</a>                                                                                         |
| 6  | Ozbreed – How to divide <i>Lomandras</i> (late winter to early spring; trim leaves)                                                                                          | <a href="https://ozbreed.com.au/how-to-divide-lomandras/">ozbreed.com.au/how-to-divide-lomandras/</a>                                                                                                   |
| 7  | Cottesloe Coastcare Association – <i>Lomandra maritima</i> (separate male and female plants)                                                                                 | <a href="https://cottesloecoastcare.org/lomandra-maritima/">cottesloecoastcare.org/lomandra-maritima/</a>                                                                                               |
| 8  | DBCA Library – graceful sun-moth breeds on <i>Lomandra maritima</i> and <i>L. hermaphrodita</i>                                                                              | <a href="https://library.dbca.wa.gov.au/static/Journals/080525/080525-41.pdf">library.dbca.wa.gov.au/static/Journals/080525/080525-41.pdf</a>                                                           |
| 9  | Coastal Restoration Trust of NZ – <i>Ficinia nodosa</i> (wīwī) technical note: division size, survival                                                                       | <a href="https://coastalrestorationtrust.org.nz/site/assets/files/1185/8.4_wiwi.pdf">coastalrestorationtrust.org.nz/site/assets/files/1185/8.4_wiwi.pdf</a>                                             |
| 10 | NZ Plant Conservation Network – <i>Ficinia nodosa</i> (easily grown from seed and division)                                                                                  | <a href="https://nzpcn.org.nz/flora/species/ficinia-nodosa/">nzpcn.org.nz/flora/species/ficinia-nodosa/</a>                                                                                             |
| 11 | US Plant Patent PP30,800 – <i>Lepidosperma gladiatum</i> 'LEP01' (first propagated by crown division in Joondalup; tissue culture)                                           | <a href="https://patents.google.com/patent/USPP30800">patents.google.com/patent/USPP30800</a>                                                                                                           |
| 12 | Australian Plants Society NSW – <i>Conostylis aculeata</i> (rooted runners after flowering; 2–3 week window)                                                                 | <a href="https://resources.austplants.com.au/plant/conostylis-aculeata/">resources.austplants.com.au/plant/conostylis-aculeata/</a>                                                                     |
| 13 | Gardening With Angus – <i>Conostylis aculeata</i> (division of rhizomes)                                                                                                     | <a href="https://gardeningwithangus.com.au/conostylis-aculeata-prickly-conostylis/">gardeningwithangus.com.au/conostylis-aculeata-prickly-conostylis/</a>                                               |
| 14 | Murdoch University CPSM (2009) – Native garden plants resistant to dieback ( <i>Conostylis aculeata</i> )                                                                    | <a href="https://cpsm-phytophthora.org/downloads/CPSM_resistanceBrochures.pdf">cpsm-phytophthora.org/downloads/CPSM_resistanceBrochures.pdf</a>                                                         |
| 15 | ANPSA – <i>Anigozanthos humilis</i> profile (divide after flowering, only vigorous plants; short-lived; rust)                                                                | <a href="https://anpsa.org.au/plant_profiles/anigozanthos-humilis/">anpsa.org.au/plant_profiles/anigozanthos-humilis/</a>                                                                               |
| 16 | Australian National Botanic Gardens – <i>Austrostipa ramosissima</i> (division in late winter – early spring has high success)                                               | <a href="https://anbg.gov.au/gnp/trainees-2016/austrostipa-ramosissima.html">anbg.gov.au/gnp/trainees-2016/austrostipa-ramosissima.html</a>                                                             |
| 17 | Peppy Plants – <i>Poa poiformis</i> (divide mature clumps in early spring; commercial grower, low authority)                                                                 | <a href="https://peppyplants.com.au/poa-poiformis-the-native-ornamental-and-adaptable-blue-tussock-grass/">peppyplants.com.au/poa-poiformis-the-native-ornamental-and-adaptable-blue-tussock-grass/</a> |
| 18 | Australian Plants Online – <i>Poa poiformis</i> (salt spray and saline soils)                                                                                                | <a href="https://australianplantsonline.com.au/poa-poiformis-coastal-tussock-grass.html">australianplantsonline.com.au/poa-poiformis-coastal-tussock-grass.html</a>                                     |

## Revision log

| Version | Date     | Changes                                                                                                                                                                                                               |
|---------|----------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| 1.0     | Sep 2026 | First edition: Anigozanthos manglesii, Austrostipa flavescens, Dianella revoluta, Spinifex hirsutus, S. longifolius. Includes a list of other species in the Cuttings and Seed guides that can be divided or layered. |
| 1.1     | Sep 2026 | Added 4 species: Conostylis aculeata, Ficinia nodosa, Lepidosperma gladiatum, Lomandra maritima.                                                                                                                      |
| 1.2     | Sep 2026 | Added 3 species: Anigozanthos humilis, Austrostipa elegantissima, Poa poiformis.                                                                                                                                      |
| 1.3     | Oct 2026 | Added Orthrosanthus laxus and Patersonia occidentalis to 'Other species that can be divided or layered'.                                                                                                              |
| 1.4     | Oct 2026 | Added Atriplex cinerea (layering) and Tricoryne elatior to 'Other species that can be divided or layered'.                                                                                                            |
|         |          |                                                                                                                                                                                                                       |
|         |          |                                                                                                                                                                                                                       |
|         |          |                                                                                                                                                                                                                       |
|         |          |                                                                                                                                                                                                                       |