NURSERY PROPAGATION REFERENCE

Native Plant Cuttings Guide

Western Australian native species – Joondalup, WA

Version 1.9 · October 2026 · 51 species

Species in this edition (alphabetical)

Full index on page 2

- *Acrotriche cordata* * 9
- *Anthocercis littorea* 11
- *Atriplex cinerea* 13
- *Atriplex isatidea* * 15
- *Baeckea robusta* * 17
- *Calothamnus quadrifidus* * 19
- *Carpobrotus virescens* 21
- *Conostephium preissii* * 23
- *Diplolaena angustifolia* * 25
- *Eremophila glabra* 27
- *Grevillea crithmifolia* * 29
- *Grevillea preissii* 31
- *Grevillea vestita* 33
- *Guichenotia ledifolia* * 35
- *Hardenbergia comptoniana* 37
- *Hemiandra pungens* 39
- *Hibbertia cuneiformis* * 41
- *Hibbertia hypericoides* 43
- *Hibbertia racemosa* * 45
- *Hibbertia spicata* * 47
- *Hypocalymma robustum* 49
- *Kennedia prostrata* 51
- *Lechenaultia linarioides* * 53
- *Leucophyta brownii* * 55
- *Leucopogon parviflorus* * 57
- *Melaleuca cardiophylla* * 59
- *Melaleuca huegelii* * 61
- *Melaleuca lanceolata* * 63
- *Melaleuca systena* * 65
- *Myoporum caprarioides* * 67
- *Myoporum insulare* 69
- *Olearia axillaris* * 71
- *Phyllanthus calycinus* * 73
- *Pimelea argentea* * 75
- *Pimelea calcicola* * 77
- *Pimelea sulphurea* * 79
- *Pithocarpa cordata* * 81
- *Rhagodia baccata* * 83
- *Scaevola canescens* * 85
- *Scaevola crassifolia* 87
- *Scaevola globulifera* * 89
- *Scaevola nitida* * 91
- *Scaevola repens* * 93
- *Scaevola thesioides* * 95
- *Spinifex hirsutus* * 97
- *Spinifex longifolius* * 99
- *Styphelia insularis* * 101
- *Synaphea spinulosa* 103
- *Templetonia retusa* * 105
- *Thomasia triphylla* * 107
- *Threlkeldia diffusa* * 109

Easy · Easy–Mod. · Moderate · Mod.–Difficult · Difficult

* Provisional rating – based on genus behaviour; confirm with your own nursery records.

Compiled from published sources (see References). Windows and ratings are starting points for local practice.

© 2026 Mitchell Donald. All rights reserved.

How to use this guide

Each species has its own page set, listed alphabetically by scientific name. Every page follows the same layout: quick facts, a cutting calendar, a plant snapshot, the full propagation method, common problems, and a blank record table so the nursery can log strike rates and refine the advice over time.

Timing windows are written for the Joondalup / Perth coastal climate: hot, dry summers and mild, wet winters. They combine general native-propagation practice with each species' flowering period from FloraBase. Rooting is fastest in warm weather and slows markedly in winter unless bottom heat is used.

Species index – at a glance

Species	Page	Difficulty	Wood type	Best months	Length	Hormone (IBA)
Acrotriche cordata Coast Ground Berry	9	Difficult *	Semi-hardwood	**Jan – Mar**	50–100 mm	3 g/L IBA
Anthocercis littorea Yellow Tailflower	11	Easy–Moderate	Semi-hardwood	**Dec – Apr**	75–100 mm	3 g/L IBA
Atriplex cinerea Grey Saltbush	13	Easy	Semi-hardwood tips	**Nov – Mar**	100–150 mm	1.5–3 g/L IBA
Atriplex isatidea Coast Saltbush	15	Easy–Moderate *	Semi-hardwood tips	**Sep – Nov**	100–150 mm	1.5–3 g/L IBA
Baeckea robusta	17	Easy–Moderate *	Semi-hardwood	**Jan – Mar**	50–75 mm	3 g/L IBA
Calothamnus quadrifidus One-sided Bottlebrush	19	Easy–Moderate *	Semi-hardwood	**Jan – Apr**	75–100 mm	3 g/L IBA
Carpobrotus virescens Coastal Pigface	21	Easy	Succulent stem pieces	**Sep – Nov; Mar – Apr**	10–15 cm (tubes)	Not needed
Conostephium preissii	23	Difficult *	Semi-hardwood	**Jan – Mar**	80–150 mm	3 g/L IBA
Diplolaena angustifolia Yanchep Rose	25	Moderate *	Semi-hardwood	**Dec – Mar**	75–100 mm	3 g/L IBA
Eremophila glabra Tar Bush / Common Emu Bush	27	Easy	Semi-hardwood	**Jan – Apr**	75–100 mm	3 g/L IBA
Grevillea crithmifolia	29	Moderate *	Semi-hardwood	**Dec – Apr**	75–100 mm	3 g/L IBA
Grevillea preissii Spider-net Grevillea	31	Moderate	Semi-hardwood	**Nov – Mar**	75–100 mm	3 g/L IBA
Grevillea vestita	33	Easy	Semi-hardwood	**Dec – Apr**	80–120 mm	3 g/L IBA
Guichenotia ledifolia	35	Easy–Moderate *	Semi-hardwood	**Dec – Apr**	50–75 mm	3 g/L IBA
Hardenbergia comptoniana Native Wisteria	37	Easy–Moderate	Semi-hardwood	**Dec – Mar**	50–75 mm	3 g/L IBA
Hemiandra pungens Snakebush	39	Easy	Softwood – semi-hardwood	**Oct – Mar**	50–75 mm	1.5–3 g/L IBA
Hibbertia cuneiformis Cutleaf Hibbertia	41	Easy–Moderate *	Semi-hardwood	**Mar – May**	75–100 mm	3 g/L IBA
Hibbertia hypericoides Yellow Buttercups	43	Difficult	Semi-hardwood	**Dec – Feb**	50–75 mm	3 g/L IBA (trial range)
Hibbertia racemosa Stalked Guinea Flower	45	Moderate *	Semi-hardwood	**Jan – Apr**	50–75 mm	1.5–3 g/L IBA

© 2026 Mitchell Donald. All rights reserved.

Species	Page	Difficulty	Wood type	Best months	Length	Hormone (IBA)
Hibbertia spicata	47	Moderate *	Semi-hardwood	**Jan – Mar**	50–75 mm	1.5–3 g/L IBA
Hypocalymma robustum Swan River Myrtle	49	Easy	Semi-hardwood	**Dec – Mar**	75–100 mm	3 g/L IBA
Kennedia prostrata Scarlet Runner / Running Postman	51	Easy–Moderate	Semi-hardwood	**Dec – Mar**	75–100 mm	3 g/L IBA
Lechenaultia linarioides Yellow Leschenaultia	53	Easy *	Softwood – semi-hardwood	**Feb – Apr**	50–75 mm	1.5–3 g/L IBA
Leucophyta brownii Cushion Bush	55	Easy–Moderate *	Semi-hardwood	**Mar – May; Sep – Oct**	50–75 mm	1.5–3 g/L IBA
Leucopogon parviflorus Coast Beard-heath	57	Difficult *	Semi-hardwood	**Nov – Jan**	80–100 mm	3 g/L IBA
Melaleuca cardiophylla Tangling Melaleuca	59	Easy *	Semi-hardwood	**Feb – Apr**	75–100 mm	3 g/L IBA
Melaleuca huegelii Chenille Honeymyrtle	61	Easy *	Semi-hardwood	**Feb – Apr**	75–100 mm	3 g/L IBA
Melaleuca lanceolata Rottnest Teatree / Moonah	63	Easy *	Semi-hardwood	**Mar – Apr**	75–100 mm	3 g/L IBA
Melaleuca systena Coastal Honeymyrtle	65	Easy *	Semi-hardwood	**Dec – Jan; Apr – May**	75–100 mm	3 g/L IBA
Myoporum caprarioides Slender Myoporum	67	Easy *	Semi-hardwood	**Nov – Mar**	75–100 mm	3 g/L IBA
Myoporum insulare Blueberry Tree / Common Boobialla	69	Easy	Semi-hardwood	**Feb – Apr**	75–100 mm	3 g/L IBA
Olearia axillaris Coastal Daisybush	71	Easy *	Semi-hardwood	**Aug – Oct**	75–100 mm	1.5–3 g/L IBA
Phyllanthus calycinus False Boronia	73	Moderate *	Semi-hardwood	**Feb – Apr**	50–75 mm	3 g/L IBA
Pimelea argentea Silvery-leaved Pimelea	75	Moderate *	Semi-hardwood	**Mar – May**	30–40 mm	3 g/L IBA
Pimelea calcicola Coastal Banjine	77	Moderate *	Semi-hardwood	**Dec – Mar**	50–75 mm	1.5–3 g/L IBA
Pimelea sulphurea Yellow Banjine	79	Moderate *	Semi-hardwood	**Dec – Mar**	50–75 mm	1.5–3 g/L IBA
Pithocarpa cordata Tangle Daisy	81	Easy–Moderate *	Semi-hardwood tips	**Sep – Nov; May**	50–75 mm	1.5–3 g/L IBA
Rhagodia baccata Berry Saltbush	83	Moderate *	Semi-hardwood tips	**Sep – Nov**	75–100 mm	3 g/L IBA
Scaevola canescens Grey Scaevola	85	Easy–Moderate *	Softwood – semi-hardwood	**Nov – Jan**	50–80 mm	1.5–3 g/L IBA
Scaevola crassifolia Thick-leaved Fan-flower	87	Easy	Semi-hardwood	**Mar – May**	75–100 mm	3 g/L IBA
Scaevola globulifera	89	Easy *	Softwood – semi-hardwood	**Jan – Apr**	50–80 mm	1.5–3 g/L IBA
Scaevola nitida Shining Fanflower	91	Easy *	Semi-hardwood	**Jan – Apr**	75–100 mm	3 g/L IBA
Scaevola repens	93	Easy *	Softwood – semi-hardwood	**Mar – May**	50–80 mm	1.5–3 g/L IBA

© 2026 Mitchell Donald. All rights reserved.

Species	Page	Difficulty	Wood type	Best months	Length	Hormone (IBA)
Scaevola thesioides	95	Easy–Moderate *	Softwood – semi-hardwood	**Apr – May**	50–75 mm	1.5–3 g/L IBA
Spinifex hirsutus Hairy Spinifex	97	Moderate *	Runner (stolon) sections	**Mar – May**	10–15 cm, 1–2 nodes	Optional (1.5 g/L)
Spinifex longifolius Beach Spinifex	99	Moderate *	Runner (stolon) sections	**Mar – May**	10–15 cm, 1–2 nodes	Optional (1.5 g/L)
Styphelia insularis	101	Difficult *	Semi-hardwood	**Oct – Dec**	50–80 mm	3 g/L IBA
Synaphea spinulosa	103	Difficult	Semi-hardwood	**Feb – Apr**	50–75 mm	3–8 g/L IBA (trial)
Templetonia retusa Cockies Tongues	105	Easy–Moderate *	Semi-hardwood	**Dec – Mar**	50–75 mm	3 g/L IBA
Thomasia triphylla	107	Easy–Moderate *	Semi-hardwood	**Dec – Apr**	50–75 mm	3 g/L IBA
Threlkeldia diffusa Coast Bonefruit	109	Moderate *	Softwood tips	**Dec – Mar**	50–75 mm	1.5 g/L IBA

* Provisional rating – based on genus behaviour; update once nursery strike rates are recorded.

© 2026 Mitchell Donald. All rights reserved.

Cutting calendar

Species	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Acrotriche cordata	Best	Best	Best	OK	OK	–	–	–	–	OK	OK	OK
Anthocercis littorea	Best	Best	Best	Best	OK	–	–	–	–	–	OK	Best
Atriplex cinerea	Best	Best	Best	OK	OK	–	–	–	OK	OK	Best	Best
Atriplex isatidea	OK	OK	–	–	–	–	–	OK	Best	Best	Best	OK
Baeckea robusta	Best	Best	Best	OK	OK	–	–	–	–	–	–	OK
Calothamnus quadrifidus	Best	Best	Best	Best	OK	–	–	–	–	–	–	OK
Carpobrotus virescens	OK	OK	Best	Best	OK	–	–	OK	Best	Best	Best	OK
Conostephium preissii	Best	Best	Best	OK	OK	–	–	–	–	–	–	OK
Diplolaena angustifolia	Best	Best	Best	OK	OK	–	–	–	–	–	OK	Best
Eremophila glabra	Best	Best	Best	Best	OK	–	–	–	–	–	OK	OK
Grevillea crithmifolia	Best	Best	Best	Best	OK	–	–	–	–	–	OK	Best
Grevillea preissii	Best	Best	Best	OK	–	–	–	–	–	OK	Best	Best
Grevillea vestita	Best	Best	Best	Best	OK	–	–	–	–	–	OK	Best
Guichenotia ledifolia	Best	Best	Best	Best	OK	–	–	–	–	–	OK	Best
Hardenbergia comptoniana	Best	Best	Best	OK	–	–	–	–	–	–	OK	Best
Hemiandra pungens	Best	Best	Best	OK	–	–	–	–	OK	Best	Best	Best
Hibbertia cuneiformis	OK	OK	Best	Best	Best	–	–	–	–	OK	OK	OK
Hibbertia hypericoides	Best	Best	OK	–	–	–	–	–	–	–	OK	Best
Hibbertia racemosa	Best	Best	Best	Best	OK	–	–	–	–	–	–	OK
Hibbertia spicata	Best	Best	Best	OK	–	–	–	–	–	–	–	OK
Hypocalymma robustum	Best	Best	Best	OK	–	–	–	–	–	–	OK	Best
Kennedia prostrata	Best	Best	Best	OK	–	–	–	–	–	–	OK	Best
Lechenaultia linarioides	OK	Best	Best	Best	OK	–	–	–	–	OK	OK	OK
Leucophyta brownii	–	OK	Best	Best	Best	OK	–	OK	Best	Best	OK	–
Leucopogon parviflorus	Best	OK	OK	OK	OK	–	–	–	–	–	Best	Best
Melaleuca cardiophylla	OK	Best	Best	Best	OK	–	–	–	–	–	–	OK
Melaleuca huegelii	OK	Best	Best	Best	OK	–	–	–	–	–	–	OK
Melaleuca lanceolata	–	OK	Best	Best	OK	–	–	–	OK	OK	–	–
Melaleuca systena	Best	OK	OK	Best	Best	OK	–	–	–	–	OK	Best
Myoporum caprarioides	Best	Best	Best	OK	–	–	–	–	–	OK	Best	Best
Myoporum insulare	OK	Best	Best	Best	OK	–	–	–	–	OK	OK	OK
Olearia axillaris	–	–	OK	OK	–	–	OK	Best	Best	Best	OK	–
Phyllanthus calycinus	OK	Best	Best	Best	OK	–	–	–	–	–	–	–
Pimelea argentea	–	OK	Best	Best	Best	OK	OK	OK	–	–	–	–

© 2026 Mitchell Donald. All rights reserved.

Pimelea calcicola	Best	Best	Best	OK	–	–	–	–	–	–	OK	Best
Pimelea sulphurea	Best	Best	Best	OK	–	–	–	–	–	–	OK	Best
Pithocarpa cordata	–	–	–	OK	Best	OK	–	OK	Best	Best	Best	OK
Rhagodia baccata	OK	OK	OK	–	–	–	–	OK	Best	Best	Best	OK
Scaevola canescens	Best	OK	OK	OK	OK	–	–	–	–	OK	Best	Best
Scaevola crassifolia	OK	OK	Best	Best	Best	–	–	–	–	OK	OK	OK
Scaevola globulifera	Best	Best	Best	Best	OK	–	–	–	–	–	OK	OK
Scaevola nitida	Best	Best	Best	Best	OK	–	–	–	–	–	OK	OK
Scaevola repens	OK	OK	Best	Best	Best	–	–	–	OK	OK	OK	OK
Scaevola thesioides	OK	OK	OK	Best	Best	–	–	–	OK	OK	OK	OK
Spinifex hirsutus	OK	OK	Best	Best	Best	–	–	–	OK	OK	OK	OK
Spinifex longifolius	OK	OK	Best	Best	Best	–	–	–	OK	OK	OK	OK
Styphelia insularis	OK	OK	–	–	–	–	–	–	OK	Best	Best	Best
Synaphea spinulosa	OK	Best	Best	Best	OK	–	–	–	–	–	–	OK
Templetonia retusa	Best	Best	Best	OK	–	–	–	–	–	–	OK	Best
Thomasia triphylla	Best	Best	Best	Best	OK	–	–	–	–	–	OK	Best
Threlkeldia diffusa	Best	Best	Best	OK	–	–	–	–	–	–	OK	Best
Perth season	Summer			Autumn			Winter			Spring		Sum.

Best window Workable – see ‘Selecting material’ in the method Avoid / low success

© 2026 Mitchell Donald. All rights reserved.

Key to terms and ratings

Wood types and the bend test

Wood type	What it looks like	Bend test	Use
Softwood	Soft, flexible, bright new growth from the current flush.	Flops or stays bent.	Roots fast but wilts and rots easily. Suits soft-stemmed plants such as *Scaevola*.
Semi-hardwood	Current season's growth that has begun to firm; bark starting to colour.	Bends about 60° and springs back.	The standard for most WA native shrubs. More forgiving than softwood.
Hardwood	Fully mature, woody stems from previous seasons.	Stiff; snaps rather than bends.	Slow to root. Rarely needed for the species in this guide.

Difficulty ratings

Rating	Meaning
Easy	Strikes reliably with the standard method.
Easy–Moderate	Usually strikes; one factor (e.g. moisture, fragile stems) needs extra care.
Moderate	Strikes, but results depend on timing, material and moisture control. Set extra cuttings.
Moderate–Difficult	Low or variable strike; needs careful timing and good facilities.
Difficult	Low or erratic strike; specialist techniques or alternative methods usually needed.

* **Provisional** – rating is based on how the genus generally behaves because little species-specific data is published. Replace it with the nursery's own figure once a few batches have been recorded.

Confidence

Level	Meaning
High	Species-specific propagation information is published and consistent.
Medium	Built from genus-level information plus the species' own traits (habit, leaf type, flowering time). Sound starting point – verify with trials.
Low	Little information available; treat as a trial protocol.

© 2026 Mitchell Donald. All rights reserved.

Standard cutting method

Species pages refer back to these steps and only note where a species differs.

Step	What to do
1. Collect	Take material early in the morning from healthy, pest-free, well-watered, **non-flowering** stock. Seal in a moist plastic bag and keep in an esky out of the sun. Process the same day.
2. Prepare	Use a sharp, clean blade (wipe with alcohol between stock plants). Cut just below a node. Strip leaves from the lower half to two-thirds without tearing the bark. Reduce large leaves by about one-third. Remove flowers and buds.
3. Wound (optional)	On firmer stems, remove a thin sliver of bark from the bottom 15–20 mm to expose more rooting area.
4. Hormone	Decant a small amount into a clean container, dip the base to about 1 cm, and discard what's left. Store hormone in the fridge – it has a limited shelf life.
5. Set	Make a hole with a dibber so the hormone isn't wiped off, insert to the bottom of the hole, and firm the mix. Single tubes make potting on easier than community pots.
6. Environment	Water in and place in a humid frame or propagation house in bright shade. Bottom heat so the base is 5–10°C warmer than the air. Keep the mix moist, not wet. Mist sparingly on hairy-leaved species.
7. Check	Most cuttings strike in 4–8 weeks in warm weather. Look for roots at the drainage holes. Remove any rotting cuttings promptly.
8. Harden off	About 2 weeks with cover closed by day and open at night, then about 2 weeks in shade, then into the open. Many struck cuttings are lost by skipping this step.
9. Pot on	After 1–3 months, pot on into larger containers. Don't let roots become pot-bound.

Rooting hormone guide

Product (example)	IBA strength	Use for
Clonex Green	1.5 g/L	Softwood and soft tip cuttings
Clonex Purple	3 g/L	Semi-hardwood – the general all-rounder
Clonex Red	8 g/L	Hardwood cuttings

Equivalent powders or liquids work too – match the strength to the wood type. Callus forming without roots usually means the hormone is too strong for that material.

Mix, hygiene and collecting

Topic	Guidance
Propagation mix	About 75–80% washed coarse river sand with 20–25% coir or peat. Add extra perlite or coarse sand for rot-prone species.
Hygiene	Fresh mix, clean tubes and benches. Many WA natives (including *Hibbertia* and *Grevillea*) are susceptible to *Phytophthora*. Discard diseased cuttings; use fungicide only if a problem persists.
Collecting	Collecting from bushland generally requires a DBCA flora licence and landholder permission. Use local-provenance stock and label every batch with its source.

© 2026 Mitchell Donald. All rights reserved.

Acrotriche cordata

Coast Ground Berry

Family **Ericaceae**

DIFFICULTY	WOOD TYPE	BEST MONTHS	CUTTING LENGTH	HORMONE (IBA)
Difficult *	Semi-hardwood	Jan – Mar	50–100 mm	3 g/L IBA

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Cuttings	Best	Best	Best	OK	OK	–	–	–	–	OK	OK	OK

Best window | Workable – see 'Selecting material' in the method | Avoid / low success

Plant snapshot

Habit	Low spreading or erect shrub, 0.05–0.6 m (sometimes to 1 m) and 0.4–2 m wide.
Flowering	Jul – Dec (small, green or green-white).
Habitat	Grey, brown or yellow sand on coastal and near-coastal limestone, cliffs and dunes.
Local note	Recorded in the Wanneroo LGA (not listed for Joondalup). Useful for erosion control, especially on calcareous soils (Tasmania).

Propagation options

Method	Suitability	Notes	Details in
Cuttings	Difficult	Usually grown from cuttings of new growth, but slow (provisional).	This guide
Seed	Difficult	Complex dormancy; sow whole fruit in autumn – slow.	–
Division	Not used	Shrub – not divided.	–

Propagation method

Item	Details
Best time to take cuttings	**Jan – Mar** (estimate) – the new growth made after flowering has half-hardened, and it's warm. Epacrid cuttings can be taken at any time of year, so Apr – May and Oct – Dec are workable with non-flowering shoots. Avoid Jun – Sep (cool, and flowering).
Wood type	**Semi-hardwood** – half-hardened new growth.
Cutting size	50–100 mm (estimate) – short side shoots, as it's a low, spreading shrub. Doig used 80–250 mm for epacrids generally.
Selecting material	Healthy, non-flowering new growth. Take plenty – strike rates are low.
Pre-treatment	• Strip leaves from the lower half. • Light wound on firmer stems. • Dip the base in rooting hormone.
Rooting hormone	Medium strength, as for other epacrids: 3 g/L IBA (**Clonex Purple**).
Propagation mix	Very free-draining sand-based cutting mix.
Environment	Humid frame in bright shade, not heavy mist.
Expected strike time	Slow. Doig's epacrid trials without mist gave at least 30% strikes over a 6–12 month holding period (a related *Acrotriche* struck in that time). Check from about 8 weeks and record the result.
Post-treatment / aftercare	• Expect fine, fragile roots – pot up with the mix ball intact. • Harden off slowly.

© 2026 Mitchell Donald. All rights reserved.

Common problems

Problem	Likely cause and fix
No roots after months	Normal for epacrids. Keep them going; try another month or hormone strength next batch.
Cuttings blacken or rot	Too wet or humid. Less mist, better drainage and airflow.
Roots break when potting up	Move the whole plug without disturbing the roots.

Notes

Difficulty	**Difficult (provisional).** Usually grown from cuttings, but epacrids are slow and strike poorly.
Other methods	Seed – difficult: morphophysiological dormancy and complex germination requirements. Tasmanian growers sow the whole fruit 3–5 mm deep in autumn, in mix inoculated with soil from around parent plants; chilling at 3–5°C for up to 3 months may help. Germination is slow.
Confidence	**Low.** Tasmanian growers usually raise it from cuttings of new growth; method details are from general epacrid experience – a trial protocol.
Sources	See References: 4, 13, 82, 99, 100

Nursery records

Log each batch here to confirm or update the timing and difficulty above.

Date set	Stock source / provenance	No. set	No. struck	Strike %	Notes

© 2026 Mitchell Donald. All rights reserved.

Anthocercis littorea

Yellow Tailflower

Family **Solanaceae**

DIFFICULTY	WOOD TYPE	BEST MONTHS	CUTTING LENGTH	HORMONE (IBA)
Easy–Moderate	Semi-hardwood	Dec – Apr	75–100 mm	3 g/L IBA

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Cuttings	Best	Best	Best	Best	OK	–	–	–	–	–	OK	Best

Best window | Workable – see 'Selecting material' in the method | Avoid / low success

Plant snapshot

Habit	Erect, rarely sprawling shrub, 0.6–3 (–4) m.
Flowering	All year, mainly Jun – Nov (yellow, star-like with long narrow petals).
Habitat	Calcareous sand; coastal limestone, sand dunes and sandplains.
Local note	Recorded in the Wanneroo LGA. **Poisonous** – all *Anthocercis* contain tropane alkaloids. Wear gloves and wash hands after handling.

Propagation options

Method	Suitability	Notes	Details in
Cuttings	Main method	Hardened current season's growth strikes reasonably well.	This guide
Seed	Possible	GA, smoke or karrikinolide raise germination (research).	–
Division	Not used	Woody shrub.	–

Propagation method

Item	Details
Best time to take cuttings	**Dec – Apr**, after the main flowering, once new growth has hardened. Nov and May are workable. Avoid Jun – Oct (main flowering).
Wood type	**Semi-hardwood** – hardened current season's growth (ANPSA).
Cutting size	75–100 mm.
Selecting material	Non-flowering shoots; the plant flowers most of the year, so snip off buds.
Pre-treatment	• Wear gloves. • Strip leaves from the lower half; reduce large leaves by a third. • Light wound on firmer stems. • Dip the base in rooting hormone.
Rooting hormone	Semi-hardwood strength, 3 g/L IBA (**Clonex Purple**).
Propagation mix	Standard free-draining cutting mix.
Environment	Humid frame, bright shade, bottom heat.
Expected strike time	Not documented – check from about 4–6 weeks and record the result.
Post-treatment / aftercare	• Standard hardening off. • Grow on in full sun; tip-prune for a bushier plant.

© 2026 Mitchell Donald. All rights reserved.

Common problems

Problem	Likely cause and fix
Cuttings flower	Material taken from flowering shoots. Remove buds; pick non-flowering growth.
Base rotting	Mix too wet. Water less; improve drainage.
Skin irritation	Handle with gloves – the plant is toxic.

Notes

Difficulty	**Easy–Moderate.** Cuttings should strike reasonably well; seed is hard to obtain and slow.
Other methods	Seed – gibberellic acid, smoke water or karrikinolide substantially increased germination in research; untreated seed gave 0% in the only Alcoa test.
Confidence	**Medium.** ANPSA says cuttings of hardened current season's growth should strike reasonably well; details follow standard practice.
Sources	See References: 4, 32, 64, 65, 68

Nursery records

Log each batch here to confirm or update the timing and difficulty above.

Date set	Stock source / provenance	No. set	No. struck	Strike %	Notes

© 2026 Mitchell Donald. All rights reserved.

Atriplex cinerea

Grey Saltbush

Family **Amaranthaceae (chenopods)**

DIFFICULTY	WOOD TYPE	BEST MONTHS	CUTTING LENGTH	HORMONE (IBA)
Easy	Semi-hardwood tips	Nov – Mar	100–150 mm	1.5–3 g/L IBA

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Cuttings	Best	Best	Best	OK	OK	–	–	–	OK	OK	Best	Best

Best window | Workable – see 'Selecting material' in the method | Avoid / low success

Plant snapshot

Habit	Grey, erect to spreading shrub, 0.2–1.5 m (to 2 m). Plants are usually male or female, sometimes both.
Flowering	Sep – Oct or Dec (cream-yellow). Male plants have dense brown flower spikes up to 5 cm long.
Habitat	Saline sand; coastal areas, limestone cliffs, salt lakes and dunes.
Local note	Recorded in the Wanneroo LGA (not listed for Joondalup). Tolerates drought, salt spray and saline soils – often the plant closest to the sea.

Propagation options

Method	Suitability	Notes	Details in
Cuttings	Main method	Strikes easily; no mist. Keeps the parent's sex.	This guide
Seed	Good	Wash out salts (4-day soak), smoke water; 7–21 days.	→ Seed Sowing Guide
Division	Possible	Low stems layer themselves – lift rooted pieces.	–

Propagation method

Item	Details
Best time to take cuttings	**Nov – Mar** (estimate), in warm weather after the spring flowering. Apr – May and Sep – Oct are workable. Avoid Jun – Aug (cool, slow growth). NZ growers say cuttings can be collected at any time.
Wood type	**Semi-hardwood terminal (tip) cuttings** of current season's growth.
Cutting size	100–150 mm (estimate, as for related saltbushes).
Selecting material	Take cuttings from **both male and female plants** and label them – a cutting keeps its parent's sex, and only females set seed. In New Zealand, most commercial stock is female.
Pre-treatment	• Strip leaves from the lower half. • No wounding needed. • Dip the base in rooting hormone.
Rooting hormone	Softwood to semi-hardwood strength (1.5–3 g/L IBA, **Clonex Green** or **Purple**).
Propagation mix	Very free-draining sandy mix – it needs good drainage (Tasmania).
Environment	**Good ventilation and no mist** – misting encourages leaf disease on saltbush cuttings (ANBG, related species). Bottom heat may help (estimate); keep the mix just moist.
Expected strike time	Not documented – check from about 4 weeks and record the result.
Post-treatment / aftercare	• Harden off, then grow on in full sun. • Water sparingly once established.

© 2026 Mitchell Donald. All rights reserved.

Common problems

Problem	Likely cause and fix
Leaf spots, leaves dropping	Too humid or misted. Stop misting, open the frame, improve airflow.
Base rotting	Mix too wet. Use a sandier mix and water less.
No seed from the planting later	All male (or all female) stock. Keep and label cuttings from both sexes.

Notes

Difficulty	**Easy.** Growers in Tasmania and New Zealand report it strikes easily from cuttings.
Other methods	Seed – wash the salts out of the fruit first (see the Seed Sowing Guide). Low stems also layer themselves – lift rooted pieces.
Confidence	**Medium.** Tasmanian and NZ growers report it strikes easily; method details follow related saltbushes.
Sources	See References: 4, 13, 36, 95, 96

Nursery records

Log each batch here to confirm or update the timing and difficulty above.

Date set	Stock source / provenance	No. set	No. struck	Strike %	Notes

© 2026 Mitchell Donald. All rights reserved.

Atriplex isatidea

Coast Saltbush

Family **Amaranthaceae (chenopods)**

DIFFICULTY	WOOD TYPE	BEST MONTHS	CUTTING LENGTH	HORMONE (IBA)
Easy–Moderate *	Semi-hardwood tips	Sep – Nov	100–150 mm	1.5–3 g/L IBA

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Cuttings	OK	OK	–	–	–	–	–	OK	Best	Best	Best	OK

Best window | Workable – see 'Selecting material' in the method | Avoid / low success

Plant snapshot

Habit	Grey-leaved, spreading shrub, 0.5–2.5 m. Male and female flowers on the same plant.
Flowering	Mainly Mar – Jun (small, inconspicuous).
Habitat	White sand on coastal dunes.
Local note	Recorded in the Wanneroo LGA. Very salt- and wind-tolerant – a useful dune stabiliser.

Propagation options

Method	Suitability	Notes	Details in
Cuttings	Good	Saltbush cuttings are generally reliable (provisional for this species).	This guide
Seed	Main method	Soak 4 days in changed water, then smoke water.	→ Seed Sowing Guide
Division	Not used	Woody shrub.	–

Propagation method

Item	Details
Best time to take cuttings	**Sep – Nov**, in active spring growth (the time recommended for the related Old Man Saltbush). Aug and Dec – Feb are workable. Avoid Mar – Jul (flowering, then cool, slow growth).
Wood type	**Semi-hardwood terminal (tip) cuttings** of current season's growth.
Cutting size	100–150 mm. Terminal cuttings of about 15 cm are used for *A. nummularia*.
Selecting material	Vigorous, non-flowering tips from healthy plants. Avoid old, woody stems.
Pre-treatment	• Strip leaves from the lower half. • No wounding needed. • Dip the base in rooting hormone.
Rooting hormone	Softwood to semi-hardwood strength (1.5–3 g/L IBA, **Clonex Green** or **Purple**). Related saltbushes root without hormone, so it mainly evens out the batch.
Propagation mix	Free-draining mix of sand and potting mix.
Environment	**Good ventilation and no mist** – misting encourages a leaf-spotting fungus on saltbush cuttings (ANBG, *A. nummularia*). Bottom heat helps; keep the mix just moist.
Expected strike time	Not documented for this species. In one trial, 20 cm unrooted *A. halimus* cuttings with no hormone gave about **49% survival at 8 weeks**. Check from about 4 weeks and record your result.
Post-treatment / aftercare	• Harden off, then grow on in full sun. • Water sparingly once established. • Tip-prune for a bushier tube.

© 2026 Mitchell Donald. All rights reserved.

Common problems

Problem	Likely cause and fix
Leaf spots, leaves dropping	Too humid or misted. Stop misting, open the frame, improve airflow.
Base rotting	Mix too wet. Use a sandier mix and water less.
Leggy, pale growth	Not enough light. Grow on in full sun.

Notes

Difficulty	**Easy–Moderate (provisional).** Saltbushes generally strike readily from tip cuttings; the main risk is leaf disease under mist. Seed is the more common nursery method.
Confidence	**Medium.** No species-specific cutting data. Method follows published results for related saltbushes (*A. nummularia*, *A. halimus*).
Sources	See References: 4, 36, 37, 38

Nursery records

Log each batch here to confirm or update the timing and difficulty above.

Date set	Stock source / provenance	No. set	No. struck	Strike %	Notes

© 2026 Mitchell Donald. All rights reserved.

Baeckea robusta

No widely used common name (a baeckea)

Also listed as *Babingtonia robusta*

Family **Myrtaceae**

DIFFICULTY	WOOD TYPE	BEST MONTHS	CUTTING LENGTH	HORMONE (IBA)
Easy–Moderate *	Semi-hardwood	Jan – Mar	50–75 mm	3 g/L IBA

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Cuttings	Best	Best	Best	OK	OK	–	–	–	–	–	–	OK

Best window | Workable – see 'Selecting material' in the method | Avoid / low success

Plant snapshot

Habit	Erect, slender shrub, 0.9–2.6 m.
Flowering	Jun – Dec (white or white-pink).
Habitat	Grey or yellow sand over laterite or limestone; sandplains and coastal limestone.
Local note	**Not a local species** – FloraBase records it only from the Geraldton Sandplains (Dandaragan and Northampton LGAs). Check the name and provenance before using it in local plantings.
Name note	Its placement in *Baeckea* is under review (Rye 2026), so the name may change.

Propagation options

Method	Suitability	Notes	Details in
Cuttings	Main method	Firm current season's growth strikes readily in the genus (provisional).	This guide
Seed	Possible	No pre-treatment in the genus, but capsules shed seed quickly.	–
Division	Not used	Shrub – not divided.	–

Propagation method

Item	Details
Best time to take cuttings	**Jan – Mar** (estimate), after the long Jun – Dec flowering, once new growth has firmed. Apr, May and Dec are workable with non-flowering shoots. Avoid Jun – Nov (flowering).
Wood type	**Semi-hardwood** – firm current season's growth (genus advice).
Cutting size	50–75 mm.
Selecting material	Non-flowering tips from healthy plants.
Pre-treatment	• Strip leaves from the lower half to two-thirds. • Light wound on firmer stems. • Dip the base in rooting hormone.
Rooting hormone	Semi-hardwood strength, 3 g/L IBA (**Clonex Purple**).
Propagation mix	Standard free-draining cutting mix.
Environment	Humid frame, bright shade, bottom heat.
Expected strike time	Not documented – check from about 6 weeks and record the result.
Post-treatment / aftercare	• Standard hardening off. • Pot on into a free-draining native mix; tip-prune for bushiness.

© 2026 Mitchell Donald. All rights reserved.

Common problems

Problem	Likely cause and fix
Callus, no roots	Trim and re-set; try a lower hormone strength on the next batch.
Cuttings flower	Taken from flowering shoots. Use non-flowering tips, after flowering.
Base rotting	Mix too wet. Water less.

Notes

Difficulty	**Easy–Moderate (provisional).** Cuttings of firm current season's growth strike readily in *Baeckea* and its former members; there is no data for this species.
Other methods	Seed – germinates without pre-treatment in the genus, but the small capsules shed seed as soon as it's ripe, so it's hard to collect.
Confidence	**Low.** No data for this species. Based on genus advice – cuttings of firm current season's growth strike readily.
Sources	See References: 4, 13, 89, 90, 97

Nursery records

Log each batch here to confirm or update the timing and difficulty above.

Date set	Stock source / provenance	No. set	No. struck	Strike %	Notes

© 2026 Mitchell Donald. All rights reserved.

Calothamnus quadrifidus

One-sided Bottlebrush

Family **Myrtaceae**

DIFFICULTY	WOOD TYPE	BEST MONTHS	CUTTING LENGTH	HORMONE (IBA)
Easy–Moderate *	Semi-hardwood	Jan – Apr	75–100 mm	3 g/L IBA

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Cuttings	Best	Best	Best	Best	OK	–	–	–	–	–	–	OK

Best window | Workable – see 'Selecting material' in the method | Avoid / low success

Plant snapshot

Habit	Erect, compact or spreading shrub, usually 0.9–2 m.
Flowering	Jun – Dec (red, one-sided brushes).
Habitat	Wide variety of soils and habitats.
Local note	Recorded in the Wanneroo LGA. Several subspecies exist – label stock by subspecies and provenance.

Propagation options

Method	Suitability	Notes	Details in
Cuttings	Good	Presumed to strike like related melaleucas (provisional).	This guide
Seed	Main method	Easy from seed held in old capsules.	→ Seed Sowing Guide
Division	Not used	Woody shrub.	–

Propagation method

Item	Details
Best time to take cuttings	**Jan – Apr**, after flowering, once the new growth has firmed. Dec and May are workable. Avoid Jun – Nov (flowering).
Wood type	**Semi-hardwood** – firm current season's growth.
Cutting size	75–100 mm, as for *Melaleuca*.
Selecting material	Non-flowering side shoots. Keep subspecies and forms separate.
Pre-treatment	• Strip the needle-like leaves from the lower half to two-thirds. • Light wound on firmer stems. • Dip the base in rooting hormone.
Rooting hormone	Semi-hardwood strength, 3 g/L IBA (**Clonex Purple**).
Propagation mix	Standard free-draining cutting mix.
Environment	Humid frame, bright shade, bottom heat.
Expected strike time	Not documented – check from about 6 weeks and record the result.
Post-treatment / aftercare	• Standard hardening off. • Pot on into a free-draining native mix.

© 2026 Mitchell Donald. All rights reserved.

Common problems

Problem	Likely cause and fix
Cuttings brown off	Too dry or too soft. Use firmer material and keep humidity up until rooted.
Callus, no roots	Trim and re-set; try a lower hormone strength.
Slow in winter	Use bottom heat, or wait for the Jan – Apr window.

Notes

Difficulty	**Easy–Moderate (provisional).** Expected to behave like related melaleucas, but untested here. Seed is easy and is the main method.
Confidence	**Low.** ANPSA notes the genus is 'presumably' grown from cuttings. Method follows related melaleucas – trial protocol.
Sources	See References: 1, 4, 13, 41

Nursery records

Log each batch here to confirm or update the timing and difficulty above.

Date set	Stock source / provenance	No. set	No. struck	Strike %	Notes

© 2026 Mitchell Donald. All rights reserved.

Carpobrotus virescens

Coastal Pigface

Family **Aizoaceae**

DIFFICULTY	WOOD TYPE	BEST MONTHS	CUTTING LENGTH	HORMONE (IBA)
Easy	Succulent stem pieces	Sep – Nov; Mar – Apr	10–15 cm (tubes)	Not needed

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Cuttings	OK	OK	Best	Best	OK	–	–	OK	Best	Best	Best	OK

Best window · Workable – see 'Selecting material' in the method · Avoid / low success

Plant snapshot

Habit	Prostrate succulent groundcover, 0.1–0.3 m high and 0.5–3 m wide, rooting along the stems.
Flowering	Jun – Dec or Jan (purple-pink, sometimes white), followed by fleshy fruit.
Habitat	Coastal limestone and sand dunes.
Local note	Recorded in the Wanneroo LGA. Collect only from plants you are sure are native – the introduced Hottentot Fig (*C. edulis*) is naturalised on some coasts.

Propagation options

Method	Suitability	Notes	Details in
Cuttings	Main method	Stem pieces root very easily; rooted runners can be lifted.	This guide
Seed	Good	Ferment fruit, sow fine seed on the surface; 6–8 days.	→ Seed Sowing Guide
Division	Possible	Lift stems that have rooted – treat as cuttings.	–

Propagation method

Item	Details
Best time to take cuttings	**Sep – Nov** and **Mar – Apr**, when plants are actively growing. Aug, Dec – Feb and May are workable. Avoid Jun – Jul (slow growth; cuttings rot in cold, wet mix).
Wood type	**Succulent stem (runner) pieces** – not woody cuttings. Use firm stems with plump leaves.
Cutting size	**Tubes:** 10–15 cm pieces with 2–3 leaf pairs. **Direct planting / layering:** about 30 cm, with at least 5 cm left above the sand (ANBG, *C. glaucescens*).
Selecting material	Healthy runners. Pieces that have already rooted at the nodes are ideal. Avoid shrivelled or damaged stems.
Pre-treatment	• Remove the leaves from the lowest 1–2 nodes. • Leave the cut ends to dry in the shade for a day or two before setting – this reduces rot. • No hormone needed.
Rooting hormone	Not needed – stems root readily at the nodes.
Propagation mix	Coarse washed sand or a very sandy, free-draining mix.
Environment	Open bench in bright light or light shade. **No mist and no humid frame** – succulent leaves rot. Water sparingly until rooted.
Expected strike time	Not documented, but usually quick – check from about 3 weeks.

© 2026 Mitchell Donald. All rights reserved.

Item	Details
Post-treatment / aftercare	• Grow on in full sun. • Pot on or plant out once roots hold the mix together. • Water sparingly – overwatering is the main cause of losses.

Common problems

Problem	Likely cause and fix
Stems go soft and rot	Too wet or humid. Dry the cut ends first, water less, move out of the humid frame.
Leaves shrivel	Too dry for too long. Water lightly once set.
Pale, stretched growth	Not enough light. Move to full sun.

Notes

Difficulty	**Easy.** Pigface roots readily from stem pieces, and rooted runners can simply be cut off and potted up.
Other methods	Layering – pin a runner onto a pot of sand beside the parent and cut it free once rooted.
Confidence	**Medium.** Method from published advice for the closely related *C. glaucescens*; pigface is widely grown from stem pieces.
Sources	See References: 4, 12, 33

Nursery records

Log each batch here to confirm or update the timing and difficulty above.

Date set	Stock source / provenance	No. set	No. struck	Strike %	Notes

© 2026 Mitchell Donald. All rights reserved.

Conostephium preissii

No widely used common name

Family **Ericaceae**

DIFFICULTY	WOOD TYPE	BEST MONTHS	CUTTING LENGTH	HORMONE (IBA)
Difficult *	Semi-hardwood	Jan – Mar	80–150 mm	3 g/L IBA

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Cuttings	Best	Best	Best	OK	OK	–	–	–	–	–	–	OK

Best window | Workable – see 'Selecting material' in the method | Avoid / low success

Plant snapshot

Habit	Erect, multi-stemmed shrub, 0.3–1.2 m.
Flowering	White and purple to red-pink. FloraBase gives Jan – Sep or Dec; other sources say Sep – Dec, or most of the year.
Habitat	Sandplains and winter-wet flats, on sand over laterite, quartzite or limestone; Banksia woodland and heath.
Local note	Recorded in the Wanneroo LGA (not listed for Joondalup).

Propagation options

Method	Suitability	Notes	Details in
Cuttings	Difficult	No data – trial protocol based on related epacrids.	This guide
Seed	Difficult	Not documented; epacrid stones need a long time in soil.	–
Division	Not used	Shrub – not divided.	–

Propagation method

Item	Details
Best time to take cuttings	**Jan – Mar** (estimate) – firm growth in warm weather. Apr, May and Dec are workable. Flowering times vary, so pick non-flowering shoots whenever you take them.
Wood type	**Semi-hardwood** – current season's growth that has firmed, as suggested for the related *C. pendulum*.
Cutting size	80–150 mm – the shorter end of the 80–250 mm used for epacrid cuttings in Sydney trials (Doig).
Selecting material	Non-flowering tips from healthy plants. Take plenty – epacrids strike poorly.
Pre-treatment	• Strip leaves from the lower half. • Light wound on firmer stems. • Dip the base in rooting hormone.
Rooting hormone	Medium strength, as for other epacrids: 3 g/L IBA (**Clonex Purple**).
Propagation mix	Very free-draining sand-based cutting mix.
Environment	Humid frame in bright shade, not heavy mist. Trial bottom heat on part of the batch.
Expected strike time	Not documented – epacrids are slow (6–12 months in trials). Check from about 8 weeks and record the result.
Post-treatment / aftercare	• Strike in **individual small pots or tubes** to limit root disturbance (advice for *C. pendulum*). • Pot on without breaking up the root ball; harden off slowly.

© 2026 Mitchell Donald. All rights reserved.

Common problems

Problem	Likely cause and fix
No roots after months	Common in epacrids. Keep them going; try another month or hormone strength next batch.
Cuttings blacken or rot	Too wet or humid. Less mist, better drainage and airflow.
Plants sulk after potting	Roots disturbed. Use individual tubes and move the whole plug.

Notes

Difficulty	**Difficult (provisional).** No data for this species; related epacrids strike poorly, so the rating is cautious.
Other methods	Seed – not documented. The fruit is a fleshy drupe with a hard stone; related epacrids need a long time in the soil before they germinate.
Confidence	**Low.** No propagation data for this species. Based on a related *Conostephium* and general epacrid experience – a trial protocol.
Sources	See References: 1, 4, 13, 82, 93, 94

Nursery records

Log each batch here to confirm or update the timing and difficulty above.

Date set	Stock source / provenance	No. set	No. struck	Strike %	Notes

© 2026 Mitchell Donald. All rights reserved.

Diplolaena angustifolia

Yanchep Rose

Family **Rutaceae**

DIFFICULTY	WOOD TYPE	BEST MONTHS	CUTTING LENGTH	HORMONE (IBA)
Moderate *	Semi-hardwood	Dec – Mar	75–100 mm	3 g/L IBA

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Cuttings	Best	Best	Best	OK	OK	–	–	–	–	–	OK	Best

Best window | Workable – see 'Selecting material' in the method | Avoid / low success

Plant snapshot

Habit	Erect, compact or spreading shrub, 0.3–1.5 m.
Flowering	Jun – Oct (red or orange).
Habitat	White or yellow sand on limestone hills, rocky ridges and dunes.
Local note	Recorded in the Wanneroo LGA (not listed for Joondalup) – it grows from Wanneroo north to near Eneabba. Dislikes humid conditions and wet, boggy soils.

Propagation options

Method	Suitability	Notes	Details in
Cuttings	Main method	Firm current season's growth works in a relative (provisional).	This guide
Seed	Possible	Slow – first seedlings at 4–7 weeks (Kings Park); smoke may help.	→ Seed Sowing Guide
Division	Not used	Shrub – not divided.	–

Propagation method

Item	Details
Best time to take cuttings	**Dec – Mar** (estimate) – after the winter–spring flowering, once new growth has firmed. Nov, Apr and May are workable. Avoid Jun – Oct (flowering).
Wood type	**Semi-hardwood** – firm current season's growth (as for *D. microcephala*).
Cutting size	75–100 mm (estimate).
Selecting material	Non-flowering tips from healthy plants.
Pre-treatment	• Strip leaves from the lower half. • Wound the base on firmer stems. • Dip the base in rooting hormone.
Rooting hormone	Semi-hardwood strength, 3 g/L IBA (**Clonex Purple**).
Propagation mix	Very free-draining cutting mix.
Environment	Humid frame in bright shade, but vent well and don't let leaves stay wet – *D. microcephala* cuttings are prone to fungal disease on the leaves.
Expected strike time	Not documented. Check from about 6 weeks and record the result.
Post-treatment / aftercare	• Harden off with good airflow. • Pot on into a free-draining native mix; don't overwater.

© 2026 Mitchell Donald. All rights reserved.

Common problems

Problem	Likely cause and fix
Leaves spot or drop	Fungal disease. Improve airflow, mist less, remove affected leaves.
Base rotting	Mix too wet. Water less; use a more open mix.
Callus, no roots	Trim and re-set; try a different month next batch.

Notes

Difficulty	**Moderate (provisional).** Cuttings work in a related species; no data for this one yet.
Other methods	Seed – slow: first seedlings at 28–50 days in Kings Park records, and smoke may help (see the Seed Sowing Guide).
Confidence	**Low.** No cuttings data for this species. Based on the related *D. microcephala* (cuttings of firm current season's growth are successful) – a trial protocol.
Sources	See References: 4, 13, 101, 102, 103

Nursery records

Log each batch here to confirm or update the timing and difficulty above.

Date set	Stock source / provenance	No. set	No. struck	Strike %	Notes

© 2026 Mitchell Donald. All rights reserved.

Eremophila glabra

Tar Bush / Common Emu Bush

Family **Scrophulariaceae**

DIFFICULTY	WOOD TYPE	BEST MONTHS	CUTTING LENGTH	HORMONE (IBA)
Easy	Semi-hardwood	Jan – Apr	75–100 mm	3 g/L IBA

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Cuttings	Best	Best	Best	Best	OK	–	–	–	–	–	OK	OK

Best window | Workable – see 'Selecting material' in the method | Avoid / low success

Plant snapshot

Habit	Very variable, from prostrate to 3 m. Perth coastal plants are mostly subsp. *albicans*: low, spreading, to about 0.4 m high and 2 m wide, with grey, felty leaves.
Flowering	Species: Mar – Dec. Subsp. *albicans*: May – Jan. Tubular flowers, yellow, orange, red or green.
Habitat	Sand and limestone; dunes, sandplains and winter-wet flats, sometimes saline.
Local note	Recorded in the Wanneroo LGA and along the Perth coast. Several subspecies and many forms – label stock by subspecies and form.

Propagation options

Method	Suitability	Notes	Details in
Cuttings	Main method	Most forms strike readily from semi-hardwood cuttings.	This guide
Seed	Difficult	Unreliable – hard fruits; slow, erratic germination.	–
Division	Not used	Woody shrub.	–

Propagation method

Item	Details
Best time to take cuttings	**Jan – Apr**, using hardened current season's growth while the weather is warm and plants are mostly out of flower. Nov – Dec and May are workable. Avoid Jun – Oct (cool weather, flowering).
Wood type	**Semi-hardwood** – hardened current season's growth.
Cutting size	75–100 mm.
Selecting material	**Non-flowering** tips or side shoots. On grey-leaved forms pick clean foliage with no sign of mould.
Pre-treatment	• Carefully remove leaves from the lower two-thirds. • Wound: remove a thin sliver of bark from the base. • Dip the base in rooting hormone.
Rooting hormone	Semi-hardwood strength, e.g. **Clonex Purple (3 g/L IBA)**.
Propagation mix	Very free-draining, sand-based mix.
Environment	**Grey, felty-leaved forms rot under heavy mist or high humidity** – keep foliage as dry as possible, water the mix, give good airflow. Green-leaved forms tolerate normal misting. Bottom heat speeds rooting.
Expected strike time	About **4–8 weeks** in warm weather.
Post-treatment / aftercare	• Standard hardening off. • Pot on into a free-draining mix and grow on in full sun. • Keep water off grey foliage; water in the morning.

© 2026 Mitchell Donald. All rights reserved.

Common problems

Problem	Likely cause and fix
Leaves blacken or grow grey mould	Too humid. Cut back misting, open the frame, remove affected leaves.
Base rotting	Mix too wet. Water less often and use a sandier mix.
Cuttings flower instead of rooting	Use non-flowering shoots and pinch off buds.

Notes

Difficulty	**Easy.** Most forms of *E. glabra* strike readily from cuttings.
Other methods	Seed – unreliable in *Eremophila* (hard fruits, slow and erratic germination). Grafting is used for difficult eremophilas but isn't needed for this species.
Confidence	**High.** Published sources agree most forms strike readily from cuttings.
Sources	See References: 1, 4, 12, 13, 20, 21, 22, 30

Nursery records

Log each batch here to confirm or update the timing and difficulty above.

Date set	Stock source / provenance	No. set	No. struck	Strike %	Notes

© 2026 Mitchell Donald. All rights reserved.

Grevillea crithmifolia

No widely used common name

Family **Proteaceae**

DIFFICULTY	WOOD TYPE	BEST MONTHS	CUTTING LENGTH	HORMONE (IBA)
Moderate *	Semi-hardwood	Dec – Apr	75–100 mm	3 g/L IBA

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Cuttings	Best	Best	Best	Best	OK	–	–	–	–	–	OK	Best

Best window | Workable – see 'Selecting material' in the method | Avoid / low success

Plant snapshot

Habit	Dense, much-branched shrub, (0.3–) 0.6–2 m. Compact shrub and low groundcover forms exist.
Flowering	Jun or Aug – Nov (white to white-pink). Some people dislike the scent.
Habitat	White or yellow sand over limestone; dunes, hillslopes and sandplains.
Local note	Recorded in the City of Joondalup. Dieback susceptible. Keep the shrub and groundcover forms separate.

Propagation options

Method	Suitability	Notes	Details in
Cuttings	Main method	The usual method, but may be slow to root.	This guide
Seed	Difficult	About 9% in one nursery test; smoke recommended.	–
Division	Not used	–	–

Propagation method

Item	Details
Best time to take cuttings	**Dec – Apr**, after flowering, once new growth has firmed. Nov and May are workable. Avoid Jun – Oct (flowering).
Wood type	**Semi-hardwood** – firm current season's growth.
Cutting size	75–100 mm (ANPSA *Grevillea* guidance).
Selecting material	Non-flowering shoots from a plant of the form you want (shrub or groundcover).
Pre-treatment	• Strip leaves from the lower half to two-thirds (cut, don't tear). • Wound the base – a thin sliver of bark from the bottom 15–20 mm. • Dip the base in rooting hormone.
Rooting hormone	Semi-hardwood strength, 3 g/L IBA (**Clonex Purple**). Wounding and hormone improve strike in grevilleas.
Propagation mix	Free-draining, clean cutting mix. Strict hygiene – the species is dieback susceptible.
Environment	Humid frame, bright shade, bottom heat. Be patient – roots can be slow.
Expected strike time	Not documented, but reported slow – allow at least **8–12 weeks** before discarding. Record your results.
Post-treatment / aftercare	• Standard hardening off. • Pot on into a **low-phosphorus** native mix; feed only with low-P fertiliser.

© 2026 Mitchell Donald. All rights reserved.

Common problems

Problem	Likely cause and fix
No roots after 8 weeks	Normal for this species – keep them going. Next time wound and use hormone.
Heavy callus, no roots	Trim the callus and re-set; try a lower hormone strength.
Plants yellow or die after potting	Phosphorus or dieback. Low-P mix, clean pots off the ground.

Notes

Difficulty	**Moderate (provisional).** Cuttings are the usual method but may be slow to produce roots.
Other methods	Seed – poor: about 9% in a single Alcoa nursery test; a seed supplier recommends smoke treatment.
Confidence	**Medium.** Grown commercially from cuttings but reported slow to root; size and treatment follow ANPSA *Grevillea* guidance.
Sources	See References: 4, 32, 56, 57, 59, 61

Nursery records

Log each batch here to confirm or update the timing and difficulty above.

Date set	Stock source / provenance	No. set	No. struck	Strike %	Notes

© 2026 Mitchell Donald. All rights reserved.

Grevillea preissii

Spider-net Grevillea

Family **Proteaceae**

DIFFICULTY	WOOD TYPE	BEST MONTHS	CUTTING LENGTH	HORMONE (IBA)
Moderate	Semi-hardwood	Nov – Mar	75–100 mm	3 g/L IBA

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Cuttings	Best	Best	Best	OK	–	–	–	–	–	OK	Best	Best

Best window | Workable – see 'Selecting material' in the method | Avoid / low success

Plant snapshot

Habit	Shrub, 0.15–1.7 m.
Flowering	Jun – Sep (red spider flowers; attract honeyeaters).
Habitat	White, grey or yellow sand, sandy clay and limestone; ridges, outcrops and seasonally wet areas.
Local note	Recorded in Joondalup and Wanneroo. The former subsp. *glabrilimba* (including 'Magic Lantern') is now a separate species, *G. glabrilimba* – keep it apart from local stock.

Propagation options

Method	Suitability	Notes	Details in
Cuttings	Main method	Firm growth; slow, hormone improves strike.	This guide
Seed	Possible	Nick the seed coat; smoke water.	–
Division	Not used	–	–

Propagation method

Item	Details
Best time to take cuttings	**Nov – Mar**, after flowering, once the current season's growth has firmed. Oct and Apr are workable. Avoid May – Sep (flowering).
Wood type	**Semi-hardwood** – firm current season's growth (ANPSA).
Cutting size	75–100 mm (ANPSA *Grevillea* guidance).
Selecting material	Non-flowering shoots from healthy, local-provenance plants.
Pre-treatment	• Strip leaves from the lower half to two-thirds. • Wound the base on firmer stems. • Dip the base in rooting hormone – ANPSA notes it clearly improves success.
Rooting hormone	Semi-hardwood strength, 3 g/L IBA (**Clonex Purple**).
Propagation mix	Free-draining, clean cutting mix.
Environment	Humid frame, bright shade, bottom heat.
Expected strike time	Not documented, but reported slow – allow at least **8–12 weeks** before discarding. Record your results.
Post-treatment / aftercare	• Standard hardening off. • Pot on into a **low-phosphorus** native mix.

© 2026 Mitchell Donald. All rights reserved.

Common problems

Problem	Likely cause and fix
Slow to strike	Normal. Use hormone and bottom heat; be patient.
Heavy callus, no roots	Trim and re-set; try a lower hormone strength.
Yellowing after potting on	Phosphorus in mix or fertiliser. Use low-P products.

Notes

Difficulty	**Moderate.** Cuttings of firm growth can be slow to strike; hormone improves success.
Other methods	Seed – nick the seed coat before sowing (ANPSA); a seed supplier recommends smoke treatment.
Confidence	**High.** ANPSA gives species-specific cutting advice; size follows ANPSA *Grevillea* guidance.
Sources	See References: 4, 58, 59, 60, 61

Nursery records

Log each batch here to confirm or update the timing and difficulty above.

Date set	Stock source / provenance	No. set	No. struck	Strike %	Notes

© 2026 Mitchell Donald. All rights reserved.

Grevillea vestita

No widely accepted common name

Family **Proteaceae**

DIFFICULTY	WOOD TYPE	BEST MONTHS	CUTTING LENGTH	HORMONE (IBA)
Easy	Semi-hardwood	Dec – Apr	80–120 mm	3 g/L IBA

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Cuttings	Best	Best	Best	Best	OK	–	–	–	–	–	OK	Best

Best window Workable – see 'Selecting material' in the method Avoid / low success

Plant snapshot

Habit	Erect, spreading, prickly shrub, usually 0.5–3 m. Hairy branchlets; lobed leaves with sharp-pointed tips.
Flowering	Mainly Jun – Nov (white to cream, often pink-tinged, scented). Odd flowers Jan – Apr.
Habitat	Sandy soils over limestone or granite. Occurs on the Swan Coastal Plain, including the Perth area.
Local note	Two subspecies exist (subsp. *vestita* and subsp. *isopogoides*). Label stock plants by subspecies and provenance.

Propagation options

Method	Suitability	Notes	Details in
Cuttings	Main method	Firm current season's growth strikes reliably.	This guide
Seed	Good	Grows reasonably easily from seed; seedlings vary from the parent.	→ Seed Sowing Guide
Division	Possible	Can sucker – lift rooted suckers.	–

Propagation method

Item	Details
Best time to take cuttings	**Dec – Apr** (late spring to autumn), once the spring flush has firmed and flowering has finished. Nov and May are workable. Avoid Jun – Oct: plants are flowering, new growth is soft and cool weather slows rooting.
Wood type	**Semi-hardwood** – firm, current season's growth. It should bend about 60° and spring back.
Cutting size	80–120 mm long with 3–5 nodes. Tip or sub-tip pieces; two cuttings can often be taken from one shoot.
Selecting material	Healthy, vigorous, **non-flowering** shoots. Skip shoots carrying buds, old woody stems and soft new tips.
Pre-treatment	• Wear gloves: leaves are prickly and grevillea sap can irritate skin. • Remove any flowers or buds. • Strip leaves from the lower half to two-thirds (cut, don't tear the bark). • Trim remaining lobed leaves by about one-third. • Wound: remove a thin sliver of bark from the bottom 15–20 mm. • Dip the base in rooting hormone.
Rooting hormone	Semi-hardwood strength, e.g. **Clonex Purple (3 g/L IBA)**. If cuttings form a heavy callus but no roots, trim the callus back to clean wood and re-set; try a lower strength on the next batch.
Propagation mix	Free-draining, low nutrient: about 75–80% washed coarse river sand + 20–25% coir or peat (or a perlite/coir mix).
Environment	Humid propagation house or frame in bright shade. Bottom heat with the base of the cutting 5–10°C warmer than the air. Light misting only – the hairy foliage can rot if kept constantly wet.

© 2026 Mitchell Donald. All rights reserved.

Item	Details
Expected strike time	About **6–8 weeks** in warm weather. Slower in the cooler months.
Post-treatment / aftercare	• Harden off: ~2 weeks with cover closed by day and open at night, then ~2 weeks in shade before full sun. • Pot on when roots show at the drainage holes, into a **low-phosphorus** native mix. • Feed only with a low-phosphorus native fertiliser (Proteaceae are phosphorus-sensitive). • Responds well to pruning – tip-prune young plants for a bushier tube.

Common problems

Problem	Likely cause and fix
Heavy callus, no roots	Trim callus back and re-set. Lower the hormone strength next time.
Base rotting	Mix too wet or material too soft. Use firmer wood and water less often.
Leaf yellowing or scorch after potting on	Check phosphorus in the potting mix and fertiliser.

Notes

Difficulty	**Easy.** Cuttings from firm current season's growth usually strike reliably.
Other methods	Seed – grows reasonably easily from seed, but seedlings are not identical to the parent. Grafting onto *G. robusta* is used on the humid east coast; not needed locally.
Confidence	**High.** Species-specific propagation information is published and consistent across sources.
Sources	See References: 1, 2, 3, 4, 13, 15, 17

Nursery records

Log each batch here to confirm or update the timing and difficulty above.

Date set	Stock source / provenance	No. set	No. struck	Strike %	Notes

© 2026 Mitchell Donald. All rights reserved.

Guichenotia ledifolia

No widely used common name

Family **Malvaceae**

DIFFICULTY	WOOD TYPE	BEST MONTHS	CUTTING LENGTH	HORMONE (IBA)
Easy–Moderate *	Semi-hardwood	Dec – Apr	50–75 mm	3 g/L IBA

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Cuttings	Best	Best	Best	Best	OK	–	–	–	–	–	OK	Best

Best window | Workable – see 'Selecting material' in the method | Avoid / low success

Plant snapshot

Habit	Spreading shrub, 0.4–1.5 (–2) m.
Flowering	Jul – Oct (blue-purple or pink).
Habitat	Sandy or rocky clay; coastal limestone, sandplains and granite rocks.
Local note	Recorded in Joondalup and Wanneroo. Tolerates drought and alkaline (limestone) soils.

Propagation options

Method	Suitability	Notes	Details in
Cuttings	Main method	Often grown from cuttings; semi-hardwood.	This guide
Seed	Good	Hard coat: boiling water over seed; variable.	→ Seed Sowing Guide
Division	Not used	Shrub – not divided.	–

Propagation method

Item	Details
Best time to take cuttings	**Dec – Apr**, after flowering, once new growth has firmed. Nov and May are workable. Avoid Jun – Oct (cool, and plants are flowering).
Wood type	**Semi-hardwood** – firm current season's growth.
Cutting size	50–75 mm.
Selecting material	Healthy, **non-flowering** side shoots.
Pre-treatment	• Strip leaves from the lower half; the leaves are small, so no trimming is needed. • Dip the base in rooting hormone.
Rooting hormone	3 g/L IBA (**Clonex Purple**).
Propagation mix	Very free-draining cutting mix.
Environment	Humid frame, bright shade, bottom heat. Moist, not wet; good airflow.
Expected strike time	Not documented – check from about 4–6 weeks and record the result.
Post-treatment / aftercare	• Standard hardening off. • Grow on in full sun in a free-draining mix. • Tip-prune for a denser plant.

© 2026 Mitchell Donald. All rights reserved.

Common problems

Problem	Likely cause and fix
Leaves go mouldy	Too humid or crowded. Ventilate the frame; don't over-mist.
Base rotting	Too wet. Coarser mix; water less.
Slow to root	Use bottom heat and take cuttings in the warm months.

Notes

Difficulty	**Easy–Moderate (provisional).** Firm growth of the related *G. macrantha* usually strikes reliably; untested here.
Other methods	Seed – hard-coated. Pour boiling water over sown seed or soak it 24 h in water just off the boil; germination is variable and takes 30+ days (see the Seed Sowing Guide).
Confidence	**Medium.** WA seed notes say the genus is often grown from cuttings; semi-hardwood cuttings are listed for this species. No published strike rates.
Sources	See References: 4, 12, 13, 74, 75, 76, 77

Nursery records

Log each batch here to confirm or update the timing and difficulty above.

Date set	Stock source / provenance	No. set	No. struck	Strike %	Notes

© 2026 Mitchell Donald. All rights reserved.

Hardenbergia comptoniana

Native Wisteria

Family **Fabaceae**

DIFFICULTY	WOOD TYPE	BEST MONTHS	CUTTING LENGTH	HORMONE (IBA)
Easy–Moderate	Semi-hardwood	Dec – Mar	50–75 mm	3 g/L IBA

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Cuttings	Best	Best	Best	OK	–	–	–	–	–	–	OK	Best

Best window | Workable – see 'Selecting material' in the method | Avoid / low success

Plant snapshot

Habit	Vigorous twining climber or scrambling shrub.
Flowering	Jul – Oct (blue-purple, occasionally white).
Habitat	Sandy soils; coastal limestone, sandplains and dunes.
Local note	Recorded in Joondalup and Wanneroo. More vigorous than *H. violacea* – can smother small shrubs.

Propagation options

Method	Suitability	Notes	Details in
Cuttings	Good	Firm current season's growth strikes well.	This guide
Seed	Main method	Hot water + smoke water; about 10 days.	→ Seed Sowing Guide
Division	Not used	–	–

Propagation method

Item	Details
Best time to take cuttings	**Dec – Mar**, once the current season's growth has firmed after spring flowering. Nov and Apr are workable. Avoid May – Oct (cool, then flowering).
Wood type	**Semi-hardwood** – firm current season's growth (ANPSA).
Cutting size	**50–75 mm**, the length ANPSA suggests for pea-family cuttings. One runner gives several pieces.
Selecting material	Healthy runners without flowers or pods.
Pre-treatment	• Remove the leaves from the lower two-thirds. • No wounding needed. • Dip the base in rooting hormone.
Rooting hormone	Semi-hardwood strength, 3 g/L IBA (**Clonex Purple**).
Propagation mix	Standard free-draining cutting mix.
Environment	Humid frame, bright shade, bottom heat.
Expected strike time	Not documented – check from about 4 weeks and record the result.
Post-treatment / aftercare	• Standard hardening off. • Stake or trim runners so tubes don't tangle; pot on promptly.

© 2026 Mitchell Donald. All rights reserved.

Common problems

Problem	Likely cause and fix
Cuttings wilt	Material too soft. Use firmer growth; keep humidity up until rooted.
Runners tangle between tubes	Space tubes; trim or stake runners.
Base rotting	Mix too wet. Water less.

Notes

Difficulty	**Easy–Moderate.** Cuttings strike well. Seed is also easy after hot-water treatment.
Confidence	**Medium.** ANPSA reports cuttings of firm current season's growth strike well; size follows ANPSA pea-family guidance.
Sources	See References: 4, 13, 43, 44

Nursery records

Log each batch here to confirm or update the timing and difficulty above.

Date set	Stock source / provenance	No. set	No. struck	Strike %	Notes

© 2026 Mitchell Donald. All rights reserved.

Hemiandra pungens

Snakebush

Family **Lamiaceae**

DIFFICULTY	WOOD TYPE	BEST MONTHS	CUTTING LENGTH	HORMONE (IBA)
Easy	Softwood – semi-hardwood	Oct – Mar	50–75 mm	1.5–3 g/L IBA

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Cuttings	Best	Best	Best	OK	–	–	–	–	OK	Best	Best	Best

Best window | Workable – see 'Selecting material' in the method | Avoid / low success

Plant snapshot

Habit	Prostrate to ascending shrub, 0.05–1 m, with sharp, prickly leaves.
Flowering	Throughout the year (white, blue-purple or pink).
Habitat	Sand, clay, loam, gravel, laterite and granite; rock outcrops.
Local note	Recorded in Joondalup and Wanneroo. Resistant to dieback. Wear gloves when collecting.

Propagation options

Method	Suitability	Notes	Details in
Cuttings	Main method	Relatively easy from soft to semi-hardwood tips.	This guide
Seed	Difficult	Rarely available; about 4% germination in nursery tests.	–
Division	Not used	Not documented.	–

Propagation method

Item	Details
Best time to take cuttings	**Oct – Mar**, in the warm months while plants are growing actively. Sep and Apr are workable. Avoid May – Aug (slow rooting).
Wood type	**Softwood to semi-hardwood** tips.
Cutting size	50–75 mm tip cuttings.
Selecting material	Plants flower all year – choose tips without open flowers and snip off any buds.
Pre-treatment	• Wear gloves – the leaves are prickly. • Strip leaves from the lower half. • No wounding. • Dip the base in rooting hormone.
Rooting hormone	1.5 g/L IBA (**Clonex Green**) for soft tips; 3 g/L (**Clonex Purple**) for firmer ones.
Propagation mix	Very free-draining mix.
Environment	Bright shade and bottom heat, with light misting and good airflow – plants are unreliable in humid conditions.
Expected strike time	Not documented – check from about 3–4 weeks and record the result.
Post-treatment / aftercare	• Harden off promptly; grow on in full sun. • Free-draining mix; don't overwater.

© 2026 Mitchell Donald. All rights reserved.

Common problems

Problem	Likely cause and fix
Cuttings rot	Too humid or wet. Mist less and improve airflow.
Cuttings flower	Taken from flowering tips. Snip off buds and choose non-flowering shoots.
Slow in winter	Use bottom heat, or wait for the warm window.

Notes

Difficulty	**Easy.** ANPSA describes it as relatively easy to propagate from cuttings.
Other methods	Seed – rarely available and germinates poorly (about 4% untreated in Alcoa nursery tests). Grafting onto *Westringia fruticosa* is used where plants struggle.
Confidence	**Medium.** ANPSA says it is relatively easy from cuttings; timing and size follow standard practice.
Sources	See References: 4, 32, 45, 46, 47

Nursery records

Log each batch here to confirm or update the timing and difficulty above.

Date set	Stock source / provenance	No. set	No. struck	Strike %	Notes

© 2026 Mitchell Donald. All rights reserved.

Hibbertia cuneiformis

Cutleaf Hibbertia

Family **Dilleniaceae**

DIFFICULTY	WOOD TYPE	BEST MONTHS	CUTTING LENGTH	HORMONE (IBA)
Easy–Moderate *	Semi-hardwood	Mar – May	75–100 mm	3 g/L IBA

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Cuttings	OK	OK	Best	Best	Best	–	–	–	–	OK	OK	OK

Best window | Workable – see 'Selecting material' in the method | Avoid / low success

Plant snapshot

Habit	Erect or sprawling shrub, 0.5–3 m (usually 1–2 m). Wedge-shaped leaves, toothed near the tip.
Flowering	Jan – Mar or Jun – Nov (yellow).
Habitat	Coastal sand dunes and swampy places near the coast.
Local note	Recorded in the City of Joondalup. One of the more vigorous hibbertias – useful for screening.

Propagation options

Method	Suitability	Notes	Details in
Cuttings	Main method	Commonly propagated by cuttings.	This guide
Seed	Difficult	0% in a nursery test; Hibbertia seed is dormant.	–
Division	Not used	–	–

Propagation method

Item	Details
Best time to take cuttings	**Mar – May**, between the summer and winter flowering periods, once new growth has firmed. Oct – Feb is workable using non-flowering shoots. Avoid Jun – Sep (cool, and plants are flowering).
Wood type	**Semi-hardwood** – current season's growth that has begun to firm.
Cutting size	75–100 mm.
Selecting material	**Non-flowering** shoots from healthy plants. Snip off any buds.
Pre-treatment	• Strip leaves from the lower half. • Reduce large leaves by about one-third. • Light wound on firmer stems. • Dip the base in rooting hormone.
Rooting hormone	Semi-hardwood strength, 3 g/L IBA (**Clonex Purple**).
Propagation mix	Very free-draining, sand-based mix. Use fresh mix and clean tubes – hibbertias are prone to *Phytophthora*.
Environment	Humid frame, bright shade, bottom heat. Moist, never wet; good airflow.
Expected strike time	Roughly **4–8 weeks** in warm weather (typical for the genus). Record your results.
Post-treatment / aftercare	• Standard hardening off. • Pot on into a free-draining native mix; avoid overwatering. • Tip-prune for a bushier plant.

© 2026 Mitchell Donald. All rights reserved.

Common problems

Problem	Likely cause and fix
Cuttings collapse at the base	Too wet, or root rot. Improve drainage, use clean mix, remove affected cuttings.
Cuttings flower instead of rooting	Material taken from flowering shoots. Use non-flowering shoots.
Slow to strike	Cool weather. Use bottom heat or wait for the Mar – May window.

Notes

Difficulty	**Easy–Moderate (provisional).** Commonly grown from cuttings. Hibbertias usually strike well, but *H. hypericoides* shows not every WA species does – set extra cuttings until your strike rate is known.
Other methods	Seed – 0% germination in the one Alcoa nursery test; *Hibbertia* seed has complex dormancy. Layering – check low, sprawling stems for roots.
Confidence	**Medium.** Commonly propagated from cuttings in the nursery trade; little published detail on timing or strike rate.
Sources	See References: 1, 4, 9, 10, 32, 39, 40

Nursery records

Log each batch here to confirm or update the timing and difficulty above.

Date set	Stock source / provenance	No. set	No. struck	Strike %	Notes

© 2026 Mitchell Donald. All rights reserved.

Hibbertia hypericoides

Yellow Buttercups

Family **Dilleniaceae**

DIFFICULTY	WOOD TYPE	BEST MONTHS	CUTTING LENGTH	HORMONE (IBA)
Difficult	**Semi-hardwood**	**Dec – Feb**	**50–75 mm**	**3 g/L IBA (trial range)**

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Cuttings	Best	Best	OK	–	–	–	–	–	–	–	OK	Best

Best window Workable – see 'Selecting material' in the method Avoid / low success

Plant snapshot

Habit	Spreading shrub, 0.3–0.8 m. Densely hairy branches; small leaves with rolled edges and white-hairy undersides.
Flowering	Apr – Nov (bright yellow, notched petals).
Habitat	Sandy soils in Banksia woodland and shrubland; one of the most common understorey plants between Perth and Albany. Resprouts from its rootstock after fire.
Local note	Recorded in the City of Joondalup.

Propagation options

Method	Suitability	Notes	Details in
Cuttings	Difficult	Known to be difficult; trial protocol on the cuttings page.	This guide
Seed	Difficult	Complex seed dormancy; rarely practical.	–
Division	Not used	Resprouts from a rootstock; not practical to divide.	–

Propagation method

Item	Details
Best time to take cuttings	**Dec – Feb**, after flowering finishes and once the new flush has started to firm. Nov and Mar are workable. Avoid Apr – Oct (flowering period).
Wood type	**Semi-hardwood** – new growth that is just firming. Soft tips wilt quickly; old wood rarely roots.
Cutting size	50–75 mm.
Selecting material	Vigorous **non-flowering** shoots. Well-watered, actively growing stock plants (e.g. nursery-held stock) usually give better material than stressed bushland plants.
Pre-treatment	• Strip lower leaves by cutting, not pulling. • Light wound on the firmer stems. • Dip the base in rooting hormone.
Rooting hormone	Start with 3 g/L IBA (**Clonex Purple**). Because results are unreliable, split trial batches across 1.5, 3 and 8 g/L (Green, Purple, Red) and record which works best.
Propagation mix	Very free-draining, fresh, clean mix. Strict hygiene – hibbertias are prone to *Phytophthora* root rot.
Environment	Humid frame, bottom heat, bright shade. Mist lightly – the hairy leaf undersides hold water. Good airflow.
Expected strike time	Slow and variable – allow **8–16 weeks**. Leave cuttings in as long as they stay green and healthy.

© 2026 Mitchell Donald. All rights reserved.

Item	Details
Post-treatment / aftercare	• Harden off slowly (extend each stage if cuttings wilt). • Pot on gently, disturbing roots as little as possible. • Avoid overwatering; keep benches clean.

Common problems

Problem	Likely cause and fix
Cuttings stay green but don't root	Leave longer; check for heavy callus and trim it; try a different hormone strength.
Cuttings collapse or rot	Too wet or root rot. Improve drainage and hygiene; remove affected cuttings.
Rooted cuttings die after potting on	Roots damaged or hardening too fast. Use single tubes and harden off more slowly.

Notes

Difficulty	**Difficult.** DBCA notes it is difficult to grow from seed or cuttings, and specialist Perth nurseries describe it as a species other producers often fail with. Set large batches and keep detailed records.
Other methods	Seed – also difficult: WA *Hibbertia* seed has complex dormancy, and treatments are still the subject of research. If numbers are needed quickly, consider buying in tubestock from a specialist grower.
Confidence	**Low.** Its difficulty is well documented (DBCA; specialist WA nurseries), but no detailed cutting protocol is published. Treat the method below as a trial protocol.
Sources	See References: 1, 4, 10, 11, 27, 30, 31

Nursery records

Log each batch here to confirm or update the timing and difficulty above.

Date set	Stock source / provenance	No. set	No. struck	Strike %	Notes

© 2026 Mitchell Donald. All rights reserved.

Hibbertia racemosa

Stalked Guinea Flower

Family **Dilleniaceae**

DIFFICULTY	WOOD TYPE	BEST MONTHS	CUTTING LENGTH	HORMONE (IBA)
Moderate *	Semi-hardwood	Jan – Apr	50–75 mm	1.5–3 g/L IBA

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Cuttings	Best	Best	Best	Best	OK	–	–	–	–	–	–	OK

Best window | Workable – see 'Selecting material' in the method | Avoid / low success

Plant snapshot

Habit	Small erect or ascending, spreading shrub, 0.1–0.75 m.
Flowering	Jul – Dec (yellow, on long stalks).
Habitat	Grey, white or yellow sand in coastal areas: dunes, plains and limestone.
Local note	Recorded in the City of Joondalup. Grey-leaved and prostrate forms are sold in the trade – keep forms separate and labelled.

Propagation options

Method	Suitability	Notes	Details in
Cuttings	Main method	Semi-hardwood cuttings (provisional rating).	This guide
Seed	Difficult	Hibbertia seed is dormant and unreliable.	–
Division	Possible	Low stems may root where they touch the soil (layering).	–

Propagation method

Item	Details
Best time to take cuttings	**Jan – Apr**, after flowering finishes and new growth has begun to firm. Dec and May are workable. Avoid Jun – Nov (cool weather, then flowering).
Wood type	**Semi-hardwood** – new growth that has started to harden. Firm tip cuttings also work.
Cutting size	50–75 mm long. Stems are fine and leaves small, so keep cuttings short.
Selecting material	**Non-flowering** side shoots from healthy plants. Snip off any flower stalks or buds.
Pre-treatment	• Strip leaves from the lower half (cut rather than pull on fine stems). • Wounding not needed on fine stems; a light wound is optional on firmer material. • Dip the base in rooting hormone.
Rooting hormone	Semi-hardwood strength (3 g/L IBA, e.g. **Clonex Purple**). For softer tips use softwood strength (1.5 g/L IBA, e.g. **Clonex Green**).
Propagation mix	Very free-draining, sand-based mix. Use fresh, clean mix and clean tubes – hibbertias are prone to *Phytophthora* root rot.
Environment	Humid frame, bright shade, bottom heat. Keep the mix moist but never wet; good airflow.
Expected strike time	Roughly **4–8 weeks** in warm weather; may be slower. Record your results.
Post-treatment / aftercare	• Standard hardening off (2 weeks closed by day/open at night, then 2 weeks shade). • Pot on into a free-draining native mix; avoid overwatering. • Keep good hygiene on benches to limit root rot.

© 2026 Mitchell Donald. All rights reserved.

Common problems

Problem	Likely cause and fix
Cuttings collapse at the base	Too wet, or root rot. Improve drainage, use clean mix, remove affected cuttings.
Few roots, cuttings flower	Material was taken from flowering shoots. Use non-flowering side shoots.
Slow to strike in winter	Use bottom heat, or wait for the Jan – Apr window.

Notes

Difficulty	**Moderate (provisional).** Hibbertias are generally easy from cuttings, but some WA species (e.g. *H. hypericoides*) are known to be difficult. Set larger batches until your own strike rate is known.
Other methods	Seed – generally unreliable in *Hibbertia* (complex dormancy). Layering – several hibbertias root where stems touch the ground; check low stems on prostrate forms and pot up any rooted pieces.
Confidence	**Medium.** Little species-specific data published. Based on how the genus behaves and this species' growth cycle.
Sources	See References: 1, 4, 8, 9, 10, 11

Nursery records

Log each batch here to confirm or update the timing and difficulty above.

Date set	Stock source / provenance	No. set	No. struck	Strike %	Notes

© 2026 Mitchell Donald. All rights reserved.

Hibbertia spicata

No widely used common name

Perth coastal plants are now *Hibbertia leptotheca* (formerly *H. spicata* subsp. *leptotheca*)

Family **Dilleniaceae**

DIFFICULTY	WOOD TYPE	BEST MONTHS	CUTTING LENGTH	HORMONE (IBA)
Moderate *	Semi-hardwood	Jan – Mar	50–75 mm	1.5–3 g/L IBA

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Cuttings	Best	Best	Best	OK	–	–	–	–	–	–	–	OK

Best window | Workable – see 'Selecting material' in the method | Avoid / low success

Plant snapshot

Habit	Small erect or spreading shrub, 0.2–0.7 m (*H. leptotheca* 0.2–0.5 m).
Flowering	Jul – Nov (yellow).
Habitat	Sand, limestone and laterite; near-coastal limestone ridges, outcrops and cliffs.
Local note	A recent revision split *H. spicata* (in the broad sense) into six species. True *H. spicata* is recorded around Geraldton – Shark Bay and Augusta – Margaret River. The Perth coastal plant is *H. leptotheca* (recorded in Wanneroo). It is a **Priority Three** species, so take cuttings only from nursery stock or under the right DBCA licence, and check provenance labels.

Propagation options

Method	Suitability	Notes	Details in
Cuttings	Main method	Most hibbertias are grown from cuttings (provisional).	This guide
Seed	Difficult	Dormant; 9% in the one Alcoa test; GA may help.	–
Division	Not used	Shrub – not divided.	–

Propagation method

Item	Details
Best time to take cuttings	**Jan – Mar**, after flowering, once new growth has firmed. *H. empetrifolia* cuttings strike readily in Jan – Feb. Dec and Apr are workable. Avoid May – Nov (cool, and plants are flowering).
Wood type	**Semi-hardwood** – current season's growth that has begun to firm.
Cutting size	50–75 mm.
Selecting material	**Non-flowering** shoots from healthy plants. Snip off any buds.
Pre-treatment	• Strip leaves from the lower half. • Dip the base in rooting hormone.
Rooting hormone	1.5 g/L IBA (**Clonex Green**) for soft tips; 3 g/L (**Clonex Purple**) for firmer stems.
Propagation mix	Very free-draining, sand-based mix. Use fresh mix and clean tubes – hibbertias are prone to root rot.
Environment	Humid frame, bright shade, bottom heat. Moist, never wet; good airflow.
Expected strike time	Not documented – check from about 6 weeks and record the result.
Post-treatment / aftercare	• Standard hardening off. • Pot on into a free-draining native mix; avoid overwatering.

© 2026 Mitchell Donald. All rights reserved.

Common problems

Problem	Likely cause and fix
Cuttings collapse at the base	Too wet, or root rot. Improve drainage; clean mix; remove affected cuttings.
Cuttings flower instead of rooting	Material from flowering shoots. Use non-flowering shoots.
Very low strike	Some WA hibbertias are hard. Try softer tips, bottom heat, and record which works.

Notes

Difficulty	**Moderate (provisional).** Untested for this species. Set extra cuttings until your strike rate is known.
Other methods	Seed – dormant: 9% in the one Alcoa nursery test (about 30 seedlings per gram). Removing the seed coat or using gibberellic acid (25 mg/L) can help (WA seed notes).
Confidence	**Low.** No species data. WA seed notes say most hibbertias are grown from cuttings, but *H. hypericoides* shows some WA species are hard.
Sources	See References: 1, 4, 9, 13, 32, 69, 70

Nursery records

Log each batch here to confirm or update the timing and difficulty above.

Date set	Stock source / provenance	No. set	No. struck	Strike %	Notes

© 2026 Mitchell Donald. All rights reserved.

Hypocalymma robustum

Swan River Myrtle

Family **Myrtaceae**

DIFFICULTY	WOOD TYPE	BEST MONTHS	CUTTING LENGTH	HORMONE (IBA)
Easy	Semi-hardwood	Dec – Mar	75–100 mm	3 g/L IBA

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Cuttings	Best	Best	Best	OK	–	–	–	–	–	–	OK	Best

Best window | Workable – see 'Selecting material' in the method | Avoid / low success

Plant snapshot

Habit	Erect shrub, 0.4–1 (–1.5) m.
Flowering	Jun – Nov (pink to pink-red, along the stems).
Habitat	Gravelly lateritic soils and sandy soils; undulating ground and ridges.
Local note	Recorded in Joondalup and Wanneroo. Prune yearly for a dense, bushy plant.

Propagation options

Method	Suitability	Notes	Details in
Cuttings	Main method	Strikes readily from hardened current season's growth.	This guide
Seed	Possible	Smoke water likely helps; about 35% in one test.	–
Division	Not used	–	–

Propagation method

Item	Details
Best time to take cuttings	**Dec – Mar** (January is suggested for the Geraldton wax group), once the current season's growth has hardened. Nov and Apr are workable. Avoid May – Oct (cool, then flowering).
Wood type	**Semi-hardwood** – hardened current season's growth (ANPSA).
Cutting size	75–100 mm.
Selecting material	Non-flowering tips from healthy plants.
Pre-treatment	• Carefully remove the leaves from the lower two-thirds. • Wound the base. • Dip the base in rooting hormone.
Rooting hormone	Semi-hardwood strength, 3 g/L IBA (**Clonex Purple**) – wounding and hormone both improve results.
Propagation mix	Standard free-draining cutting mix. Clean mix and tubes – the genus includes dieback-affected species.
Environment	Humid frame, bright shade, bottom heat.
Expected strike time	Not documented – usually quick; check from about 4 weeks.
Post-treatment / aftercare	• Standard hardening off. • Pot on into a free-draining native mix; tip-prune for bushiness.

© 2026 Mitchell Donald. All rights reserved.

Common problems

Problem	Likely cause and fix
Cuttings brown off	Too soft or too dry. Use hardened growth; keep humidity up.
Base rotting	Mix too wet. Water less, improve drainage.
Leggy plants	Not pruned. Tip-prune young plants.

Notes

Difficulty	**Easy.** Cuttings are the preferred method and strike readily (ANPSA).
Other methods	Seed – possible; smoke water is likely to help (about 35% in a single Alcoa test with smoke).
Confidence	**Medium.** ANPSA says cuttings are the preferred method and strike readily; size and treatment follow the related Geraldton wax group.
Sources	See References: 4, 32, 53, 54, 55

Nursery records

Log each batch here to confirm or update the timing and difficulty above.

Date set	Stock source / provenance	No. set	No. struck	Strike %	Notes

© 2026 Mitchell Donald. All rights reserved.

Kennedia prostrata

Scarlet Runner / Running Postman

Family **Fabaceae**

DIFFICULTY	WOOD TYPE	BEST MONTHS	CUTTING LENGTH	HORMONE (IBA)
Easy–Moderate	Semi-hardwood	Dec – Mar	75–100 mm	3 g/L IBA

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Cuttings	Best	Best	Best	OK	–	–	–	–	–	–	OK	Best

Best window | Workable – see 'Selecting material' in the method | Avoid / low success

Plant snapshot

Habit	Prostrate or twining groundcover, spreading to about 2.5 m.
Flowering	Apr – Nov (red pea flowers).
Habitat	Usually sandy, gravelly soils.
Local note	Recorded in Joondalup and Wanneroo.

Propagation options

Method	Suitability	Notes	Details in
Cuttings	Good	Firm current season's growth strikes well.	This guide
Seed	Main method	Hot water + smoke water; about 8 days.	→ Seed Sowing Guide
Division	Not used	–	–

Propagation method

Item	Details
Best time to take cuttings	**Dec – Mar**, when the current season's runners have firmed. Nov and Apr are workable. Avoid May – Oct (flowering and cool weather).
Wood type	**Semi-hardwood** – firm current season's growth (ANPSA).
Cutting size	75–100 mm with 2–3 nodes. One runner can be cut into several pieces.
Selecting material	Healthy runners without flowers or pods.
Pre-treatment	• Remove the leaves from the lower nodes; cut remaining leaflets back by about half. • No wounding needed. • Dip the base in rooting hormone.
Rooting hormone	Semi-hardwood strength, 3 g/L IBA (**Clonex Purple**).
Propagation mix	Standard free-draining cutting mix.
Environment	Humid frame, bright shade, bottom heat.
Expected strike time	Not documented – check from about 4 weeks and record the result.
Post-treatment / aftercare	• Standard hardening off. • Pot on promptly – runners tangle across the bench. Trim runners to keep tubes tidy.

© 2026 Mitchell Donald. All rights reserved.

Common problems

Problem	Likely cause and fix
Cuttings wilt and collapse	Material too soft. Use firmer runners and keep humidity up.
Runners tangled between tubes	Space tubes out and trim runners regularly.
Base rotting	Mix too wet. Water less, improve drainage.

Notes

Difficulty	**Easy–Moderate.** Cuttings of firm growth strike well. Seed is quick and is the usual nursery method.
Confidence	**Medium.** ANPSA reports cuttings of firm current season's growth strike well; timing and size follow standard practice.
Sources	See References: 4, 13, 42

Nursery records

Log each batch here to confirm or update the timing and difficulty above.

Date set	Stock source / provenance	No. set	No. struck	Strike %	Notes

© 2026 Mitchell Donald. All rights reserved.

Lechenaultia linarioides

Yellow Leschenaultia

Family **Goodeniaceae**

DIFFICULTY	WOOD TYPE	BEST MONTHS	CUTTING LENGTH	HORMONE (IBA)
Easy *	Softwood – semi-hardwood	Feb – Apr	50–75 mm	1.5–3 g/L IBA

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Cuttings	OK	Best	Best	Best	OK	–	–	–	–	OK	OK	OK

Best window | Workable – see 'Selecting material' in the method | Avoid / low success

Plant snapshot

Habit	Sprawling, open shrub to 1.5 m, with narrow, linear leaves.
Flowering	Can flower in any month. Flowers yellow with red or pink tones, sometimes cream.
Habitat	Sand (often over limestone) and sandy clay; sandplains, low dunes, ridges and winter-wet flats.
Local note	Recorded in the Wanneroo LGA.

Propagation options

Method	Suitability	Notes	Details in
Cuttings	Main method	Cuttings strike easily in this genus.	This guide
Seed	Difficult	Slow; trays may need a second season.	–
Division	Possible	Some leschenaultias sucker – pot up rooted suckers.	–

Propagation method

Item	Details
Best time to take cuttings	**Feb – Apr** (late summer to autumn) while it is warm. One grower reports roots in 4–6 weeks from late-February leschenaultia cuttings. Oct – Jan and May are workable. Avoid Jun – Sep.
Wood type	**Softwood to semi-hardwood** tip cuttings.
Cutting size	50–75 mm.
Selecting material	**Non-flowering** tips. This species can flower in any month, so check each cutting and remove buds.
Pre-treatment	• Strip leaves from the lower half (narrow leaves strip easily). • No wounding needed. • Dip the base in rooting hormone.
Rooting hormone	1.5 g/L IBA (**Clonex Green**) for soft tips; 3 g/L (**Clonex Purple**) for firmer material.
Propagation mix	Very free-draining – leschenaultias dislike wet roots.
Environment	Humid frame, bottom heat, bright shade. Keep the mix moist, never wet.
Expected strike time	About **4–6 weeks** in warm weather.
Post-treatment / aftercare	• Standard hardening off. • Pot on into a free-draining mix; water sparingly. • Tip-prune to counter the sprawling habit.

© 2026 Mitchell Donald. All rights reserved.

Common problems

Problem	Likely cause and fix
Base rotting	Overwatering. Water less often; add coarse sand or perlite to the mix.
Wilting in the frame	Material too soft or humidity too low. Use slightly firmer tips; keep the frame closed.
Cuttings flowering	Remove buds; pick vegetative shoots.

Notes

Difficulty	**Easy (provisional).** *Lechenaultia* cuttings generally strike very easily; seed is difficult.
Other methods	Seed – slow and unreliable. The Geraldton community nursery soaked seed in smoke water, saw germination at about 46 days, and kept trays for a second year. Division – some leschenaultias sucker; pot up rooted suckers.
Confidence	**Medium.** No species-specific cutting data found. Based on the genus, whose cuttings strike very easily.
Sources	See References: 1, 4, 12, 13, 24, 25

Nursery records

Log each batch here to confirm or update the timing and difficulty above.

Date set	Stock source / provenance	No. set	No. struck	Strike %	Notes

© 2026 Mitchell Donald. All rights reserved.

Leucophyta brownii

Cushion Bush

Family **Asteraceae**

DIFFICULTY	WOOD TYPE	BEST MONTHS	CUTTING LENGTH	HORMONE (IBA)
Easy–Moderate *	Semi-hardwood	Mar – May; Sep – Oct	50–75 mm	1.5–3 g/L IBA

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Cuttings	–	OK	Best	Best	Best	OK	–	OK	Best	Best	OK	–

Best window | Workable – see 'Selecting material' in the method | Avoid / low success

Plant snapshot

Habit	Compact, rounded shrub to about 1 m, densely covered in silvery-white woolly hairs.
Flowering	Dec or Jan – Feb (small yellow button heads).
Habitat	Coastal foredunes and cliffs.
Local note	Recorded in the Wanneroo LGA. Very salt-tolerant; grows poorly in high humidity.

Propagation options

Method	Suitability	Notes	Details in
Cuttings	Main method	Semi-hardened cuttings are the preferred method.	This guide
Seed	Good	No pre-treatment; 10–30 days.	→ Seed Sowing Guide
Division	Not used	–	–

Propagation method

Item	Details
Best time to take cuttings	**Mar – May**, after flowering, and **Sep – Oct** as spring growth firms. Feb, Jun, Aug and Nov are workable. Avoid Dec – Jan (flowering and heat) and Jul (cold).
Wood type	**Semi-hardwood** – ANBG notes cuttings should come from semi-hardened branches.
Cutting size	50–75 mm tip cuttings.
Selecting material	Non-flowering tips from compact, healthy plants. Avoid soft, sappy growth.
Pre-treatment	• Rub the small leaves off the lower half. • No wounding. • Dip the base in rooting hormone.
Rooting hormone	1.5 g/L IBA (**Clonex Green**) for firm tips; 3 g/L for firmer material.
Propagation mix	Very free-draining – add extra coarse sand or perlite.
Environment	**Keep it airy.** The woolly foliage rots in humidity: use a well-ventilated frame in bright light with little or no mist. Bottom heat helps. Keep the mix only just moist.
Expected strike time	Not documented – check from about 4 weeks and record the result.
Post-treatment / aftercare	• Harden off promptly and grow on in full sun. • Free-draining mix; never let pots sit wet – plants die easily if waterlogged. • Tip-prune for a dense cushion.

© 2026 Mitchell Donald. All rights reserved.

Common problems

Problem	Likely cause and fix
Cuttings turn grey-brown and mushy	Too humid or wet. Reduce mist, open the frame, improve airflow.
Slow to root in winter	Use bottom heat, or wait for the autumn or spring window.
Loses its dense shape	Tip-prune young plants.

Notes

Difficulty	**Easy–Moderate (provisional).** Cuttings are the preferred method; the main risk is rot from too much moisture or humidity.
Confidence	**Medium.** ANBG growing notes say semi-hardened cuttings are the preferred method; timing and details follow general daisy practice.
Sources	See References: 4, 34, 35

Nursery records

Log each batch here to confirm or update the timing and difficulty above.

Date set	Stock source / provenance	No. set	No. struck	Strike %	Notes

© 2026 Mitchell Donald. All rights reserved.

Leucopogon parviflorus

Coast Beard-heath

Family **Ericaceae**

DIFFICULTY	WOOD TYPE	BEST MONTHS	CUTTING LENGTH	HORMONE (IBA)
Difficult *	Semi-hardwood	Nov – Jan	80–100 mm	3 g/L IBA

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Cuttings	Best	OK	OK	OK	OK	–	–	–	–	–	Best	Best

Best window | Workable – see 'Selecting material' in the method | Avoid / low success

Plant snapshot

Habit	Erect, densely branched shrub or small tree, usually 0.3–3 m (to 5 m).
Flowering	Feb – Mar or Jun – Oct (white), followed by round green fruit that turn translucent white when ripe.
Habitat	Sandy soils over limestone or granite; coastal dunes and limestone.
Local note	Recorded in Joondalup and Wanneroo. A frontline coastal shrub that buffers wind; birds and lizards eat the fruit (State Flora SA).

Propagation options

Method	Suitability	Notes	Details in
Cuttings	Difficult	Low strike and fine roots – trial protocol (provisional).	This guide
Seed	Difficult	Dormant stones; peroxide + smoke water worked best in SA trials.	→ Seed Sowing Guide
Division	Not used	Shrub – not divided.	–

Propagation method

Item	Details
Best time to take cuttings	**Nov – Jan** (estimate), once the spring growth has firmed and before the summer heat peaks. Feb – May are workable with non-flowering shoots. Avoid Jun – Oct (main flowering, and cool).
Wood type	**Semi-hardwood** – firm current season's growth gives the best results (ANPSA).
Cutting size	80–100 mm – the short end of the 80–250 mm range used for epacrid cuttings in Sydney trials (Doig).
Selecting material	Non-flowering tips from healthy, vigorous plants. Take plenty – strike rates are low.
Pre-treatment	• Strip leaves from the lower half. • Light wound on firmer stems. • Dip the base in rooting hormone – ANPSA advises using one.
Rooting hormone	Doig used a medium-strength hormone powder for epacrids; use 3 g/L IBA (**Clonex Purple**) as the equivalent.
Propagation mix	Very free-draining sand-based cutting mix.
Environment	Humid frame in bright shade, not heavy mist. Doig got at least 30% strike with epacrids in a cold, non-misting frame. Bottom heat is worth trialling on part of the batch.
Expected strike time	**Slow** – epacrid cuttings were held for 6–12 months in trials. Check from about 8 weeks and record the result.
Post-treatment / aftercare	• The roots are **very fine and easily damaged** (Tasmania) – pot up with the mix ball intact; don't tease the roots. • Harden off slowly; keep potted plants in light shade at first.

© 2026 Mitchell Donald. All rights reserved.

Common problems

Problem	Likely cause and fix
Cuttings sit for months without roots	Normal for epacrids. Keep them going; try bottom heat or a different month on the next batch.
Roots break when potting up	Pot up before roots fill the cell, and move the whole plug without disturbing it.
Cuttings blacken or rot	Too wet or humid. Less mist, better drainage and airflow.

Notes

Difficulty	**Difficult (provisional).** ANPSA calls cuttings unreliable, and Tasmanian growers report low strike rates even though cuttings are their usual method.
Other methods	Seed – difficult (dormant stones). South Australian trials reached 80–100% with special treatments; see the Seed Sowing Guide.
Confidence	**Low.** Sources agree cuttings strike poorly (low strike, very fine roots); no WA trial data. Treat as a trial protocol and record results.
Sources	See References: 1, 4, 13, 79, 80, 81, 82, 86

Nursery records

Log each batch here to confirm or update the timing and difficulty above.

Date set	Stock source / provenance	No. set	No. struck	Strike %	Notes

© 2026 Mitchell Donald. All rights reserved.

Melaleuca cardiophylla

Tangling Melaleuca

Family **Myrtaceae**

DIFFICULTY	WOOD TYPE	BEST MONTHS	CUTTING LENGTH	HORMONE (IBA)
Easy *	Semi-hardwood	Feb – Apr	75–100 mm	3 g/L IBA

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Cuttings	OK	Best	Best	Best	OK	–	–	–	–	–	–	OK

Best window | Workable – see 'Selecting material' in the method | Avoid / low success

Plant snapshot

Habit	Dense, prickly, erect to spreading shrub, 0.2–2 m (to 4 m), with tangled branches.
Flowering	Aug – Dec or Jan (white-cream).
Habitat	White, grey or brown sand on limestone ridges, outcrops and dunes.
Local note	Recorded in Joondalup and Wanneroo. Regrows after fire from epicormic shoots and the base.

Propagation options

Method	Suitability	Notes	Details in
Cuttings	Good	Reliable in this genus; keeps a chosen form.	This guide
Seed	Main method	Surface-sow fine seed from old capsules; no pre-treatment.	→ Seed Sowing Guide
Division	Not used	Shrub – not divided.	–

Propagation method

Item	Details
Best time to take cuttings	**Feb – Apr**, after flowering, once new growth has firmed. May, Dec and Jan are workable with non-flowering shoots. Avoid Jun – Jul (cool) and Aug – Nov (main flowering).
Wood type	**Semi-hardwood** – firm current season's growth.
Cutting size	75–100 mm (ANPSA).
Selecting material	Straight, non-flowering tips. **Wear gloves** – the leaves are sharp and branches tangle.
Pre-treatment	• Strip leaves from the lower half to two-thirds. • Wound the base on firmer stems. • Dip the base in rooting hormone.
Rooting hormone	Semi-hardwood strength, 3 g/L IBA (**Clonex Purple**).
Propagation mix	Standard free-draining cutting mix.
Environment	Humid frame, bright shade, bottom heat.
Expected strike time	Est. **4–8 weeks** in warm weather, as for other melaleucas. Record your results.
Post-treatment / aftercare	• Standard hardening off. • Pot on into a free-draining native mix; tip-prune for a dense plant.

© 2026 Mitchell Donald. All rights reserved.

Common problems

Problem	Likely cause and fix
Callus, no roots	Trim and re-set; lower the hormone strength next batch.
Hard to get neat material	Tangled, prickly stems. Take straight outer tips and trim leaves carefully.
Base rotting	Mix too wet. Water less.

Notes

Difficulty	**Easy (provisional).** Melaleucas are generally reliable from cuttings.
Other methods	Seed – easy; capsules hold seed on the plant (see the Seed Sowing Guide).
Confidence	**Medium.** Based on ANPSA *Melaleuca* genus guidance; Australian Plants Society lists cuttings for this species.
Sources	See References: 4, 13, 41, 83

Nursery records

Log each batch here to confirm or update the timing and difficulty above.

Date set	Stock source / provenance	No. set	No. struck	Strike %	Notes

© 2026 Mitchell Donald. All rights reserved.

Melaleuca huegelii

Chenille Honeymyrtle

Family **Myrtaceae**

DIFFICULTY	WOOD TYPE	BEST MONTHS	CUTTING LENGTH	HORMONE (IBA)
Easy *	Semi-hardwood	Feb – Apr	75–100 mm	3 g/L IBA

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Cuttings	OK	Best	Best	Best	OK	–	–	–	–	–	–	OK

Best window | Workable – see 'Selecting material' in the method | Avoid / low success

Plant snapshot

Habit	Shrub or small tree, 0.5–5 m.
Flowering	Sep – Dec or Jan (pink, white or pink-purple).
Habitat	Sand; limestone cliffs, coastal plains and dunes.
Local note	Recorded in Joondalup and Wanneroo. Cuttings keep a chosen flower colour.

Propagation options

Method	Suitability	Notes	Details in
Cuttings	Good	Reliable in this genus; keeps a chosen form.	This guide
Seed	Main method	Surface-sow fine seed; no pre-treatment.	→ Seed Sowing Guide
Division	Not used	Woody shrub.	–

Propagation method

Item	Details
Best time to take cuttings	**Feb – Apr**, after flowering, once new growth has firmed. Jan, May and Dec are workable. Avoid Jun – Nov.
Wood type	**Semi-hardwood** – firm current season's growth.
Cutting size	75–100 mm (ANPSA).
Selecting material	Non-flowering tips. Take from a plant with the flower colour you want.
Pre-treatment	• Strip leaves from the lower half to two-thirds. • Wound the base on firmer stems. • Dip the base in rooting hormone.
Rooting hormone	Semi-hardwood strength, 3 g/L IBA (**Clonex Purple**).
Propagation mix	Standard free-draining cutting mix.
Environment	Humid frame, bright shade, bottom heat.
Expected strike time	Typically **4–8 weeks** in warm weather. Record your results.
Post-treatment / aftercare	• Standard hardening off. • Pot on into a free-draining native mix; tip-prune for bushiness.

© 2026 Mitchell Donald. All rights reserved.

Common problems

Problem	Likely cause and fix
Callus, no roots	Trim and re-set; lower the hormone strength next batch.
Leaves drop in the frame	Too dry or too hot. Shade more and keep humidity steady.
Base rotting	Mix too wet. Water less.

Notes

Difficulty	**Easy (provisional).** Melaleucas are generally reliable from cuttings.
Confidence	**Medium.** Based on ANPSA *Melaleuca* genus guidance – cuttings are reliable in the genus.
Sources	See References: 4, 13, 41

Nursery records

Log each batch here to confirm or update the timing and difficulty above.

Date set	Stock source / provenance	No. set	No. struck	Strike %	Notes

© 2026 Mitchell Donald. All rights reserved.

Melaleuca lanceolata

Rottnest Teatree / Moonah

Family **Myrtaceae**

DIFFICULTY	WOOD TYPE	BEST MONTHS	CUTTING LENGTH	HORMONE (IBA)
Easy *	Semi-hardwood	Mar – Apr	75–100 mm	3 g/L IBA

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Cuttings	–	OK	Best	Best	OK	–	–	–	OK	OK	–	–

Best window | Workable – see 'Selecting material' in the method | Avoid / low success

Plant snapshot

Habit	Shrub or tree, 1–8 m.
Flowering	Mainly in summer – FloraBase's calendar shows Nov – Mar (white-cream).
Habitat	Limestone ridges, coastal cliffs and dunes, salt flats and the edges of salt lakes.
Local note	Recorded in the Wanneroo LGA (not listed for Joondalup). Resists salt spray – a good windbreak or shelterbelt plant.

Propagation options

Method	Suitability	Notes	Details in
Cuttings	Good	Reliable in this genus (provisional).	This guide
Seed	Main method	Surface-sow; no pre-treatment; about 55% (Alcoa).	→ Seed Sowing Guide
Division	Not used	Shrub – not divided.	–

Propagation method

Item	Details
Best time to take cuttings	**Mar – Apr** (estimate) – after the main summer flowering, while it's still warm and new growth has firmed. Feb, May, Sep and Oct are workable with non-flowering shoots. Avoid Jun – Aug (cool) and Nov – Jan (peak flowering).
Wood type	**Semi-hardwood** – firm current season's growth.
Cutting size	75–100 mm (ANPSA).
Selecting material	Straight, non-flowering tips from healthy plants.
Pre-treatment	• Strip leaves from the lower half to two-thirds. • Wound the base on firmer stems. • Dip the base in rooting hormone.
Rooting hormone	Semi-hardwood strength, 3 g/L IBA (**Clonex Purple**).
Propagation mix	Standard free-draining cutting mix.
Environment	Humid frame, bright shade, bottom heat.
Expected strike time	Est. **4–8 weeks** in warm weather, as for other melaleucas. Record your results.
Post-treatment / aftercare	• Standard hardening off. • Pot on into a free-draining native mix; tip-prune for a bushy plant.

© 2026 Mitchell Donald. All rights reserved.

Common problems

Problem	Likely cause and fix
Callus, no roots	Trim and re-set; lower the hormone strength next batch.
Cuttings flower, not root	Flowering shoots taken. Use non-flowering tips.
Base rotting	Mix too wet. Water less.

Notes

Difficulty	**Easy (provisional).** Melaleucas are generally reliable from cuttings.
Other methods	Seed – easy; about 55% germination with no pre-treatment (see the Seed Sowing Guide).
Confidence	**Medium.** Based on ANPSA *Melaleuca* genus guidance – no cuttings data found for this species.
Sources	See References: 4, 13, 41, 98

Nursery records

Log each batch here to confirm or update the timing and difficulty above.

Date set	Stock source / provenance	No. set	No. struck	Strike %	Notes

© 2026 Mitchell Donald. All rights reserved.

Melaleuca systena

Coastal Honeymyrtle

Family **Myrtaceae**

DIFFICULTY	WOOD TYPE	BEST MONTHS	CUTTING LENGTH	HORMONE (IBA)
Easy *	Semi-hardwood	Dec – Jan; Apr – May	75–100 mm	3 g/L IBA

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Cuttings	Best	OK	OK	Best	Best	OK	–	–	–	–	OK	Best

Best window Workable – see 'Selecting material' in the method Avoid / low success

Plant snapshot

Habit	Erect to spreading shrub, usually 0.5–2 m.
Flowering	Feb – Mar or Aug – Dec (yellow-cream).
Habitat	Sand over laterite or limestone; stabilised coastal dunes and rocky limestone.
Local note	Recorded in Joondalup and Wanneroo. Seed germination is low (about 14% in nursery tests), so cuttings are worth using.

Propagation options

Method	Suitability	Notes	Details in
Cuttings	Good	Reliable in this genus; a good option given low seed germination.	This guide
Seed	Main method	Surface-sow fine seed; no pre-treatment.	→ Seed Sowing Guide
Division	Not used	Woody shrub.	–

Propagation method

Item	Details
Best time to take cuttings	**Dec – Jan** and **Apr – May** – the gaps between its two flowering periods. Feb – Mar, Jun and Nov are workable with non-flowering shoots. Avoid Jul – Oct.
Wood type	**Semi-hardwood** – firm current season's growth.
Cutting size	75–100 mm (ANPSA).
Selecting material	Non-flowering tips from healthy plants.
Pre-treatment	• Strip leaves from the lower half to two-thirds. • Wound the base on firmer stems. • Dip the base in rooting hormone.
Rooting hormone	Semi-hardwood strength, 3 g/L IBA (**Clonex Purple**).
Propagation mix	Standard free-draining cutting mix.
Environment	Humid frame, bright shade, bottom heat. Autumn batches root more slowly.
Expected strike time	Typically **4–8 weeks** in warm weather; longer in May. Record your results.
Post-treatment / aftercare	• Standard hardening off. • Pot on into a free-draining native mix.

© 2026 Mitchell Donald. All rights reserved.

Common problems

Problem	Likely cause and fix
Callus, no roots	Trim and re-set; lower the hormone strength next batch.
Cuttings flower	Taken from flowering shoots. Use non-flowering tips, between flowering periods.
Base rotting	Mix too wet. Water less.

Notes

Difficulty	**Easy (provisional).** Melaleucas are generally reliable from cuttings.
Confidence	**Medium.** Based on ANPSA *Melaleuca* genus guidance – cuttings are reliable in the genus.
Sources	See References: 4, 13, 41

Nursery records

Log each batch here to confirm or update the timing and difficulty above.

Date set	Stock source / provenance	No. set	No. struck	Strike %	Notes

© 2026 Mitchell Donald. All rights reserved.

Myoporum caprarioides

Slender Myoporum

Family **Scrophulariaceae**

DIFFICULTY	WOOD TYPE	BEST MONTHS	CUTTING LENGTH	HORMONE (IBA)
Easy *	Semi-hardwood	Nov – Mar	75–100 mm	3 g/L IBA

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Cuttings	Best	Best	Best	OK	–	–	–	–	–	OK	Best	Best

Best window | Workable – see 'Selecting material' in the method | Avoid / low success

Plant snapshot

Habit	Erect, slender, compact or sprawling shrub, (0.3–) 0.5–3 m.
Flowering	Variable – from Feb, Apr or Jun to Dec (white, sometimes blue-pink tinged).
Habitat	White-grey to black sand, peaty sand and clay loam; seasonally wet flats, swamps, dunes and limestone ridges.
Local note	Recorded in Joondalup and Wanneroo. Suits sites that stay damp in winter. Small berries attract birds.

Propagation options

Method	Suitability	Notes	Details in
Cuttings	Main method	Hardened current season's growth strikes readily in the genus.	This guide
Seed	Possible	Not documented; the related M. insulare is fermented + smoke.	–
Division	Not used	Woody shrub.	–

Propagation method

Item	Details
Best time to take cuttings	**Nov – Mar**, in the warm months once current growth has hardened. Oct and Apr are workable. Avoid May – Sep (slow rooting).
Wood type	**Semi-hardwood** – hardened current season's growth.
Cutting size	75–100 mm.
Selecting material	Non-flowering tips from a plant with the form you want (upright or sprawling).
Pre-treatment	• Strip leaves from the lower half. • Light wound on firmer stems. • Dip the base in rooting hormone.
Rooting hormone	Semi-hardwood strength, 3 g/L IBA (**Clonex Purple**).
Propagation mix	Standard free-draining cutting mix.
Environment	Humid frame, bright shade, bottom heat.
Expected strike time	Not documented – usually quick in the genus; check from about 4 weeks.
Post-treatment / aftercare	• Standard hardening off. • Pot on into a native mix that holds some moisture.

© 2026 Mitchell Donald. All rights reserved.

Common problems

Problem	Likely cause and fix
Base rotting	Mix too wet in the frame. Water less; improve drainage.
Cuttings wilt	Material too soft. Use hardened growth.
Leggy plants	Tip-prune young plants.

Notes

Difficulty	**Easy (provisional).** Most myoporums strike readily; this species is untested here.
Other methods	Seed – not documented for this species. The related *M. insulare* is fermented, then soaked in smoke water (Geraldton).
Confidence	**Medium.** No species data. ANPSA: most *Myoporum* strike readily from hardened current season's growth, the preferred method.
Sources	See References: 4, 12, 62, 63

Nursery records

Log each batch here to confirm or update the timing and difficulty above.

Date set	Stock source / provenance	No. set	No. struck	Strike %	Notes

© 2026 Mitchell Donald. All rights reserved.

Myoporum insulare

Blueberry Tree / Common Boobialla

Family **Scrophulariaceae**

DIFFICULTY	WOOD TYPE	BEST MONTHS	CUTTING LENGTH	HORMONE (IBA)
Easy	Semi-hardwood	Feb – Apr	75–100 mm	3 g/L IBA

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Cuttings	OK	Best	Best	Best	OK	–	–	–	–	OK	OK	OK

Best window | Workable – see 'Selecting material' in the method | Avoid / low success

Plant snapshot

Habit	Dense, low spreading to erect shrub or small tree, 0.25–5 m, with thick, fleshy leaves.
Flowering	Jul – Dec, also Jan – Feb (white, purple-spotted), followed by purple berries.
Habitat	White or grey sand on coastal dunes and limestone.
Local note	Recorded in Joondalup and Wanneroo. Forms range from prostrate to tree-like – take cuttings from a plant with the form you want.

Propagation options

Method	Suitability	Notes	Details in
Cuttings	Main method	Firm young growth roots readily.	This guide
Seed	Good	Ferment berries, smoke water; 15–30 days. Seedling form varies.	→ Seed Sowing Guide
Division	Not used	Woody shrub.	–

Propagation method

Item	Details
Best time to take cuttings	**Feb – Apr** – warm, and after the main flowering. Oct – Jan and May are workable. Avoid Jun – Sep.
Wood type	**Semi-hardwood** – firm young growth.
Cutting size	75–100 mm.
Selecting material	Firm, **non-flowering** young shoots from a stock plant with the growth form you need (prostrate for groundcover, erect for screening).
Pre-treatment	• Strip leaves from the lower half. • Reduce the remaining thick leaves by about one-third. • Dip the base in rooting hormone.
Rooting hormone	3 g/L IBA (**Clonex Purple**). Strikes readily without it, but hormone gives faster, more even rooting.
Propagation mix	Standard sand/coir cutting mix.
Environment	Standard humid frame with bottom heat. Hairless leaves tolerate normal misting.
Expected strike time	About **4–8 weeks** in warm weather.
Post-treatment / aftercare	• Standard hardening off. • Fast-growing – pot on promptly before roots circle. • Tip-prune for a bushier plant.

© 2026 Mitchell Donald. All rights reserved.

Common problems

Problem	Likely cause and fix
Leaf drop in the frame	Too dark or too wet. Give brighter shade and water less.
Roots circling in tubes	Pot on sooner – it grows quickly.
Plants not the expected shape	Seed-grown or mixed stock. Propagate from a known-form stock plant.

Notes

Difficulty	**Easy.** Firm young growth roots readily; even branchlets placed straight into soil can strike.
Other methods	Seed – the Geraldton community nursery ferments ripe (darkest purple) berries for 4–5 days, washes off the pulp and soaks in smoke water; germination took 15–30 days. Seedlings vary in form.
Confidence	**High.** Published sources agree it strikes readily from firm young growth.
Sources	See References: 1, 4, 12, 13, 23

Nursery records

Log each batch here to confirm or update the timing and difficulty above.

Date set	Stock source / provenance	No. set	No. struck	Strike %	Notes

© 2026 Mitchell Donald. All rights reserved.

Olearia axillaris

Coastal Daisybush

Family **Asteraceae**

DIFFICULTY	WOOD TYPE	BEST MONTHS	CUTTING LENGTH	HORMONE (IBA)
Easy *	Semi-hardwood	Aug – Oct	75–100 mm	1.5–3 g/L IBA

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Cuttings	–	–	OK	OK	–	–	OK	Best	Best	Best	OK	–

Best window | Workable – see 'Selecting material' in the method | Avoid / low success

Plant snapshot

Habit	Erect, much-branched grey shrub, 0.2–3 m.
Flowering	Jan – Jul or Nov – Dec (small white to yellow daisies).
Habitat	White, grey or red sand and loam; coastal limestone, sand dunes and rocky hillsides.
Local note	Recorded in Joondalup and Wanneroo.

Propagation options

Method	Suitability	Notes	Details in
Cuttings	Good	Perennial daisies strike quickly from vigorous growth.	This guide
Seed	Main method	Fresh seed, aired 3 months; surface-sow.	→ Seed Sowing Guide
Division	Not used	Woody shrub.	–

Propagation method

Item	Details
Best time to take cuttings	**Aug – Oct**, on vigorous spring growth between flowering periods. Mar – Apr, Jul and Nov are workable with non-flowering shoots. Avoid Dec – Feb and May – Jun.
Wood type	**Semi-hardwood** – vigorous current growth that has begun to firm.
Cutting size	75–100 mm.
Selecting material	Vigorous, healthy, non-flowering shoots (ANPSA).
Pre-treatment	• Strip leaves from the lower half. • No wounding needed. • Dip the base in rooting hormone.
Rooting hormone	1.5 g/L IBA (**Clonex Green**) for softer tips; 3 g/L for firmer material.
Propagation mix	Free-draining: about 3 parts sand to 1 part peat or coir (ANPSA daisy mix).
Environment	Humid frame, bright shade, bottom heat. The grey, hairy leaves rot if kept wet – mist lightly.
Expected strike time	Usually quick – check from about 3–4 weeks.
Post-treatment / aftercare	• Standard hardening off. • Grow on in full sun; tip-prune for a compact plant.

© 2026 Mitchell Donald. All rights reserved.

Common problems

Problem	Likely cause and fix
Leaves go grey-brown and soft	Too wet or humid. Mist less, improve airflow.
Cuttings flower	Taken from flowering shoots. Use vigorous non-flowering growth.
Leggy plants	Too much shade. Move to full sun after hardening off.

Notes

Difficulty	**Easy (provisional).** Perennial daisies usually strike quickly and easily.
Confidence	**Medium.** ANPSA: stem cuttings of most perennial Australian daisies strike quickly from vigorous growth.
Sources	See References: 4, 13, 35

Nursery records

Log each batch here to confirm or update the timing and difficulty above.

Date set	Stock source / provenance	No. set	No. struck	Strike %	Notes

© 2026 Mitchell Donald. All rights reserved.

Phyllanthus calycinus

False Boronia

Current WA Herbarium name: *Lysiandra calycina*

Family **Phyllanthaceae**

DIFFICULTY	WOOD TYPE	BEST MONTHS	CUTTING LENGTH	HORMONE (IBA)
Moderate *	Semi-hardwood	Feb – Apr	50–75 mm	3 g/L IBA

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Cuttings	OK	Best	Best	Best	OK	–	–	–	–	–	–	–

Best window | Workable – see 'Selecting material' in the method | Avoid / low success

Plant snapshot

Habit	Small erect shrub, 0.2–1.2 m, with smooth branchlets.
Flowering	Jun – Dec or Jan (white-cream to pink). Sepals enlarge to enclose the fruit.
Habitat	Usually sandy soils, including coastal alkaline sands.
Name note	FloraBase now lists this plant as *Lysiandra calycina*; many nurseries still sell it as *Phyllanthus calycinus*. Kept under *Phyllanthus* here to match your plant list.
Local note	Recorded in Joondalup and Wanneroo.

Propagation options

Method	Suitability	Notes	Details in
Cuttings	Trial	No published cutting data; trial protocol.	This guide
Seed	Possible	Fresh seed is recommended; little published detail.	–
Division	Not used	Woody shrub.	–

Propagation method

Item	Details
Best time to take cuttings	**Feb – Apr** – warm, and after the long flowering season ends. Jan and May are workable. Avoid Jun – Dec (flowering, and cool in winter).
Wood type	**Semi-hardwood**.
Cutting size	50–75 mm.
Selecting material	**Non-flowering** shoots from healthy plants.
Pre-treatment	• Strip leaves from the lower half. • Light wound on firmer stems. • Dip the base in rooting hormone.
Rooting hormone	3 g/L IBA (**Clonex Purple**); trial 1.5 g/L (**Clonex Green**) on softer tips.
Propagation mix	Free-draining sand-based mix.
Environment	Standard humid frame, bottom heat, bright shade.
Expected strike time	Not documented – start checking from about 6 weeks and record the result.
Post-treatment / aftercare	• Standard hardening off. • Pot on into a free-draining native mix.

© 2026 Mitchell Donald. All rights reserved.

Common problems

Problem	Likely cause and fix
No roots after 10+ weeks	Try a different hormone strength or timing on the next batch; record what you tried.
Rotting	Mix too wet. Water less often; improve drainage.
Cuttings flowering	Use non-flowering shoots only.

Notes

Difficulty	**Moderate (provisional).** No cutting data found for this species; the rating is a cautious starting point.
Other methods	Seed – the method most guides mention; use fresh seed. Few published details on pre-treatment.
Confidence	**Low.** No published cutting data found. Treat as a trial protocol and record results.
Sources	See References: 1, 4, 29, 30

Nursery records

Log each batch here to confirm or update the timing and difficulty above.

Date set	Stock source / provenance	No. set	No. struck	Strike %	Notes

© 2026 Mitchell Donald. All rights reserved.

Pimelea argentea

Silvery-leaved Pimelea

Family **Thymelaeaceae**

DIFFICULTY	WOOD TYPE	BEST MONTHS	CUTTING LENGTH	HORMONE (IBA)
Moderate *	Semi-hardwood	Mar – May	30–40 mm	3 g/L IBA

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Cuttings	–	OK	Best	Best	Best	OK	OK	OK	–	–	–	–

Best window | Workable – see 'Selecting material' in the method | Avoid / low success

Plant snapshot

Habit	Erect shrub, 0.4–2 m, with silvery leaves.
Flowering	Jul – Nov (cream-yellow to white-green). **Male and female flowers are on separate plants.**
Habitat	Sand or sandy clay, usually on coastal dunes, often with granite or near watercourses and lakes.
Local note	Recorded in Joondalup and Wanneroo.

Propagation options

Method	Suitability	Notes	Details in
Cuttings	Main method	Firm growth, autumn – winter; cut leaves off (provisional).	This guide
Seed	Difficult	Poor germination, about 21 days; smoke helps in the genus.	–
Division	Not used	Shrub – not divided.	–

Propagation method

Item	Details
Best time to take cuttings	**Mar – May**. Kings Park found WA pimeleas are best taken from autumn through to late winter, before new growth starts. Feb and Jun – Aug are workable (non-flowering shoots). Avoid Sep – Jan (flowering, then soft new growth).
Wood type	**Semi-hardwood**.
Cutting size	30–40 mm (Keighery and Dixon).
Selecting material	Take cuttings from **both male and female plants** and label them. A cutting keeps its parent's sex – the reason cuttings are preferred for this group – and only females set seed.
Pre-treatment	• Cut the lower leaves off with a sharp blade – **don't strip them**, or the bark tears. • Dip the base in rooting hormone.
Rooting hormone	Semi-hardwood strength: 3 g/L IBA (**Clonex Purple**).
Propagation mix	Free-draining sand-based cutting mix.
Environment	Humid frame, bright shade, bottom heat.
Expected strike time	Not documented for this species – strike varies widely in the genus (90% or more in *P. ferruginea*, low in *P. rosea*). Check from about 6 weeks and record the result.
Post-treatment / aftercare	• Standard hardening off. • Pot on into a free-draining native mix; keep the sex label with each plant.

© 2026 Mitchell Donald. All rights reserved.

Common problems

Problem	Likely cause and fix
Bark tears	Leaves pulled off. Cut them off with a sharp blade.
Cuttings rot	Too wet. Water less; improve drainage.
Low strike	Try earlier in autumn on the next batch, and record the month.

Notes

Difficulty	**Moderate (provisional).** Pimeleas strike from firm growth, but the bark is fragile and results vary between species.
Other methods	Seed – poor germination, about 21 days, and seedlings damp off (Keighery and Dixon). Wash the fruit off in running water and sow in autumn; smoke helps in the genus.
Confidence	**Medium.** Method from Kings Park's work on WA pimeleas (Keighery and Dixon); no strike data for this species.
Sources	See References: 4, 13, 87, 88

Nursery records

Log each batch here to confirm or update the timing and difficulty above.

Date set	Stock source / provenance	No. set	No. struck	Strike %	Notes

© 2026 Mitchell Donald. All rights reserved.

Pimelea calcicola

Coastal Banjine

Family **Thymelaeaceae**

DIFFICULTY	WOOD TYPE	BEST MONTHS	CUTTING LENGTH	HORMONE (IBA)
Moderate *	Semi-hardwood	Dec – Mar	50–75 mm	1.5–3 g/L IBA

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Cuttings	Best	Best	Best	OK	–	–	–	–	–	–	OK	Best

Best window | Workable – see 'Selecting material' in the method | Avoid / low success

Plant snapshot

Habit	Erect to spreading shrub, 0.2–1 m.
Flowering	Sep – Nov (pink heads).
Habitat	Sand on coastal limestone ridges.
Local note	Recorded in Joondalup and Wanneroo. **Priority Three** conservation species – take cuttings only from nursery stock or under the right DBCA licence.

Propagation options

Method	Suitability	Notes	Details in
Cuttings	Main method	Firm growth strikes; snip leaves so the bark doesn't tear (provisional).	This guide
Seed	Difficult	Pimeleas are hard from seed; smoke helps some.	–
Division	Not used	–	–

Propagation method

Item	Details
Best time to take cuttings	**Dec – Mar**, after flowering, once new growth has firmed. Nov and Apr are workable. Avoid May – Oct.
Wood type	**Semi-hardwood** – firm current season's growth.
Cutting size	50–75 mm.
Selecting material	Healthy, non-flowering side shoots.
Pre-treatment	• **Snip leaves off with fine scissors** – pulling them tears the thin bark. • No wounding. • Dip the base in rooting hormone.
Rooting hormone	1.5 g/L IBA (**Clonex Green**) for softer tips; 3 g/L for firmer stems.
Propagation mix	Very free-draining cutting mix.
Environment	Humid frame, bright shade, bottom heat. Moist, not wet.
Expected strike time	Not documented – check from about 4–6 weeks and record the result.
Post-treatment / aftercare	• Standard hardening off. • Grow on in full sun in a free-draining mix.

© 2026 Mitchell Donald. All rights reserved.

Common problems

Problem	Likely cause and fix
Bark torn at the base	Leaves were pulled off. Snip them with scissors.
Base rotting	Too wet. Coarser mix; water less.
Slow to root	Use bottom heat; take cuttings in the warm months.

Notes

Difficulty	**Moderate (provisional).** Pimeleas strike from cuttings, but the bark damages easily; untested for this species.
Other methods	Seed – pimeleas are difficult from seed; smoke water helps some species. The only Alcoa test of this species gave 0%.
Confidence	**Medium.** No species data. ANPSA: *Pimelea* cuttings are successful and firm growth usually strikes, but the bark tears easily.
Sources	See References: 4, 32, 66, 67

Nursery records

Log each batch here to confirm or update the timing and difficulty above.

Date set	Stock source / provenance	No. set	No. struck	Strike %	Notes

© 2026 Mitchell Donald. All rights reserved.

Pimelea sulphurea

Yellow Banjine

Family **Thymelaeaceae**

DIFFICULTY	WOOD TYPE	BEST MONTHS	CUTTING LENGTH	HORMONE (IBA)
Moderate *	Semi-hardwood	Dec – Mar	50–75 mm	1.5–3 g/L IBA

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Cuttings	Best	Best	Best	OK	–	–	–	–	–	–	OK	Best

Best window | Workable – see 'Selecting material' in the method | Avoid / low success

Plant snapshot

Habit	Erect, spindly or open shrub, 0.15–0.7 m.
Flowering	Jul – Nov (yellow heads).
Habitat	White, grey or yellow sand, sometimes with gravel or laterite; sandplains and ridges.
Local note	Recorded in Joondalup and Wanneroo.

Propagation options

Method	Suitability	Notes	Details in
Cuttings	Main method	Firm growth; snip leaves – bark tears (provisional).	This guide
Seed	Difficult	Winter sowing with smoke water, and patience.	–
Division	Not used	Shrub – not divided.	–

Propagation method

Item	Details
Best time to take cuttings	**Dec – Mar**, after flowering, once new growth has firmed. Nov and Apr are workable. Avoid May – Oct.
Wood type	**Semi-hardwood** – firm current season's growth.
Cutting size	50–75 mm. Stems are thin – take extra.
Selecting material	Healthy, non-flowering side shoots.
Pre-treatment	• **Snip leaves off with fine scissors** – pulling them tears the thin bark. • No wounding. • Dip the base in rooting hormone.
Rooting hormone	1.5 g/L IBA (**Clonex Green**) for softer tips; 3 g/L for firmer stems.
Propagation mix	Very free-draining cutting mix.
Environment	Humid frame, bright shade, bottom heat. Moist, not wet.
Expected strike time	Not documented – check from about 4–6 weeks and record the result.
Post-treatment / aftercare	• Standard hardening off. • Grow on in full sun in a free-draining mix.

© 2026 Mitchell Donald. All rights reserved.

Common problems

Problem	Likely cause and fix
Bark torn at the base	Leaves were pulled off. Snip them with scissors.
Base rotting	Too wet. Coarser mix; water less.
Slow to root	Use bottom heat; take cuttings in the warm months.

Notes

Difficulty	**Moderate (provisional).** Pimeleas strike from cuttings, but the bark damages easily; untested for this species.
Other methods	Seed – pimeleas are difficult from seed. Sowing in winter with smoke water, and patience, is the best bet. Not in the Alcoa nursery records.
Confidence	**Low.** No species data. Other WA pimeleas strike from firm growth, but the bark tears easily.
Sources	See References: 4, 13, 66, 67, 73

Nursery records

Log each batch here to confirm or update the timing and difficulty above.

Date set	Stock source / provenance	No. set	No. struck	Strike %	Notes

© 2026 Mitchell Donald. All rights reserved.

Pithocarpa cordata

Tangle Daisy

Previously *Ozothamnus cordatus / Helichrysum cordatum*

Family **Asteraceae**

DIFFICULTY	WOOD TYPE	BEST MONTHS	CUTTING LENGTH	HORMONE (IBA)
Easy–Moderate *	Semi-hardwood tips	Sep – Nov; May	50–75 mm	1.5–3 g/L IBA

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Cuttings	–	–	–	OK	Best	OK	–	OK	Best	Best	Best	OK

Best window | Workable – see 'Selecting material' in the method | Avoid / low success

Plant snapshot

Habit	Scrambling, tangled shrub, 1–2 m, with small heart-shaped leaves.
Flowering	Jan – Apr (white to yellow).
Habitat	Coastal limestone and dunes.
Local note	Recorded in the Wanneroo LGA. Older books list it as *Ozothamnus cordatus* or *Helichrysum cordatum*.

Propagation options

Method	Suitability	Notes	Details in
Cuttings	Good	Perennial daisy – expected to strike well (provisional).	This guide
Seed	Possible	Not documented; daisy seed often needs light.	–
Division	Not used	–	–

Propagation method

Item	Details
Best time to take cuttings	**Sep – Nov** on vigorous spring growth, and **May** after flowering. Apr, Jun, Aug and Dec are workable. Avoid Jan – Mar (flowering and heat) and Jul (cold).
Wood type	**Semi-hardwood tips** – vigorous growth that has started to firm.
Cutting size	50–75 mm.
Selecting material	Vigorous, healthy, non-flowering shoots.
Pre-treatment	• Strip leaves from the lower half. • No wounding. • Dip the base in rooting hormone.
Rooting hormone	1.5 g/L IBA (**Clonex Green**); 3 g/L for firmer stems.
Propagation mix	Free-draining: about 3 parts sand to 1 part peat or coir.
Environment	Humid frame, bright shade, bottom heat. Moist, not wet.
Expected strike time	Not documented – check from about 3–4 weeks and record the result.
Post-treatment / aftercare	• Standard hardening off. • Grow on in full sun; tip-prune to control the scrambling habit.

© 2026 Mitchell Donald. All rights reserved.

Common problems

Problem	Likely cause and fix
Cuttings rot	Too wet or humid. Mist less, improve airflow.
No roots after 8 weeks	Try firmer tips, a stronger hormone, or a different season, and record the result.
Tangled, sprawling tubes	Tip-prune and space tubes.

Notes

Difficulty	**Easy–Moderate (provisional).** Most perennial daisies strike quickly from vigorous growth, but this species is untested. Start with a trial batch.
Other methods	Seed – not documented. Daisy-family seed often needs light, so surface-sow if trialling.
Confidence	**Low.** No published propagation data. Based on perennial Australian daisies, which usually strike quickly – trial protocol.
Sources	See References: 4, 35

Nursery records

Log each batch here to confirm or update the timing and difficulty above.

Date set	Stock source / provenance	No. set	No. struck	Strike %	Notes

© 2026 Mitchell Donald. All rights reserved.

Rhagodia baccata

Berry Saltbush

Also listed as *Chenopodium baccatum*

Family **Amaranthaceae (chenopods)**

DIFFICULTY	WOOD TYPE	BEST MONTHS	CUTTING LENGTH	HORMONE (IBA)
Moderate *	Semi-hardwood tips	Sep – Nov	75–100 mm	3 g/L IBA

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Cuttings	OK	OK	OK	–	–	–	–	OK	Best	Best	Best	OK

Best window | Workable – see 'Selecting material' in the method | Avoid / low success

Plant snapshot

Habit	Spreading shrub, 0.3–2 m, with grey-green leaves.
Flowering	Variable – Feb, Apr – May or Oct – Dec (tiny), followed by red berries eaten by birds and bobtails.
Habitat	Sand dunes and coastal rock.
Local note	Recorded in Joondalup and Wanneroo. Salt-tolerant; a breeding plant for the saltbush blue butterfly.

Propagation options

Method	Suitability	Notes	Details in
Cuttings	Good	Chenopod shrubs generally strike well (provisional).	This guide
Seed	Main method	Ferment berries, then smoke water.	→ Seed Sowing Guide
Division	Not used	Woody shrub.	–

Propagation method

Item	Details
Best time to take cuttings	**Sep – Nov**, in active spring growth. Aug and Dec – Mar are workable with non-flowering shoots. Avoid Apr – Jul.
Wood type	**Semi-hardwood tips** of current season's growth.
Cutting size	75–100 mm.
Selecting material	Vigorous tips without flowers or berries.
Pre-treatment	• Strip leaves from the lower half. • Light wound on firmer stems. • Dip the base in rooting hormone.
Rooting hormone	Semi-hardwood strength, 3 g/L IBA (**Clonex Purple**).
Propagation mix	Free-draining mix of sand and potting mix.
Environment	Bright shade with **good ventilation and little or no mist** (misting encourages leaf disease on saltbush cuttings). Bottom heat helps.
Expected strike time	Not documented – check from about 4–6 weeks and record the result.
Post-treatment / aftercare	• Harden off, then grow on in full sun. • Water sparingly once established.

© 2026 Mitchell Donald. All rights reserved.

Common problems

Problem	Likely cause and fix
Leaf spots, leaves dropping	Too humid. Stop misting and improve airflow.
Base rotting	Mix too wet. Use a sandier mix, water less.
No roots	Try firmer material or the seed method, and record the result.

Notes

Difficulty	**Moderate (provisional).** Untested here; related saltbushes strike from tip cuttings. Seed is the documented method.
Confidence	**Low.** No published cutting data found. Method follows related saltbushes – trial protocol.
Sources	See References: 4, 20, 36, 37

Nursery records

Log each batch here to confirm or update the timing and difficulty above.

Date set	Stock source / provenance	No. set	No. struck	Strike %	Notes

© 2026 Mitchell Donald. All rights reserved.

Scaevola canescens

Grey Scaevola

Family **Goodeniaceae**

DIFFICULTY	WOOD TYPE	BEST MONTHS	CUTTING LENGTH	HORMONE (IBA)
Easy–Moderate *	Softwood – semi-hardwood	Nov – Jan	50–80 mm	1.5–3 g/L IBA

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Cuttings	Best	OK	OK	OK	OK	–	–	–	–	OK	Best	Best

Best window | Workable – see 'Selecting material' in the method | Avoid / low success

Plant snapshot

Habit	Prostrate to ascending shrub, 0.1–0.5 m. Densely hairy grey leaves; stems are brittle.
Flowering	Mar – Oct (white to white-purple), in spikes in the leaf axils.
Habitat	Sand, often with lateritic gravel or limestone.
Local note	Recorded in the northern Perth area (Wanneroo and Stirling LGAs).

Propagation options

Method	Suitability	Notes	Details in
Cuttings	Main method	Soft to semi-hardwood tips; keep hairy foliage dry.	This guide
Seed	Difficult	Scaevola seed is generally difficult.	–
Division	Possible	Only if stems have rooted or suckered.	–

Propagation method

Item	Details
Best time to take cuttings	**Nov – Jan**, on new growth after flowering ends, while it is warm. Oct and Feb – May are workable if you choose shoots without flower spikes. Avoid Jun – Sep (cool, peak flowering).
Wood type	**Softwood to semi-hardwood** tip cuttings.
Cutting size	50–80 mm tips.
Selecting material	Vigorous **non-flowering** tips. Flowers form in the leaf axils along the stem, so check the whole cutting and pinch off any spikes.
Pre-treatment	• Handle gently – stems snap easily. Carry in a rigid tray or box. • Strip lower leaves by cutting, not tearing. • Dip the base in rooting hormone.
Rooting hormone	Softwood strength (1.5 g/L IBA, e.g. **Clonex Green**) for soft tips; semi-hardwood strength (3 g/L, **Clonex Purple**) for firmer material.
Propagation mix	Extra free-draining: standard sand/coir mix with extra coarse sand or perlite.
Environment	**Keep foliage as dry as possible.** Dense hairy leaves rot under heavy mist. Water the mix rather than the leaves, give good airflow, and use bottom heat (root zone about 21–24°C).
Expected strike time	About **4–8 weeks** in warm weather.
Post-treatment / aftercare	• Standard hardening off. • Pot on into a free-draining mix; keep water off the foliage. • Remove any leaves showing grey mould promptly.

© 2026 Mitchell Donald. All rights reserved.

Common problems

Problem	Likely cause and fix
Grey fuzz or blackened leaves	Too humid/wet. Cut back misting, remove affected material, improve airflow.
Snapped stems	Brittle stems – transport in rigid containers and use a dibber when setting.
Cuttings flowering, not rooting	Take shoots without axillary flower spikes.

Notes

Difficulty	**Easy–Moderate (provisional).** *Scaevola* cuttings generally strike very readily; the hairy foliage makes rot the main risk.
Other methods	Seed – *Scaevola* seed is generally difficult. Division – possible if a plant has suckered or rooted along its stems.
Confidence	**Medium.** Little species-specific data published. Based on the genus (strikes readily) plus this species' hairy foliage.
Sources	See References: 1, 4, 5, 6, 7, 12, 13, 14, 16

Nursery records

Log each batch here to confirm or update the timing and difficulty above.

Date set	Stock source / provenance	No. set	No. struck	Strike %	Notes

© 2026 Mitchell Donald. All rights reserved.

Scaevola crassifolia

Thick-leaved Fan-flower

Family **Goodeniaceae**

DIFFICULTY	WOOD TYPE	BEST MONTHS	CUTTING LENGTH	HORMONE (IBA)
Easy	Semi-hardwood	Mar – May	75–100 mm	3 g/L IBA

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Cuttings	OK	OK	Best	Best	Best	–	–	–	–	OK	OK	OK

Best window | Workable – see 'Selecting material' in the method | Avoid / low success

Plant snapshot

Habit	Erect or spreading shrub, 0.1–1.5 m and up to 3 m wide. Young growth is sticky; thick, toothed, egg-shaped to round leaves.
Flowering	Jul – Dec, also Jan – Feb (blue, mauve or white fans).
Habitat	Sand on coastal dunes, plains and limestone cliffs.
Local note	Recorded in Joondalup and Wanneroo. Flower colour and habit vary – cuttings keep the parent's form.

Propagation options

Method	Suitability	Notes	Details in
Cuttings	Main method	Standard method; keeps flower colour and form.	This guide
Seed	Possible	Large seeds sometimes germinate naturally.	–
Division	Not used	–	–

Propagation method

Item	Details
Best time to take cuttings	**Mar – May** – warm, and outside the main flowering period. Oct – Feb is workable with non-flowering shoots (shade well in summer). Avoid Jun – Sep.
Wood type	**Semi-hardwood**.
Cutting size	75–100 mm.
Selecting material	**Non-flowering** tips. Flower spikes carry leafy bracts – check closely and pinch them off.
Pre-treatment	• New growth is sticky – wipe the blade often. • Strip lower leaves; cut the large remaining leaves back by a third to a half. • Dip the base in rooting hormone.
Rooting hormone	3 g/L IBA (**Clonex Purple**).
Propagation mix	Free-draining sand/coir mix.
Environment	Standard humid frame with bottom heat, bright shade.
Expected strike time	About **4–8 weeks** in warm weather.
Post-treatment / aftercare	• Standard hardening off. • Pot on into a free-draining mix. • Tip-prune young plants for dense growth.

© 2026 Mitchell Donald. All rights reserved.

Common problems

Problem	Likely cause and fix
Wilting in the first week	Too much leaf area. Trim leaves harder; keep the frame closed.
Rotting	Mix too wet. Water less often; improve drainage.
Cuttings flowering	Use non-flowering tips or wait for the Mar – May window.

Notes

Difficulty	**Easy.** Semi-hardwood cuttings in the warmer months are the standard method.
Other methods	Seed – large seeds are produced and sometimes germinate naturally; cuttings are preferred to keep flower colour and habit.
Confidence	**High.** Cuttings are the standard method in WA nurseries (e.g. Geraldton community nursery).
Sources	See References: 1, 4, 5, 12, 13, 26, 30

Nursery records

Log each batch here to confirm or update the timing and difficulty above.

Date set	Stock source / provenance	No. set	No. struck	Strike %	Notes

© 2026 Mitchell Donald. All rights reserved.

Scaevola globulifera

No widely used common name

Family **Goodeniaceae**

DIFFICULTY	WOOD TYPE	BEST MONTHS	CUTTING LENGTH	HORMONE (IBA)
Easy *	Softwood – semi-hardwood	Jan – Apr	50–80 mm	1.5–3 g/L IBA

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Cuttings	Best	Best	Best	Best	OK	–	–	–	–	–	OK	OK

Best window | Workable – see 'Selecting material' in the method | Avoid / low success

Plant snapshot

Habit	Spreading, low-lying to ascending shrub, 0.2–1.2 m. Hairless.
Flowering	Aug – Dec (blue to white).
Habitat	Sand, sandhills and limestone ridges.
Local note	Recorded in Joondalup and Wanneroo.

Propagation options

Method	Suitability	Notes	Details in
Cuttings	Main method	Soft to semi-hardwood tips (provisional rating).	This guide
Seed	Difficult	Scaevola seed is generally difficult.	–
Division	Possible	Check stock plants for rooted stems.	–

Propagation method

Item	Details
Best time to take cuttings	**Jan – Apr** – warm, and after flowering finishes in Dec. Nov – Dec and May are workable. Avoid Jun – Oct.
Wood type	**Softwood to semi-hardwood** tip cuttings.
Cutting size	50–80 mm.
Selecting material	**Non-flowering** tips; pinch off any buds.
Pre-treatment	• Strip lower leaves. • Trim larger leaves by about one-third. • Dip the base in rooting hormone.
Rooting hormone	1.5 g/L IBA (**Clonex Green**) for soft tips; 3 g/L (**Clonex Purple**) for firmer material.
Propagation mix	Free-draining sand/coir mix.
Environment	Humid frame, bottom heat, bright shade. Hairless leaves tolerate normal misting.
Expected strike time	About **4–8 weeks** in warm weather.
Post-treatment / aftercare	• Standard hardening off. • Pot on into a free-draining native mix. • Pinch tips to encourage branching.

© 2026 Mitchell Donald. All rights reserved.

Common problems

Problem	Likely cause and fix
Wilting	Low humidity or too much leaf. Trim leaves; keep the frame closed.
Rotting	Mix too wet. Water less often.
Cuttings flowering	Use non-flowering tips, or wait for the Jan – Apr window.

Notes

Difficulty	**Easy (provisional).** *Scaevola* cuttings generally strike very readily.
Other methods	Seed – *Scaevola* seed is generally difficult. Division – check stock plants for rooted stems.
Confidence	**Medium.** No species-specific data found. Based on the genus, which strikes readily from cuttings.
Sources	See References: 1, 4, 5, 6, 7, 13, 14

Nursery records

Log each batch here to confirm or update the timing and difficulty above.

Date set	Stock source / provenance	No. set	No. struck	Strike %	Notes

© 2026 Mitchell Donald. All rights reserved.

Scaevola nitida

Shining Fanflower

Family
Goodeniaceae

DIFFICULTY	WOOD TYPE	BEST MONTHS	CUTTING LENGTH	HORMONE (IBA)
Easy *	Semi-hardwood	Jan – Apr	75–100 mm	3 g/L IBA

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Cuttings	Best	Best	Best	Best	OK	–	–	–	–	–	OK	OK

Best window | Workable – see 'Selecting material' in the method | Avoid / low success

Plant snapshot

Habit	Erect shrub, 0.3–3 m.
Flowering	Aug – Dec (blue-purple fans).
Habitat	White or grey sand and clay; coastal limestone cliffs and dunes.
Local note	Recorded in Joondalup and Wanneroo. Tolerates full coastal exposure, sand blast and salt spray.

Propagation options

Method	Suitability	Notes	Details in
Cuttings	Main method	Scaevola cuttings strike very readily.	This guide
Seed	Possible	Smoke; about 21–42 days (seed suppliers).	–
Division	Not used	Shrub – not divided.	–

Propagation method

Item	Details
Best time to take cuttings	**Jan – Apr**, after the main flowering, while it's warm. Nov – Dec and May are workable with non-flowering shoots. Avoid Jun – Oct.
Wood type	**Semi-hardwood** – firm current season's growth.
Cutting size	75–100 mm.
Selecting material	**Non-flowering** tips. Pinch off any buds.
Pre-treatment	• Strip lower leaves; cut large remaining leaves back by a third to a half. • Dip the base in rooting hormone.
Rooting hormone	3 g/L IBA (**Clonex Purple**).
Propagation mix	Free-draining sand/coir mix.
Environment	Standard humid frame with bottom heat, bright shade.
Expected strike time	About **4–8 weeks** in warm weather (typical for the genus). Record your results.
Post-treatment / aftercare	• Standard hardening off. • Pot on into a free-draining mix. • Tip-prune young plants for dense growth.

© 2026 Mitchell Donald. All rights reserved.

Common problems

Problem	Likely cause and fix
Wilting in the first week	Too much leaf area. Trim leaves harder; keep the frame closed.
Rotting	Mix too wet. Water less often; improve drainage.
Cuttings flowering	Use non-flowering tips, or wait for the Jan – Apr window.

Notes

Difficulty	**Easy (provisional).** Scaevolas strike very readily; untested for this species.
Other methods	Seed – smoke treatment is recommended by seed suppliers; germination in about 21–42 days. Not in the Alcoa records.
Confidence	**Medium.** No species-specific cutting data. Based on the genus – ANPSA says cuttings of *Scaevola* species strike very readily.
Sources	See References: 1, 4, 5, 13, 51, 71, 72, 78

Nursery records

Log each batch here to confirm or update the timing and difficulty above.

Date set	Stock source / provenance	No. set	No. struck	Strike %	Notes

© 2026 Mitchell Donald. All rights reserved.

Scaevola repens

No widely used common name ('repens' = creeping)

Family **Goodeniaceae**

DIFFICULTY	WOOD TYPE	BEST MONTHS	CUTTING LENGTH	HORMONE (IBA)
Easy *	Softwood – semi-hardwood	Mar – May	50–80 mm	1.5–3 g/L IBA

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Cuttings	OK	OK	Best	Best	Best	–	–	–	OK	OK	OK	OK

Best window Workable – see 'Selecting material' in the method Avoid / low success

Plant snapshot

Habit	Prostrate to ascending or erect shrub, 0.05–1 m; trailing branches to about 50 cm.
Flowering	Jun – Dec, sometimes Jan – Feb (white or cream, occasionally blue-purple; golden hairs on the outside of the flower).
Habitat	Sand dunes and plains on sand, gravel, laterite or limestone.
Local note	Recorded in the City of Joondalup. A variable species with informal subspecies – use local provenance and keep it labelled.

Propagation options

Method	Suitability	Notes	Details in
Cuttings	Main method	Soft to semi-hardwood tips (provisional rating).	This guide
Seed	Difficult	Scaevola seed is generally difficult.	–
Division	Possible	Some scaevolas sucker; check for rooted pieces.	–

Propagation method

Item	Details
Best time to take cuttings	**Mar – May** – warm autumn weather, mostly outside the main flowering period. Sep – Feb is workable using non-flowering shoots (shade well in summer). Avoid Jun – Aug.
Wood type	**Softwood to semi-hardwood** tip cuttings.
Cutting size	50–80 mm tips from the trailing stems.
Selecting material	**Non-flowering** tips. Flower spikes sit in the leaf axils – pinch off any you find.
Pre-treatment	• Strip lower leaves. • Leaves can be long (up to 90 mm) – trim remaining leaves by about one-third to cut water loss. • Dip the base in rooting hormone.
Rooting hormone	Softwood strength (1.5 g/L IBA, **Clonex Green**) for soft tips; 3 g/L (**Clonex Purple**) for firmer material.
Propagation mix	Free-draining sand/coir mix.
Environment	Humid frame with bottom heat, bright shade. Keep the mix moist but on the drier side rather than wet.
Expected strike time	About **4–8 weeks** in warm weather.
Post-treatment / aftercare	• Standard hardening off. • Pot on into a free-draining native mix. • Pinch tips once growing to encourage branching.

© 2026 Mitchell Donald. All rights reserved.

Common problems

Problem	Likely cause and fix
Wilting in the first week	Too much leaf area or low humidity. Trim leaves harder; keep the frame closed.
Rotting	Mix too wet. Water less often; improve drainage.
Cuttings flowering, not rooting	Take non-flowering tips, or wait for the Mar – May window.

Notes

Difficulty	**Easy (provisional).** *Scaevola* cuttings generally strike very readily.
Other methods	Seed – *Scaevola* seed is generally difficult. Division – some scaevolas sucker; check stock plants for rooted pieces.
Confidence	**Medium.** Little species-specific data published. Based on the genus, which strikes readily from cuttings.
Sources	See References: 1, 4, 5, 6, 7, 12, 13, 14, 16

Nursery records

Log each batch here to confirm or update the timing and difficulty above.

Date set	Stock source / provenance	No. set	No. struck	Strike %	Notes

© 2026 Mitchell Donald. All rights reserved.

Scaevola thesioides

No widely used common name

Family **Goodeniaceae**

DIFFICULTY	WOOD TYPE	BEST MONTHS	CUTTING LENGTH	HORMONE (IBA)
Easy–Moderate *	Softwood – semi-hardwood	Apr – May	50–75 mm	1.5–3 g/L IBA

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Cuttings	OK	OK	OK	Best	Best	–	–	–	OK	OK	OK	OK

Best window | Workable – see 'Selecting material' in the method | Avoid / low success

Plant snapshot

Habit	Spreading subshrub or perennial herb, 0.15–1 m. Mostly hairless; thread-like to narrow leaves; ribbed stems.
Flowering	Aug – Dec and Jan – Mar (pale blue to white), in spikes at the ends of branches.
Habitat	Coastal sand and limestone.
Local note	The coastal Perth plant is subsp. *thesioides*; recorded in the Wanneroo LGA.

Propagation options

Method	Suitability	Notes	Details in
Cuttings	Main method	Non-flowering side shoots (provisional rating).	This guide
Seed	Difficult	Scaevola seed is generally difficult.	–
Division	Possible	Only if stock plants have suckered.	–

Propagation method

Item	Details
Best time to take cuttings	**Apr – May** – warm autumn weather between flowering periods. Sep – Mar is workable using non-flowering side shoots. Avoid Jun – Aug.
Wood type	**Softwood to semi-hardwood** – firm but still green tip or side-shoot cuttings.
Cutting size	50–75 mm.
Selecting material	**Non-flowering side shoots.** Flowers form at the branch tips, so main tips often carry buds.
Pre-treatment	• Strip leaves from the lower half; leaves are narrow so little trimming is needed. • Stems are slender – always use a dibber hole. • Dip the base in rooting hormone.
Rooting hormone	Softwood strength (1.5 g/L IBA, **Clonex Green**); 3 g/L (**Clonex Purple**) for firmer material.
Propagation mix	Free-draining sand/coir mix.
Environment	Hairless leaves tolerate misting better than *S. canescens*, but fine stems collapse quickly if the mix is waterlogged or dries out. Keep evenly moist, with bottom heat and bright shade.
Expected strike time	About **4–8 weeks** in warm weather.
Post-treatment / aftercare	• Standard hardening off. • Pot on gently – fine roots break easily. Single cuttings per tube make this easier. • Pinch tips once growing to encourage branching.

© 2026 Mitchell Donald. All rights reserved.

Common problems

Problem	Likely cause and fix
Fine stems wilt or collapse	Mix drying out or waterlogged. Keep evenly moist; check daily.
Cuttings flower instead of rooting	Pinch off terminal flower buds; use side shoots.
Root damage when potting on	Strike one cutting per tube instead of community pots.

Notes

Difficulty	**Easy–Moderate (provisional).** The genus strikes readily; the fine stems are less forgiving of watering errors.
Other methods	Seed – *Scaevola* seed is generally difficult. Division – possible if stock plants have suckered.
Confidence	**Medium.** Little species-specific data published. Based on the genus plus this species' fine, slender growth.
Sources	See References: 1, 4, 5, 6, 7, 12, 13, 14, 16

Nursery records

Log each batch here to confirm or update the timing and difficulty above.

Date set	Stock source / provenance	No. set	No. struck	Strike %	Notes

© 2026 Mitchell Donald. All rights reserved.

Spinifex hirsutus

Hairy Spinifex

Family **Poaceae**

DIFFICULTY	WOOD TYPE	BEST MONTHS	CUTTING LENGTH	HORMONE (IBA)
Moderate *	Runner (stolon) sections	Mar – May	10–15 cm, 1–2 nodes	Optional (1.5 g/L)

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Cuttings	OK	OK	Best	Best	Best	–	–	–	OK	OK	OK	OK

Best window Workable – see 'Selecting material' in the method Avoid / low success

Plant snapshot

Habit	Rhizomatous, tussock-forming grass, 0.3–0.9 m, sending long runners across the sand. Broad, flat, silvery leaves, hairy on both sides.
Flowering	Oct – Jan. Male and female flowers are on separate plants; female heads are large spiky balls that roll in the wind.
Habitat	White sand on coastal dunes; an important stabiliser at the toe of unstable dunes.
Local note	Recorded in the City of Joondalup. Less common than *S. longifolius* on the Cottesloe foreshore.

Propagation options

Method	Suitability	Notes	Details in
Cuttings	Good	Runner (stolon) sections with 1–2 nodes.	This guide
Seed	Main method	The usual nursery method; cover seed (needs dark).	→ Seed Sowing Guide
Division	Possible	Tussock divisions work but survived less well than rooted runners.	→ Division Guide

Propagation method

Item	Details
Best time to take cuttings	**Mar – May**, when runners are young, green and actively growing, giving plants ready for winter planting. Sep – Feb is workable. Avoid Jun – Aug (slow growth).
Wood type	**Runner (stolon) sections** – not woody cuttings. Use young, green runners, not old brown ones.
Cutting size	**Nursery tubes:** 10–15 cm sections with at least one (ideally two) nodes. **Direct planting on site:** 40–50 cm runners, buried about halfway in damp sand.
Selecting material	Actively growing runners from the dune edge. Plants are male or female, so collect from several plants if you want seed-producing plantings later.
Pre-treatment	• Keep runners moist and shaded after collecting – they dry out fast. • Cut just below a node; trim leaves back by about half. • Hormone optional – roots form naturally at the nodes.
Rooting hormone	Optional. If used, softwood strength (1.5 g/L IBA, **Clonex Green**) on the node.
Propagation mix	Coarse washed sand or a very sandy mix. Deep tubes suit the root system.
Environment	Bottom heat and misting gave good rooting at the nodes in *S. sericeus* trials. The hairy leaves of this species can hold water, so mist lightly.
Expected strike time	Nodes root readily, but plants are **slow to reach planting size** – in one trial about 80% of cuttings were ready to plant within 18 months.

© 2026 Mitchell Donald. All rights reserved.

Item	Details
Post-treatment / aftercare	• Harden off, then grow on in full sun. • Pot on into deep tubes or larger pots. • Plant out in late autumn or winter when rain is reliable.

Common problems

Problem	Likely cause and fix
Runners dry out before setting	Carry in damp bags in an esky; set the same day.
Nodes rot	Mix too wet or too much mist. Use coarser sand; mist less.
Slow growth after rooting	Normal for spinifex – allow time; feed lightly and give full sun.

Notes

Difficulty	**Moderate (provisional).** Runner sections root well under bottom heat and mist, but it's labour-intensive and slow. Seed is the more common nursery method.
Other methods	Seed – the usual nursery method. The Geraldton community nursery advises collecting from female plants growing near males. *S. sericeus* research found seed germinates far better in darkness (56%) than light (14%), so cover seed. Division of tussocks also works but survived less well than rooted cuttings in trials (31% vs 40%).
Confidence	**Medium.** Method based on published trials with the closely related *S. sericeus* and general spinifex practice.
Sources	See References: 4, 12, 18, 19, 20

Nursery records

Log each batch here to confirm or update the timing and difficulty above.

Date set	Stock source / provenance	No. set	No. struck	Strike %	Notes

© 2026 Mitchell Donald. All rights reserved.

Spinifex longifolius

Beach Spinifex

Family **Poaceae**

DIFFICULTY	WOOD TYPE	BEST MONTHS	CUTTING LENGTH	HORMONE (IBA)
Moderate *	Runner (stolon) sections	Mar – May	10–15 cm, 1–2 nodes	Optional (1.5 g/L)

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Cuttings	OK	OK	Best	Best	Best	–	–	–	OK	OK	OK	OK

Best window · Workable – see 'Selecting material' in the method · Avoid / low success

Plant snapshot

Habit	Rhizomatous, tussock-forming grass, 0.3–1 m high and up to 2 m wide, with long runners through the sand.
Flowering	Apr – Dec or Jan (green-brown). Male and female plants are separate; female heads up to 250 mm across roll in the wind.
Habitat	White sand on coastal dunes; a key foredune stabiliser.
Local note	Recorded in the Wanneroo LGA; the more common of the two spinifex species on the Cottesloe foreshore.

Propagation options

Method	Suitability	Notes	Details in
Cuttings	Good	Runner (stolon) sections with 1–2 nodes.	This guide
Seed	Main method	The usual nursery method; smoke water, about 10 days.	→ Seed Sowing Guide
Division	Possible	Tussock divisions work but survived less well than rooted runners.	→ Division Guide

Propagation method

Item	Details
Best time to take cuttings	**Mar – May**, when runners are young, green and actively growing, giving plants ready for winter planting. Sep – Feb is workable. Avoid Jun – Aug (slow growth).
Wood type	**Runner (stolon) sections** – not woody cuttings. Use young, green runners, not old brown ones.
Cutting size	**Nursery tubes:** 10–15 cm sections with at least one (ideally two) nodes. **Direct planting on site:** 40–50 cm runners, buried about halfway in damp sand.
Selecting material	Actively growing runners from the dune edge. Plants are male or female, so collect from several plants if you want seed-producing plantings later.
Pre-treatment	• Keep runners moist and shaded after collecting – they dry out fast. • Cut just below a node; trim leaves back by about half. • Hormone optional – roots form naturally at the nodes.
Rooting hormone	Optional. If used, softwood strength (1.5 g/L IBA, **Clonex Green**) on the node.
Propagation mix	Coarse washed sand or a very sandy mix. Deep tubes suit the root system.
Environment	Bottom heat and misting gave good rooting at the nodes in *S. sericeus* trials. Keep the sand moist, not wet.
Expected strike time	Nodes root readily, but plants are **slow to reach planting size** – in one trial about 80% of cuttings were ready to plant within 18 months.

© 2026 Mitchell Donald. All rights reserved.

Item	Details
Post-treatment / aftercare	• Harden off, then grow on in full sun. • Pot on into deep tubes or larger pots. • Plant out in late autumn or winter when rain is reliable.

Common problems

Problem	Likely cause and fix
Runners dry out before setting	Carry in damp bags in an esky; set the same day.
Nodes rot	Mix too wet or too much mist. Use coarser sand; mist less.
Slow growth after rooting	Normal for spinifex – allow time; feed lightly and give full sun.

Notes

Difficulty	**Moderate (provisional).** Runner sections root well under bottom heat and mist, but it's labour-intensive and slow. Seed is the more common nursery method.
Other methods	Seed – the usual nursery method. The Geraldton community nursery collects from female plants growing close to males, separates the seed, cuts off the awns and sows en masse with smoke water (about 10 days to germinate). *S. sericeus* research found seed germinates far better in darkness, so cover seed. Division of tussocks also works.
Confidence	**Medium.** Method based on published trials with the closely related *S. sericeus* and general spinifex practice.
Sources	See References: 4, 12, 18, 19, 20

Nursery records

Log each batch here to confirm or update the timing and difficulty above.

Date set	Stock source / provenance	No. set	No. struck	Strike %	Notes

© 2026 Mitchell Donald. All rights reserved.

Styphelia insularis

No widely used common name (a beard-heath)

Formerly *Leucopogon insularis*

Family **Ericaceae**

DIFFICULTY	WOOD TYPE	BEST MONTHS	CUTTING LENGTH	HORMONE (IBA)
Difficult *	**Semi-hardwood**	**Oct – Dec**	**50–80 mm**	**3 g/L IBA**

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Cuttings	OK	OK	–	–	–	–	–	–	OK	Best	Best	Best

Best window | Workable – see 'Selecting material' in the method | Avoid / low success

Plant snapshot

Habit	Erect, spreading shrub, usually 0.25–0.9 m.
Flowering	Mar – Jul (white).
Habitat	Calcareous white or grey sand; coastal dunes, limestone and sandplains.
Name note	Moved from *Leucopogon* to *Styphelia* in 2020 (FloraBase) – older lists and labels use *Leucopogon insularis*.
Local note	Recorded in Joondalup and Wanneroo.

Propagation options

Method	Suitability	Notes	Details in
Cuttings	Difficult	No data – trial protocol based on related beard-heaths.	This guide
Seed	Difficult	Not documented; woody epacrid fruits need soil ageing + smoke.	–
Division	Not used	Shrub – not divided.	–

Propagation method

Item	Details
Best time to take cuttings	**Oct – Dec** (estimate) – new growth made after the autumn–winter flowering has firmed, and it's warm. Sep, Jan and Feb are workable. Avoid Mar – Aug (flowering, and cool).
Wood type	**Semi-hardwood** – firm current season's growth.
Cutting size	50–80 mm – a small shrub, so shorter than for the Coast Beard-heath.
Selecting material	Non-flowering tips from healthy plants. Take plenty.
Pre-treatment	• Strip leaves from the lower half. • Light wound on firmer stems. • Dip the base in rooting hormone.
Rooting hormone	Medium strength, as for other epacrids: 3 g/L IBA (**Clonex Purple**).
Propagation mix	Very free-draining sand-based cutting mix.
Environment	Humid frame in bright shade, not heavy mist. Trial bottom heat on part of the batch.
Expected strike time	Not documented – epacrids are slow (6–12 months in trials). Check from about 8 weeks and record the result.
Post-treatment / aftercare	• Expect fine, fragile roots – pot up with the mix ball intact. • Harden off slowly.

© 2026 Mitchell Donald. All rights reserved.

Common problems

Problem	Likely cause and fix
No roots after months	Common in epacrids. Keep them going; try another month or hormone strength next batch.
Cuttings blacken or rot	Too wet or humid. Less mist, better drainage and airflow.
Roots break when potting up	Move the whole plug without disturbing the roots.

Notes

Difficulty	**Difficult (provisional).** No data for this species; related beard-heaths strike poorly, so the rating is cautious.
Other methods	Seed – not documented for this species. Woody epacrid fruits don't respond to smoke alone; Kings Park found that ageing seed in soil for 1–2 years with repeated smoke worked for smoke-resistant Ericaceae.
Confidence	**Low.** No propagation data found for this species. Based on the related Coast Beard-heath and general epacrid experience – a trial protocol.
Sources	See References: 1, 4, 13, 80, 82, 84, 85

Nursery records

Log each batch here to confirm or update the timing and difficulty above.

Date set	Stock source / provenance	No. set	No. struck	Strike %	Notes

© 2026 Mitchell Donald. All rights reserved.

Synaphea spinulosa

No widely used common name (a synaphea)

Family **Proteaceae**

DIFFICULTY	WOOD TYPE	BEST MONTHS	CUTTING LENGTH	HORMONE (IBA)
Difficult	**Semi-hardwood**	**Feb – Apr**	**50–75 mm**	**3–8 g/L IBA (trial)**

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Cuttings	OK	Best	Best	Best	OK	–	–	–	–	–	–	OK

Best window · Workable – see 'Selecting material' in the method · Avoid / low success

Plant snapshot

Habit	Shrub, 0.3–1.2 m (subsp. *spinulosa*), with divided leaves whose lobes end in sharp points.
Flowering	Jul – Nov (yellow).
Habitat	Sandy soils, sometimes gravelly, over laterite.
Local note	Subsp. *spinulosa* is recorded in Joondalup and Wanneroo; the other two subspecies grow inland or further north – check your source. Wear gloves.

Propagation options

Method	Suitability	Notes	Details in
Cuttings	Difficult	Kings Park has struck it, at an extremely low rate.	This guide
Seed	Difficult	Low seed set; untested – one seed of a relative came up after nick + GA.	–
Division	Not used	Shrub – not divided.	–

Propagation method

Item	Details
Best time to take cuttings	**Feb – Apr** (estimate), once summer growth has firmed, well after the Jul – Nov flowering. Jan, May and Dec are workable. Avoid Jun – Nov.
Wood type	**Semi-hardwood**.
Cutting size	50–75 mm.
Selecting material	Vigorous, non-flowering shoots. Take plenty – expect very few to strike.
Pre-treatment	• Strip leaves from the lower half (gloves – the lobes are sharp). • Wound the base. • Dip the base in rooting hormone.
Rooting hormone	No published rate. Trial two strengths on split batches: 3 g/L (**Clonex Purple**) and 8 g/L (**Clonex Red**).
Propagation mix	Very free-draining sand-based cutting mix.
Environment	Humid frame, bright shade, bottom heat; keep the mix just moist.
Expected strike time	Not documented – check from about 8 weeks. Record every batch; there's almost no published data.
Post-treatment / aftercare	• Harden off slowly. • Use a low-phosphorus native mix and fertiliser, as for other Proteaceae.

© 2026 Mitchell Donald. All rights reserved.

Common problems

Problem	Likely cause and fix
No roots	Expected for most cuttings. Record the month and hormone, and compare batches.
Cuttings rot	Too wet. Water less; improve drainage.
Hard to handle	Prickly leaf lobes. Wear gloves.

Notes

Difficulty	**Difficult.** Kings Park's nursery has struck it, but at an extremely low rate.
Other methods	Seed – seed set is low in *Synaphea*, and two related species germinated under 2% (0.7% and 1.7%) in nursery tests. A single seed of a related threatened species germinated about 10 weeks after nicking and gibberellic acid (the seedling later died). Kings Park has used tissue culture for another species.
Confidence	**Low.** Kings Park's nursery has grown this species from cuttings, but with an extremely low strike rate; no method details published.
Sources	See References: 4, 13, 32, 59, 91, 92

Nursery records

Log each batch here to confirm or update the timing and difficulty above.

Date set	Stock source / provenance	No. set	No. struck	Strike %	Notes

© 2026 Mitchell Donald. All rights reserved.

Templetonia retusa

Cockies Tongues

Family **Fabaceae**

DIFFICULTY	WOOD TYPE	BEST MONTHS	CUTTING LENGTH	HORMONE (IBA)
Easy–Moderate *	Semi-hardwood	Dec – Mar	50–75 mm	3 g/L IBA

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Cuttings	Best	Best	Best	OK	–	–	–	–	–	–	OK	Best

Best window | Workable – see 'Selecting material' in the method | Avoid / low success

Plant snapshot

Habit	Much-branched shrub, 0.3–4 m.
Flowering	Apr – Nov (red, sometimes white-yellow).
Habitat	White sand, red clay, gravel and limestone; coastal cliffs, hillsides and plateaus.
Local note	Recorded in the Wanneroo LGA. Cuttings keep a chosen flower colour.

Propagation options

Method	Suitability	Notes	Details in
Cuttings	Good	Firm current season's growth is successful.	This guide
Seed	Main method	Hot water + smoke water; 8–12 days.	→ Seed Sowing Guide
Division	Not used	Woody shrub.	–

Propagation method

Item	Details
Best time to take cuttings	**Dec – Mar**, after flowering, once the new growth has firmed. Nov and Apr are workable. Avoid May – Oct (flowering).
Wood type	**Semi-hardwood** – firm current season's growth (ANPSA).
Cutting size	**50–75 mm** (ANPSA pea-family guidance).
Selecting material	Non-flowering side shoots from healthy plants.
Pre-treatment	• Remove the leaves from the lower two-thirds. • Light wound on firmer stems. • Dip the base in rooting hormone.
Rooting hormone	Semi-hardwood strength, 3 g/L IBA (**Clonex Purple**).
Propagation mix	Standard free-draining cutting mix.
Environment	Humid frame, bright shade, bottom heat.
Expected strike time	Not documented – check from about 6 weeks and record the result.
Post-treatment / aftercare	• Standard hardening off. • Pot on into a free-draining mix.

© 2026 Mitchell Donald. All rights reserved.

Common problems

Problem	Likely cause and fix
Callus, no roots	Trim and re-set; try a lower hormone strength.
Cuttings brown off	Too dry or too soft. Firmer material; keep humidity up.
Base rotting	Mix too wet. Water less.

Notes

Difficulty	**Easy–Moderate (provisional).** Cuttings are reported successful; no strike rates published. Seed is easy after hot water.
Confidence	**Medium.** ANPSA reports cuttings of firm current season's growth are successful; size follows ANPSA pea-family guidance.
Sources	See References: 4, 13, 43, 48

Nursery records

Log each batch here to confirm or update the timing and difficulty above.

Date set	Stock source / provenance	No. set	No. struck	Strike %	Notes

© 2026 Mitchell Donald. All rights reserved.

Thomasia triphylla

No widely used common name

Family **Malvaceae**

DIFFICULTY	WOOD TYPE	BEST MONTHS	CUTTING LENGTH	HORMONE (IBA)
Easy–Moderate *	Semi-hardwood	Dec – Apr	50–75 mm	3 g/L IBA

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Cuttings	Best	Best	Best	Best	OK	–	–	–	–	–	OK	Best

Best window | Workable – see 'Selecting material' in the method | Avoid / low success

Plant snapshot

Habit	Multi-stemmed shrub, 0.3–1.2 m.
Flowering	Jul – Nov (pink-purple or white).
Habitat	Calcareous sand; coastal limestone and dunes.
Local note	Recorded in the Wanneroo LGA. Suits alkaline limestone sand and dry coastal sites.

Propagation options

Method	Suitability	Notes	Details in
Cuttings	Main method	Cuttings usually strike reliably in the genus (provisional).	This guide
Seed	Difficult	Unreliable in the genus; a supplier recommends smoke.	–
Division	Not used	–	–

Propagation method

Item	Details
Best time to take cuttings	**Dec – Apr**, after flowering, once new growth has firmed. Nov and May are workable. Avoid Jun – Oct (cool, then flowering).
Wood type	**Semi-hardwood** – firm current season's growth.
Cutting size	50–75 mm.
Selecting material	Non-flowering side shoots from healthy plants.
Pre-treatment	• Strip leaves from the lower half; reduce large leaves by a third. • Light wound on firmer stems. • Dip the base in rooting hormone.
Rooting hormone	Semi-hardwood strength, 3 g/L IBA (**Clonex Purple**).
Propagation mix	Standard free-draining cutting mix.
Environment	Humid frame, bright shade, bottom heat. Keep the mix moist, not wet, and mist lightly.
Expected strike time	Not documented – check from about 4–6 weeks and record the result.
Post-treatment / aftercare	• Standard hardening off. • Grow on in full sun; tip-prune for a bushier plant.

© 2026 Mitchell Donald. All rights reserved.

Common problems

Problem	Likely cause and fix
Leaves go brown and soft	Too wet or humid. Mist less, improve airflow.
Callus, no roots	Trim and re-set; try a lower hormone strength.
Slow in winter	Use bottom heat, or wait for the summer window.

Notes

Difficulty	**Easy–Moderate (provisional).** *Thomasia* cuttings usually strike reliably, but this species is untested – start with a trial batch.
Other methods	Seed – unreliable in *Thomasia*; a seed supplier recommends smoke treatment.
Confidence	**Medium.** No species data. ANPSA says cuttings of firm growth usually strike reliably in *Thomasia*, though at least one species is difficult.
Sources	See References: 4, 49, 50, 51, 52

Nursery records

Log each batch here to confirm or update the timing and difficulty above.

Date set	Stock source / provenance	No. set	No. struck	Strike %	Notes

© 2026 Mitchell Donald. All rights reserved.

Threlkeldia diffusa

Coast Bonefruit

Family **Amaranthaceae (chenopods)**

DIFFICULTY	WOOD TYPE	BEST MONTHS	CUTTING LENGTH	HORMONE (IBA)
Moderate *	Softwood tips	Dec – Mar	50–75 mm	1.5 g/L IBA

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Cuttings	Best	Best	Best	OK	–	–	–	–	–	–	OK	Best

Best window | Workable – see 'Selecting material' in the method | Avoid / low success

Plant snapshot

Habit	Much-branched, prostrate to erect perennial, 0.1–0.4 m, forming a spreading mat. Semi-succulent leaves.
Flowering	Oct – Nov (tiny, green), followed by small red to black fruits.
Habitat	White or grey sand over limestone, coastal areas and saline flats. Salt- and drought-tolerant.
Local note	Recorded in the City of Joondalup.

Propagation options

Method	Suitability	Notes	Details in
Cuttings	Trial	No published cutting data; trial protocol.	This guide
Seed	Main method	Smoke-water soak; 10–15 days in cool weather.	→ Seed Sowing Guide
Division	Not used	Not documented.	–

Propagation method

Item	Details
Best time to take cuttings	**Dec – Mar** – warm, with plants actively growing after flowering. Nov and Apr are workable. Avoid May – Oct.
Wood type	**Softwood tip cuttings** of the semi-succulent stems.
Cutting size	50–75 mm.
Selecting material	Healthy tips without developing fruit.
Pre-treatment	• Strip lower leaves. • No wounding. • Dip the base in rooting hormone.
Rooting hormone	Softwood strength, 1.5 g/L IBA (**Clonex Green**).
Propagation mix	Very free-draining sandy mix.
Environment	Semi-succulent leaves rot if kept wet – mist very little, keep the mix on the dry side, give good airflow and bright light.
Expected strike time	Not documented – start checking from about 4 weeks and record the result.
Post-treatment / aftercare	• Standard hardening off. • Grow on in full sun; water sparingly.

© 2026 Mitchell Donald. All rights reserved.

Common problems

Problem	Likely cause and fix
Cuttings turn soft and rot	Too wet or humid. Water less, mist less, improve airflow.
Weak, stretched growth	Not enough light. Move to a brighter position.
No roots	Try firmer tips or the seed method.

Notes

Difficulty	**Moderate (provisional).** No cutting data found; seed is the proven method (see Other methods).
Other methods	Seed – the documented method. The Geraldton community nursery soaks seed in smoke water (1:9) for 24 hours; germination takes 10–15 days and happens in cooler weather, so sow in autumn.
Confidence	**Low.** No published cutting data found. Seed is the documented method; cuttings are a trial.
Sources	See References: 1, 4, 12, 28

Nursery records

Log each batch here to confirm or update the timing and difficulty above.

Date set	Stock source / provenance	No. set	No. struck	Strike %	Notes

© 2026 Mitchell Donald. All rights reserved.

References

Numbers match the source list on each species page. Web addresses shortened; accessed September–October 2026.

#	Source	Address
1	Australian Native Plants Society (ANPSA) – Plant Propagation: cuttings, hormones, mixes, hardening off	anpsa.org.au/plant-propagation/
2	ANPSA – Grevillea vestita plant profile	anpsa.org.au/plant_profiles/grevillea-vestita/
3	Australian Plants Society NSW – Grevillea vestita	resources.austplants.com.au/plant/grevillea-vestita/
4	FloraBase, WA Herbarium (DBCA) – species profiles for every species in this guide (e.g. 2463, 2798, 5117, 5162, 7603, 11341, 12585, 12824, 13152, 16449, 42281, 3961, 4256, 5105, 6839, 1982, 5825, 8839, 5237, 6949, 7289, 5011, 5171, 5268, 7626, 49637, 6427, 5887, 49808, 2452, 5365, 6349, 5232, 2329, 15532, 5922, 6295, 4453; others as cited in the Seed Sowing Guide)	florabase.dbca.wa.gov.au
5	ANPSA – Scaevola ramosissima profile (genus note: cuttings strike very readily, seed difficult)	anpsa.org.au/plant_profiles/scaevola-ramosissima/
6	Sustainable Gardening Australia – Fan Flower: take cuttings during the growing season	sgaonline.org.au/fan-flower-fabulous-native-ground-cover/
7	Greenhouse Product News – Scaevola propagation (IBA rates, flower-free cuttings, bottom heat)	gpnmag.com/article/scaevola-outback-fan-flower/
8	ANPSA – Hibbertia vestita profile (genus note: easy from cuttings, seed unreliable)	anpsa.org.au/plant_profiles/hibbertia-vestita/
9	Australian National Botanic Gardens – Hibbertia pedunculata and H. empetrifolia (cutting timing, root rot)	anbg.gov.au/gnp/gnp11/hibbertia-pedunculata.html
10	Trees and Shrubs Online – Hibbertia (nodal cuttings while in growth; seed dormancy)	treesandshrubsonline.org/articles/hibbertia/
11	DBCA Library – wildflower booklet: H. hypericoides difficult from seed or cuttings	library.dbca.wa.gov.au/FullTextFiles/072542.pdf
12	City of Greater Geraldton Community Nursery – Native Plant Propagation Guide (2019)	cgg.wa.gov.au
13	Growth Technology – Clonex rooting gel strengths (Green 1.5, Purple 3, Red 8 g/L IBA)	growthtechnology.com.au
14	EverydayWits – Propagating Scaevola (small-nursery practice, seasonal success rates)	everydaywits.com/propagate-scaevola/
15	Minimalist Gardener – Propagating Australian natives from cuttings (grevillea strike times, sap irritant)	minimalistgardener.com.au
16	Wikipedia – Scaevola canescens, S. repens, S. thesioides (descriptions)	en.wikipedia.org
17	CompleteGrow – How to propagate Grevillea (low-phosphorus feeding)	completegrow.com.au
18	NZ Department of Conservation – Rehabilitation of coastal foredunes using indigenous sand-binding species (Spinifex sericeus stolon cuttings, seed)	doc.govt.nz/Documents/science-and-technical/sfc122.pdf
19	ABC Gardening Australia – Spinifex (40–50 cm runner cuttings)	abc.net.au/gardening/how-to/spinifex/9432152
20	Cottesloe Coastcare Association – plant pages: Spinifex longifolius, S. hirsutus, Eremophila glabra, Rhagodia baccata	cottesloecoastcare.org
21	ANPSA – Eremophila glabra plant profile	anpsa.org.au/plant_profiles/eremophila-glabra/
22	Australian Bushfoods – The Eremophila Page (cutting length, wounding, hormone)	ausbushfoods.com/bushfoodsonline/reports/Plants/EREMO.TXT
23	Australian National Botanic Gardens – Myoporum insulare	anbg.gov.au/gnp/interns-2010/myoporum-insulare.html
24	ANPSA – Lechenaultia biloba profile (cuttings strike very easily; seed difficult)	anpsa.org.au/plant_profiles/lechenaultia-biloba/

© 2026 Mitchell Donald. All rights reserved.

#	Source	Address
25	Better Homes and Gardens – fastest and easiest native plants to propagate (leschenaultia timing)	bhg.com.au/how-to-propagate-natives
26	Nurseries Online – Scaevola crassifolia (semi-hardwood cuttings in warmer months)	nurseriesonline.com.au/plant-index/australian-native-plants/scaevola-crassifolia/
27	Natural Area Nursery – WA native plant propagation (H. hypericoides difficult to propagate)	naturalarea.com.au/revegetation-plant-production/wa-native-plant-propagation/
28	APACE WA – Threlkeldia diffusa	apacewa.org.au/plant/threlkeldia-diffusa/
29	Geographe Community Landcare Nursery – Phyllanthus calycinus (Lysiandra calycina)	gcln.org.au/product/phyllanthus-calycinus/
30	Wikipedia – Hibbertia hypericoides, Eremophila glabra subsp. albicans, Phyllanthus calycinus, Scaevola crassifolia	en.wikipedia.org
31	PMC – Sympatric species of Hibbertia vary in dormancy break and germination requirements	pmc.ncbi.nlm.nih.gov/articles/PMC3336950
32	Cromer E.L. (2007) Seed Germination and Research Records from Alcoa's Marrinup Nursery (seed test figures quoted under Other methods)	library.dbca.wa.gov.au/FullTextFiles/924754.pdf
33	Australian National Botanic Gardens – Carpobrotus glaucescens (layering, 30 cm stem pieces, division)	anbg.gov.au/gnp/interns-2005/carpobrotus-glaucescens.html
34	Australian National Botanic Gardens – Leucophyta brownii (semi-hardened cuttings; dislikes humidity)	anbg.gov.au/gnp/interns-2014/leucophyta-brownii.html
35	ANPSA – Australian Daisies (perennial stem cuttings strike quickly; 3:1 sand:peat mix)	anpsa.org.au/genera/daisies/
36	Australian National Botanic Gardens – Atriplex nummularia (ventilate, avoid mist on cuttings)	anbg.gov.au/gnp/interns-2007/atriplex-nummularia.html
37	Nurseries Online – Atriplex nummularia (15 cm terminal cuttings in spring)	nurseriesonline.com.au/plant-index/australian-native-plants/atriplex-nummularia/
38	Applied Sciences (MDPI, 2025) – Propagation of Atriplex halimus by unrooted cuttings	mdpi.com/2076-3417/15/13/7027
39	Peppy Plants – Hibbertia cuneiformis (propagated by cuttings)	peppyplants.com.au/hibbertia-cuneiformis-unveiling-the-charms-of-the-cutleaf-hibbertia/
40	Wikipedia – Hibbertia cuneiformis (description, flowering, habitat)	en.wikipedia.org/wiki/Hibbertia_cuneiformis
41	ANPSA – Melaleuca genus page (75–100 mm cuttings, strip, wound, hormone)	anpsa.org.au/genera/melaleuca/
42	ANPSA – Kennedia prostrata profile (cuttings of firm current season's growth)	anpsa.org.au/plant_profiles/kennedia-prostrata/
43	ANPSA – Pea family (Fabaceae) genus page (50–75 mm cuttings, leaves off the lower two-thirds)	anpsa.org.au/genera/pea-family/
44	ANPSA – Hardenbergia comptoniana profile (cuttings of firm current season's growth strike well)	anpsa.org.au/plant_profiles/hardenbergia-comptoniana/
45	ANPSA – Hemiandra pungens profile (relatively easy from cuttings; unreliable in humid conditions; grafting)	anpsa.org.au/plant_profiles/hemiandra-pungens/
46	Gardening With Angus – Hemiandra pungens (softwood and semi-hardwood cuttings)	gardeningwithangus.com.au/hemiandra-pungens-snake-plant/
47	APACE WA – Hemiandra pungens (dieback resistant)	apacewa.org.au/plant/hemiandra-pungens/
48	ANPSA – Templetonia retusa profile (cuttings of firm current season's growth are successful)	anpsa.org.au/plant_profiles/templetonia-retusa/
49	ANPSA – Thomasia grandiflora profile (genus: cuttings of firm growth usually strike reliably; seed unreliable)	anpsa.org.au/plant_profiles/thomasia-grandiflora/
50	Geographe Community Landcare Nursery – Thomasia triphylla (coastal, drought tolerant, alkaline sand)	gcln.org.au/product/thomasia-triphylla/
51	Nindethana Australian Seed – Thomasia triphylla and Scaevola nitida seed (smoke treatment recommended)	nindethana.net.au

© 2026 Mitchell Donald. All rights reserved.

#	Source	Address
52	ANPSA – Australian Plants Online (2007): Thomasia pygmaea, cuttings difficult to strike	anpsa.org.au/APOL2007/jul07-2.html
53	ANPSA – Hypocalymma robustum profile (cuttings preferred, strike readily from hardened growth)	anpsa.org.au/plant_profiles/hypocalymma-robustum/
54	ANPSA – Geraldton wax and relatives (75–100 mm cuttings in January, strip lower two-thirds, wound, hormone)	anpsa.org.au/genera/geraldton-wax-and-relatives/
55	Gardening With Angus – Hypocalymma robustum (softwood and semi-hardwood cuttings)	gardeningwithangus.com.au/hypocalymma-robustum-swan-river-myrtle/
56	Australian Plants Society NSW – Grevillea crithmifolia (cuttings may be slow to root; shrub and groundcover forms)	resources.austplants.com.au/plant/grevillea-crithmifolia/
57	APACE WA – Grevillea crithmifolia (dieback susceptible)	apacewa.org.au/plant/grevillea-crithmifolia/
58	ANPSA – Grevillea preissii profile (cuttings slow; hormone improves success; nick seed)	anpsa.org.au/plant_profiles/grevillea-preissii/
59	ANPSA – Grevillea genus page (75–100 mm cuttings, strip lower half to two-thirds, wounding, hormone; phosphorus)	anpsa.org.au/genera/grevillea/
60	Olde P.M. (2022) Telopea – Grevillea preissii subsp. glabrilimba raised to species rank as G. glabrilimba	openjournals.library.sydney.edu.au/TEL/article/view/15872
61	Nindethana Australian Seed – Grevillea crithmifolia and G. preissii seed (smoke treatment recommended)	nindethana.net.au
62	ANPSA – Eremophila and relatives (Myoporum: cuttings of hardened current season's growth strike readily)	anpsa.org.au/genera/eremophila-and-relatives/
63	APACE WA – Myoporum caprarioides (winter-wet sites; berries attract birds)	apacewa.org.au/plant/myoporum-caprarioides/
64	ANPSA – Anthocercis littorea profile (cuttings of hardened growth strike reasonably well)	anpsa.org.au/plant_profiles/anthocercis-littorea/
65	Wikipedia – Anthocercis (tropane alkaloids; poisoning)	en.wikipedia.org/wiki/Anthocercis
66	ANPSA – Pimelea physodes profile (genus: cuttings successful but bark tears; seed difficult, smoke helps)	anpsa.org.au/plant_profiles/pimelea-physodes/
67	ANPSA – Pimelea ferruginea profile (firm current season's growth usually strikes reliably)	anpsa.org.au/plant_profiles/pimelea-ferruginea/
68	Commander L.E. et al. (2009) Journal of Arid Environments – GA, smoke water and karrikinolide raise germination (incl. Anthocercis littorea)	sciencedirect.com/science/article/abs/pii/S0140196309000287
69	Cochrane A. (2007) Seed Notes 17 – Hibbertia (most plants grown from cuttings; dormant seed; GA 25 mg/L). Wildflower Society of WA / DEC	library.dbca.wa.gov.au/FullTextFiles/PAM02004.pdf
70	Thiele K.R. – Nuytsia 30: the Hibbertia spicata group split into six species, incl. H. leptotheca	florabase.dbca.wa.gov.au/nuytsia/article/924
71	ANPSA – Scaevola aemula profile (cuttings strike very easily from firm current season's growth)	anpsa.org.au/plant_profiles/scaevola-aemula/
72	Australian Seed – Scaevola nitida (smoke pre-treatment; 21–42 days; tolerates full coastal exposure)	australianseed.com/shop/item/scaevola-nitda
73	ANGAIR – Propagating the impossibles (Pimelea: winter sowing with smoke water)	angair.org.au/propagating-the-impossibles/
74	Cochrane A. (2007) Seed Notes 20 – Lasiopetalum, Thomasia and Guichenotia (often grown from cuttings; hard seed). Wildflower Society of WA / DEC	library.dbca.wa.gov.au/FullTextFiles/PAM02007.pdf
75	Gardening With Angus – Guichenotia ledifolia (semi-hardwood cuttings)	gardeningwithangus.com.au/guichenotia-ledifolia/
76	ANPSA – Guichenotia macrantha profile (cuttings of firm current season's growth usually strike reliably)	anpsa.org.au/plant_profiles/guichenotia-macrantha/
77	Botanic Gardens of South Australia Plant Selector – Guichenotia ledifolia (drought and lime tolerant)	plantselector.botanicgardens.sa.gov.au

© 2026 Mitchell Donald. All rights reserved.

#	Source	Address
78	ANPSA – Scaevola calendulacea profile (cuttings of this and other Scaevola species strike very readily)	anpsa.org.au/plant_profiles/scaevola-calendulacea/
79	ANPSA – Leucopogon parviflorus profile (cuttings unreliable; firm current season's growth and a root-promoting hormone)	anpsa.org.au/?p=11543
80	Understorey Network (Tasmania) – Leucopogon parviflorus (usually grown from cuttings; low strike; very fine roots)	database.understorey-network.org.au/plant/leucopogon-parviflorus
81	NZ Plant Conservation Network – Leucopogon parviflorus (semi-hardwood cuttings, slow to strike)	nzpcn.org.nz/flora/species/leucopogon-parviflorus/
82	Doig – Epacrids from cuttings, ANPSA Australian Plants Online (Dec 1999): half-hardened wood 80–250 mm, medium-strength hormone, cold non-misting frame, 6–12 months	anpsa.org.au/APOL16/dec99-6.html
83	Australian Plants Society NSW – Melaleuca cardiophylla (propagate from seed and cuttings; regrows after fire)	resources.austplants.com.au/plant/melaleuca-cardiophylla/
84	Dixon K. & Roche S., ANPSA Australian Plants Online (Jun 1996) – smoke germination: drupaceous Epacridaceae don't respond	anpsa.org.au/APOL2/jun96-6.html
85	ANPSA – Smoke germination of Australian plants (ageing seed in soil 1–2 years plus repeat smoke for Ericaceae)	anpsa.org.au/smoke-germination-of-australian-plants/
86	State Flora SA – coastal garden plants (Leucopogon parviflorus buffers coastal wind; birds and lizards eat the fruit)	stateflora.sa.gov.au/news/coastal-garden-plants
87	Keighery G.J. & Dixon K. (1984) Australian Plants 12 – Pimelea in WA: separate male and female plants, P. argentea seed (poor germination, 21 days), cuttings 3–4 cm from autumn to late winter, bark strips	anpsa.org.au/wp-content/uploads/Australian-Plants/Australian-Plants-Vol12-99.pdf
88	Wikipedia – Pimelea argentea (male and female flowers on separate plants)	en.wikipedia.org/wiki/Pimelea_argentea
89	Australian Plants Society NSW – Baeckea linifolia (genus: seed hard to collect; cuttings of firm current season's growth strike readily)	resources.austplants.com.au/plant/baeckea-linifolia/
90	ANPSA – Euryomyrtus ramosissima profile (former Baeckea; strikes readily from firm current season's growth)	anpsa.org.au/?p=1767
91	DBCA Interim Recovery Plan 128 – Synaphea quartzitica (notes that S. spinulosa has been grown from cuttings at the BGPA nursery, extremely low strike rate; GA and nicking)	dbca.wa.gov.au/media/2652/download
92	DBCA recovery plans – Synaphea sp. Pinjarra and S. stenoloba (low seed set; mild dieback susceptibility; tissue culture)	dbca.wa.gov.au/media/2654/download
93	ANPSA – Conostephium pendulum profile (cuttings of current season's growth; individual tubes to limit root disturbance)	anpsa.org.au/plant_profiles/conostephium-pendulum/
94	Hislop M. (2013) Nuytsia 23 – Conostephium revision (habitat; fleshy drupe; fire-resistant rootstock)	florabase.dbca.wa.gov.au/science/nuytsia/680.pdf
95	NZ Plant Conservation Network – Atriplex cinerea fact sheet (easy from cuttings; layers itself; most stock female)	nzpcn.org.nz
96	Understorey Network (Tasmania) – Atriplex cinerea (strikes easily from cuttings; needs good drainage; male and female plants)	database.understorey-network.org.au/plant/atriplex-cinerea
97	Rye B.L. (2026) Nuytsia 37 – notes on Baeckea in WA (B. robusta's placement in Baeckea questioned)	florabase.dbca.wa.gov.au/science/nuytsia/1104.pdf
98	Wikipedia – Melaleuca lanceolata (resistant to salt spray; windbreak and shelterbelt use; synonyms)	en.wikipedia.org/wiki/Melaleuca_lanceolata
99	Understorey Network (Tasmania) – Acrotriche cordata (usually grown from cuttings of new growth; difficult from seed; erosion control)	database.understorey-network.org.au/plant/acrotriche-cordata
100	Seeds of South Australia – Acrotriche cordata (morphophysiological dormancy; collection and cleaning)	seedsofsa.au (read via syzygium.xyz mirror)
101	ANPSA – Diplolaena microcephala profile (cuttings of firm current season's growth successful; seed difficult; leaf fungal disease)	anpsa.org.au/plant_profiles/diplolaena-microcephala/

© 2026 Mitchell Donald. All rights reserved.

#	Source	Address
102	The Seed Collection – Yanchep Rose, Diplolaena angustifolia (dislikes humid conditions and wet, boggy soils)	theseedcollection.com.au/Yanchep-Rose
103	Flora of Australia online – Diplolaena angustifolia (Wanneroo north to near Eneabba)	profiles.ala.org.au/opus/foa/profile/Diplolaena%20angustifolia

Revision log

Version	Date	Changes
1.0	Sep 2026	First edition: Grevillea vestita, Hibbertia racemosa, Scaevola canescens, S. repens, S. thesioides.
1.1	Sep 2026	Added 10 species: Eremophila glabra, Hibbertia hypericoides, Lechenaultia linarioides, Myoporum insulare, Phyllanthus calycinus, Scaevola crassifolia, S. globulifera, Spinifex hirsutus, S. longifolius, Threlkeldia diffusa. Added page numbers to the species index.
1.2	Sep 2026	Added 11 species: Atriplex isatidea, Calothamnus quadrifidus, Carpobrotus virescens, Hibbertia cuneiformis, Kennedia prostrata, Leucophyta brownii, Melaleuca huegelii, M. systena, Olearia axillaris, Pithocarpa cordata, Rhagodia baccata. Added a ‘Propagation options’ table (cuttings, seed, division) to every species page.
1.3	Sep 2026	Added 4 species: Hardenbergia comptoniana, Hemiandra pungens, Templetonia retusa, Thomasia triphylla.
1.4	Sep 2026	Added 3 species: Grevillea crithmifolia, G. preissii, Hypocalymma robustum.
1.5	Sep 2026	Added 3 species: Anthocercis littorea, Myoporum caprarioides, Pimelea calcicola.
1.6	Sep 2026	Added 4 species: Guichenotia ledifolia, Hibbertia spicata (incl. H. leptotheca), Pimelea sulphurea, Scaevola nitida.
1.7	Oct 2026	Added 3 species: Leucopogon parviflorus, Melaleuca cardiophylla, Styphelia insularis (formerly Leucopogon insularis).
1.8	Oct 2026	Added 5 species: Atriplex cinerea, Baeckea robusta, Conostephium preissii, Pimelea argentea, Synaphea spinulosa.
1.9	Oct 2026	Added 3 species: Acrotriche cordata, Diplolaena angustifolia, Melaleuca lanceolata.

© 2026 Mitchell Donald. All rights reserved.