

Native Plant Cuttings Guide

Western Australian native species – Joondalup, WA

Version 1.3 · September 2026 · 30 species

Species in this edition (alphabetical)

Full index on page 2

● <i>Atriplex isatidea</i>	7	● <i>Kennedia prostrata</i>	27	● <i>Scaevola canescens</i>	47
● <i>Calothamnus quadrifidus</i>	9	● <i>Lechenaultia linarioides</i>	29	● <i>Scaevola crassifolia</i>	49
● <i>Carpobrotus virescens</i>	11	● <i>Leucophyta brownii</i>	31	● <i>Scaevola globulifera</i>	51
● <i>Eremophila glabra</i>	13	● <i>Melaleuca huegelii</i>	33	● <i>Scaevola repens</i>	53
● <i>Grevillea vestita</i>	15	● <i>Melaleuca systema</i>	35	● <i>Scaevola thesioides</i>	55
● <i>Hardenbergia comptoniana</i>	17	● <i>Myoporum insulare</i>	37	● <i>Spinifex hirsutus</i>	57
● <i>Hemiandra pungens</i>	19	● <i>Olearia axillaris</i>	39	● <i>Spinifex longifolius</i>	59
● <i>Hibbertia cuneiformis</i>	21	● <i>Phyllanthus calycinus</i>	41	● <i>Templetonia retusa</i>	61
● <i>Hibbertia hypericoides</i>	23	● <i>Pithocarpa cordata</i>	43	● <i>Thomasia triphylla</i>	63
● <i>Hibbertia racemosa</i>	25	● <i>Rhagodia baccata</i>	45	● <i>Threlkeldia diffusa</i>	65

● Easy ● Easy-Mod. ● Moderate ● Mod.-Difficult ● Difficult

* Provisional rating – based on genus behaviour; confirm with your own nursery records.

Compiled from published sources (see References). Windows and ratings are starting points for local practice.

How to use this guide

Each species has its own page set, listed alphabetically by scientific name. Every page follows the same layout: quick facts, a cutting calendar, a plant snapshot, the full propagation method, common problems, and a blank record table so the nursery can log strike rates and refine the advice over time.

Timing windows are written for the Joondalup / Perth coastal climate: hot, dry summers and mild, wet winters. They combine general native-propagation practice with each species' flowering period from FloraBase. Rooting is fastest in warm weather and slows markedly in winter unless bottom heat is used.

Species index – at a glance

Species	Page	Difficulty	Wood type	Best months	Length	Hormone (IBA)
<i>Atriplex isatidea</i> Coast Saltbush	7	Easy-Moderate *	Semi-hardwood tips	Sep – Nov	100–150 mm	1.5–3 g/L IBA
<i>Calothamnus quadrifidus</i> One-sided Bottlebrush	9	Easy-Moderate *	Semi-hardwood	Jan – Apr	75–100 mm	3 g/L IBA
<i>Carpobrotus virescens</i> Coastal Pigface	11	Easy	Succulent stem pieces	Sep – Nov; Mar – Apr	10–15 cm (tubes)	Not needed
<i>Eremophila glabra</i> Tar Bush / Common Emu Bush	13	Easy	Semi-hardwood	Jan – Apr	75–100 mm	3 g/L IBA
<i>Grevillea vestita</i>	15	Easy	Semi-hardwood	Dec – Apr	80–120 mm	3 g/L IBA
<i>Hardenbergia comptoniana</i> Native Wisteria	17	Easy-Moderate	Semi-hardwood	Dec – Mar	50–75 mm	3 g/L IBA
<i>Hemiandra pungens</i> Snakebush	19	Easy	Softwood – semi-hardwood	Oct – Mar	50–75 mm	1.5–3 g/L IBA
<i>Hibbertia cuneiformis</i> Cutleaf Hibbertia	21	Easy-Moderate *	Semi-hardwood	Mar – May	75–100 mm	3 g/L IBA
<i>Hibbertia hypericoides</i> Yellow Buttercups	23	Difficult	Semi-hardwood	Dec – Feb	50–75 mm	3 g/L IBA (trial range)
<i>Hibbertia racemosa</i> Stalked Guinea Flower	25	Moderate *	Semi-hardwood	Jan – Apr	50–75 mm	1.5–3 g/L IBA
<i>Kennedia prostrata</i> Scarlet Runner / Running Postman	27	Easy-Moderate	Semi-hardwood	Dec – Mar	75–100 mm	3 g/L IBA
<i>Lechenaultia linarioides</i> Yellow Leschenaultia	29	Easy *	Softwood – semi-hardwood	Feb – Apr	50–75 mm	1.5–3 g/L IBA
<i>Leucophyta brownii</i> Cushion Bush	31	Easy-Moderate *	Semi-hardwood	Mar – May; Sep – Oct	50–75 mm	1.5–3 g/L IBA
<i>Melaleuca huegelii</i> Chenille Honeymyrtle	33	Easy *	Semi-hardwood	Feb – Apr	75–100 mm	3 g/L IBA
<i>Melaleuca systema</i> Coastal Honeymyrtle	35	Easy *	Semi-hardwood	Dec – Jan; Apr – May	75–100 mm	3 g/L IBA
<i>Myoporum insulare</i> Blueberry Tree / Common Boobialla	37	Easy	Semi-hardwood	Feb – Apr	75–100 mm	3 g/L IBA
<i>Olearia axillaris</i> Coastal Daisybush	39	Easy *	Semi-hardwood	Aug – Oct	75–100 mm	1.5–3 g/L IBA
<i>Phyllanthus calycinus</i> False Boronia	41	Moderate *	Semi-hardwood	Feb – Apr	50–75 mm	3 g/L IBA
<i>Pithocarpa cordata</i> Tangle Daisy	43	Easy-Moderate *	Semi-hardwood tips	Sep – Nov; May	50–75 mm	1.5–3 g/L IBA

Species	Page	Difficulty	Wood type	Best months	Length	Hormone (IBA)
<i>Rhagodia baccata</i> Berry Saltbush	45	Moderate *	Semi-hardwood tips	Sep – Nov	75–100 mm	3 g/L IBA
<i>Scaevola canescens</i> Grey Scaevola	47	Easy–Moderate *	Softwood – semi-hardwood	Nov – Jan	50–80 mm	1.5–3 g/L IBA
<i>Scaevola crassifolia</i> Thick-leaved Fan-flower	49	Easy	Semi-hardwood	Mar – May	75–100 mm	3 g/L IBA
<i>Scaevola globulifera</i>	51	Easy *	Softwood – semi-hardwood	Jan – Apr	50–80 mm	1.5–3 g/L IBA
<i>Scaevola repens</i>	53	Easy *	Softwood – semi-hardwood	Mar – May	50–80 mm	1.5–3 g/L IBA
<i>Scaevola thesioides</i>	55	Easy–Moderate *	Softwood – semi-hardwood	Apr – May	50–75 mm	1.5–3 g/L IBA
<i>Spinifex hirsutus</i> Hairy Spinifex	57	Moderate *	Runner (stolon) sections	Mar – May	10–15 cm, 1–2 nodes	Optional (1.5 g/L)
<i>Spinifex longifolius</i> Beach Spinifex	59	Moderate *	Runner (stolon) sections	Mar – May	10–15 cm, 1–2 nodes	Optional (1.5 g/L)
<i>Templetonia retusa</i> Cockies Tongues	61	Easy–Moderate *	Semi-hardwood	Dec – Mar	50–75 mm	3 g/L IBA
<i>Thomasia triphylla</i>	63	Easy–Moderate *	Semi-hardwood	Dec – Apr	50–75 mm	3 g/L IBA
<i>Threlkeldia diffusa</i> Coast Bonefruit	65	Moderate *	Softwood tips	Dec – Mar	50–75 mm	1.5 g/L IBA

* Provisional rating – based on genus behaviour; update once nursery strike rates are recorded.

Cutting calendar

Species	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
<i>Atriplex isatidea</i>	OK	OK	–	–	–	–	–	OK	Best	Best	Best	OK
<i>Calothamnus quadrifidus</i>	Best	Best	Best	Best	OK	–	–	–	–	–	–	OK
<i>Carpobrotus virescens</i>	OK	OK	Best	Best	OK	–	–	OK	Best	Best	Best	OK
<i>Eremophila glabra</i>	Best	Best	Best	Best	OK	–	–	–	–	–	OK	OK
<i>Grevillea vestita</i>	Best	Best	Best	Best	OK	–	–	–	–	–	OK	Best
<i>Hardenbergia comptoniana</i>	Best	Best	Best	OK	–	–	–	–	–	–	OK	Best
<i>Hemianandra pungens</i>	Best	Best	Best	OK	–	–	–	–	OK	Best	Best	Best
<i>Hibbertia cuneiformis</i>	OK	OK	Best	Best	Best	–	–	–	–	OK	OK	OK
<i>Hibbertia hypericoides</i>	Best	Best	OK	–	–	–	–	–	–	–	OK	Best
<i>Hibbertia racemosa</i>	Best	Best	Best	Best	OK	–	–	–	–	–	–	OK
<i>Kennedia prostrata</i>	Best	Best	Best	OK	–	–	–	–	–	–	OK	Best
<i>Lechenaultia linarioides</i>	OK	Best	Best	Best	OK	–	–	–	–	OK	OK	OK
<i>Leucophyta brownii</i>	–	OK	Best	Best	Best	OK	–	OK	Best	Best	OK	–
<i>Melaleuca huegelii</i>	OK	Best	Best	Best	OK	–	–	–	–	–	–	OK
<i>Melaleuca systema</i>	Best	OK	OK	Best	Best	OK	–	–	–	–	OK	Best
<i>Myoporum insulare</i>	OK	Best	Best	Best	OK	–	–	–	–	OK	OK	OK
<i>Olearia axillaris</i>	–	–	OK	OK	–	–	OK	Best	Best	Best	OK	–
<i>Phyllanthus calycinus</i>	OK	Best	Best	Best	OK	–	–	–	–	–	–	–
<i>Pithocarpa cordata</i>	–	–	–	OK	Best	OK	–	OK	Best	Best	Best	OK
<i>Rhagodia baccata</i>	OK	OK	OK	–	–	–	–	OK	Best	Best	Best	OK
<i>Scaevola canescens</i>	Best	OK	OK	OK	OK	–	–	–	–	OK	Best	Best
<i>Scaevola crassifolia</i>	OK	OK	Best	Best	Best	–	–	–	–	OK	OK	OK
<i>Scaevola globulifera</i>	Best	Best	Best	Best	OK	–	–	–	–	–	OK	OK
<i>Scaevola repens</i>	OK	OK	Best	Best	Best	–	–	–	OK	OK	OK	OK
<i>Scaevola thesioides</i>	OK	OK	OK	Best	Best	–	–	–	OK	OK	OK	OK
<i>Spinifex hirsutus</i>	OK	OK	Best	Best	Best	–	–	–	OK	OK	OK	OK
<i>Spinifex longifolius</i>	OK	OK	Best	Best	Best	–	–	–	OK	OK	OK	OK
<i>Templetonia retusa</i>	Best	Best	Best	OK	–	–	–	–	–	–	OK	Best
<i>Thomasia triphylla</i>	Best	Best	Best	Best	OK	–	–	–	–	–	OK	Best
<i>Threlkeldia diffusa</i>	Best	Best	Best	OK	–	–	–	–	–	–	OK	Best

Perth season

Summer

Autumn

Winter

Spring

Sum.



Best window



Workable – see ‘Selecting material’ in the method



Avoid / low success

Key to terms and ratings

Wood types and the bend test

Wood type	What it looks like	Bend test	Use
Softwood	Soft, flexible, bright new growth from the current flush.	Flops or stays bent.	Roots fast but wilts and rots easily. Suits soft-stemmed plants such as <i>Scaevola</i> .
Semi-hardwood	Current season's growth that has begun to firm; bark starting to colour.	Bends about 60° and springs back.	The standard for most WA native shrubs. More forgiving than softwood.
Hardwood	Fully mature, woody stems from previous seasons.	Stiff; snaps rather than bends.	Slow to root. Rarely needed for the species in this guide.

Difficulty ratings

Rating	Meaning
Easy	Strikes reliably with the standard method.
Easy-Moderate	Usually strikes; one factor (e.g. moisture, fragile stems) needs extra care.
Moderate	Strikes, but results depend on timing, material and moisture control. Set extra cuttings.
Moderate-Difficult	Low or variable strike; needs careful timing and good facilities.
Difficult	Low or erratic strike; specialist techniques or alternative methods usually needed.

* **Provisional** – rating is based on how the genus generally behaves because little species-specific data is published. Replace it with the nursery's own figure once a few batches have been recorded.

Confidence

Level	Meaning
High	Species-specific propagation information is published and consistent.
Medium	Built from genus-level information plus the species' own traits (habit, leaf type, flowering time). Sound starting point – verify with trials.
Low	Little information available; treat as a trial protocol.

Standard cutting method

Species pages refer back to these steps and only note where a species differs.

Step	What to do
1. Collect	Take material early in the morning from healthy, pest-free, well-watered, non-flowering stock. Seal in a moist plastic bag and keep in an esky out of the sun. Process the same day.
2. Prepare	Use a sharp, clean blade (wipe with alcohol between stock plants). Cut just below a node. Strip leaves from the lower half to two-thirds without tearing the bark. Reduce large leaves by about one-third. Remove flowers and buds.
3. Wound (optional)	On firmer stems, remove a thin sliver of bark from the bottom 15–20 mm to expose more rooting area.
4. Hormone	Decant a small amount into a clean container, dip the base to about 1 cm, and discard what's left. Store hormone in the fridge – it has a limited shelf life.
5. Set	Make a hole with a dibber so the hormone isn't wiped off, insert to the bottom of the hole, and firm the mix. Single tubes make potting on easier than community pots.
6. Environment	Water in and place in a humid frame or propagation house in bright shade. Bottom heat so the base is 5–10°C warmer than the air. Keep the mix moist, not wet. Mist sparingly on hairy-leaved species.
7. Check	Most cuttings strike in 4–8 weeks in warm weather. Look for roots at the drainage holes. Remove any rotting cuttings promptly.
8. Harden off	About 2 weeks with cover closed by day and open at night, then about 2 weeks in shade, then into the open. Many struck cuttings are lost by skipping this step.
9. Pot on	After 1–3 months, pot on into larger containers. Don't let roots become pot-bound.

Rooting hormone guide

Product (example)	IBA strength	Use for
Clonex Green	1.5 g/L	Softwood and soft tip cuttings
Clonex Purple	3 g/L	Semi-hardwood – the general all-rounder
Clonex Red	8 g/L	Hardwood cuttings

Equivalent powders or liquids work too – match the strength to the wood type. Callus forming without roots usually means the hormone is too strong for that material.

Mix, hygiene and collecting

Topic	Guidance
Propagation mix	About 75–80% washed coarse river sand with 20–25% coir or peat. Add extra perlite or coarse sand for rot-prone species.
Hygiene	Fresh mix, clean tubes and benches. Many WA natives (including <i>Hibbertia</i> and <i>Grevillea</i>) are susceptible to <i>Phytophthora</i> . Discard diseased cuttings; use fungicide only if a problem persists.
Collecting	Collecting from bushland generally requires a DBCA flora licence and landholder permission. Use local-provenance stock and label every batch with its source.

Atriplex isatidea

Coast Saltbush

Family
Amaranthaceae
(chenopods)

DIFFICULTY	WOOD TYPE	BEST MONTHS	CUTTING LENGTH	HORMONE (IBA)
Easy-Moderate *	Semi-hardwood tips	Sep – Nov	100–150 mm	1.5–3 g/L IBA

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Cuttings	OK	OK	–	–	–	–	–	OK	Best	Best	Best	OK
	Best window	Workable – see ‘Selecting material’ in the method						Avoid / low success				

Plant snapshot

Habit	Grey-leaved, spreading shrub, 0.5–2.5 m. Male and female flowers on the same plant.
Flowering	Mainly Mar – Jun (small, inconspicuous).
Habitat	White sand on coastal dunes.
Local note	Recorded in the Wanneroo LGA. Very salt- and wind-tolerant – a useful dune stabiliser.

Propagation options

Method	Suitability	Notes	Details in
Cuttings	Good	Saltbush cuttings are generally reliable (provisional for this species).	This guide
Seed	Main method	Soak 4 days in changed water, then smoke water.	→ Seed Sowing Guide
Division	Not used	Woody shrub.	–

Propagation method

Item	Details
Best time to take cuttings	Sep – Nov, in active spring growth (the time recommended for the related Old Man Saltbush). Aug and Dec – Feb are workable. Avoid Mar – Jul (flowering, then cool, slow growth).
Wood type	Semi-hardwood terminal (tip) cuttings of current season's growth.
Cutting size	100–150 mm. Terminal cuttings of about 15 cm are used for <i>A. nummularia</i> .
Selecting material	Vigorous, non-flowering tips from healthy plants. Avoid old, woody stems.
Pre-treatment	- Strip leaves from the lower half. - No wounding needed. - Dip the base in rooting hormone.
Rooting hormone	Softwood to semi-hardwood strength (1.5–3 g/L IBA, Clonex Green or Purple). Related saltbushes root without hormone, so it mainly evens out the batch.
Propagation mix	Free-draining mix of sand and potting mix.
Environment	Good ventilation and no mist – misting encourages a leaf-spotting fungus on saltbush cuttings (ANBG, <i>A. nummularia</i>). Bottom heat helps; keep the mix just moist.
Expected strike time	Not documented for this species. In one trial, 20 cm unrooted <i>A. halimus</i> cuttings with no hormone gave about 49% survival at 8 weeks . Check from about 4 weeks and record your result.
Post-treatment / aftercare	- Harden off, then grow on in full sun. - Water sparingly once established. - Tip-prune for a bushier tube.

Common problems

Problem	Likely cause and fix
Leaf spots, leaves dropping	Too humid or misted. Stop misting, open the frame, improve airflow.
Base rotting	Mix too wet. Use a sandier mix and water less.
Leggy, pale growth	Not enough light. Grow on in full sun.

Notes

Difficulty	Easy-Moderate (provisional). Saltbushes generally strike readily from tip cuttings; the main risk is leaf disease under mist. Seed is the more common nursery method.
Confidence	Medium. No species-specific cutting data. Method follows published results for related saltbushes (<i>A. nummularia</i> , <i>A. halimus</i>).
Sources	See References: 4, 36, 37, 38

Nursery records

Log each batch here to confirm or update the timing and difficulty above.

Date set	Stock source / provenance	No. set	No. struck	Strike %	Notes

Calothamnus quadrifidus

One-sided Bottlebrush

Family
Myrtaceae

DIFFICULTY	WOOD TYPE	BEST MONTHS	CUTTING LENGTH	HORMONE (IBA)
Easy-Moderate *	Semi-hardwood	Jan – Apr	75–100 mm	3 g/L IBA

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Cuttings	Best	Best	Best	Best	OK	–	–	–	–	–	–	OK

Best window	Workable – see 'Selecting material' in the method	Avoid / low success
-------------	---	---------------------

Plant snapshot

Habit	Erect, compact or spreading shrub, usually 0.9–2 m.
Flowering	Jun – Dec (red, one-sided brushes).
Habitat	Wide variety of soils and habitats.
Local note	Recorded in the Wanneroo LGA. Several subspecies exist – label stock by subspecies and provenance.

Propagation options

Method	Suitability	Notes	Details in
Cuttings	Good	Presumed to strike like related melaleucas (provisional).	This guide
Seed	Main method	Easy from seed held in old capsules.	→ Seed Sowing Guide
Division	Not used	Woody shrub.	–

Propagation method

Item	Details
Best time to take cuttings	Jan – Apr, after flowering, once the new growth has firmed. Dec and May are workable. Avoid Jun – Nov (flowering).
Wood type	Semi-hardwood – firm current season's growth.
Cutting size	75–100 mm, as for <i>Melaleuca</i> .
Selecting material	Non-flowering side shoots. Keep subspecies and forms separate.
Pre-treatment	- Strip the needle-like leaves from the lower half to two-thirds. - Light wound on firmer stems. - Dip the base in rooting hormone.
Rooting hormone	Semi-hardwood strength, 3 g/L IBA (Clonex Purple).
Propagation mix	Standard free-draining cutting mix.
Environment	Humid frame, bright shade, bottom heat.
Expected strike time	Not documented – check from about 6 weeks and record the result.
Post-treatment / aftercare	- Standard hardening off. - Pot on into a free-draining native mix.

Common problems

Problem	Likely cause and fix
Cuttings brown off	Too dry or too soft. Use firmer material and keep humidity up until rooted.
Callus, no roots	Trim and re-set; try a lower hormone strength.
Slow in winter	Use bottom heat, or wait for the Jan – Apr window.

Notes

Difficulty	Easy-Moderate (provisional). Expected to behave like related melaleucas, but untested here. Seed is easy and is the main method.
Confidence	Low. ANPSA notes the genus is 'presumably' grown from cuttings. Method follows related melaleucas – trial protocol.
Sources	See References: 1, 4, 13, 41

Nursery records

Log each batch here to confirm or update the timing and difficulty above.

Date set	Stock source / provenance	No. set	No. struck	Strike %	Notes

Carpobrotus virescens

Coastal Pigface

Family
Aizoaceae

DIFFICULTY	WOOD TYPE		BEST MONTHS			CUTTING LENGTH			HORMONE (IBA)			
Easy	Succulent stem pieces		Sep – Nov; Mar – Apr			10–15 cm (tubes)			Not needed			
	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Cuttings	OK	OK	Best	Best	OK	–	–	OK	Best	Best	Best	OK
	Best window		Workable – see 'Selecting material' in the method						Avoid / low success			

Plant snapshot

Habit	Prostrate succulent groundcover, 0.1–0.3 m high and 0.5–3 m wide, rooting along the stems.
Flowering	Jun – Dec or Jan (purple-pink, sometimes white), followed by fleshy fruit.
Habitat	Coastal limestone and sand dunes.
Local note	Recorded in the Wanneroo LGA. Collect only from plants you are sure are native – the introduced Hottentot Fig (<i>C. edulis</i>) is naturalised on some coasts.

Propagation options

Method	Suitability	Notes	Details in
Cuttings	Main method	Stem pieces root very easily; rooted runners can be lifted.	This guide
Seed	Good	Ferment fruit, sow fine seed on the surface; 6–8 days.	→ Seed Sowing Guide
Division	Possible	Lift stems that have rooted – treat as cuttings.	–

Propagation method

Item	Details
Best time to take cuttings	Sep – Nov and Mar – Apr, when plants are actively growing. Aug, Dec – Feb and May are workable. Avoid Jun – Jul (slow growth; cuttings rot in cold, wet mix).
Wood type	Succulent stem (runner) pieces – not woody cuttings. Use firm stems with plump leaves.
Cutting size	Tubes: 10–15 cm pieces with 2–3 leaf pairs. Direct planting / layering: about 30 cm, with at least 5 cm left above the sand (ANBG, <i>C. glaucescens</i>).
Selecting material	Healthy runners. Pieces that have already rooted at the nodes are ideal. Avoid shrivelled or damaged stems.
Pre-treatment	- Remove the leaves from the lowest 1–2 nodes. - Leave the cut ends to dry in the shade for a day or two before setting – this reduces rot. - No hormone needed.
Rooting hormone	Not needed – stems root readily at the nodes.
Propagation mix	Coarse washed sand or a very sandy, free-draining mix.
Environment	Open bench in bright light or light shade. No mist and no humid frame – succulent leaves rot. Water sparingly until rooted.
Expected strike time	Not documented, but usually quick – check from about 3 weeks.

Item	Details
Post-treatment / aftercare	• Grow on in full sun. • Pot on or plant out once roots hold the mix together. • Water sparingly – overwatering is the main cause of losses.

Common problems

Problem	Likely cause and fix
Stems go soft and rot	Too wet or humid. Dry the cut ends first, water less, move out of the humid frame.
Leaves shrivel	Too dry for too long. Water lightly once set.
Pale, stretched growth	Not enough light. Move to full sun.

Notes

Difficulty	Easy. Pigface roots readily from stem pieces, and rooted runners can simply be cut off and potted up.
Other methods	Layering – pin a runner onto a pot of sand beside the parent and cut it free once rooted.
Confidence	Medium. Method from published advice for the closely related <i>C. glaucescens</i> ; pigface is widely grown from stem pieces.
Sources	See References: 4, 12, 33

Nursery records

Log each batch here to confirm or update the timing and difficulty above.

Date set	Stock source / provenance	No. set	No. struck	Strike %	Notes

Eremophila glabra

Tar Bush / Common Emu Bush

Family
Scrophulariaceae

DIFFICULTY	WOOD TYPE	BEST MONTHS	CUTTING LENGTH	HORMONE (IBA)
Easy	Semi-hardwood	Jan – Apr	75–100 mm	3 g/L IBA

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Cuttings	Best	Best	Best	Best	OK	–	–	–	–	–	OK	OK

Best window
Workable – see 'Selecting material' in the method
Avoid / low success

Plant snapshot

Habit	Very variable, from prostrate to 3 m. Perth coastal plants are mostly subsp. <i>albicans</i> : low, spreading, to about 0.4 m high and 2 m wide, with grey, felty leaves.
Flowering	Species: Mar – Dec. Subsp. <i>albicans</i> : May – Jan. Tubular flowers, yellow, orange, red or green.
Habitat	Sand and limestone; dunes, sandplains and winter-wet flats, sometimes saline.
Local note	Recorded in the Wanneroo LGA and along the Perth coast. Several subspecies and many forms – label stock by subspecies and form.

Propagation options

Method	Suitability	Notes	Details in
Cuttings	Main method	Most forms strike readily from semi-hardwood cuttings.	This guide
Seed	Difficult	Unreliable – hard fruits; slow, erratic germination.	–
Division	Not used	Woody shrub.	–

Propagation method

Item	Details
Best time to take cuttings	Jan – Apr, using hardened current season's growth while the weather is warm and plants are mostly out of flower. Nov – Dec and May are workable. Avoid Jun – Oct (cool weather, flowering).
Wood type	Semi-hardwood – hardened current season's growth.
Cutting size	75–100 mm.
Selecting material	Non-flowering tips or side shoots. On grey-leaved forms pick clean foliage with no sign of mould.
Pre-treatment	- Carefully remove leaves from the lower two-thirds. - Wound: remove a thin sliver of bark from the base. - Dip the base in rooting hormone.
Rooting hormone	Semi-hardwood strength, e.g. Clonex Purple (3 g/L IBA).
Propagation mix	Very free-draining, sand-based mix.
Environment	Grey, felty-leaved forms rot under heavy mist or high humidity – keep foliage as dry as possible, water the mix, give good airflow. Green-leaved forms tolerate normal misting. Bottom heat speeds rooting.
Expected strike time	About 4–8 weeks in warm weather.
Post-treatment / aftercare	- Standard hardening off. - Pot on into a free-draining mix and grow on in full sun. - Keep water off grey foliage; water in the morning.

Common problems

Problem	Likely cause and fix
Leaves blacken or grow grey mould	Too humid. Cut back misting, open the frame, remove affected leaves.
Base rotting	Mix too wet. Water less often and use a sandier mix.
Cuttings flower instead of rooting	Use non-flowering shoots and pinch off buds.

Notes

Difficulty	Easy. Most forms of <i>E. glabra</i> strike readily from cuttings.
Other methods	Seed – unreliable in <i>Eremophila</i> (hard fruits, slow and erratic germination). Grafting is used for difficult eremophilas but isn't needed for this species.
Confidence	High. Published sources agree most forms strike readily from cuttings.
Sources	See References: 1, 4, 12, 13, 20, 21, 22, 30

Nursery records

Log each batch here to confirm or update the timing and difficulty above.

Date set	Stock source / provenance	No. set	No. struck	Strike %	Notes

Grevillea vestita

No widely accepted common name

Family
Proteaceae

DIFFICULTY	WOOD TYPE	BEST MONTHS	CUTTING LENGTH	HORMONE (IBA)
Easy	Semi-hardwood	Dec – Apr	80–120 mm	3 g/L IBA

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Cuttings	Best	Best	Best	Best	OK	–	–	–	–	–	OK	Best

Best window
Workable – see 'Selecting material' in the method
Avoid / low success

Plant snapshot

Habit	Erect, spreading, prickly shrub, usually 0.5–3 m. Hairy branchlets; lobed leaves with sharp-pointed tips.
Flowering	Mainly Jun – Nov (white to cream, often pink-tinged, scented). Odd flowers Jan – Apr.
Habitat	Sandy soils over limestone or granite. Occurs on the Swan Coastal Plain, including the Perth area.
Local note	Two subspecies exist (subsp. <i>vestita</i> and subsp. <i>isopogoides</i>). Label stock plants by subspecies and provenance.

Propagation options

Method	Suitability	Notes	Details in
Cuttings	Main method	Firm current season's growth strikes reliably.	This guide
Seed	Good	Grows reasonably easily from seed; seedlings vary from the parent.	→ Seed Sowing Guide
Division	Possible	Can sucker – lift rooted suckers.	–

Propagation method

Item	Details
Best time to take cuttings	Dec – Apr (late spring to autumn), once the spring flush has firmed and flowering has finished. Nov and May are workable. Avoid Jun – Oct: plants are flowering, new growth is soft and cool weather slows rooting.
Wood type	Semi-hardwood – firm, current season's growth. It should bend about 60° and spring back.
Cutting size	80–120 mm long with 3–5 nodes. Tip or sub-tip pieces; two cuttings can often be taken from one shoot.
Selecting material	Healthy, vigorous, non-flowering shoots. Skip shoots carrying buds, old woody stems and soft new tips.
Pre-treatment	- Wear gloves: leaves are prickly and grevillea sap can irritate skin. - Remove any flowers or buds. - Strip leaves from the lower half to two-thirds (cut, don't tear the bark). - Trim remaining lobed leaves by about one-third. - Wound: remove a thin sliver of bark from the bottom 15–20 mm. - Dip the base in rooting hormone.
Rooting hormone	Semi-hardwood strength, e.g. Clonex Purple (3 g/L IBA) . If cuttings form a heavy callus but no roots, trim the callus back to clean wood and re-set; try a lower strength on the next batch.
Propagation mix	Free-draining, low nutrient: about 75–80% washed coarse river sand + 20–25% coir or peat (or a perlite/coir mix).
Environment	Humid propagation house or frame in bright shade. Bottom heat with the base of the cutting 5–10°C warmer than the air. Light misting only – the hairy foliage can rot if kept constantly wet.

Item	Details
Expected strike time	About 6–8 weeks in warm weather. Slower in the cooler months.
Post-treatment / aftercare	• Harden off: ~2 weeks with cover closed by day and open at night, then ~2 weeks in shade before full sun. • Pot on when roots show at the drainage holes, into a low-phosphorus native mix. • Feed only with a low-phosphorus native fertiliser (Proteaceae are phosphorus-sensitive). • Responds well to pruning – tip-prune young plants for a bushier tube.

Common problems

Problem	Likely cause and fix
Heavy callus, no roots	Trim callus back and re-set. Lower the hormone strength next time.
Base rotting	Mix too wet or material too soft. Use firmer wood and water less often.
Leaf yellowing or scorch after potting on	Check phosphorus in the potting mix and fertiliser.

Notes

Difficulty	Easy. Cuttings from firm current season's growth usually strike reliably.
Other methods	Seed – grows reasonably easily from seed, but seedlings are not identical to the parent. Grafting onto <i>G. robusta</i> is used on the humid east coast; not needed locally.
Confidence	High. Species-specific propagation information is published and consistent across sources.
Sources	See References: 1, 2, 3, 4, 13, 15, 17

Nursery records

Log each batch here to confirm or update the timing and difficulty above.

Date set	Stock source / provenance	No. set	No. struck	Strike %	Notes

Hardenbergia comptoniana

Native Wisteria

Family
Fabaceae

DIFFICULTY	WOOD TYPE	BEST MONTHS	CUTTING LENGTH	HORMONE (IBA)
Easy-Moderate	Semi-hardwood	Dec - Mar	50-75 mm	3 g/L IBA

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Cuttings	Best	Best	Best	OK	-	-	-	-	-	-	OK	Best
	Best window		Workable - see 'Selecting material' in the method									Avoid / low success

Plant snapshot

Habit	Vigorous twining climber or scrambling shrub.
Flowering	Jul - Oct (blue-purple, occasionally white).
Habitat	Sandy soils; coastal limestone, sandplains and dunes.
Local note	Recorded in Joondalup and Wanneroo. More vigorous than <i>H. violacea</i> - can smother small shrubs.

Propagation options

Method	Suitability	Notes	Details in
Cuttings	Good	Firm current season's growth strikes well.	This guide
Seed	Main method	Hot water + smoke water; about 10 days.	→ Seed Sowing Guide
Division	Not used	-	-

Propagation method

Item	Details
Best time to take cuttings	Dec - Mar, once the current season's growth has firmed after spring flowering. Nov and Apr are workable. Avoid May - Oct (cool, then flowering).
Wood type	Semi-hardwood - firm current season's growth (ANPSA).
Cutting size	50-75 mm, the length ANPSA suggests for pea-family cuttings. One runner gives several pieces.
Selecting material	Healthy runners without flowers or pods.
Pre-treatment	- Remove the leaves from the lower two-thirds. - No wounding needed. - Dip the base in rooting hormone.
Rooting hormone	Semi-hardwood strength, 3 g/L IBA (Clonex Purple).
Propagation mix	Standard free-draining cutting mix.
Environment	Humid frame, bright shade, bottom heat.
Expected strike time	Not documented - check from about 4 weeks and record the result.
Post-treatment / aftercare	- Standard hardening off. - Stake or trim runners so tubes don't tangle; pot on promptly.

Common problems

Problem	Likely cause and fix
Cuttings wilt	Material too soft. Use firmer growth; keep humidity up until rooted.
Runners tangle between tubes	Space tubes; trim or stake runners.
Base rotting	Mix too wet. Water less.

Notes

Difficulty	Easy-Moderate. Cuttings strike well. Seed is also easy after hot-water treatment.
Confidence	Medium. ANPSA reports cuttings of firm current season's growth strike well; size follows ANPSA pea-family guidance.
Sources	See References: 4, 13, 43, 44

Nursery records

Log each batch here to confirm or update the timing and difficulty above.

Date set	Stock source / provenance	No. set	No. struck	Strike %	Notes

Hemiandra pungens

Snakebush

Family
Lamiaceae

DIFFICULTY	WOOD TYPE	BEST MONTHS	CUTTING LENGTH	HORMONE (IBA)
Easy	Softwood – semi-hardwood	Oct – Mar	50–75 mm	1.5–3 g/L IBA

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Cuttings	Best	Best	Best	OK	–	–	–	–	OK	Best	Best	Best

Best window	Workable – see 'Selecting material' in the method	Avoid / low success
-------------	---	---------------------

Plant snapshot

Habit	Prostrate to ascending shrub, 0.05–1 m, with sharp, prickly leaves.
Flowering	Throughout the year (white, blue-purple or pink).
Habitat	Sand, clay, loam, gravel, laterite and granite; rock outcrops.
Local note	Recorded in Joondalup and Wanneroo. Resistant to dieback. Wear gloves when collecting.

Propagation options

Method	Suitability	Notes	Details in
Cuttings	Main method	Relatively easy from soft to semi-hardwood tips.	This guide
Seed	Difficult	Rarely available; about 4% germination in nursery tests.	–
Division	Not used	Not documented.	–

Propagation method

Item	Details
Best time to take cuttings	Oct – Mar, in the warm months while plants are growing actively. Sep and Apr are workable. Avoid May – Aug (slow rooting).
Wood type	Softwood to semi-hardwood tips.
Cutting size	50–75 mm tip cuttings.
Selecting material	Plants flower all year – choose tips without open flowers and snip off any buds.
Pre-treatment	- Wear gloves – the leaves are prickly. - Strip leaves from the lower half. - No wounding. - Dip the base in rooting hormone.
Rooting hormone	1.5 g/L IBA (Clonex Green) for soft tips; 3 g/L (Clonex Purple) for firmer ones.
Propagation mix	Very free-draining mix.
Environment	Bright shade and bottom heat, with light misting and good airflow – plants are unreliable in humid conditions.
Expected strike time	Not documented – check from about 3–4 weeks and record the result.
Post-treatment / aftercare	- Harden off promptly; grow on in full sun. - Free-draining mix; don't overwater.

Common problems

Problem	Likely cause and fix
Cuttings rot	Too humid or wet. Mist less and improve airflow.
Cuttings flower	Taken from flowering tips. Snip off buds and choose non-flowering shoots.
Slow in winter	Use bottom heat, or wait for the warm window.

Notes

Difficulty	Easy. ANPSA describes it as relatively easy to propagate from cuttings.
Other methods	Seed – rarely available and germinates poorly (about 4% untreated in Alcoa nursery tests). Grafting onto <i>Westringia fruticosa</i> is used where plants struggle.
Confidence	Medium. ANPSA says it is relatively easy from cuttings; timing and size follow standard practice.
Sources	See References: 4, 32, 45, 46, 47

Nursery records

Log each batch here to confirm or update the timing and difficulty above.

Date set	Stock source / provenance	No. set	No. struck	Strike %	Notes

Hibbertia cuneiformis

Cutleaf Hibbertia

Family
Dilleniaceae

DIFFICULTY	WOOD TYPE				BEST MONTHS		CUTTING LENGTH			HORMONE (IBA)		
Easy-Moderate *	Semi-hardwood				Mar - May		75-100 mm			3 g/L IBA		
	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Cuttings	OK	OK	Best	Best	Best	-	-	-	-	OK	OK	OK
Best window	Workable – see ‘Selecting material’ in the method				Avoid / low success							

Plant snapshot

Habit	Erect or sprawling shrub, 0.5-3 m (usually 1-2 m). Wedge-shaped leaves, toothed near the tip.
Flowering	Jan - Mar or Jun - Nov (yellow).
Habitat	Coastal sand dunes and swampy places near the coast.
Local note	Recorded in the City of Joondalup. One of the more vigorous hibbertias - useful for screening.

Propagation options

Method	Suitability	Notes	Details in
Cuttings	Main method	Commonly propagated by cuttings.	This guide
Seed	Difficult	0% in a nursery test; Hibbertia seed is dormant.	-
Division	Not used	-	-

Propagation method

Item	Details
Best time to take cuttings	Mar - May, between the summer and winter flowering periods, once new growth has firmed. Oct - Feb is workable using non-flowering shoots. Avoid Jun - Sep (cool, and plants are flowering).
Wood type	Semi-hardwood - current season's growth that has begun to firm.
Cutting size	75-100 mm.
Selecting material	Non-flowering shoots from healthy plants. Snip off any buds.
Pre-treatment	- Strip leaves from the lower half. - Reduce large leaves by about one-third. - Light wound on firmer stems. - Dip the base in rooting hormone.
Rooting hormone	Semi-hardwood strength, 3 g/L IBA (Clonex Purple).
Propagation mix	Very free-draining, sand-based mix. Use fresh mix and clean tubes - hibbertias are prone to Phytophthora.
Environment	Humid frame, bright shade, bottom heat. Moist, never wet; good airflow.
Expected strike time	Roughly 4-8 weeks in warm weather (typical for the genus). Record your results.
Post-treatment / aftercare	- Standard hardening off. - Pot on into a free-draining native mix; avoid overwatering. - Tip-prune for a bushier plant.

Common problems

Problem	Likely cause and fix
Cuttings collapse at the base	Too wet, or root rot. Improve drainage, use clean mix, remove affected cuttings.
Cuttings flower instead of rooting	Material taken from flowering shoots. Use non-flowering shoots.
Slow to strike	Cool weather. Use bottom heat or wait for the Mar – May window.

Notes

Difficulty	Easy-Moderate (provisional). Commonly grown from cuttings. Hibbertias usually strike well, but <i>H. hypericoides</i> shows not every WA species does – set extra cuttings until your strike rate is known.
Other methods	Seed – 0% germination in the one Alcoa nursery test; <i>Hibbertia</i> seed has complex dormancy. Layering – check low, sprawling stems for roots.
Confidence	Medium. Commonly propagated from cuttings in the nursery trade; little published detail on timing or strike rate.
Sources	See References: 1, 4, 9, 10, 32, 39, 40

Nursery records

Log each batch here to confirm or update the timing and difficulty above.

Date set	Stock source / provenance	No. set	No. struck	Strike %	Notes

Hibbertia hypericoides

Yellow Buttercups

Family
Dilleniaceae

DIFFICULTY	WOOD TYPE	BEST MONTHS	CUTTING LENGTH	HORMONE (IBA)
Difficult	Semi-hardwood	Dec – Feb	50–75 mm	3 g/L IBA (trial range)

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Cuttings	Best	Best	OK	–	–	–	–	–	–	–	OK	Best
 Best window	 Workable – see 'Selecting material' in the method						 Avoid / low success					

Plant snapshot

Habit	Spreading shrub, 0.3–0.8 m. Densely hairy branches; small leaves with rolled edges and white-hairy undersides.
Flowering	Apr – Nov (bright yellow, notched petals).
Habitat	Sandy soils in Banksia woodland and shrubland; one of the most common understorey plants between Perth and Albany. Resprouts from its rootstock after fire.
Local note	Recorded in the City of Joondalup.

Propagation options

Method	Suitability	Notes	Details in
Cuttings	Difficult	Known to be difficult; trial protocol on the cuttings page.	This guide
Seed	Difficult	Complex seed dormancy; rarely practical.	–
Division	Not used	Resprouts from a rootstock; not practical to divide.	–

Propagation method

Item	Details
Best time to take cuttings	Dec – Feb , after flowering finishes and once the new flush has started to firm. Nov and Mar are workable. Avoid Apr – Oct (flowering period).
Wood type	Semi-hardwood – new growth that is just firming. Soft tips wilt quickly; old wood rarely roots.
Cutting size	50–75 mm.
Selecting material	Vigorous non-flowering shoots. Well-watered, actively growing stock plants (e.g. nursery-held stock) usually give better material than stressed bushland plants.
Pre-treatment	- Strip lower leaves by cutting, not pulling. - Light wound on the firmer stems. - Dip the base in rooting hormone.
Rooting hormone	Start with 3 g/L IBA (Clonex Purple). Because results are unreliable, split trial batches across 1.5, 3 and 8 g/L (Green, Purple, Red) and record which works best.
Propagation mix	Very free-draining, fresh, clean mix. Strict hygiene – hibbertias are prone to <i>Phytophthora</i> root rot.
Environment	Humid frame, bottom heat, bright shade. Mist lightly – the hairy leaf undersides hold water. Good airflow.
Expected strike time	Slow and variable – allow 8–16 weeks . Leave cuttings in as long as they stay green and healthy.

Item	Details
Post-treatment / aftercare	• Harden off slowly (extend each stage if cuttings wilt). • Pot on gently, disturbing roots as little as possible. • Avoid overwatering; keep benches clean.

Common problems

Problem	Likely cause and fix
Cuttings stay green but don't root	Leave longer; check for heavy callus and trim it; try a different hormone strength.
Cuttings collapse or rot	Too wet or root rot. Improve drainage and hygiene; remove affected cuttings.
Rooted cuttings die after potting on	Roots damaged or hardening too fast. Use single tubes and harden off more slowly.

Notes

Difficulty	Difficult. DBCA notes it is difficult to grow from seed or cuttings, and specialist Perth nurseries describe it as a species other producers often fail with. Set large batches and keep detailed records.
Other methods	Seed – also difficult: WA <i>Hibbertia</i> seed has complex dormancy, and treatments are still the subject of research. If numbers are needed quickly, consider buying in tubestock from a specialist grower.
Confidence	Low. Its difficulty is well documented (DBCA; specialist WA nurseries), but no detailed cutting protocol is published. Treat the method below as a trial protocol.
Sources	See References: 1, 4, 10, 11, 27, 30, 31

Nursery records

Log each batch here to confirm or update the timing and difficulty above.

Date set	Stock source / provenance	No. set	No. struck	Strike %	Notes

Hibbertia racemosa

Stalked Guinea Flower

Family
Dilleniaceae

DIFFICULTY	WOOD TYPE				BEST MONTHS			CUTTING LENGTH			HORMONE (IBA)		
Moderate *	Semi-hardwood				Jan - Apr			50-75 mm			1.5-3 g/L IBA		
	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	
Cuttings	Best	Best	Best	Best	OK	-	-	-	-	-	-	OK	
	Best window				Workable - see 'Selecting material' in the method				Avoid / low success				

Plant snapshot

Habit	Small erect or ascending, spreading shrub, 0.1–0.75 m.
Flowering	Jul – Dec (yellow, on long stalks).
Habitat	Grey, white or yellow sand in coastal areas: dunes, plains and limestone.
Local note	Recorded in the City of Joondalup. Grey-leaved and prostrate forms are sold in the trade – keep forms separate and labelled.

Propagation options

Method	Suitability	Notes	Details in
Cuttings	Main method	Semi-hardwood cuttings (provisional rating).	This guide
Seed	Difficult	Hibbertia seed is dormant and unreliable.	–
Division	Possible	Low stems may root where they touch the soil (layering).	–

Propagation method

Item	Details
Best time to take cuttings	Jan – Apr, after flowering finishes and new growth has begun to firm. Dec and May are workable. Avoid Jun – Nov (cool weather, then flowering).
Wood type	Semi-hardwood – new growth that has started to harden. Firm tip cuttings also work.
Cutting size	50–75 mm long. Stems are fine and leaves small, so keep cuttings short.
Selecting material	Non-flowering side shoots from healthy plants. Snip off any flower stalks or buds.
Pre-treatment	- Strip leaves from the lower half (cut rather than pull on fine stems). - Wounding not needed on fine stems; a light wound is optional on firmer material. - Dip the base in rooting hormone.
Rooting hormone	Semi-hardwood strength (3 g/L IBA, e.g. Clonex Purple). For softer tips use softwood strength (1.5 g/L IBA, e.g. Clonex Green).
Propagation mix	Very free-draining, sand-based mix. Use fresh, clean mix and clean tubes – hibbertias are prone to <i>Phytophthora</i> root rot.
Environment	Humid frame, bright shade, bottom heat. Keep the mix moist but never wet; good airflow.
Expected strike time	Roughly 4–8 weeks in warm weather; may be slower. Record your results.
Post-treatment / aftercare	- Standard hardening off (2 weeks closed by day/open at night, then 2 weeks shade). - Pot on into a free-draining native mix; avoid overwatering. - Keep good hygiene on benches to limit root rot.

Common problems

Problem	Likely cause and fix
Cuttings collapse at the base	Too wet, or root rot. Improve drainage, use clean mix, remove affected cuttings.
Few roots, cuttings flower	Material was taken from flowering shoots. Use non-flowering side shoots.
Slow to strike in winter	Use bottom heat, or wait for the Jan – Apr window.

Notes

Difficulty	Moderate (provisional). Hibbertias are generally easy from cuttings, but some WA species (e.g. <i>H. hypericoides</i>) are known to be difficult. Set larger batches until your own strike rate is known.
Other methods	Seed – generally unreliable in <i>Hibbertia</i> (complex dormancy). Layering – several hibbertias root where stems touch the ground; check low stems on prostrate forms and pot up any rooted pieces.
Confidence	Medium. Little species-specific data published. Based on how the genus behaves and this species' growth cycle.
Sources	See References: 1, 4, 8, 9, 10, 11

Nursery records

Log each batch here to confirm or update the timing and difficulty above.

Date set	Stock source / provenance	No. set	No. struck	Strike %	Notes

Kennedia prostrata

Scarlet Runner / Running Postman

Family
Fabaceae

DIFFICULTY	WOOD TYPE	BEST MONTHS	CUTTING LENGTH	HORMONE (IBA)
Easy-Moderate	Semi-hardwood	Dec - Mar	75-100 mm	3 g/L IBA

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Cuttings	Best	Best	Best	OK	-	-	-	-	-	-	OK	Best
	Best window		Workable - see 'Selecting material' in the method									Avoid / low success

Plant snapshot

Habit	Prostrate or twining groundcover, spreading to about 2.5 m.
Flowering	Apr - Nov (red pea flowers).
Habitat	Usually sandy, gravelly soils.
Local note	Recorded in Joondalup and Wanneroo.

Propagation options

Method	Suitability	Notes	Details in
Cuttings	Good	Firm current season's growth strikes well.	This guide
Seed	Main method	Hot water + smoke water; about 8 days.	→ Seed Sowing Guide
Division	Not used	-	-

Propagation method

Item	Details
Best time to take cuttings	Dec - Mar, when the current season's runners have firmed. Nov and Apr are workable. Avoid May - Oct (flowering and cool weather).
Wood type	Semi-hardwood - firm current season's growth (ANPSA).
Cutting size	75-100 mm with 2-3 nodes. One runner can be cut into several pieces.
Selecting material	Healthy runners without flowers or pods.
Pre-treatment	- Remove the leaves from the lower nodes; cut remaining leaflets back by about half. - No wounding needed. - Dip the base in rooting hormone.
Rooting hormone	Semi-hardwood strength, 3 g/L IBA (Clonex Purple).
Propagation mix	Standard free-draining cutting mix.
Environment	Humid frame, bright shade, bottom heat.
Expected strike time	Not documented - check from about 4 weeks and record the result.
Post-treatment / aftercare	- Standard hardening off. - Pot on promptly - runners tangle across the bench. Trim runners to keep tubes tidy.

Common problems

Problem	Likely cause and fix
Cuttings wilt and collapse	Material too soft. Use firmer runners and keep humidity up.
Runners tangled between tubes	Space tubes out and trim runners regularly.
Base rotting	Mix too wet. Water less, improve drainage.

Notes

Difficulty	Easy-Moderate. Cuttings of firm growth strike well. Seed is quick and is the usual nursery method.
Confidence	Medium. ANPSA reports cuttings of firm current season's growth strike well; timing and size follow standard practice.
Sources	See References: 4, 13, 42

Nursery records

Log each batch here to confirm or update the timing and difficulty above.

Date set	Stock source / provenance	No. set	No. struck	Strike %	Notes

Lechenaultia linarioides

Yellow Leschenaultia

Family
Goodeniaceae

DIFFICULTY	WOOD TYPE	BEST MONTHS	CUTTING LENGTH	HORMONE (IBA)
Easy *	Softwood - semi-hardwood	Feb - Apr	50-75 mm	1.5-3 g/L IBA

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Cuttings	OK	Best	Best	Best	OK	-	-	-	-	OK	OK	OK
	Best window	Workable - see 'Selecting material' in the method										

Plant snapshot

Habit	Sprawling, open shrub to 1.5 m, with narrow, linear leaves.
Flowering	Can flower in any month. Flowers yellow with red or pink tones, sometimes cream.
Habitat	Sand (often over limestone) and sandy clay; sandplains, low dunes, ridges and winter-wet flats.
Local note	Recorded in the Wanneroo LGA.

Propagation options

Method	Suitability	Notes	Details in
Cuttings	Main method	Cuttings strike easily in this genus.	This guide
Seed	Difficult	Slow; trays may need a second season.	-
Division	Possible	Some leschenaultias sucker - pot up rooted suckers.	-

Propagation method

Item	Details
Best time to take cuttings	Feb - Apr (late summer to autumn) while it is warm. One grower reports roots in 4-6 weeks from late-February leschenaultia cuttings. Oct - Jan and May are workable. Avoid Jun - Sep.
Wood type	Softwood to semi-hardwood tip cuttings.
Cutting size	50-75 mm.
Selecting material	Non-flowering tips. This species can flower in any month, so check each cutting and remove buds.
Pre-treatment	- Strip leaves from the lower half (narrow leaves strip easily). - No wounding needed. - Dip the base in rooting hormone.
Rooting hormone	1.5 g/L IBA (Clonex Green) for soft tips; 3 g/L (Clonex Purple) for firmer material.
Propagation mix	Very free-draining - leschenaultias dislike wet roots.
Environment	Humid frame, bottom heat, bright shade. Keep the mix moist, never wet.
Expected strike time	About 4-6 weeks in warm weather.
Post-treatment / aftercare	- Standard hardening off. - Pot on into a free-draining mix; water sparingly. - Tip-prune to counter the sprawling habit.

Common problems

Problem	Likely cause and fix
Base rotting	Overwatering. Water less often; add coarse sand or perlite to the mix.
Wilting in the frame	Material too soft or humidity too low. Use slightly firmer tips; keep the frame closed.
Cuttings flowering	Remove buds; pick vegetative shoots.

Notes

Difficulty	Easy (provisional). <i>Lechenaultia</i> cuttings generally strike very easily; seed is difficult.
Other methods	Seed – slow and unreliable. The Geraldton community nursery soaked seed in smoke water, saw germination at about 46 days, and kept trays for a second year. Division – some <i>leschenaultias</i> sucker; pot up rooted suckers.
Confidence	Medium. No species-specific cutting data found. Based on the genus, whose cuttings strike very easily.
Sources	See References: 1, 4, 12, 13, 24, 25

Nursery records

Log each batch here to confirm or update the timing and difficulty above.

Date set	Stock source / provenance	No. set	No. struck	Strike %	Notes

Leucophyta brownii

Cushion Bush

Family
Asteraceae

DIFFICULTY	WOOD TYPE	BEST MONTHS	CUTTING LENGTH	HORMONE (IBA)
Easy-Moderate *	Semi-hardwood	Mar – May; Sep – Oct	50–75 mm	1.5–3 g/L IBA

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Cuttings	–	OK	Best	Best	Best	OK	–	OK	Best	Best	OK	–
	Best window	Workable – see 'Selecting material' in the method										Avoid / low success

Plant snapshot

Habit	Compact, rounded shrub to about 1 m, densely covered in silvery-white woolly hairs.
Flowering	Dec or Jan – Feb (small yellow button heads).
Habitat	Coastal foredunes and cliffs.
Local note	Recorded in the Wanneroo LGA. Very salt-tolerant; grows poorly in high humidity.

Propagation options

Method	Suitability	Notes	Details in
Cuttings	Main method	Semi-hardened cuttings are the preferred method.	This guide
Seed	Good	No pre-treatment; 10–30 days.	→ Seed Sowing Guide
Division	Not used	–	–

Propagation method

Item	Details
Best time to take cuttings	Mar – May, after flowering, and Sep – Oct as spring growth firms. Feb, Jun, Aug and Nov are workable. Avoid Dec – Jan (flowering and heat) and Jul (cold).
Wood type	Semi-hardwood – ANBG notes cuttings should come from semi-hardened branches.
Cutting size	50–75 mm tip cuttings.
Selecting material	Non-flowering tips from compact, healthy plants. Avoid soft, sappy growth.
Pre-treatment	• Rub the small leaves off the lower half. • No wounding. • Dip the base in rooting hormone.
Rooting hormone	1.5 g/L IBA (Clonex Green) for firm tips; 3 g/L for firmer material.
Propagation mix	Very free-draining – add extra coarse sand or perlite.
Environment	Keep it airy. The woolly foliage rots in humidity: use a well-ventilated frame in bright light with little or no mist. Bottom heat helps. Keep the mix only just moist.
Expected strike time	Not documented – check from about 4 weeks and record the result.
Post-treatment / aftercare	• Harden off promptly and grow on in full sun. • Free-draining mix; never let pots sit wet – plants die easily if waterlogged. • Tip-prune for a dense cushion.

Common problems

Problem	Likely cause and fix
Cuttings turn grey-brown and mushy	Too humid or wet. Reduce mist, open the frame, improve airflow.
Slow to root in winter	Use bottom heat, or wait for the autumn or spring window.
Loses its dense shape	Tip-prune young plants.

Notes

Difficulty	Easy-Moderate (provisional). Cuttings are the preferred method; the main risk is rot from too much moisture or humidity.
Confidence	Medium. ANBG growing notes say semi-hardened cuttings are the preferred method; timing and details follow general daisy practice.
Sources	See References: 4, 34, 35

Nursery records

Log each batch here to confirm or update the timing and difficulty above.

Date set	Stock source / provenance	No. set	No. struck	Strike %	Notes

Melaleuca huegelii

Chenille Honeymyrtle

Family
Myrtaceae

DIFFICULTY Easy *	WOOD TYPE Semi-hardwood	BEST MONTHS Feb – Apr	CUTTING LENGTH 75–100 mm	HORMONE (IBA) 3 g/L IBA
----------------------	----------------------------	--------------------------	-----------------------------	----------------------------

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Cuttings	OK	Best	Best	Best	OK	–	–	–	–	–	–	OK
	Best window	Workable – see ‘Selecting material’ in the method							Avoid / low success			

Plant snapshot

Habit	Shrub or small tree, 0.5–5 m.
Flowering	Sep – Dec or Jan (pink, white or pink-purple).
Habitat	Sand; limestone cliffs, coastal plains and dunes.
Local note	Recorded in Joondalup and Wanneroo. Cuttings keep a chosen flower colour.

Propagation options

Method	Suitability	Notes	Details in
Cuttings	Good	Reliable in this genus; keeps a chosen form.	This guide
Seed	Main method	Surface-sow fine seed; no pre-treatment.	→ Seed Sowing Guide
Division	Not used	Woody shrub.	–

Propagation method

Item	Details
Best time to take cuttings	Feb – Apr, after flowering, once new growth has firmed. Jan, May and Dec are workable. Avoid Jun – Nov.
Wood type	Semi-hardwood – firm current season's growth.
Cutting size	75–100 mm (ANPSA).
Selecting material	Non-flowering tips. Take from a plant with the flower colour you want.
Pre-treatment	- Strip leaves from the lower half to two-thirds. - Wound the base on firmer stems. - Dip the base in rooting hormone.
Rooting hormone	Semi-hardwood strength, 3 g/L IBA (Clonex Purple).
Propagation mix	Standard free-draining cutting mix.
Environment	Humid frame, bright shade, bottom heat.
Expected strike time	Typically 4–8 weeks in warm weather. Record your results.
Post-treatment / aftercare	- Standard hardening off. - Pot on into a free-draining native mix; tip-prune for bushiness.

Common problems

Problem	Likely cause and fix
Callus, no roots	Trim and re-set; lower the hormone strength next batch.
Leaves drop in the frame	Too dry or too hot. Shade more and keep humidity steady.
Base rotting	Mix too wet. Water less.

Notes

Difficulty	Easy (provisional). Melaleucas are generally reliable from cuttings.
Confidence	Medium. Based on ANPSA <i>Melaleuca</i> genus guidance – cuttings are reliable in the genus.
Sources	See References: 4, 13, 41

Nursery records

Log each batch here to confirm or update the timing and difficulty above.

Date set	Stock source / provenance	No. set	No. struck	Strike %	Notes

Melaleuca systema

Coastal Honeymyrtle

Family
Myrtaceae

DIFFICULTY	WOOD TYPE	BEST MONTHS	CUTTING LENGTH	HORMONE (IBA)
Easy *	Semi-hardwood	Dec – Jan; Apr – May	75–100 mm	3 g/L IBA

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Cuttings	Best	OK	OK	Best	Best	OK	–	–	–	–	OK	Best

Best window

Workable – see ‘Selecting material’ in the method

Avoid / low success

Plant snapshot

Habit	Erect to spreading shrub, usually 0.5–2 m.
Flowering	Feb – Mar or Aug – Dec (yellow-cream).
Habitat	Sand over laterite or limestone; stabilised coastal dunes and rocky limestone.
Local note	Recorded in Joondalup and Wanneroo. Seed germination is low (about 14% in nursery tests), so cuttings are worth using.

Propagation options

Method	Suitability	Notes	Details in
Cuttings	Good	Reliable in this genus; a good option given low seed germination.	This guide
Seed	Main method	Surface-sow fine seed; no pre-treatment.	→ Seed Sowing Guide
Division	Not used	Woody shrub.	–

Propagation method

Item	Details
Best time to take cuttings	Dec – Jan and Apr – May – the gaps between its two flowering periods. Feb – Mar, Jun and Nov are workable with non-flowering shoots. Avoid Jul – Oct.
Wood type	Semi-hardwood – firm current season's growth.
Cutting size	75–100 mm (ANPSA).
Selecting material	Non-flowering tips from healthy plants.
Pre-treatment	- Strip leaves from the lower half to two-thirds. - Wound the base on firmer stems. - Dip the base in rooting hormone.
Rooting hormone	Semi-hardwood strength, 3 g/L IBA (Clonex Purple).
Propagation mix	Standard free-draining cutting mix.
Environment	Humid frame, bright shade, bottom heat. Autumn batches root more slowly.
Expected strike time	Typically 4–8 weeks in warm weather; longer in May. Record your results.
Post-treatment / aftercare	- Standard hardening off. - Pot on into a free-draining native mix.

Common problems

Problem	Likely cause and fix
Callus, no roots	Trim and re-set; lower the hormone strength next batch.
Cuttings flower	Taken from flowering shoots. Use non-flowering tips, between flowering periods.
Base rotting	Mix too wet. Water less.

Notes

Difficulty	Easy (provisional). Melaleucas are generally reliable from cuttings.
Confidence	Medium. Based on ANPSA <i>Melaleuca</i> genus guidance – cuttings are reliable in the genus.
Sources	See References: 4, 13, 41

Nursery records

Log each batch here to confirm or update the timing and difficulty above.

Date set	Stock source / provenance	No. set	No. struck	Strike %	Notes

Myoporum insulare

Blueberry Tree / Common Boobialla

Family
Scrophulariaceae

DIFFICULTY		WOOD TYPE				BEST MONTHS		CUTTING LENGTH			HORMONE (IBA)		
Easy		Semi-hardwood				Feb – Apr		75–100 mm			3 g/L IBA		
		Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Cuttings		OK	Best	Best	Best	OK	–	–	–	–	OK	OK	OK
Best window		Workable – see ‘Selecting material’ in the method						Avoid / low success					

Plant snapshot

Habit	Dense, low spreading to erect shrub or small tree, 0.25–5 m, with thick, fleshy leaves.
Flowering	Jul – Dec, also Jan – Feb (white, purple-spotted), followed by purple berries.
Habitat	White or grey sand on coastal dunes and limestone.
Local note	Recorded in Joondalup and Wanneroo. Forms range from prostrate to tree-like – take cuttings from a plant with the form you want.

Propagation options

Method	Suitability	Notes	Details in
Cuttings	Main method	Firm young growth roots readily.	This guide
Seed	Good	Ferment berries, smoke water; 15–30 days. Seedling form varies.	→ Seed Sowing Guide
Division	Not used	Woody shrub.	–

Propagation method

Item	Details
Best time to take cuttings	Feb – Apr – warm, and after the main flowering. Oct – Jan and May are workable. Avoid Jun – Sep.
Wood type	Semi-hardwood – firm young growth.
Cutting size	75–100 mm.
Selecting material	Firm, non-flowering young shoots from a stock plant with the growth form you need (prostrate for groundcover, erect for screening).
Pre-treatment	- Strip leaves from the lower half. - Reduce the remaining thick leaves by about one-third. - Dip the base in rooting hormone.
Rooting hormone	3 g/L IBA (Clonex Purple). Strikes readily without it, but hormone gives faster, more even rooting.
Propagation mix	Standard sand/coir cutting mix.
Environment	Standard humid frame with bottom heat. Hairless leaves tolerate normal misting.
Expected strike time	About 4–8 weeks in warm weather.
Post-treatment / aftercare	- Standard hardening off. - Fast-growing – pot on promptly before roots circle. - Tip-prune for a bushier plant.

Common problems

Problem	Likely cause and fix
Leaf drop in the frame	Too dark or too wet. Give brighter shade and water less.
Roots circling in tubes	Pot on sooner – it grows quickly.
Plants not the expected shape	Seed-grown or mixed stock. Propagate from a known-form stock plant.

Notes

Difficulty	Easy. Firm young growth roots readily; even branchlets placed straight into soil can strike.
Other methods	Seed – the Geraldton community nursery ferments ripe (darkest purple) berries for 4–5 days, washes off the pulp and soaks in smoke water; germination took 15–30 days. Seedlings vary in form.
Confidence	High. Published sources agree it strikes readily from firm young growth.
Sources	See References: 1, 4, 12, 13, 23

Nursery records

Log each batch here to confirm or update the timing and difficulty above.

Date set	Stock source / provenance	No. set	No. struck	Strike %	Notes

Olearia axillaris

Coastal Daisybush

Family
Asteraceae

DIFFICULTY Easy *	WOOD TYPE Semi-hardwood	BEST MONTHS Aug – Oct	CUTTING LENGTH 75–100 mm	HORMONE (IBA) 1.5–3 g/L IBA
----------------------	----------------------------	--------------------------	-----------------------------	--------------------------------

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Cuttings	–	–	OK	OK	–	–	OK	Best	Best	Best	OK	–
Best window												
Workable – see ‘Selecting material’ in the method												
Avoid / low success												

Plant snapshot

Habit	Erect, much-branched grey shrub, 0.2–3 m.
Flowering	Jan – Jul or Nov – Dec (small white to yellow daisies).
Habitat	White, grey or red sand and loam; coastal limestone, sand dunes and rocky hillsides.
Local note	Recorded in Joondalup and Wanneroo.

Propagation options

Method	Suitability	Notes	Details in
Cuttings	Good	Perennial daisies strike quickly from vigorous growth.	This guide
Seed	Main method	Fresh seed, aired 3 months; surface-sow.	→ Seed Sowing Guide
Division	Not used	Woody shrub.	–

Propagation method

Item	Details
Best time to take cuttings	Aug – Oct, on vigorous spring growth between flowering periods. Mar – Apr, Jul and Nov are workable with non-flowering shoots. Avoid Dec – Feb and May – Jun.
Wood type	Semi-hardwood – vigorous current growth that has begun to firm.
Cutting size	75–100 mm.
Selecting material	Vigorous, healthy, non-flowering shoots (ANPSA).
Pre-treatment	- Strip leaves from the lower half. - No wounding needed. - Dip the base in rooting hormone.
Rooting hormone	1.5 g/L IBA (Clonex Green) for softer tips; 3 g/L for firmer material.
Propagation mix	Free-draining: about 3 parts sand to 1 part peat or coir (ANPSA daisy mix).
Environment	Humid frame, bright shade, bottom heat. The grey, hairy leaves rot if kept wet – mist lightly.
Expected strike time	Usually quick – check from about 3–4 weeks.
Post-treatment / aftercare	- Standard hardening off. - Grow on in full sun; tip-prune for a compact plant.

Common problems

Problem	Likely cause and fix
Leaves go grey-brown and soft	Too wet or humid. Mist less, improve airflow.
Cuttings flower	Taken from flowering shoots. Use vigorous non-flowering growth.
Leggy plants	Too much shade. Move to full sun after hardening off.

Notes

Difficulty	Easy (provisional). Perennial daisies usually strike quickly and easily.
Confidence	Medium. ANPSA: stem cuttings of most perennial Australian daisies strike quickly from vigorous growth.
Sources	See References: 4, 13, 35

Nursery records

Log each batch here to confirm or update the timing and difficulty above.

Date set	Stock source / provenance	No. set	No. struck	Strike %	Notes

Phyllanthus calycinus

False Boronia

Current WA Herbarium name: *Lysiandra calycina*Family
Phyllanthaceae

DIFFICULTY Moderate *	WOOD TYPE Semi-hardwood	BEST MONTHS Feb – Apr	CUTTING LENGTH 50–75 mm	HORMONE (IBA) 3 g/L IBA
---------------------------------	-----------------------------------	---------------------------------	-----------------------------------	-----------------------------------

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Cuttings	OK	Best	Best	Best	OK	–	–	–	–	–	–	–
	Best window	Workable – see 'Selecting material' in the method										

Plant snapshot

Habit	Small erect shrub, 0.2–1.2 m, with smooth branchlets.
Flowering	Jun – Dec or Jan (white-cream to pink). Sepals enlarge to enclose the fruit.
Habitat	Usually sandy soils, including coastal alkaline sands.
Name note	FloraBase now lists this plant as <i>Lysiandra calycina</i> ; many nurseries still sell it as <i>Phyllanthus calycinus</i> . Kept under <i>Phyllanthus</i> here to match your plant list.
Local note	Recorded in Joondalup and Wanneroo.

Propagation options

Method	Suitability	Notes	Details in
Cuttings	Trial	No published cutting data; trial protocol.	This guide
Seed	Possible	Fresh seed is recommended; little published detail.	–
Division	Not used	Woody shrub.	–

Propagation method

Item	Details
Best time to take cuttings	Feb – Apr – warm, and after the long flowering season ends. Jan and May are workable. Avoid Jun – Dec (flowering, and cool in winter).
Wood type	Semi-hardwood.
Cutting size	50–75 mm.
Selecting material	Non-flowering shoots from healthy plants.
Pre-treatment	- Strip leaves from the lower half. - Light wound on firmer stems. - Dip the base in rooting hormone.
Rooting hormone	3 g/L IBA (Clonex Purple); trial 1.5 g/L (Clonex Green) on softer tips.
Propagation mix	Free-draining sand-based mix.
Environment	Standard humid frame, bottom heat, bright shade.
Expected strike time	Not documented – start checking from about 6 weeks and record the result.
Post-treatment / aftercare	- Standard hardening off. - Pot on into a free-draining native mix.

Common problems

Problem	Likely cause and fix
No roots after 10+ weeks	Try a different hormone strength or timing on the next batch; record what you tried.
Rotting	Mix too wet. Water less often; improve drainage.
Cuttings flowering	Use non-flowering shoots only.

Notes

Difficulty	Moderate (provisional). No cutting data found for this species; the rating is a cautious starting point.
Other methods	Seed – the method most guides mention; use fresh seed. Few published details on pre-treatment.
Confidence	Low. No published cutting data found. Treat as a trial protocol and record results.
Sources	See References: 1, 4, 29, 30

Nursery records

Log each batch here to confirm or update the timing and difficulty above.

Date set	Stock source / provenance	No. set	No. struck	Strike %	Notes

Pithocarpa cordata

Tangle Daisy

Previously *Ozothamnus cordatus* / *Helichrysum cordatum*Family
Asteraceae

DIFFICULTY	WOOD TYPE	BEST MONTHS	CUTTING LENGTH	HORMONE (IBA)
Easy-Moderate *	Semi-hardwood tips	Sep – Nov; May	50–75 mm	1.5–3 g/L IBA

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Cuttings	–	–	–	OK	Best	OK	–	OK	Best	Best	Best	OK
	Best window	Workable – see 'Selecting material' in the method										

Plant snapshot

Habit	Scrambling, tangled shrub, 1–2 m, with small heart-shaped leaves.
Flowering	Jan – Apr (white to yellow).
Habitat	Coastal limestone and dunes.
Local note	Recorded in the Wanneroo LGA. Older books list it as <i>Ozothamnus cordatus</i> or <i>Helichrysum cordatum</i> .

Propagation options

Method	Suitability	Notes	Details in
Cuttings	Good	Perennial daisy – expected to strike well (provisional).	This guide
Seed	Possible	Not documented; daisy seed often needs light.	–
Division	Not used	–	–

Propagation method

Item	Details
Best time to take cuttings	Sep – Nov on vigorous spring growth, and May after flowering. Apr, Jun, Aug and Dec are workable. Avoid Jan – Mar (flowering and heat) and Jul (cold).
Wood type	Semi-hardwood tips – vigorous growth that has started to firm.
Cutting size	50–75 mm.
Selecting material	Vigorous, healthy, non-flowering shoots.
Pre-treatment	- Strip leaves from the lower half. - No wounding. - Dip the base in rooting hormone.
Rooting hormone	1.5 g/L IBA (Clonex Green); 3 g/L for firmer stems.
Propagation mix	Free-draining: about 3 parts sand to 1 part peat or coir.
Environment	Humid frame, bright shade, bottom heat. Moist, not wet.
Expected strike time	Not documented – check from about 3–4 weeks and record the result.
Post-treatment / aftercare	- Standard hardening off. - Grow on in full sun; tip-prune to control the scrambling habit.

Common problems

Problem	Likely cause and fix
Cuttings rot	Too wet or humid. Mist less, improve airflow.
No roots after 8 weeks	Try firmer tips, a stronger hormone, or a different season, and record the result.
Tangled, sprawling tubes	Tip-prune and space tubes.

Notes

Difficulty	Easy-Moderate (provisional). Most perennial daisies strike quickly from vigorous growth, but this species is untested. Start with a trial batch.
Other methods	Seed – not documented. Daisy-family seed often needs light, so surface-sow if trialling.
Confidence	Low. No published propagation data. Based on perennial Australian daisies, which usually strike quickly – trial protocol.
Sources	See References: 4, 35

Nursery records

Log each batch here to confirm or update the timing and difficulty above.

Date set	Stock source / provenance	No. set	No. struck	Strike %	Notes

Rhagodia baccata

Berry Saltbush

Also listed as *Chenopodium baccatum*

Family
Amaranthaceae
(chenopods)

DIFFICULTY	WOOD TYPE	BEST MONTHS	CUTTING LENGTH	HORMONE (IBA)
Moderate *	Semi-hardwood tips	Sep – Nov	75–100 mm	3 g/L IBA

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Cuttings	OK	OK	OK	–	–	–	–	OK	Best	Best	Best	OK
	Best window	Workable – see ‘Selecting material’ in the method							Avoid / low success			

Plant snapshot

Habit	Spreading shrub, 0.3–2 m, with grey-green leaves.
Flowering	Variable – Feb, Apr – May or Oct – Dec (tiny), followed by red berries eaten by birds and bobtails.
Habitat	Sand dunes and coastal rock.
Local note	Recorded in Joondalup and Wanneroo. Salt-tolerant; a breeding plant for the saltbush blue butterfly.

Propagation options

Method	Suitability	Notes	Details in
Cuttings	Good	Chenopod shrubs generally strike well (provisional).	This guide
Seed	Main method	Ferment berries, then smoke water.	→ Seed Sowing Guide
Division	Not used	Woody shrub.	–

Propagation method

Item	Details
Best time to take cuttings	Sep – Nov, in active spring growth. Aug and Dec – Mar are workable with non-flowering shoots. Avoid Apr – Jul.
Wood type	Semi-hardwood tips of current season's growth.
Cutting size	75–100 mm.
Selecting material	Vigorous tips without flowers or berries.
Pre-treatment	- Strip leaves from the lower half. - Light wound on firmer stems. - Dip the base in rooting hormone.
Rooting hormone	Semi-hardwood strength, 3 g/L IBA (Clonex Purple).
Propagation mix	Free-draining mix of sand and potting mix.
Environment	Bright shade with good ventilation and little or no mist (misting encourages leaf disease on saltbush cuttings). Bottom heat helps.
Expected strike time	Not documented – check from about 4–6 weeks and record the result.
Post-treatment / aftercare	- Harden off, then grow on in full sun. - Water sparingly once established.

Common problems

Problem	Likely cause and fix
Leaf spots, leaves dropping	Too humid. Stop misting and improve airflow.
Base rotting	Mix too wet. Use a sandier mix, water less.
No roots	Try firmer material or the seed method, and record the result.

Notes

Difficulty	Moderate (provisional). Untested here; related saltbushes strike from tip cuttings. Seed is the documented method.
Confidence	Low. No published cutting data found. Method follows related saltbushes – trial protocol.
Sources	See References: 4, 20, 36, 37

Nursery records

Log each batch here to confirm or update the timing and difficulty above.

Date set	Stock source / provenance	No. set	No. struck	Strike %	Notes

Scaevola canescens

Grey Scaevola

Family
Goodeniaceae

DIFFICULTY	WOOD TYPE	BEST MONTHS	CUTTING LENGTH	HORMONE (IBA)
Easy-Moderate *	Softwood – semi-hardwood	Nov – Jan	50–80 mm	1.5–3 g/L IBA

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Cuttings	Best	OK	OK	OK	OK	–	–	–	–	OK	Best	Best

Best window	Workable – see 'Selecting material' in the method	Avoid / low success
-------------	---	---------------------

Plant snapshot

Habit	Prostrate to ascending shrub, 0.1–0.5 m. Densely hairy grey leaves; stems are brittle.
Flowering	Mar – Oct (white to white-purple), in spikes in the leaf axils.
Habitat	Sand, often with lateritic gravel or limestone.
Local note	Recorded in the northern Perth area (Wanneroo and Stirling LGAs).

Propagation options

Method	Suitability	Notes	Details in
Cuttings	Main method	Soft to semi-hardwood tips; keep hairy foliage dry.	This guide
Seed	Difficult	Scaevola seed is generally difficult.	–
Division	Possible	Only if stems have rooted or suckered.	–

Propagation method

Item	Details
Best time to take cuttings	Nov – Jan, on new growth after flowering ends, while it is warm. Oct and Feb – May are workable if you choose shoots without flower spikes. Avoid Jun – Sep (cool, peak flowering).
Wood type	Softwood to semi-hardwood tip cuttings.
Cutting size	50–80 mm tips.
Selecting material	Vigorous non-flowering tips. Flowers form in the leaf axils along the stem, so check the whole cutting and pinch off any spikes.
Pre-treatment	- Handle gently – stems snap easily. Carry in a rigid tray or box. - Strip lower leaves by cutting, not tearing. - Dip the base in rooting hormone.
Rooting hormone	Softwood strength (1.5 g/L IBA, e.g. Clonex Green) for soft tips; semi-hardwood strength (3 g/L, Clonex Purple) for firmer material.
Propagation mix	Extra free-draining: standard sand/coir mix with extra coarse sand or perlite.
Environment	Keep foliage as dry as possible. Dense hairy leaves rot under heavy mist. Water the mix rather than the leaves, give good airflow, and use bottom heat (root zone about 21–24°C).
Expected strike time	About 4–8 weeks in warm weather.
Post-treatment / aftercare	- Standard hardening off. - Pot on into a free-draining mix; keep water off the foliage. - Remove any leaves showing grey mould promptly.

Common problems

Problem	Likely cause and fix
Grey fuzz or blackened leaves	Too humid/wet. Cut back misting, remove affected material, improve airflow.
Snapped stems	Brittle stems – transport in rigid containers and use a dibber when setting.
Cuttings flowering, not rooting	Take shoots without axillary flower spikes.

Notes

Difficulty	Easy-Moderate (provisional). <i>Scaevola</i> cuttings generally strike very readily; the hairy foliage makes rot the main risk.
Other methods	Seed – <i>Scaevola</i> seed is generally difficult. Division – possible if a plant has suckered or rooted along its stems.
Confidence	Medium. Little species-specific data published. Based on the genus (strikes readily) plus this species' hairy foliage.
Sources	See References: 1, 4, 5, 6, 7, 12, 13, 14, 16

Nursery records

Log each batch here to confirm or update the timing and difficulty above.

Date set	Stock source / provenance	No. set	No. struck	Strike %	Notes

Scaevola crassifolia

Thick-leaved Fan-flower

Family
Goodeniaceae

DIFFICULTY	WOOD TYPE				BEST MONTHS		CUTTING LENGTH			HORMONE (IBA)		
Easy	Semi-hardwood				Mar - May		75-100 mm			3 g/L IBA		
	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Cuttings	OK	OK	Best	Best	Best	-	-	-	-	OK	OK	OK
Best window		Workable – see ‘Selecting material’ in the method						Avoid / low success				

Plant snapshot

Habit	Erect or spreading shrub, 0.1–1.5 m and up to 3 m wide. Young growth is sticky; thick, toothed, egg-shaped to round leaves.
Flowering	Jul – Dec, also Jan – Feb (blue, mauve or white fans).
Habitat	Sand on coastal dunes, plains and limestone cliffs.
Local note	Recorded in Joondalup and Wanneroo. Flower colour and habit vary – cuttings keep the parent's form.

Propagation options

Method	Suitability	Notes	Details in
Cuttings	Main method	Standard method; keeps flower colour and form.	This guide
Seed	Possible	Large seeds sometimes germinate naturally.	–
Division	Not used	–	–

Propagation method

Item	Details
Best time to take cuttings	Mar – May – warm, and outside the main flowering period. Oct – Feb is workable with non-flowering shoots (shade well in summer). Avoid Jun – Sep.
Wood type	Semi-hardwood.
Cutting size	75–100 mm.
Selecting material	Non-flowering tips. Flower spikes carry leafy bracts – check closely and pinch them off.
Pre-treatment	• New growth is sticky – wipe the blade often. • Strip lower leaves; cut the large remaining leaves back by a third to a half. • Dip the base in rooting hormone.
Rooting hormone	3 g/L IBA (Clonex Purple).
Propagation mix	Free-draining sand/coir mix.
Environment	Standard humid frame with bottom heat, bright shade.
Expected strike time	About 4–8 weeks in warm weather.
Post-treatment / aftercare	• Standard hardening off. • Pot on into a free-draining mix. • Tip-prune young plants for dense growth.

Common problems

Problem	Likely cause and fix
Wilting in the first week	Too much leaf area. Trim leaves harder; keep the frame closed.
Rotting	Mix too wet. Water less often; improve drainage.
Cuttings flowering	Use non-flowering tips or wait for the Mar – May window.

Notes

Difficulty	Easy. Semi-hardwood cuttings in the warmer months are the standard method.
Other methods	Seed – large seeds are produced and sometimes germinate naturally; cuttings are preferred to keep flower colour and habit.
Confidence	High. Cuttings are the standard method in WA nurseries (e.g. Geraldton community nursery).
Sources	See References: 1, 4, 5, 12, 13, 26, 30

Nursery records

Log each batch here to confirm or update the timing and difficulty above.

Date set	Stock source / provenance	No. set	No. struck	Strike %	Notes

Scaevola globulifera

No widely used common name

Family
Goodeniaceae

DIFFICULTY	WOOD TYPE				BEST MONTHS			CUTTING LENGTH			HORMONE (IBA)	
Easy *	Softwood - semi-hardwood				Jan - Apr			50-80 mm			1.5-3 g/L IBA	

Common problems

Problem	Likely cause and fix
Wilting	Low humidity or too much leaf. Trim leaves; keep the frame closed.
Rotting	Mix too wet. Water less often.
Cuttings flowering	Use non-flowering tips, or wait for the Jan – Apr window.

Notes

Difficulty	Easy (provisional). <i>Scaevola</i> cuttings generally strike very readily.
Other methods	Seed – <i>Scaevola</i> seed is generally difficult. Division – check stock plants for rooted stems.
Confidence	Medium. No species-specific data found. Based on the genus, which strikes readily from cuttings.
Sources	See References: 1, 4, 5, 6, 7, 13, 14

Nursery records

Log each batch here to confirm or update the timing and difficulty above.

Date set	Stock source / provenance	No. set	No. struck	Strike %	Notes

Scaevola repens

No widely used common name ('repens' = creeping)

Family
Goodeniaceae

DIFFICULTY	WOOD TYPE	BEST MONTHS	CUTTING LENGTH	HORMONE (IBA)
Easy *	Softwood – semi-hardwood	Mar – May	50–80 mm	1.5–3 g/L IBA

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Cuttings	OK	OK	Best	Best	Best	–	–	–	OK	OK	OK	OK
	Best window	Workable – see 'Selecting material' in the method						Avoid / low success				

Plant snapshot

Habit	Prostrate to ascending or erect shrub, 0.05–1 m; trailing branches to about 50 cm.
Flowering	Jun – Dec, sometimes Jan – Feb (white or cream, occasionally blue-purple; golden hairs on the outside of the flower).
Habitat	Sand dunes and plains on sand, gravel, laterite or limestone.
Local note	Recorded in the City of Joondalup. A variable species with informal subspecies – use local provenance and keep it labelled.

Propagation options

Method	Suitability	Notes	Details in
Cuttings	Main method	Soft to semi-hardwood tips (provisional rating).	This guide
Seed	Difficult	Scaevola seed is generally difficult.	–
Division	Possible	Some scaevolae sucker; check for rooted pieces.	–

Propagation method

Item	Details
Best time to take cuttings	Mar – May – warm autumn weather, mostly outside the main flowering period. Sep – Feb is workable using non-flowering shoots (shade well in summer). Avoid Jun – Aug.
Wood type	Softwood to semi-hardwood tip cuttings.
Cutting size	50–80 mm tips from the trailing stems.
Selecting material	Non-flowering tips. Flower spikes sit in the leaf axils – pinch off any you find.
Pre-treatment	- Strip lower leaves. - Leaves can be long (up to 90 mm) – trim remaining leaves by about one-third to cut water loss. - Dip the base in rooting hormone.
Rooting hormone	Softwood strength (1.5 g/L IBA, Clonex Green) for soft tips; 3 g/L (Clonex Purple) for firmer material.
Propagation mix	Free-draining sand/coir mix.
Environment	Humid frame with bottom heat, bright shade. Keep the mix moist but on the drier side rather than wet.
Expected strike time	About 4–8 weeks in warm weather.
Post-treatment / aftercare	- Standard hardening off. - Pot on into a free-draining native mix. - Pinch tips once growing to encourage branching.

Common problems

Problem	Likely cause and fix
Wilting in the first week	Too much leaf area or low humidity. Trim leaves harder; keep the frame closed.
Rotting	Mix too wet. Water less often; improve drainage.
Cuttings flowering, not rooting	Take non-flowering tips, or wait for the Mar – May window.

Notes

Difficulty	Easy (provisional). <i>Scaevola</i> cuttings generally strike very readily.
Other methods	Seed – <i>Scaevola</i> seed is generally difficult. Division – some <i>scaevolas</i> sucker; check stock plants for rooted pieces.
Confidence	Medium. Little species-specific data published. Based on the genus, which strikes readily from cuttings.
Sources	See References: 1, 4, 5, 6, 7, 12, 13, 14, 16

Nursery records

Log each batch here to confirm or update the timing and difficulty above.

Date set	Stock source / provenance	No. set	No. struck	Strike %	Notes

Scaevola thesioides

No widely used common name

Family
Goodeniaceae

DIFFICULTY	WOOD TYPE			BEST MONTHS			CUTTING LENGTH			HORMONE (IBA)		
Easy-Moderate *	Softwood – semi-hardwood			Apr – May			50–75 mm			1.5–3 g/L IBA		

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Cuttings	OK	OK	OK	Best	Best	–	–	–	OK	OK	OK	OK

Best window

Workable – see ‘Selecting material’ in the method

Avoid / low success

Plant snapshot

Habit	Spreading subshrub or perennial herb, 0.15–1 m. Mostly hairless; thread-like to narrow leaves; ribbed stems.
Flowering	Aug – Dec and Jan – Mar (pale blue to white), in spikes at the ends of branches.
Habitat	Coastal sand and limestone.
Local note	The coastal Perth plant is subsp. <i>thesioides</i> ; recorded in the Wanneroo LGA.

Propagation options

Method	Suitability	Notes	Details in
Cuttings	Main method	Non-flowering side shoots (provisional rating).	This guide
Seed	Difficult	Scaevola seed is generally difficult.	–
Division	Possible	Only if stock plants have suckered.	–

Propagation method

Item	Details
Best time to take cuttings	Apr – May – warm autumn weather between flowering periods. Sep – Mar is workable using non-flowering side shoots. Avoid Jun – Aug.
Wood type	Softwood to semi-hardwood – firm but still green tip or side-shoot cuttings.
Cutting size	50–75 mm.
Selecting material	Non-flowering side shoots. Flowers form at the branch tips, so main tips often carry buds.
Pre-treatment	- Strip leaves from the lower half; leaves are narrow so little trimming is needed. - Stems are slender – always use a dibber hole. - Dip the base in rooting hormone.
Rooting hormone	Softwood strength (1.5 g/L IBA, Clonex Green); 3 g/L (Clonex Purple) for firmer material.
Propagation mix	Free-draining sand/coir mix.
Environment	Hairless leaves tolerate misting better than <i>S. canescens</i> , but fine stems collapse quickly if the mix is waterlogged or dries out. Keep evenly moist, with bottom heat and bright shade.
Expected strike time	About 4–8 weeks in warm weather.
Post-treatment / aftercare	- Standard hardening off. - Pot on gently – fine roots break easily. Single cuttings per tube make this easier. - Pinch tips once growing to encourage branching.

Common problems

Problem	Likely cause and fix
Fine stems wilt or collapse	Mix drying out or waterlogged. Keep evenly moist; check daily.
Cuttings flower instead of rooting	Pinch off terminal flower buds; use side shoots.
Root damage when potting on	Strike one cutting per tube instead of community pots.

Notes

Difficulty	Easy-Moderate (provisional). The genus strikes readily; the fine stems are less forgiving of watering errors.
Other methods	Seed – <i>Scaevola</i> seed is generally difficult. Division – possible if stock plants have suckered.
Confidence	Medium. Little species-specific data published. Based on the genus plus this species' fine, slender growth.
Sources	See References: 1, 4, 5, 6, 7, 12, 13, 14, 16

Nursery records

Log each batch here to confirm or update the timing and difficulty above.

Date set	Stock source / provenance	No. set	No. struck	Strike %	Notes

Spinifex hirsutus

Hairy Spinifex

Family
Poaceae

DIFFICULTY	WOOD TYPE	BEST MONTHS	CUTTING LENGTH	HORMONE (IBA)
Moderate *	Runner (stolon) sections	Mar – May	10–15 cm, 1–2 nodes	Optional (1.5 g/L)

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Cuttings	OK	OK	Best	Best	Best	–	–	–	OK	OK	OK	OK

Best window
Workable – see 'Selecting material' in the method
Avoid / low success

Plant snapshot

Habit	Rhizomatous, tussock-forming grass, 0.3–0.9 m, sending long runners across the sand. Broad, flat, silvery leaves, hairy on both sides.
Flowering	Oct – Jan. Male and female flowers are on separate plants; female heads are large spiky balls that roll in the wind.
Habitat	White sand on coastal dunes; an important stabiliser at the toe of unstable dunes.
Local note	Recorded in the City of Joondalup. Less common than <i>S. longifolius</i> on the Cottesloe foreshore.

Propagation options

Method	Suitability	Notes	Details in
Cuttings	Good	Runner (stolon) sections with 1–2 nodes.	This guide
Seed	Main method	The usual nursery method; cover seed (needs dark).	→ Seed Sowing Guide
Division	Possible	Tussock divisions work but survived less well than rooted runners.	→ Division Guide

Propagation method

Item	Details
Best time to take cuttings	Mar – May, when runners are young, green and actively growing, giving plants ready for winter planting. Sep – Feb is workable. Avoid Jun – Aug (slow growth).
Wood type	Runner (stolon) sections – not woody cuttings. Use young, green runners, not old brown ones.
Cutting size	Nursery tubes: 10–15 cm sections with at least one (ideally two) nodes. Direct planting on site: 40–50 cm runners, buried about halfway in damp sand.
Selecting material	Actively growing runners from the dune edge. Plants are male or female, so collect from several plants if you want seed-producing plantings later.
Pre-treatment	- Keep runners moist and shaded after collecting – they dry out fast. - Cut just below a node; trim leaves back by about half. - Hormone optional – roots form naturally at the nodes.
Rooting hormone	Optional. If used, softwood strength (1.5 g/L IBA, Clonex Green) on the node.
Propagation mix	Coarse washed sand or a very sandy mix. Deep tubes suit the root system.
Environment	Bottom heat and misting gave good rooting at the nodes in <i>S. sericeus</i> trials. The hairy leaves of this species can hold water, so mist lightly.
Expected strike time	Nodes root readily, but plants are slow to reach planting size – in one trial about 80% of cuttings were ready to plant within 18 months.

Item	Details
Post-treatment / aftercare	• Harden off, then grow on in full sun. • Pot on into deep tubes or larger pots. • Plant out in late autumn or winter when rain is reliable.

Common problems

Problem	Likely cause and fix
Runners dry out before setting	Carry in damp bags in an esky; set the same day.
Nodes rot	Mix too wet or too much mist. Use coarser sand; mist less.
Slow growth after rooting	Normal for spinifex – allow time; feed lightly and give full sun.

Notes

Difficulty	Moderate (provisional). Runner sections root well under bottom heat and mist, but it's labour-intensive and slow. Seed is the more common nursery method.
Other methods	Seed – the usual nursery method. The Geraldton community nursery advises collecting from female plants growing near males. <i>S. sericeus</i> research found seed germinates far better in darkness (56%) than light (14%), so cover seed. Division of tussocks also works but survived less well than rooted cuttings in trials (31% vs 40%).
Confidence	Medium. Method based on published trials with the closely related <i>S. sericeus</i> and general spinifex practice.
Sources	See References: 4, 12, 18, 19, 20

Nursery records

Log each batch here to confirm or update the timing and difficulty above.

Date set	Stock source / provenance	No. set	No. struck	Strike %	Notes

Spinifex longifolius

Beach Spinifex

Family
Poaceae

DIFFICULTY	WOOD TYPE	BEST MONTHS	CUTTING LENGTH	HORMONE (IBA)
Moderate *	Runner (stolon) sections	Mar – May	10–15 cm, 1-2 nodes	Optional (1.5 g/L)

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Cuttings	OK	OK	Best	Best	Best	–	–	–	OK	OK	OK	OK
Best window	Workable – see ‘Selecting material’ in the method							Avoid / low success				

Plant snapshot

Habit	Rhizomatous, tussock-forming grass, 0.3–1 m high and up to 2 m wide, with long runners through the sand.
Flowering	Apr – Dec or Jan (green-brown). Male and female plants are separate; female heads up to 250 mm across roll in the wind.
Habitat	White sand on coastal dunes; a key foredune stabiliser.
Local note	Recorded in the Wanneroo LGA; the more common of the two spinifex species on the Cottesloe foreshore.

Propagation options

Method	Suitability	Notes	Details in
Cuttings	Good	Runner (stolon) sections with 1–2 nodes.	This guide
Seed	Main method	The usual nursery method; smoke water, about 10 days.	→ Seed Sowing Guide
Division	Possible	Tussock divisions work but survived less well than rooted runners.	→ Division Guide

Propagation method

Item	Details
Best time to take cuttings	Mar – May , when runners are young, green and actively growing, giving plants ready for winter planting. Sep – Feb is workable. Avoid Jun – Aug (slow growth).
Wood type	Runner (stolon) sections – not woody cuttings. Use young, green runners, not old brown ones.
Cutting size	Nursery tubes: 10–15 cm sections with at least one (ideally two) nodes. Direct planting on site: 40–50 cm runners, buried about halfway in damp sand.
Selecting material	Actively growing runners from the dune edge. Plants are male or female, so collect from several plants if you want seed-producing plantings later.
Pre-treatment	• Keep runners moist and shaded after collecting – they dry out fast. • Cut just below a node; trim leaves back by about half. • Hormone optional – roots form naturally at the nodes.
Rooting hormone	Optional. If used, softwood strength (1.5 g/L IBA, Clonex Green) on the node.
Propagation mix	Coarse washed sand or a very sandy mix. Deep tubes suit the root system.
Environment	Bottom heat and misting gave good rooting at the nodes in <i>S. sericeus</i> trials. Keep the sand moist, not wet.
Expected strike time	Nodes root readily, but plants are slow to reach planting size – in one trial about 80% of cuttings were ready to plant within 18 months.

Item	Details
Post-treatment / aftercare	• Harden off, then grow on in full sun. • Pot on into deep tubes or larger pots. • Plant out in late autumn or winter when rain is reliable.

Common problems

Problem	Likely cause and fix
Runners dry out before setting	Carry in damp bags in an esky; set the same day.
Nodes rot	Mix too wet or too much mist. Use coarser sand; mist less.
Slow growth after rooting	Normal for spinifex – allow time; feed lightly and give full sun.

Notes

Difficulty	Moderate (provisional). Runner sections root well under bottom heat and mist, but it's labour-intensive and slow. Seed is the more common nursery method.
Other methods	Seed – the usual nursery method. The Geraldton community nursery collects from female plants growing close to males, separates the seed, cuts off the awns and sows en masse with smoke water (about 10 days to germinate). <i>S. sericeus</i> research found seed germinates far better in darkness, so cover seed. Division of tussocks also works.
Confidence	Medium. Method based on published trials with the closely related <i>S. sericeus</i> and general spinifex practice.
Sources	See References: 4, 12, 18, 19, 20

Nursery records

Log each batch here to confirm or update the timing and difficulty above.

Date set	Stock source / provenance	No. set	No. struck	Strike %	Notes

Templetonia retusa

Cookies Tongues

Family
Fabaceae

DIFFICULTY	WOOD TYPE	BEST MONTHS	CUTTING LENGTH	HORMONE (IBA)
Easy-Moderate *	Semi-hardwood	Dec - Mar	50-75 mm	3 g/L IBA

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Cuttings	Best	Best	Best	OK	-	-	-	-	-	-	OK	Best

Best window
Workable - see 'Selecting material' in the method
Avoid / low success

Plant snapshot

Habit	Much-branched shrub, 0.3-4 m.
Flowering	Apr - Nov (red, sometimes white-yellow).
Habitat	White sand, red clay, gravel and limestone; coastal cliffs, hillsides and plateaus.
Local note	Recorded in the Wanneroo LGA. Cuttings keep a chosen flower colour.

Propagation options

Method	Suitability	Notes	Details in
Cuttings	Good	Firm current season's growth is successful.	This guide
Seed	Main method	Hot water + smoke water; 8-12 days.	→ Seed Sowing Guide
Division	Not used	Woody shrub.	-

Propagation method

Item	Details
Best time to take cuttings	Dec - Mar, after flowering, once the new growth has firmed. Nov and Apr are workable. Avoid May - Oct (flowering).
Wood type	Semi-hardwood - firm current season's growth (ANPSA).
Cutting size	50-75 mm (ANPSA pea-family guidance).
Selecting material	Non-flowering side shoots from healthy plants.
Pre-treatment	- Remove the leaves from the lower two-thirds. - Light wound on firmer stems. - Dip the base in rooting hormone.
Rooting hormone	Semi-hardwood strength, 3 g/L IBA (Clonex Purple).
Propagation mix	Standard free-draining cutting mix.
Environment	Humid frame, bright shade, bottom heat.
Expected strike time	Not documented - check from about 6 weeks and record the result.
Post-treatment / aftercare	- Standard hardening off. - Pot on into a free-draining mix.

Common problems

Problem	Likely cause and fix
Callus, no roots	Trim and re-set; try a lower hormone strength.
Cuttings brown off	Too dry or too soft. Firmer material; keep humidity up.
Base rotting	Mix too wet. Water less.

Notes

Difficulty	Easy-Moderate (provisional). Cuttings are reported successful; no strike rates published. Seed is easy after hot water.
Confidence	Medium. ANPSA reports cuttings of firm current season's growth are successful; size follows ANPSA pea-family guidance.
Sources	See References: 4, 13, 43, 48

Nursery records

Log each batch here to confirm or update the timing and difficulty above.

Date set	Stock source / provenance	No. set	No. struck	Strike %	Notes

Thomasia triphylla

No widely used common name

Family
Malvaceae

DIFFICULTY	WOOD TYPE	BEST MONTHS	CUTTING LENGTH	HORMONE (IBA)
Easy-Moderate *	Semi-hardwood	Dec – Apr	50–75 mm	3 g/L IBA

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Cuttings	Best	Best	Best	Best	OK	–	–	–	–	–	OK	Best
	Best window	Workable – see ‘Selecting material’ in the method									Avoid / low success	

Plant snapshot

Habit	Multi-stemmed shrub, 0.3–1.2 m.
Flowering	Jul – Nov (pink-purple or white).
Habitat	Calcareous sand; coastal limestone and dunes.
Local note	Recorded in the Wanneroo LGA. Suits alkaline limestone sand and dry coastal sites.

Propagation options

Method	Suitability	Notes	Details in
Cuttings	Main method	Cuttings usually strike reliably in the genus (provisional).	This guide
Seed	Difficult	Unreliable in the genus; a supplier recommends smoke.	–
Division	Not used	–	–

Propagation method

Item	Details
Best time to take cuttings	Dec – Apr, after flowering, once new growth has firmed. Nov and May are workable. Avoid Jun – Oct (cool, then flowering).
Wood type	Semi-hardwood – firm current season's growth.
Cutting size	50–75 mm.
Selecting material	Non-flowering side shoots from healthy plants.
Pre-treatment	- Strip leaves from the lower half; reduce large leaves by a third. - Light wound on firmer stems. - Dip the base in rooting hormone.
Rooting hormone	Semi-hardwood strength, 3 g/L IBA (Clonex Purple).
Propagation mix	Standard free-draining cutting mix.
Environment	Humid frame, bright shade, bottom heat. Keep the mix moist, not wet, and mist lightly.
Expected strike time	Not documented – check from about 4–6 weeks and record the result.
Post-treatment / aftercare	- Standard hardening off. - Grow on in full sun; tip-prune for a bushier plant.

Common problems

Problem	Likely cause and fix
Leaves go brown and soft	Too wet or humid. Mist less, improve airflow.
Callus, no roots	Trim and re-set; try a lower hormone strength.
Slow in winter	Use bottom heat, or wait for the summer window.

Notes

Difficulty	Easy-Moderate (provisional). <i>Thomasia</i> cuttings usually strike reliably, but this species is untested – start with a trial batch.
Other methods	Seed – unreliable in <i>Thomasia</i> ; a seed supplier recommends smoke treatment.
Confidence	Medium. No species data. ANPSA says cuttings of firm growth usually strike reliably in <i>Thomasia</i> , though at least one species is difficult.
Sources	See References: 4, 49, 50, 51, 52

Nursery records

Log each batch here to confirm or update the timing and difficulty above.

Date set	Stock source / provenance	No. set	No. struck	Strike %	Notes

Threlkeldia diffusa

Coast Bonefruit

Family
Amaranthaceae
(chenopods)

DIFFICULTY Moderate *	WOOD TYPE Softwood tips	BEST MONTHS Dec – Mar	CUTTING LENGTH 50–75 mm	HORMONE (IBA) 1.5 g/L IBA
--------------------------	----------------------------	--------------------------	----------------------------	------------------------------

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Cuttings	Best	Best	Best	OK	–	–	–	–	–	–	OK	Best
Best window												
Workable – see ‘Selecting material’ in the method												
Avoid / low success												

Plant snapshot

Habit	Much-branched, prostrate to erect perennial, 0.1–0.4 m, forming a spreading mat. Semi-succulent leaves.
Flowering	Oct – Nov (tiny, green), followed by small red to black fruits.
Habitat	White or grey sand over limestone, coastal areas and saline flats. Salt- and drought-tolerant.
Local note	Recorded in the City of Joondalup.

Propagation options

Method	Suitability	Notes	Details in
Cuttings	Trial	No published cutting data; trial protocol.	This guide
Seed	Main method	Smoke-water soak; 10–15 days in cool weather.	→ Seed Sowing Guide
Division	Not used	Not documented.	–

Propagation method

Item	Details
Best time to take cuttings	Dec – Mar – warm, with plants actively growing after flowering. Nov and Apr are workable. Avoid May – Oct.
Wood type	Softwood tip cuttings of the semi-succulent stems.
Cutting size	50–75 mm.
Selecting material	Healthy tips without developing fruit.
Pre-treatment	- Strip lower leaves. - No wounding. - Dip the base in rooting hormone.
Rooting hormone	Softwood strength, 1.5 g/L IBA (Clonex Green).
Propagation mix	Very free-draining sandy mix.
Environment	Semi-succulent leaves rot if kept wet – mist very little, keep the mix on the dry side, give good airflow and bright light.
Expected strike time	Not documented – start checking from about 4 weeks and record the result.
Post-treatment / aftercare	- Standard hardening off. - Grow on in full sun; water sparingly.

Common problems

Problem	Likely cause and fix
Cuttings turn soft and rot	Too wet or humid. Water less, mist less, improve airflow.
Weak, stretched growth	Not enough light. Move to a brighter position.
No roots	Try firmer tips or the seed method.

Notes

Difficulty	Moderate (provisional). No cutting data found; seed is the proven method (see Other methods).
Other methods	Seed – the documented method. The Geraldton community nursery soaks seed in smoke water (1:9) for 24 hours; germination takes 10–15 days and happens in cooler weather, so sow in autumn.
Confidence	Low. No published cutting data found. Seed is the documented method; cuttings are a trial.
Sources	See References: 1, 4, 12, 28

Nursery records

Log each batch here to confirm or update the timing and difficulty above.

Date set	Stock source / provenance	No. set	No. struck	Strike %	Notes

References

Numbers match the source list on each species page. Web addresses shortened; accessed September 2026.

#	Source	Address
1	Australian Native Plants Society (ANPSA) – Plant Propagation: cuttings, hormones, mixes, hardening off	anpsa.org.au/plant-propagation/
2	ANPSA – Grevillea vestita plant profile	anpsa.org.au/plant_profiles/grevillea-vestita/
3	Australian Plants Society NSW – Grevillea vestita	resources.austplants.com.au/plant/grevillea-vestita/
4	FloraBase, WA Herbarium (DBCA) – species profiles for every species in this guide (e.g. 2463, 2798, 5117, 5162, 7603, 11341, 12585, 12824, 13152, 16449, 42281, 3961, 4256, 5105, 6839; others as cited in the Seed Sowing Guide)	florabase.dbca.wa.gov.au
5	ANPSA – Scaevola ramosissima profile (genus note: cuttings strike very readily, seed difficult)	anpsa.org.au/plant_profiles/scaevola-ramosissima/
6	Sustainable Gardening Australia – Fan Flower: take cuttings during the growing season	sgaonline.org.au/fan-flower-fabulous-native-ground-cover/
7	Greenhouse Product News – Scaevola propagation (IBA rates, flower-free cuttings, bottom heat)	gpnmag.com/article/scaevola-outback-fan-flower/
8	ANPSA – Hibbertia vestita profile (genus note: easy from cuttings, seed unreliable)	anpsa.org.au/plant_profiles/hibbertia-vestita/
9	Australian National Botanic Gardens – Hibbertia pedunculata and H. empetrifolia (cutting timing, root rot)	anbg.gov.au/gnp/gnp11/hibbertia-pedunculata.html
10	Trees and Shrubs Online – Hibbertia (nodal cuttings while in growth; seed dormancy)	treesandshrubsonline.org/articles/hibbertia/
11	DBCA Library – wildflower booklet: H. hypericoides difficult from seed or cuttings	library.dbca.wa.gov.au/FullTextFiles/072542.pdf
12	City of Greater Geraldton Community Nursery – Native Plant Propagation Guide (2019)	cgg.wa.gov.au
13	Growth Technology – Clonex rooting gel strengths (Green 1.5, Purple 3, Red 8 g/L IBA)	growthtechnology.com.au
14	EverydayWits – Propagating Scaevola (small-nursery practice, seasonal success rates)	everydaywits.com/propagate-scaevola/
15	Minimalist Gardener – Propagating Australian natives from cuttings (grevillea strike times, sap irritant)	minimalistgardener.com.au
16	Wikipedia – Scaevola canescens, S. repens, S. thesioides (descriptions)	en.wikipedia.org
17	CompleteGrow – How to propagate Grevillea (low-phosphorus feeding)	completegrow.com.au
18	NZ Department of Conservation – Rehabilitation of coastal foredunes using indigenous sand-binding species (Spinifex sericeus stolon cuttings, seed)	doc.govt.nz/Documents/science-and-technical/sfc122.pdf
19	ABC Gardening Australia – Spinifex (40–50 cm runner cuttings)	abc.net.au/gardening/how-to/spinifex/9432152
20	Cottesloe Coastcare Association – plant pages: Spinifex longifolius, S. hirsutus, Eremophila glabra, Rhagodia baccata	cottesloecoastcare.org
21	ANPSA – Eremophila glabra plant profile	anpsa.org.au/plant_profiles/eremophila-glabra/
22	Australian Bushfoods – The Eremophila Page (cutting length, wounding, hormone)	ausbushfoods.com/bushfoodsonline/reports/Plants/EREMO.TXT
23	Australian National Botanic Gardens – Myoporum insulare	anbg.gov.au/gnp/interns-2010/myoporum-insulare.html
24	ANPSA – Lechenaultia biloba profile (cuttings strike very easily; seed difficult)	anpsa.org.au/plant_profiles/lechenaultia-biloba/
25	Better Homes and Gardens – fastest and easiest native plants to propagate (leschenaultia timing)	bhg.com.au/how-to-propagate-natives

#	Source	Address
26	Nurseries Online – <i>Scaevola crassifolia</i> (semi-hardwood cuttings in warmer months)	nurseriesonline.com.au/plant-index/australian-native-plants/scaevola-crassifolia/
27	Natural Area Nursery – WA native plant propagation (<i>H. hypericoides</i> difficult to propagate)	naturalarea.com.au/revegetation-plant-production/wa-native-plant-propagation/
28	APACE WA – <i>Threlkeldia diffusa</i>	apacewa.org.au/plant/threlkeldia-diffusa/
29	Geographe Community Landcare Nursery – <i>Phyllanthus calycinus</i> (<i>Lysiandra calycina</i>)	gcln.org.au/product/phyllanthus-calycinus/
30	Wikipedia – <i>Hibbertia hypericoides</i> , <i>Eremophila glabra</i> subsp. <i>albicans</i> , <i>Phyllanthus calycinus</i> , <i>Scaevola crassifolia</i>	en.wikipedia.org
31	PMC – Sympatric species of <i>Hibbertia</i> vary in dormancy break and germination requirements	pmc.ncbi.nlm.nih.gov/articles/PMC3336950
32	Cromer E.L. (2007) Seed Germination and Research Records from Alcoa's Marrinup Nursery (seed test figures quoted under Other methods)	library.dbca.wa.gov.au/static/FullTextFiles/924754.pdf
33	Australian National Botanic Gardens – <i>Carpobrotus glaucescens</i> (layering, 30 cm stem pieces, division)	anbg.gov.au/gnp/interns-2005/carpobrotus-glaucescens.html
34	Australian National Botanic Gardens – <i>Leucophyta brownii</i> (semi-hardened cuttings; dislikes humidity)	anbg.gov.au/gnp/interns-2014/leucophyta-brownii.html
35	ANPSA – Australian Daisies (perennial stem cuttings strike quickly; 3:1 sand:peat mix)	anpsa.org.au/genera/daisies/
36	Australian National Botanic Gardens – <i>Atriplex nummularia</i> (ventilate, avoid mist on cuttings)	anbg.gov.au/gnp/interns-2007/atriples-nummularia.html
37	Nurseries Online – <i>Atriplex nummularia</i> (15 cm terminal cuttings in spring)	nurseriesonline.com.au/plant-index/australian-native-plants/atriples-nummularia/
38	Applied Sciences (MDPI, 2025) – Propagation of <i>Atriplex halimus</i> by unrooted cuttings	mdpi.com/2076-3417/15/13/7027
39	Peppy Plants – <i>Hibbertia cuneiformis</i> (propagated by cuttings)	peppyplants.com.au/hibbertia-cuneiformis-unveiling-the-charms-of-the-cutleaf-hibbertia/
40	Wikipedia – <i>Hibbertia cuneiformis</i> (description, flowering, habitat)	en.wikipedia.org/wiki/Hibbertia_cuneiformis
41	ANPSA – <i>Melaleuca</i> genus page (75–100 mm cuttings, strip, wound, hormone)	anpsa.org.au/genera/melaleuca/
42	ANPSA – <i>Kennedia prostrata</i> profile (cuttings of firm current season's growth)	anpsa.org.au/plant_profiles/kennedia-prostrata/
43	ANPSA – Pea family (Fabaceae) genus page (50–75 mm cuttings, leaves off the lower two-thirds)	anpsa.org.au/genera/pea-family/
44	ANPSA – <i>Hardenbergia comptoniana</i> profile (cuttings of firm current season's growth strike well)	anpsa.org.au/plant_profiles/hardenbergia-comptoniana/
45	ANPSA – <i>Hemiandra pungens</i> profile (relatively easy from cuttings; unreliable in humid conditions; grafting)	anpsa.org.au/plant_profiles/hemiandra-pungens/
46	Gardening With Angus – <i>Hemiandra pungens</i> (softwood and semi-hardwood cuttings)	gardeningwithangus.com.au/hemiandra-pungens-snake-plant/
47	APACE WA – <i>Hemiandra pungens</i> (dieback resistant)	apacewa.org.au/plant/hemiandra-pungens/
48	ANPSA – <i>Templetonia retusa</i> profile (cuttings of firm current season's growth are successful)	anpsa.org.au/plant_profiles/templetonia-retusa/
49	ANPSA – <i>Thomasia grandiflora</i> profile (genus: cuttings of firm growth usually strike reliably; seed unreliable)	anpsa.org.au/plant_profiles/thomasia-grandiflora/
50	Geographe Community Landcare Nursery – <i>Thomasia triphylla</i> (coastal, drought tolerant, alkaline sand)	gcln.org.au/product/thomasia-triphylla/
51	Nindethana Australian Seed – <i>Thomasia triphylla</i> (smoke treatment recommended for seed)	nindethana.net.au/thomasia-triphylla-2716
52	ANPSA – Australian Plants Online (2007): <i>Thomasia pygmaea</i> , cuttings difficult to strike	anpsa.org.au/APOL2007/jul07-2.html

Revision log

Version	Date	Changes
1.0	Sep 2026	First edition: <i>Grevillea vestita</i> , <i>Hibbertia racemosa</i> , <i>Scaevola canescens</i> , <i>S. repens</i> , <i>S. thesioides</i> .
1.1	Sep 2026	Added 10 species: <i>Eremophila glabra</i> , <i>Hibbertia hypericoides</i> , <i>Lechenaultia linarioides</i> , <i>Myoporum insulare</i> , <i>Phyllanthus calycinus</i> , <i>Scaevola crassifolia</i> , <i>S. globulifera</i> , <i>Spinifex hirsutus</i> , <i>S. longifolius</i> , <i>Threlkeldia diffusa</i> . Added page numbers to the species index.
1.2	Sep 2026	Added 11 species: <i>Atriplex isatidea</i> , <i>Calothamnus quadrifidus</i> , <i>Carpobrotus virescens</i> , <i>Hibbertia cuneiformis</i> , <i>Kennedia prostrata</i> , <i>Leucophyta brownii</i> , <i>Melaleuca huegelii</i> , <i>M. systema</i> , <i>Olearia axillaris</i> , <i>Pithocarpa cordata</i> , <i>Rhagodia baccata</i> . Added a 'Propagation options' table (cuttings, seed, division) to every species page.
1.3	Sep 2026	Added 4 species: <i>Hardenbergia comptoniana</i> , <i>Hemiandra pungens</i> , <i>Templetonia retusa</i> , <i>Thomasia triphylla</i> .