

Native Plant Division Guide

Western Australian native species – Joondalup, WA

Version 1.3 · October 2026 · 12 species

Species in this edition (alphabetical)

Full index on page 2

Anigozanthos humilis Catspaw	p. 6	Moderate-Difficult *	Ficinia nodosa Knotted Club Rush	p. 18	Easy
Anigozanthos manglesii Red and Green Kangaroo Paw	p. 8	Moderate	Lepidosperma gladiatum Coast Sword-sedge	p. 20	Moderate *
Austrostipa elegantissima Feather Speargrass	p. 10	Easy-Moderate *	Lomandra maritima Maritime Mat Rush	p. 22	Easy-Moderate *
Austrostipa flavescens Coast Speargrass	p. 12	Moderate *	Poa poiformis Coastal Poa	p. 24	Easy-Moderate *
Conostylis aculeata Prickly Conostylis	p. 14	Moderate	Spinifex hirsutus Hairy Spinifex	p. 26	Moderate-Difficult *
Dianella revoluta Blueberry Lily / Flax Lily	p. 16	Easy	Spinifex longifolius Beach Spinifex	p. 28	Moderate-Difficult *

* Provisional rating – based on related plants; confirm with your own nursery records.

Compiled from published sources (see References). Windows and ratings are starting points for local practice.

© 2026 Mitchell Donald. All rights reserved.

How to use this guide

Division means splitting one plant into several pieces that each already have roots. It is the quickest way to multiply clump-forming plants such as kangaroo paws, flax lilies and tufted grasses, and every new plant is identical to its parent. Each species has its own page set, listed alphabetically: quick facts, a division calendar, a plant snapshot, the method, common problems and a record table.

Divide nursery stock plants, not plants from bushland. Timing is written for the Joondalup / Perth coastal climate: most clumps are best divided in autumn so the pieces root before winter.

Species index – at a glance

Species	Page	Difficulty	Division type	Best months	Division size
<i>Anigozanthos humilis</i> Catspaw	6	Moderate-Difficult *	Rhizome clump division	Oct – Nov	3+ shoots with roots
<i>Anigozanthos manglesii</i> Red and Green Kangaroo Paw	8	Moderate	Rhizome clump division	Mar – May	3+ shoots with roots
<i>Austrostipa elegantissima</i> Feather Speargrass	10	Easy-Moderate *	Rhizome / tussock division	Aug – Sep	Clumps of 3–5 tillers
<i>Austrostipa flavescens</i> Coast Speargrass	12	Moderate *	Tussock division	Apr – Jun	Small clumps, 3–5 tillers
<i>Conostylis aculeata</i> Prickly Conostylis	14	Moderate	Clump division / rooted runners	Mar – May	Larger clumps, 5+ shoots
<i>Dianella revoluta</i> Blueberry Lily / Flax Lily	16	Easy	Rhizome clump division	Mar – May	3–4 shoots with rhizome
<i>Ficinia nodosa</i> Knotted Club Rush	18	Easy	Rhizome clump division	Mar – May	6–10 stems with rhizome
<i>Lepidosperma gladiatum</i> Coast Sword-sedge	20	Moderate *	Crown (clump) division	Mar – May	Large pieces with rhizome & roots
<i>Lomandra maritima</i> Maritime Mat Rush	22	Easy-Moderate *	Clump division	Aug – Sep	3–4 shoots with roots
<i>Poa poiformis</i> Coastal Poa	24	Easy-Moderate *	Tussock division	Aug – Sep	Clumps of 3–5 tillers
<i>Spinifex hirsutus</i> Hairy Spinifex	26	Moderate-Difficult *	Tussock division	Mar – May	Rooted clumps with runners
<i>Spinifex longifolius</i> Beach Spinifex	28	Moderate-Difficult *	Tussock division	Mar – May	Rooted clumps with runners

* Provisional rating – based on related plants; update once your own results are recorded.

Division calendar

Species	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
<i>Anigozanthos humilis</i>	–	–	OK	OK	–	–	–	–	–	Best	Best	–
<i>Anigozanthos manglesii</i>	–	OK	Best	Best	Best	OK	–	–	OK	OK	–	–
<i>Austrostipa elegantissima</i>	–	–	OK	OK	OK	OK	OK	Best	Best	OK	–	–
<i>Austrostipa flavescens</i>	–	–	OK	Best	Best	Best	OK	–	–	–	–	–
<i>Conostylis aculeata</i>	–	OK	Best	Best	Best	OK	–	–	–	OK	OK	OK
<i>Dianella revoluta</i>	–	OK	Best	Best	Best	OK	–	OK	OK	OK	–	–
<i>Ficinia nodosa</i>	–	OK	Best	Best	Best	OK	OK	OK	OK	OK	–	–
<i>Lepidosperma gladiatum</i>	–	OK	Best	Best	Best	OK	OK	–	–	–	–	–

<i>Lomandra maritima</i>	–	–	OK	OK	OK	OK	OK	Best	Best	OK	–	–
<i>Poa poiformis</i>	–	–	OK	OK	OK	–	OK	Best	Best	OK	–	–
<i>Spinifex hirsutus</i>	OK	OK	Best	Best	Best	–	–	–	OK	OK	OK	OK
<i>Spinifex longifolius</i>	OK	OK	Best	Best	Best	–	–	–	OK	OK	OK	OK
Perth season	Summer		Autumn			Winter			Spring		Sum.	
Best window	Workable – slower to re-establish or needs more care					Avoid / low success						

Other species that can be divided or layered

These have no page in this guide because another method works better, but they can be split, or rooted pieces lifted, when stock plants allow.

Species	How	Main method
<i>Acanthocarpus preissii</i>	Rhizomatous clumps; division not documented.	Seed
<i>Carpobrotus virescens</i>	Lift stems that have rooted – treat as cuttings.	Cuttings
<i>Conostylis candicans</i>	Less easy than kangaroo paws – large pieces (3+ big shoots), autumn.	Seed
<i>Conostylis setigera</i>	Large pieces in autumn, as for other <i>Conostylis</i> .	Seed
<i>Grevillea vestita</i>	Can sucker – lift rooted suckers.	Cuttings
<i>Hibbertia racemosa</i>	Low stems may root where they touch the soil (layering).	Cuttings
<i>Lechenaultia linarioides</i>	Some <i>lechenaultias</i> sucker – pot up rooted suckers.	Cuttings
<i>Opercularia vaginata</i>	May sucker; untested.	–
<i>Orthrosanthus laxus</i>	Clumps can be divided in spring (RHS); untested here.	Seed
<i>Patersonia occidentalis</i>	Large pieces in autumn; ANBG calls division unreliable.	Seed
<i>Scaevola canescens</i>	Only if stems have rooted or suckered.	Cuttings
<i>Scaevola globulifera</i>	Check stock plants for rooted stems.	Cuttings
<i>Scaevola repens</i>	Some <i>scaevolae</i> sucker; check for rooted pieces.	Cuttings
<i>Scaevola thesioides</i>	Only if stock plants have suckered.	Cuttings

Types of division

Type	What it is	Examples
Clump / rhizome	Grass-like clumps with a creeping underground stem. Cut into pieces of rhizome, each with shoots and roots.	<i>Anigozanthos</i> , <i>Dianella</i> , <i>Lomandra</i> , <i>Patersonia</i>
Tussock	True grasses. Split a clump into smaller clumps of shoots (tillers) with roots.	<i>Austrostipa</i> , <i>Spinifex</i>
Suckers	New shoots arising from roots a short way from the parent. Dig up with roots attached and pot individually.	Some <i>Lechenaultia</i> , <i>Scaevola</i> , <i>Grevillea</i>
Layering	Stems that root where they touch the soil. Cut into rooted pieces and pot up, or pin a stem into a pot beside the parent.	<i>Hibbertia</i> , <i>Carpobrotus</i> , spinifex runners

Based on ANPSA's propagation notes (source 1).

Difficulty ratings

Rating	Meaning
Easy	Divisions re-establish reliably with the standard method.
Easy-Moderate	Usually re-establish; one factor (e.g. moisture, size of piece) needs care.
Moderate	Re-establish, but results depend on timing, piece size and aftercare.
Moderate-Difficult	Many divisions fail; another method is usually better.
Difficult	Rarely successful; use another method.

Confidence

Level	Meaning
High	A published division method exists for the species or genus.
Medium	Based on closely related plants or general practice for this plant type. Verify with trials.
Low	Little information available; treat as a trial protocol.

Standard division method

Species pages note where a species differs.

Step	What to do
1. Choose stock	Healthy, vigorous, pest- and disease-free nursery plants with plenty of shoots. Label each stock plant's provenance.
2. Time it	Warmer months give the best results; late-summer to autumn divisions root well before winter (ANPSA). Summer divisions need extra shelter.
3. Water first	Water the stock plant the day before so roots and shoots are full of water.
4. Lift and clean	Lift the clump or knock it out of its pot. Shake or wash off the mix so you can see where to cut.
5. Divide	Cut with a clean, sharp knife or spade. Aim for pieces with three or four leaf shoots , a good section of rhizome and healthy roots (ANPSA). Smaller pieces need a humid house.
6. Trim	Cut off damaged roots. Cut leaves back by about one-third to reduce water loss.
7. Pot	Pot into a free-draining mix with shoots at the same level as before . Firm in and water well.
8. Shelter	Keep in a sheltered, shaded spot (or humid house for small pieces) until new leaves grow and roots show at the drainage holes. Small divisions typically need 4–6 weeks.
9. Harden off	About a week with nights exposed, then a week in a sheltered spot, then into the open (ANPSA).
10. Pot on	Pot on or plant out once roots fill the pot.

Hygiene

Topic	Guidance
Tools	Clean, sharp knives and spades. Wipe with alcohol between stock plants.
Disease	Don't divide plants showing disease. Kangaroo paws are prone to ink disease (black leaf spots); many WA natives are susceptible to <i>Phytophthora</i> – keep pots off the ground and use fresh mix.
Collecting	Digging plants from bushland needs a DBCA licence and landholder permission. Divide nursery stock instead, and label every batch with its source.

Anigozanthos humilis

Catspaw

Family
Haemodoraceae

DIFFICULTY	DIVISION TYPE	BEST MONTHS	DIVISION SIZE	TIME TO ESTABLISH
Moderate-Difficult *	Rhizome clump division	Oct – Nov	3+ shoots with roots	4–6 weeks sheltered

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Division	–	–	OK	OK	–	–	–	–	–	Best	Best	–
	Best window		Workable – slower to re-establish or needs more care									Avoid / low success

Plant snapshot

Habit	Low rhizomatous herb, usually 0.1–0.3 m locally. Often short-lived – about five years at most in humid-summer areas.
Flowering	Jul – Oct (yellow, orange and red furry paws).
Habitat	Sand, sandy loam, clay, laterite and limestone; well-drained ground to winter-wet swamps.
Local note	Recorded in Joondalup and Wanneroo (subsp. <i>humilis</i>). Prone to rust.

Propagation options

Method	Suitability	Notes	Details in
Cuttings	Not used	Kangaroo paws are not grown from cuttings.	–
Seed	Main method	No pre-treatment needed; smoke lifts germination.	→ Seed Sowing Guide
Division	Possible	After flowering, vigorous plants only.	This guide

Division method

Item	Details
Best time to divide	Oct – Nov, just after flowering (ANPSA). Mar – Apr is workable, as for other kangaroo paws. Avoid Dec – Feb (heat) and May – Sep (plants are growing and flowering).
Division type	Rhizome clump division.
Division size	At least 3 shoots with a piece of rhizome and roots. Plants are small, so you may only get two or three pieces.
Selecting stock	Only vigorous plants – weak or older plants rarely recover. Choose plants free of rust.
Preparation	- Water the stock plant the day before. - Lift the clump and cut the rhizome apart with a clean, sharp knife. - Trim the leaves back by about a third.
Potting	Pot into a very free-draining mix at the original depth.
Environment	Sheltered, shaded spot until new growth starts. Keep the leaves dry – water the mix, not the plant.
Time to establish	About 4–6 weeks (record yours).
Aftercare	- Harden off gradually. - Grow on in full sun with good airflow.

Common problems

Problem	Likely cause and fix
Divisions fail	Stock plant too weak. Divide only vigorous plants.
Orange pustules on leaves	Rust. Remove affected leaves; keep foliage dry; don't divide infected plants.
Base rots	Too wet. Very free-draining mix; water less.

Notes

Difficulty	Moderate–Difficult (provisional). Works on vigorous plants, but the species is small and short-lived. Seed is the main method.
Confidence	Medium. ANPSA: divide after flowering and only if the plant is vigorous. Timing and size follow the kangaroo paw genus method.
Sources	See References: 2, 3, 15

Nursery records

Log each batch of divisions here to confirm or update the timing and difficulty above.

Date divided	Stock plant / provenance	No. pieces	No. established	Success %	Notes

Anigozanthos manglesii

Red and Green Kangaroo Paw

Family
Haemodoraceae

DIFFICULTY	DIVISION TYPE				BEST MONTHS			DIVISION SIZE			TIME TO ESTABLISH	
Moderate	Rhizome clump division				Mar – May			3+ shoots with roots			4–6 weeks sheltered	
	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Division	–	OK	Best	Best	Best	OK	–	–	OK	OK	–	–
	Best window	Workable – slower to re-establish or needs more care						Avoid / low success				

Plant snapshot

Habit	Rhizomatous perennial herb, 0.2–1.1 m (flower stems). Often short-lived – best treated as a 2–3 year plant.
Flowering	Aug – Nov (green and red). WA's floral emblem.
Habitat	White, yellow or grey sand and sandy loam.
Local note	Recorded in the Wanneroo LGA. Less vigorous and shorter-lived than most garden hybrids.

Propagation options

Method	Suitability	Notes	Details in
Cuttings	Not used	Clumping herb – no stems to cut.	–
Seed	Main method	Dry heat or aged seed; autumn sowing.	→ Seed Sowing Guide
Division	Good	Split vigorous clumps into rhizome pieces with shoots and roots.	This guide

Division method

Item	Details
Best time to divide	Mar – May – ANPSA says clumps are best divided in autumn, though other times also work. Feb, Jun and Sep – Oct are workable; summer divisions need protection until re-established. Avoid Jul – Aug (cold, wet) and Nov – Jan (heat).
Division type	Rhizome clump division – the clump is cut into pieces of rhizome, each with leaf shoots and roots.
Division size	At least three large shoots per piece gives the best chance of success (ANPSA). Single-shoot pieces with some roots can work but need a humid glasshouse until new roots form.
Selecting stock	Vigorous, healthy clumps, ideally in their second year – this species is often short-lived. Don't divide plants with black leaf spotting (ink disease).
Preparation	- Water the stock plant the day before. - Lift the clump (or knock it out of its pot) and shake or wash off the mix so you can see the rhizome. - Cut through the rhizome with a clean, sharp knife. - Trim damaged roots. - Cut all leaves back by about one-third to reduce water loss (ANPSA).
Potting	Pot into a well-drained potting mix with the shoots at the same level as before (ANPSA). Don't bury the crown. Water in.
Environment	Sheltered, shaded spot until new leaf growth is well advanced and roots show at the drainage holes. Small divisions go in a humid house.
Time to establish	Small divisions need about 4–6 weeks in humid conditions (ANPSA); larger pieces re-establish in the shelter over a similar time.

Item	Details
Aftercare	• Harden off gradually – about a week with nights exposed, then a week in a sheltered spot, then into the open. • Keep foliage dry and airflow good to limit ink disease. • Pot on once roots fill the pot.

Common problems

Problem	Likely cause and fix
Division rots at the base	Too wet, too cold, or planted too deep. Pot at the original level; water less; divide in autumn.
Black spots spreading on leaves	Ink disease. Discard affected pieces; keep foliage dry; improve airflow.
Division wilts and dies	Too few roots or leaves not trimmed. Use larger pieces, trim leaves by a third, keep sheltered.

Notes

Difficulty	Moderate. Kangaroo paws divide well, but this species is shorter-lived than many hybrids, so divide young, healthy clumps. Seed is the main method.
Confidence	High. ANPSA publishes a detailed division method for kangaroo paws.
Sources	See References: 1, 2, 3, 5

Nursery records

Log each batch of divisions here to confirm or update the timing and difficulty above.

Date divided	Stock plant / provenance	No. pieces	No. established	Success %	Notes

Austrostipa elegantissima

Feather Speargrass

Previously *Stipa elegantissima*Family
Poaceae

DIFFICULTY	DIVISION TYPE	BEST MONTHS	DIVISION SIZE	TIME TO ESTABLISH
Easy-Moderate *	Rhizome / tussock division	Aug – Sep	Clumps of 3-5 tillers	4-6 weeks

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Division	–	–	OK	OK	OK	OK	OK	Best	Best	OK	–	–
	Best window		Workable – slower to re-establish or needs more care							Avoid / low success		

Plant snapshot

Habit	Rhizomatous, tufted, often sprawling perennial grass, 0.35–2 (–3) m.
Flowering	Aug – Dec or Jan (fluffy, feathery green heads).
Habitat	Sand, loam, clay and granite, occasionally saline.
Local note	Recorded around Perth (Cottesloe, Cambridge, Cockburn); FloraBase has no Joondalup or Wanneroo specimens.

Propagation options

Method	Suitability	Notes	Details in
Cuttings	Not used	Grasses are not grown from cuttings.	–
Seed	Main method	1–2 year old seed; smoke-water soak.	→ Seed Sowing Guide
Division	Good	Late winter – early spring (genus).	This guide

Division method

Item	Details
Best time to divide	Aug – Sep – late winter to early spring gives high success for <i>Austrostipa</i> . Mar – Jul and Oct are workable. Avoid Nov – Feb (heat).
Division type	Rhizome / tussock division – split the clump into pieces of shoots (tillers), each with roots.
Division size	Clumps of 3–5 tillers with a good root mass. Very small pieces establish poorly.
Selecting stock	Healthy, established pot-grown clumps that are actively growing.
Preparation	- Water the stock plant the day before. - Knock the clump out and cut it apart with a sharp knife or spade. - Cut the stems and leaves back to about 15 cm to reduce water loss.
Potting	Pot into a free-draining mix at the original depth. Firm in and water well.
Environment	Shade house until new leaves appear. Keep the mix moist, not wet.
Time to establish	Roughly 4–6 weeks. Record your results.
Aftercare	- Move gradually to full sun once new growth shows. - Give tubes room – plants sprawl.

Common problems

Problem	Likely cause and fix
Divisions brown off	Too small, or roots dried out. Keep 3–5 tillers per piece; work in the shade.
Base rots	Too wet or planted too deep.
No new growth	Divided in summer. Divide in late winter – early spring.

Notes

Difficulty	Easy–Moderate (provisional). Division works well in the genus; untested for this species.
Confidence	Medium. No species method published. ANBG reports high success from division of <i>Austrostipa</i> in late winter – early spring.
Sources	See References: 1, 3, 16

Nursery records

Log each batch of divisions here to confirm or update the timing and difficulty above.

Date divided	Stock plant / provenance	No. pieces	No. established	Success %	Notes

Austrostipa flavescens

Coast Speargrass

Family
Poaceae

DIFFICULTY	DIVISION TYPE			BEST MONTHS			DIVISION SIZE			TIME TO ESTABLISH		
Moderate *	Tussock division			Apr – Jun			Small clumps, 3-5 tillers			4-6 weeks		
	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Division	–	–	OK	Best	Best	Best	OK	–	–	–	–	–
	Best window		Workable – slower to re-establish or needs more care					Avoid / low success				

Plant snapshot

Habit	Tufted perennial grass, 0.2–1.2 m.
Flowering	Sep – Oct.
Habitat	Grey, red or calcareous sand, sandy loam and limestone; sand dunes.
Local note	Coastal species. Use nursery stock plants rather than digging from bushland.

Propagation options

Method	Suitability	Notes	Details in
Cuttings	Not used	Tufted grass.	–
Seed	Main method	Aged seed and smoke water; sow shallow.	→ Seed Sowing Guide
Division	Good	Divide established tussocks into small clumps.	This guide

Division method

Item	Details
Best time to divide	Apr – Jun, as autumn rain restarts growth. Mar and Jul are workable. Avoid Aug – Feb (flowering, seed set and summer heat).
Division type	Tussock division – a clump is split into smaller clumps of shoots (tillers), each with roots.
Division size	Small clumps of 3-5 tillers with a good root mass. Very small pieces establish poorly.
Selecting stock	Healthy, established pot-grown clumps that are actively growing.
Preparation	- Water the stock plant the day before. - Knock the clump out and tease or cut it apart with a sharp knife or spade. - Trim damaged roots. - Cut the leaves back to about 10–15 cm to reduce water loss.
Potting	Pot into a free-draining mix at the original depth. Firm in and water well.
Environment	Shade house until new leaves appear. Keep the mix moist, not wet.
Time to establish	Roughly 4-6 weeks in autumn. Record your results.
Aftercare	- Move gradually to full sun once new growth shows. - Pot on or plant out in winter.

Common problems

Problem	Likely cause and fix
Divisions brown off	Too small, or roots dried out. Keep 3–5 tillers per piece; work quickly in the shade.
Base rots	Too wet or planted too deep. Pot at the original level; water less.
No new growth	Divided in the wrong season. Divide as autumn growth starts.

Notes

Difficulty	Moderate (provisional). Grass clumps usually divide readily, but this species hasn't been tested here. Seed is the main method.
Confidence	Low. Division of established clumps works for this grass, but no species-specific method is published. Follows general tussock-grass practice – trial protocol.
Sources	See References: 1, 3

Nursery records

Log each batch of divisions here to confirm or update the timing and difficulty above.

Date divided	Stock plant / provenance	No. pieces	No. established	Success %	Notes

Conostylis aculeata

Prickly Conostylis

Family
Haemodoraceae

DIFFICULTY	DIVISION TYPE				BEST MONTHS		DIVISION SIZE			TIME TO ESTABLISH		
Moderate	Clump division / rooted runners				Mar – May		Larger clumps, 5+ shoots			4-8 weeks sheltered		
	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Division	–	OK	Best	Best	Best	OK	–	–	–	OK	OK	OK
Best window		Workable – slower to re-establish or needs more care						Avoid / low success				

Plant snapshot

Habit	Rhizomatous, tufted perennial, 0.06–0.5 m, with prickly-edged leaves.
Flowering	Aug – Nov (yellow).
Habitat	Sand, loam, clay, gravel, limestone or laterite; winter-wet areas and sand dunes.
Local note	Recorded in Joondalup and Wanneroo (subsp. <i>cygnorum</i> on the coastal plain). Dieback resistant. Keep subspecies separate.

Propagation options

Method	Suitability	Notes	Details in
Cuttings	Not used	Tufted herb.	–
Seed	Main method	Smoke water; about 19% in nursery tests.	→ Seed Sowing Guide
Division	Good	Larger divisions, or rooted runners after flowering.	This guide

Division method

Item	Details
Best time to divide	Mar – May for clump division. Rooted runners are ready just after flowering (about Oct – Dec). Feb and Jun workable. Avoid Jul – Sep for clumps (flowering, cold).
Division type	Clump division of the rhizome, or rooted runners : after flowering the runners form aerial roots at the leaf bases.
Division size	Keep divisions larger than for kangaroo paws – ANPSA notes <i>Conostylis</i> divide less readily and larger pieces do best. Aim for 5 or more shoots with a good piece of rhizome.
Selecting stock	Vigorous, established nursery clumps. Watch runners after flowering for new roots.
Preparation	- Water the stock plant the day before. - Lift and cut the clump with a sharp knife, keeping pieces large. Rooted runners: there is only a 2–3 week window after the roots appear – cut and pot them promptly. - Trim leaves back by about a third.
Potting	Pot into a free-draining mix at the original depth.
Environment	Sheltered, shaded spot (humid house for runners and small pieces) until new growth starts.
Time to establish	About 4–8 weeks (record yours).
Aftercare	- Harden off gradually. - Grow on in full sun; don't overwater.

Common problems

Problem	Likely cause and fix
Small divisions die	Pieces too small. Keep divisions larger.
Runners don't root in the pot	Cut too late or too early. Take them in the 2–3 weeks after aerial roots appear.
Base rots	Too wet. Free-draining mix; water less.

Notes

Difficulty	Moderate. Divides less readily than kangaroo paws; use larger pieces or rooted runners. Seed with smoke is the other option.
Confidence	Medium. ANPSA notes <i>Conostylis</i> divide less readily than kangaroo paws (larger divisions best); APS NSW describes potting rooted runners.
Sources	See References: 2, 3, 12, 13, 14

Nursery records

Log each batch of divisions here to confirm or update the timing and difficulty above.

Date divided	Stock plant / provenance	No. pieces	No. established	Success %	Notes

Dianella revoluta

Blueberry Lily / Flax Lily

Family
Asphodelaceae
(Hemerocallidoideae)

DIFFICULTY		DIVISION TYPE				BEST MONTHS		DIVISION SIZE		TIME TO ESTABLISH			
Easy		Rhizome clump division				Mar – May		3-4 shoots with rhizome		4-6 weeks			
		Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Division		-	OK	Best	Best	Best	OK	-	OK	OK	OK	-	-
Best window		Workable – slower to re-establish or needs more care						Avoid / low success					

Plant snapshot

Habit	Rhizomatous, tufted perennial herb, 0.3–1.5 m.
Flowering	Sep – Dec (blue-purple), followed by blue berries.
Habitat	White or yellow sand, sandy clay, laterite, loam and clay.
Local note	Recorded in Joondalup and Wanneroo. Division is much quicker than seed, which is slow and erratic.

Propagation options

Method	Suitability	Notes	Details in
Cuttings	Not used	Clumping herb.	–
Seed	Difficult	GA + smoke water; 8–11 weeks.	→ Seed Sowing Guide
Division	Main method	The quickest way to produce plants.	This guide

Division method

Item	Details
Best time to divide	Mar – May, so divisions root before winter. Feb, Jun and Aug – Oct are workable. Avoid Nov – Jan (heat) and Jul (cold).
Division type	Rhizome clump division.
Division size	Three or four leaf shoots with a good section of rhizome and healthy roots (ANPSA). Single shoots need a humid house.
Selecting stock	Vigorous clumps with plenty of shoots. Keep forms and provenances separate.
Preparation	- Water the stock plant the day before. - Lift the clump and shake or wash off the mix. - Cut the rhizome into pieces with a clean, sharp knife. - Trim damaged roots and cut leaves back by about one-third to one-half.
Potting	Pot into a free-draining native mix at the original depth. Water in.
Environment	Sheltered, shaded spot (humid house for small pieces) until new growth starts.
Time to establish	About 4–6 weeks in autumn.
Aftercare	- Harden off gradually – about a week with nights exposed, then a sheltered week. - Pot on once roots fill the pot.

Common problems

Problem	Likely cause and fix
Divisions wilt	Leaves not trimmed or too few roots. Trim harder; use bigger pieces.
Rhizome rots	Too wet or planted too deep. Water less; pot at the original level.
Slow to re-shoot	Divided in cold weather. Divide in autumn or spring.

Notes

Difficulty	Easy. Clump-forming lilies divide readily, and this is the quickest way to produce plants.
Confidence	Medium. ANPSA lists <i>Dianella</i> among the grass-like clumps suited to division; details follow the general method.
Sources	See References: 1, 3

Nursery records

Log each batch of divisions here to confirm or update the timing and difficulty above.

Date divided	Stock plant / provenance	No. pieces	No. established	Success %	Notes

Ficinia nodosa

Knotted Club Rush

Family
Cyperaceae (sedges)

DIFFICULTY		DIVISION TYPE				BEST MONTHS		DIVISION SIZE		TIME TO ESTABLISH			
Easy		Rhizome clump division				Mar – May		6–10 stems with rhizome		Plantable in a few months			
		Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Division		–	OK	Best	Best	Best	OK	OK	OK	OK	OK	–	–
Best window		Workable – slower to re-establish or needs more care						Avoid / low success					

Plant snapshot

Habit	Erect, clumping, rhizomatous sedge to 1 m high and 0.8 m wide.
Flowering	Oct – Dec or Jan (brown knot-like heads).
Habitat	Calcareous sand and limestone; coastal dunes, seasonally wet swamps and salt-lake shores.
Local note	Recorded in Joondalup and Wanneroo. Salt tolerant, fast growing, a strong sand binder.

Propagation options

Method	Suitability	Notes	Details in
Cuttings	Not used	Sedge.	–
Seed	Main method	No pre-treatment; about 55% germination.	→ Seed Sowing Guide
Division	Main method	Divides readily; plantable in a few months.	This guide

Division method

Item	Details
Best time to divide	Mar – May, so divisions root before winter. Feb and Jun – Oct are workable. Avoid Nov – Jan (heat).
Division type	Rhizome clump division.
Division size	Anything from a short piece of rhizome with 6–10 stems up to a 10 × 10 cm clod (NZ restoration guidance).
Selecting stock	Vigorous nursery clumps or tray-grown stock.
Preparation	- Water the stock plant the day before. - Lift the clump and cut it into pieces with a sharp knife or spade. - Trim stems back by about a third to cut water loss.
Potting	Pot into tubes or pots of free-draining mix at the original depth. Water well.
Environment	Shade house until new stems appear; keep the mix moist – it tolerates wetter mix than most.
Time to establish	Divisions are ready for planting within a few months .
Aftercare	- Harden off, then grow on in full sun. - Plant out in winter.

Common problems

Problem	Likely cause and fix
Divisions brown off	Pieces dried out. Keep moist and shaded while working.
Slow to re-shoot	Divided in cold weather. Divide in autumn.
Base rots	Waterlogged in heavy mix. Use a free-draining mix.

Notes

Difficulty	Easy. Divides readily – transplanted divisions had 100% survival in one NZ planting trial.
Confidence	Medium. Coastal restoration guidance for this species (NZ); timing follows ANPSA's general division advice.
Sources	See References: 1, 3, 9, 10

Nursery records

Log each batch of divisions here to confirm or update the timing and difficulty above.

Date divided	Stock plant / provenance	No. pieces	No. established	Success %	Notes

Lepidosperma gladiatum

Coast Sword-sedge

Family
Cyperaceae (sedges)

DIFFICULTY		DIVISION TYPE				BEST MONTHS		DIVISION SIZE		TIME TO ESTABLISH			
Moderate *		Crown (clump) division				Mar – May		Large pieces with rhizome & roots		Slow – several weeks			
		Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Division		–	OK	Best	Best	Best	OK	OK	–	–	–	–	–
Best window		Workable – slower to re-establish or needs more care						Avoid / low success					

Plant snapshot

Habit	Robust, rhizomatous, tufted sedge, 0.5–1.5 (–3) m, clumps to 1.5 m wide. Knife-sharp leaf edges.
Flowering	Nov – Dec or Jan – May (brown).
Habitat	White, grey or calcareous sand, limestone and loam; dunes and creeklines.
Local note	Recorded in the Wanneroo LGA. Wear heavy gloves and long sleeves when dividing.

Propagation options

Method	Suitability	Notes	Details in
Cuttings	Not used	Sedge.	–
Seed	Difficult	Variable; 41% in one test, 53–72 days.	→ Seed Sowing Guide
Division	Possible	Crown division is used commercially (trial here).	This guide

Division method

Item	Details
Best time to divide	Mar – May (trial). Feb, Jun – Jul workable. Avoid Aug – Jan.
Division type	Crown (clump) division of the rhizome.
Division size	Keep pieces large, with a good section of rhizome and plenty of roots (trial).
Selecting stock	Vigorous nursery clumps. Mark any selected forms.
Preparation	- Water the day before. - Lift the clump; cut through the tough rhizome with a sharp spade or pruning saw. - Trim leaves back by about half to cut water loss.
Potting	Pot into deep pots of free-draining mix at the original depth. Water well.
Environment	Shade house until new leaves appear.
Time to establish	Slow – allow several weeks; record the result.
Aftercare	- Harden off, then grow on in full sun. - Plant out in late autumn or winter.

Common problems

Problem	Likely cause and fix
Divisions die back	Too few roots or too small. Keep pieces large.
Hands cut	Leaf edges are sharp – heavy gloves and sleeves.
Slow establishment	Normal. Be patient; full sun once rooted.

Notes

Difficulty	Moderate (provisional). Used commercially, but untested here. Tissue-cultured stock is also available.
Confidence	Low. Crown division is used commercially (a Joondalup-selected form was first propagated this way) but no method or survival data are published. Trial protocol.
Sources	See References: 3, 11

Nursery records

Log each batch of divisions here to confirm or update the timing and difficulty above.

Date divided	Stock plant / provenance	No. pieces	No. established	Success %	Notes

Lomandra maritima

Maritime Mat Rush

Family
Asparagaceae
(Lomandroideae)

DIFFICULTY	DIVISION TYPE	BEST MONTHS	DIVISION SIZE	TIME TO ESTABLISH
Easy-Moderate *	Clump division	Aug – Sep	3–4 shoots with roots	4–6 weeks

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Division	–	–	OK	OK	OK	OK	OK	Best	Best	OK	–	–
Best window												

Plant snapshot

Habit	Rhizomatous, tufted perennial, 0.2–0.6 m, clumps to 0.2 m wide. Male and female plants are separate (dioecious).
Flowering	Aug – Nov (purple and yellow).
Habitat	White or grey sand and limestone; coastal dunes and hills.
Local note	Recorded in Joondalup and Wanneroo. A breeding plant for the threatened graceful sun-moth.

Propagation options

Method	Suitability	Notes	Details in
Cuttings	Not used	Tufted herb.	–
Seed	Difficult	Lomandra seed is strongly dormant; not documented for this species.	–
Division	Main method	Clump division, late winter to early spring (provisional).	This guide

Division method

Item	Details
Best time to divide	Aug – Sep – late winter to early spring is usually best for lomandras. Mar – Jul and Oct are workable. Avoid Nov – Feb.
Division type	Clump division.
Division size	Three or four shoots with a piece of rhizome and healthy roots (ANPSA general guidance).
Selecting stock	Vigorous nursery clumps. Plants are male or female – label stock plants when they flower so plantings include both.
Preparation	- Water the day before. - Lift the clump and cut it apart with a sharp knife. - Trim the leaves back to about a third of their length.
Potting	Pot into a free-draining mix at the original depth.
Environment	Sheltered, shaded spot until new growth starts.
Time to establish	About 4–6 weeks (record yours).
Aftercare	- Harden off gradually. - Pot on once roots fill the pot.

Common problems

Problem	Likely cause and fix
Divisions wilt	Leaves not trimmed or too few roots. Trim harder; use bigger pieces.
Base rots	Too wet or planted too deep.
All plants one sex	Stock came from a single plant. Divide from several labelled male and female plants.

Notes

Difficulty	Easy-Moderate (provisional). Lomandras generally divide well; untested for this species.
Confidence	Medium. No species method published. Follows general <i>Lomandra</i> and ANPSA clump-division guidance.
Sources	See References: 1, 3, 6, 7, 8

Nursery records

Log each batch of divisions here to confirm or update the timing and difficulty above.

Date divided	Stock plant / provenance	No. pieces	No. established	Success %	Notes

Poa poiformis

Coastal Poa

Family
Poaceae

DIFFICULTY	DIVISION TYPE	BEST MONTHS	DIVISION SIZE	TIME TO ESTABLISH
Easy-Moderate *	Tussock division	Aug – Sep	Clumps of 3-5 tillers	4-6 weeks

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Division	–	–	OK	OK	OK	–	OK	Best	Best	OK	–	–
Best window												
Workable – slower to re-establish or needs more care												
Avoid / low success												

Plant snapshot

Habit	Densely tufted perennial tussock grass, 0.15–0.9 m, with fine blue-green leaves.
Flowering	Oct – Nov (green to yellow).
Habitat	Sand, granite and sandy loam; coastal areas, estuaries and dunes.
Local note	Recorded in the Perth area (Cottesloe, Stirling); FloraBase has no Joondalup or Wanneroo specimens. Tolerates salt spray and saline soils once established.

Propagation options

Method	Suitability	Notes	Details in
Cuttings	Not used	Grasses are not grown from cuttings.	–
Seed	Main method	Chill 1-2 weeks; sow at 15°C or more.	→ Seed Sowing Guide
Division	Good	Divide mature clumps in early spring.	This guide

Division method

Item	Details
Best time to divide	Aug – Sep (early spring). Mar – May, Jul and Oct are workable. Avoid Nov – Feb (heat).
Division type	Tussock division.
Division size	Clumps of 3-5 tillers with roots. Very small pieces establish poorly.
Selecting stock	Mature, crowded nursery clumps – these benefit from dividing.
Preparation	- Water the stock plant the day before. - Tease or cut the clump apart with a sharp knife. - Trim the leaves back by about half.
Potting	Pot into a free-draining mix at the original depth.
Environment	Shade house until new leaves appear.
Time to establish	Roughly 4-6 weeks (record yours).
Aftercare	- Harden off, then grow on in full sun. - Plant out in winter.

Common problems

Problem	Likely cause and fix
Divisions brown off	Pieces too small or dried out. Bigger pieces; work in the shade.
Base rots	Too wet or planted too deep.
Slow to re-shoot	Divided in the heat. Divide in early spring.

Notes

Difficulty	Easy-Moderate (provisional). Tussock grasses usually divide readily; untested for this species.
Confidence	Low. Only a commercial grower's note: divide mature clumps in early spring. Method follows general tussock-grass practice.
Sources	See References: 1, 3, 17, 18

Nursery records

Log each batch of divisions here to confirm or update the timing and difficulty above.

Date divided	Stock plant / provenance	No. pieces	No. established	Success %	Notes

Spinifex hirsutus

Hairy Spinifex

Family
Poaceae

DIFFICULTY	DIVISION TYPE	BEST MONTHS	DIVISION SIZE	TIME TO ESTABLISH
Moderate-Difficult *	Tussock division	Mar – May	Rooted clumps with runners	Slow – several weeks

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Division	OK	OK	Best	Best	Best	–	–	–	OK	OK	OK	OK
	Best window	Workable – slower to re-establish or needs more care						Avoid / low success				

Plant snapshot

Habit	Rhizomatous, tussock-forming grass, 0.3–0.9 m, with long runners and silvery, hairy leaves.
Flowering	Oct – Jan. Male and female flowers on separate plants.
Habitat	White sand on coastal dunes.
Local note	Recorded in the City of Joondalup. Dune plants are protected – use nursery stock or approved salvage.

Propagation options

Method	Suitability	Notes	Details in
Cuttings	Good	Runner (stolon) sections with 1–2 nodes.	→ Cuttings Guide
Seed	Main method	The usual nursery method; cover seed (needs dark).	→ Seed Sowing Guide
Division	Possible	Tussock divisions work but survived less well than rooted runners.	This guide

Division method

Item	Details
Best time to divide	Mar – May, while growing actively, giving plants ready for winter planting. Sep – Feb workable. Avoid Jun – Aug.
Division type	Tussock division – small clumps with rooted runners.
Division size	Small clumps with as much root and rooted runner as possible.
Selecting stock	Vigorous nursery stock plants. Collect male and female plants if you want seed-producing plantings later.
Preparation	- Keep pieces moist and shaded – they dry out fast. - Cut clumps apart with a sharp spade or knife. - Trim leaves back by about half.
Potting	Pot into deep tubes of coarse sand or a very sandy mix. Firm in and water well.
Environment	Shade house or light shade until new growth shows; keep the sand moist, not wet.
Time to establish	Slow – allow several weeks for new roots and shoots. Plants take many months to reach planting size.
Aftercare	- Harden off, then grow on in full sun. - Plant out in late autumn or winter when rain is reliable.

Common problems

Problem	Likely cause and fix
Divisions dry out and die	Roots exposed too long. Work in the shade; keep pieces in damp bags.
Base rots	Mix too wet. Use coarse sand, water less.
Slow establishment	Normal for spinifex. Full sun once hardened; deep tubes.

Notes

Difficulty	Moderate-Difficult (provisional). Divisions survived less well than rooted runner cuttings in trials. Runner cuttings or seed are usually better choices.
Confidence	Medium. Based on <i>S. sericeus</i> trials, where divided tussocks survived less well (31%) than rooted runner cuttings (40%).
Sources	See References: 3, 4

Nursery records

Log each batch of divisions here to confirm or update the timing and difficulty above.

Date divided	Stock plant / provenance	No. pieces	No. established	Success %	Notes

Spinifex longifolius

Beach Spinifex

Family
Poaceae

DIFFICULTY Moderate–Difficult *	DIVISION TYPE Tussock division	BEST MONTHS Mar – May	DIVISION SIZE Rooted clumps with runners	TIME TO ESTABLISH Slow – several weeks
--	--	---	--	--

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Division	OK	OK	Best	Best	Best	–	–	–	OK	OK	OK	OK
	Best window	Workable – slower to re-establish or needs more care						Avoid / low success				

Plant snapshot

Habit	Rhizomatous, tussock-forming grass, 0.3–1 m high and up to 2 m wide, with long runners.
Flowering	Apr – Dec or Jan. Male and female plants are separate.
Habitat	White sand on coastal dunes; a key foredune stabiliser.
Local note	Recorded in the Wanneroo LGA. Dune plants are protected – use nursery stock or approved salvage.

Propagation options

Method	Suitability	Notes	Details in
Cuttings	Good	Runner (stolon) sections with 1–2 nodes.	→ Cuttings Guide
Seed	Main method	The usual nursery method; smoke water, about 10 days.	→ Seed Sowing Guide
Division	Possible	Tussock divisions work but survived less well than rooted runners.	This guide

Division method

Item	Details
Best time to divide	Mar – May, while growing actively, giving plants ready for winter planting. Sep – Feb workable. Avoid Jun – Aug.
Division type	Tussock division – small clumps with rooted runners.
Division size	Small clumps with as much root and rooted runner as possible .
Selecting stock	Vigorous nursery stock plants.
Preparation	- Keep pieces moist and shaded – they dry out fast. - Cut clumps apart with a sharp spade or knife. - Trim leaves back by about half.
Potting	Pot into deep tubes of coarse sand or a very sandy mix. Firm in and water well.
Environment	Shade house or light shade until new growth shows; keep the sand moist, not wet.
Time to establish	Slow – allow several weeks for new roots and shoots.
Aftercare	- Harden off, then grow on in full sun. - Plant out in late autumn or winter when rain is reliable.

Common problems

Problem	Likely cause and fix
Divisions dry out and die	Roots exposed too long. Work in the shade; keep pieces in damp bags.
Base rots	Mix too wet. Use coarse sand, water less.
Slow establishment	Normal for spinifex. Full sun once hardened; deep tubes.

Notes

Difficulty	Moderate-Difficult (provisional). Divisions survived less well than rooted runner cuttings in trials. Runner cuttings or seed are usually better choices.
Confidence	Medium. Based on <i>S. sericeus</i> trials, where divided tussocks survived less well (31%) than rooted runner cuttings (40%).
Sources	See References: 3, 4

Nursery records

Log each batch of divisions here to confirm or update the timing and difficulty above.

Date divided	Stock plant / provenance	No. pieces	No. established	Success %	Notes

References

Numbers match the source list on each species page. Web addresses shortened; accessed September–October 2026.

#	Source	Address
1	Australian Native Plants Society (ANPSA) – Plant Propagation: division and layering (grass-like clumps, suckers, self-layering plants; division size, timing, hardening off)	anpsa.org.au/plant-propagation/
2	ANPSA – Kangaroo Paws & Relatives: propagation by division (autumn, 3+ shoots, trim leaves by a third)	anpsa.org.au/genera/kangaroo-paw-family/
3	FloraBase, WA Herbarium (DBCA) – species profiles (descriptions, flowering, habitat, local records)	florabase.dbca.wa.gov.au
4	NZ Department of Conservation – Rehabilitation of coastal foredunes using indigenous sand-binding species (<i>Spinifex sericeus</i> division vs stolon cuttings)	doc.govt.nz/Documents/science-and-technical/sfc122.pdf
5	Wikipedia – <i>Anigozanthos manglesii</i> (short-lived habit, ink disease)	en.wikipedia.org/wiki/Anigozanthos_manglesii
6	Ozbreed – How to divide <i>Lomandras</i> (late winter to early spring; trim leaves)	ozbreed.com.au/how-to-divide-lomandras/
7	Cottesloe Coastcare Association – <i>Lomandra maritima</i> (separate male and female plants)	cottesloecoastcare.org/lomandra-maritima/
8	DBCA Library – graceful sun-moth breeds on <i>Lomandra maritima</i> and <i>L. hermaphrodita</i>	library.dbca.wa.gov.au/static/Journals/080525/080525-41.pdf
9	Coastal Restoration Trust of NZ – <i>Ficinia nodosa</i> (wīwī) technical note: division size, survival	coastalrestorationtrust.org.nz/site/assets/files/1185/8.4_wiwi.pdf
10	NZ Plant Conservation Network – <i>Ficinia nodosa</i> (easily grown from seed and division)	nzpcn.org.nz/flora/species/ficinia-nodosa/
11	US Plant Patent PP30,800 – <i>Lepidosperma gladiatum</i> 'LEP01' (first propagated by crown division in Joondalup; tissue culture)	patents.google.com/patent/USPP30800
12	Australian Plants Society NSW – <i>Conostylis aculeata</i> (rooted runners after flowering; 2–3 week window)	resources.austplants.com.au/plant/conostylis-aculeata/
13	Gardening With Angus – <i>Conostylis aculeata</i> (division of rhizomes)	gardeningwithangus.com.au/conostylis-aculeata-prickly-conostylis/
14	Murdoch University CPSM (2009) – Native garden plants resistant to dieback (<i>Conostylis aculeata</i>)	cpsm-phytophthora.org/downloads/CPSM_resistanceBrochures.pdf
15	ANPSA – <i>Anigozanthos humilis</i> profile (divide after flowering, only vigorous plants; short-lived; rust)	anpsa.org.au/plant_profiles/anigozanthos-humilis/
16	Australian National Botanic Gardens – <i>Austrostipa ramosissima</i> (division in late winter – early spring has high success)	anbg.gov.au/gnp/trainees-2016/austrostipa-ramosissima.html
17	Peppy Plants – <i>Poa poiformis</i> (divide mature clumps in early spring; commercial grower, low authority)	peppyplants.com.au/poa-poiformis-the-native-ornamental-and-adaptable-blue-tussock-grass/
18	Australian Plants Online – <i>Poa poiformis</i> (salt spray and saline soils)	australianplantsonline.com.au/poa-poiformis-coastal-tussock-grass.html

Revision log

Version	Date	Changes
1.0	Sep 2026	First edition: Anigozanthos manglesii, Austrostipa flavescens, Dianella revoluta, Spinifex hirsutus, S. longifolius. Includes a list of other species in the Cuttings and Seed guides that can be divided or layered.
1.1	Sep 2026	Added 4 species: Conostylis aculeata, Ficinia nodosa, Lepidosperma gladiatum, Lomandra maritima.
1.2	Sep 2026	Added 3 species: Anigozanthos humilis, Austrostipa elegantissima, Poa poiformis.
1.3	Oct 2026	Added Orthrosanthus laxus and Patersonia occidentalis to 'Other species that can be divided or layered'.