

Native Plant Cuttings Guide

Western Australian native species – Joondalup, WA

Version 1.1 · September 2026 · 15 species

Species in this edition (alphabetical)

Full index on page 2

Eremophila glabra
Tar Bush / Common Emu Bush

p. 6

Easy

Grevillea vestita

p. 8

Easy

Hibbertia hypericoides
Yellow Buttercups

p. 10

Difficult

Hibbertia racemosa
Stalked Guinea Flower

p. 12

Moderate *

Lechenaultia linarioides
Yellow Leschenaultia

p. 14

Easy *

Myoporum insulare
Blueberry Tree / Common Boobialla

p. 16

Easy

Phyllanthus calycinus
False Boronia

p. 18

Moderate *

Scaevola canescens
Grey Scaevola

p. 20

Easy-Moderate *

Scaevola crassifolia
Thick-leaved Fan-flower

p. 22

Easy

Scaevola globulifera

p. 24

Easy *

Scaevola repens

p. 26

Easy *

Scaevola thesioides

p. 28

Easy-Moderate *

Spinifex hirsutus
Hairy Spinifex

p. 30

Moderate *

Spinifex longifolius
Beach Spinifex

p. 32

Moderate *

Threlkeldia diffusa
Coast Bonefruit

p. 34

Moderate *

* Provisional rating – based on genus behaviour; confirm with your own nursery records.

Compiled from published sources (see References). Windows and ratings are starting points for local practice.

How to use this guide

Each species has its own page set, listed alphabetically by scientific name. Every page follows the same layout: quick facts, a cutting calendar, a plant snapshot, the full propagation method, common problems, and a blank record table so the nursery can log strike rates and refine the advice over time.

Timing windows are written for the Joondalup / Perth coastal climate: hot, dry summers and mild, wet winters. They combine general native-propagation practice with each species' flowering period from FloraBase. Rooting is fastest in warm weather and slows markedly in winter unless bottom heat is used.

Species index – at a glance

Species	Page	Difficulty	Wood type	Best months	Length	Hormone (IBA)
<i>Eremophila glabra</i> Tar Bush / Common Emu Bush	6	Easy	Semi-hardwood	Jan – Apr	75–100 mm	3 g/L IBA
<i>Grevillea vestita</i>	8	Easy	Semi-hardwood	Dec – Apr	80–120 mm	3 g/L IBA
<i>Hibbertia hypericoides</i> Yellow Buttercups	10	Difficult	Semi-hardwood	Dec – Feb	50–75 mm	3 g/L IBA (trial range)
<i>Hibbertia racemosa</i> Stalked Guinea Flower	12	Moderate *	Semi-hardwood	Jan – Apr	50–75 mm	1.5–3 g/L IBA
<i>Lechenaultia linarioides</i> Yellow Leschenaultia	14	Easy *	Softwood – semi-hardwood	Feb – Apr	50–75 mm	1.5–3 g/L IBA
<i>Myoporum insulare</i> Blueberry Tree / Common Boobialla	16	Easy	Semi-hardwood	Feb – Apr	75–100 mm	3 g/L IBA
<i>Phyllanthus calycinus</i> False Boronia	18	Moderate *	Semi-hardwood	Feb – Apr	50–75 mm	3 g/L IBA
<i>Scaevola canescens</i> Grey Scaevola	20	Easy–Moderate *	Softwood – semi-hardwood	Nov – Jan	50–80 mm	1.5–3 g/L IBA
<i>Scaevola crassifolia</i> Thick-leaved Fan-flower	22	Easy	Semi-hardwood	Mar – May	75–100 mm	3 g/L IBA
<i>Scaevola globulifera</i>	24	Easy *	Softwood – semi-hardwood	Jan – Apr	50–80 mm	1.5–3 g/L IBA
<i>Scaevola repens</i>	26	Easy *	Softwood – semi-hardwood	Mar – May	50–80 mm	1.5–3 g/L IBA
<i>Scaevola thesioides</i>	28	Easy–Moderate *	Softwood – semi-hardwood	Apr – May	50–75 mm	1.5–3 g/L IBA
<i>Spinifex hirsutus</i> Hairy Spinifex	30	Moderate *	Runner (stolon) sections	Mar – May	10–15 cm, 1–2 nodes	Optional (1.5 g/L)
<i>Spinifex longifolius</i> Beach Spinifex	32	Moderate *	Runner (stolon) sections	Mar – May	10–15 cm, 1–2 nodes	Optional (1.5 g/L)
<i>Threlkeldia diffusa</i> Coast Bonefruit	34	Moderate *	Softwood tips	Dec – Mar	50–75 mm	1.5 g/L IBA

* Provisional rating – based on genus behaviour; update once nursery strike rates are recorded.

Cutting calendar

Species	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
<i>Eremophila glabra</i>	Best	Best	Best	Best	OK	–	–	–	–	–	OK	OK
<i>Grevillea vestita</i>	Best	Best	Best	Best	OK	–	–	–	–	–	OK	Best
<i>Hibbertia hypericoides</i>	Best	Best	OK	–	–	–	–	–	–	–	OK	Best
<i>Hibbertia racemosa</i>	Best	Best	Best	Best	OK	–	–	–	–	–	–	OK
<i>Lechenaultia linarioides</i>	OK	Best	Best	Best	OK	–	–	–	–	OK	OK	OK
<i>Myoporum insulare</i>	OK	Best	Best	Best	OK	–	–	–	–	OK	OK	OK
<i>Phyllanthus calycinus</i>	OK	Best	Best	Best	OK	–	–	–	–	–	–	–
<i>Scaevola canescens</i>	Best	OK	OK	OK	OK	–	–	–	–	OK	Best	Best
<i>Scaevola crassifolia</i>	OK	OK	Best	Best	Best	–	–	–	–	OK	OK	OK
<i>Scaevola globulifera</i>	Best	Best	Best	Best	OK	–	–	–	–	–	OK	OK
<i>Scaevola repens</i>	OK	OK	Best	Best	Best	–	–	–	OK	OK	OK	OK
<i>Scaevola thesioides</i>	OK	OK	OK	Best	Best	–	–	–	OK	OK	OK	OK
<i>Spinifex hirsutus</i>	OK	OK	Best	Best	Best	–	–	–	OK	OK	OK	OK
<i>Spinifex longifolius</i>	OK	OK	Best	Best	Best	–	–	–	OK	OK	OK	OK
<i>Threlkeldia diffusa</i>	Best	Best	Best	OK	–	–	–	–	–	–	OK	Best

Perth season

Summer

Autumn

Winter

Spring

Sum.

 Best window

 Workable – see ‘Selecting material’ in the method

 Avoid / low success

Key to terms and ratings

Wood types and the bend test

Wood type	What it looks like	Bend test	Use
Softwood	Soft, flexible, bright new growth from the current flush.	Flops or stays bent.	Roots fast but wilts and rots easily. Suits soft-stemmed plants such as <i>Scaevola</i> .
Semi-hardwood	Current season's growth that has begun to firm; bark starting to colour.	Bends about 60° and springs back.	The standard for most WA native shrubs. More forgiving than softwood.
Hardwood	Fully mature, woody stems from previous seasons.	Stiff; snaps rather than bends.	Slow to root. Rarely needed for the species in this guide.

Difficulty ratings

Rating	Meaning
Easy	Strikes reliably with the standard method.
Easy-Moderate	Usually strikes; one factor (e.g. moisture, fragile stems) needs extra care.
Moderate	Strikes, but results depend on timing, material and moisture control. Set extra cuttings.
Moderate-Difficult	Low or variable strike; needs careful timing and good facilities.
Difficult	Low or erratic strike; specialist techniques or alternative methods usually needed.

* **Provisional** – rating is based on how the genus generally behaves because little species-specific data is published. Replace it with the nursery's own figure once a few batches have been recorded.

Confidence

Level	Meaning
High	Species-specific propagation information is published and consistent.
Medium	Built from genus-level information plus the species' own traits (habit, leaf type, flowering time). Sound starting point – verify with trials.
Low	Little information available; treat as a trial protocol.

Standard cutting method

Species pages refer back to these steps and only note where a species differs.

Step	What to do
1. Collect	Take material early in the morning from healthy, pest-free, well-watered, non-flowering stock. Seal in a moist plastic bag and keep in an esky out of the sun. Process the same day.
2. Prepare	Use a sharp, clean blade (wipe with alcohol between stock plants). Cut just below a node. Strip leaves from the lower half to two-thirds without tearing the bark. Reduce large leaves by about one-third. Remove flowers and buds.
3. Wound (optional)	On firmer stems, remove a thin sliver of bark from the bottom 15–20 mm to expose more rooting area.
4. Hormone	Decant a small amount into a clean container, dip the base to about 1 cm, and discard what's left. Store hormone in the fridge – it has a limited shelf life.
5. Set	Make a hole with a dibber so the hormone isn't wiped off, insert to the bottom of the hole, and firm the mix. Single tubes make potting on easier than community pots.
6. Environment	Water in and place in a humid frame or propagation house in bright shade. Bottom heat so the base is 5–10°C warmer than the air. Keep the mix moist, not wet. Mist sparingly on hairy-leaved species.
7. Check	Most cuttings strike in 4–8 weeks in warm weather. Look for roots at the drainage holes. Remove any rotting cuttings promptly.
8. Harden off	About 2 weeks with cover closed by day and open at night, then about 2 weeks in shade, then into the open. Many struck cuttings are lost by skipping this step.
9. Pot on	After 1–3 months, pot on into larger containers. Don't let roots become pot-bound.

Rooting hormone guide

Product (example)	IBA strength	Use for
Clonex Green	1.5 g/L	Softwood and soft tip cuttings
Clonex Purple	3 g/L	Semi-hardwood – the general all-rounder
Clonex Red	8 g/L	Hardwood cuttings

Equivalent powders or liquids work too – match the strength to the wood type. Callus forming without roots usually means the hormone is too strong for that material.

Mix, hygiene and collecting

Topic	Guidance
Propagation mix	About 75–80% washed coarse river sand with 20–25% coir or peat. Add extra perlite or coarse sand for rot-prone species.
Hygiene	Fresh mix, clean tubes and benches. Many WA natives (including <i>Hibbertia</i> and <i>Grevillea</i>) are susceptible to <i>Phytophthora</i> . Discard diseased cuttings; use fungicide only if a problem persists.
Collecting	Collecting from bushland generally requires a DBCA flora licence and landholder permission. Use local-provenance stock and label every batch with its source.

Eremophila glabra

Tar Bush / Common Emu Bush

Family
Scrophulariaceae

DIFFICULTY		WOOD TYPE				BEST MONTHS		CUTTING LENGTH			HORMONE (IBA)		
Easy		Semi-hardwood				Jan - Apr		75-100 mm			3 g/L IBA		
		Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Cuttings		Best	Best	Best	Best	OK	-	-	-	-	-	OK	OK
Best window		Workable – see ‘Selecting material’ in the method						Avoid / low success					

Plant snapshot

Habit	Very variable, from prostrate to 3 m. Perth coastal plants are mostly subsp. <i>albicans</i> : low, spreading, to about 0.4 m high and 2 m wide, with grey, felty leaves.
Flowering	Species: Mar – Dec. Subsp. <i>albicans</i> : May – Jan. Tubular flowers, yellow, orange, red or green.
Habitat	Sand and limestone; dunes, sandplains and winter-wet flats, sometimes saline.
Local note	Recorded in the Wanneroo LGA and along the Perth coast. Several subspecies and many forms – label stock by subspecies and form.

Propagation method

Item	Details
Best time to take cuttings	Jan – Apr, using hardened current season's growth while the weather is warm and plants are mostly out of flower. Nov – Dec and May are workable. Avoid Jun – Oct (cool weather, flowering).
Wood type	Semi-hardwood – hardened current season's growth.
Cutting size	75–100 mm.
Selecting material	Non-flowering tips or side shoots. On grey-leaved forms pick clean foliage with no sign of mould.
Pre-treatment	- Carefully remove leaves from the lower two-thirds. - Wound: remove a thin sliver of bark from the base. - Dip the base in rooting hormone.
Rooting hormone	Semi-hardwood strength, e.g. Clonex Purple (3 g/L IBA).
Propagation mix	Very free-draining, sand-based mix.
Environment	Grey, felty-leaved forms rot under heavy mist or high humidity – keep foliage as dry as possible, water the mix, give good airflow. Green-leaved forms tolerate normal misting. Bottom heat speeds rooting.
Expected strike time	About 4–8 weeks in warm weather.
Post-treatment / aftercare	- Standard hardening off. - Pot on into a free-draining mix and grow on in full sun. - Keep water off grey foliage; water in the morning.

Common problems

Problem	Likely cause and fix
Leaves blacken or grow grey mould	Too humid. Cut back misting, open the frame, remove affected leaves.
Base rotting	Mix too wet. Water less often and use a sandier mix.
Cuttings flower instead of rooting	Use non-flowering shoots and pinch off buds.

Notes

Difficulty	Easy. Most forms of <i>E. glabra</i> strike readily from cuttings.
Other methods	Seed – unreliable in <i>Eremophila</i> (hard fruits, slow and erratic germination). Grafting is used for difficult eremophilas but isn't needed for this species.
Confidence	High. Published sources agree most forms strike readily from cuttings.
Sources	See References: 1, 4, 12, 13, 20, 21, 22, 30

Nursery records

Log each batch here to confirm or update the timing and difficulty above.

Date set	Stock source / provenance	No. set	No. struck	Strike %	Notes

Grevillea vestita

No widely accepted common name

Family
Proteaceae

DIFFICULTY	WOOD TYPE	BEST MONTHS	CUTTING LENGTH	HORMONE (IBA)
Easy	Semi-hardwood	Dec – Apr	80–120 mm	3 g/L IBA

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Cuttings	Best	Best	Best	Best	OK	–	–	–	–	–	OK	Best

Best window
Workable – see 'Selecting material' in the method
Avoid / low success

Plant snapshot

Habit	Erect, spreading, prickly shrub, usually 0.5–3 m. Hairy branchlets; lobed leaves with sharp-pointed tips.
Flowering	Mainly Jun – Nov (white to cream, often pink-tinged, scented). Odd flowers Jan – Apr.
Habitat	Sandy soils over limestone or granite. Occurs on the Swan Coastal Plain, including the Perth area.
Local note	Two subspecies exist (subsp. <i>vestita</i> and subsp. <i>isopogoides</i>). Label stock plants by subspecies and provenance.

Propagation method

Item	Details
Best time to take cuttings	Dec – Apr (late spring to autumn), once the spring flush has firmed and flowering has finished. Nov and May are workable. Avoid Jun – Oct: plants are flowering, new growth is soft and cool weather slows rooting.
Wood type	Semi-hardwood – firm, current season's growth. It should bend about 60° and spring back.
Cutting size	80–120 mm long with 3–5 nodes. Tip or sub-tip pieces; two cuttings can often be taken from one shoot.
Selecting material	Healthy, vigorous, non-flowering shoots. Skip shoots carrying buds, old woody stems and soft new tips.
Pre-treatment	- Wear gloves: leaves are prickly and grevillea sap can irritate skin. - Remove any flowers or buds. - Strip leaves from the lower half to two-thirds (cut, don't tear the bark). - Trim remaining lobed leaves by about one-third. - Wound: remove a thin sliver of bark from the bottom 15–20 mm. - Dip the base in rooting hormone.
Rooting hormone	Semi-hardwood strength, e.g. Clonex Purple (3 g/L IBA) . If cuttings form a heavy callus but no roots, trim the callus back to clean wood and re-set; try a lower strength on the next batch.
Propagation mix	Free-draining, low nutrient: about 75–80% washed coarse river sand + 20–25% coir or peat (or a perlite/coir mix).
Environment	Humid propagation house or frame in bright shade. Bottom heat with the base of the cutting 5–10°C warmer than the air. Light misting only – the hairy foliage can rot if kept constantly wet.
Expected strike time	About 6–8 weeks in warm weather. Slower in the cooler months.
Post-treatment / aftercare	- Harden off: ~2 weeks with cover closed by day and open at night, then ~2 weeks in shade before full sun. - Pot on when roots show at the drainage holes, into a low-phosphorus native mix. - Feed only with a low-phosphorus native fertiliser (Proteaceae are phosphorus-sensitive). - Responds well to pruning – tip-prune young plants for a bushier tube.

Common problems

Problem	Likely cause and fix
Heavy callus, no roots	Trim callus back and re-set. Lower the hormone strength next time.
Base rotting	Mix too wet or material too soft. Use firmer wood and water less often.
Leaf yellowing or scorch after potting on	Check phosphorus in the potting mix and fertiliser.

Notes

Difficulty	Easy. Cuttings from firm current season's growth usually strike reliably.
Other methods	Seed – grows reasonably easily from seed, but seedlings are not identical to the parent. Grafting onto <i>G. robusta</i> is used on the humid east coast; not needed locally.
Confidence	High. Species-specific propagation information is published and consistent across sources.
Sources	See References: 1, 2, 3, 4, 13, 15, 17

Nursery records

Log each batch here to confirm or update the timing and difficulty above.

Date set	Stock source / provenance	No. set	No. struck	Strike %	Notes

Hibbertia hypericoides

Yellow Buttercups

Family
Dilleniaceae

DIFFICULTY	WOOD TYPE	BEST MONTHS	CUTTING LENGTH	HORMONE (IBA)
Difficult	Semi-hardwood	Dec – Feb	50–75 mm	3 g/L IBA (trial range)

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Cuttings	Best	Best	OK	–	–	–	–	–	–	–	OK	Best

Best window	Workable – see ‘Selecting material’ in the method	Avoid / low success
-------------	---	---------------------

Plant snapshot

Habit	Spreading shrub, 0.3–0.8 m. Densely hairy branches; small leaves with rolled edges and white-hairy undersides.
Flowering	Apr – Nov (bright yellow, notched petals).
Habitat	Sandy soils in Banksia woodland and shrubland; one of the most common understorey plants between Perth and Albany. Resprouts from its rootstock after fire.
Local note	Recorded in the City of Joondalup.

Propagation method

Item	Details
Best time to take cuttings	Dec – Feb, after flowering finishes and once the new flush has started to firm. Nov and Mar are workable. Avoid Apr – Oct (flowering period).
Wood type	Semi-hardwood – new growth that is just firming. Soft tips wilt quickly; old wood rarely roots.
Cutting size	50–75 mm.
Selecting material	Vigorous non-flowering shoots. Well-watered, actively growing stock plants (e.g. nursery-held stock) usually give better material than stressed bushland plants.
Pre-treatment	- Strip lower leaves by cutting, not pulling. - Light wound on the firmer stems. - Dip the base in rooting hormone.
Rooting hormone	Start with 3 g/L IBA (Clonex Purple). Because results are unreliable, split trial batches across 1.5, 3 and 8 g/L (Green, Purple, Red) and record which works best.
Propagation mix	Very free-draining, fresh, clean mix. Strict hygiene – hibbertias are prone to <i>Phytophthora</i> root rot.
Environment	Humid frame, bottom heat, bright shade. Mist lightly – the hairy leaf undersides hold water. Good airflow.
Expected strike time	Slow and variable – allow 8–16 weeks . Leave cuttings in as long as they stay green and healthy.
Post-treatment / aftercare	- Harden off slowly (extend each stage if cuttings wilt). - Pot on gently, disturbing roots as little as possible. - Avoid overwatering; keep benches clean.

Common problems

Problem	Likely cause and fix
Cuttings stay green but don't root	Leave longer; check for heavy callus and trim it; try a different hormone strength.
Cuttings collapse or rot	Too wet or root rot. Improve drainage and hygiene; remove affected cuttings.
Rooted cuttings die after potting on	Roots damaged or hardening too fast. Use single tubes and harden off more slowly.

Notes

Difficulty	Difficult. DBCA notes it is difficult to grow from seed or cuttings, and specialist Perth nurseries describe it as a species other producers often fail with. Set large batches and keep detailed records.
Other methods	Seed – also difficult: WA <i>Hibbertia</i> seed has complex dormancy, and treatments are still the subject of research. If numbers are needed quickly, consider buying in tubestock from a specialist grower.
Confidence	Low. Its difficulty is well documented (DBCA; specialist WA nurseries), but no detailed cutting protocol is published. Treat the method below as a trial protocol.
Sources	See References: 1, 4, 10, 11, 27, 30, 31

Nursery records

Log each batch here to confirm or update the timing and difficulty above.

Date set	Stock source / provenance	No. set	No. struck	Strike %	Notes

Hibbertia racemosa

Stalked Guinea Flower

Family
Dilleniaceae

DIFFICULTY
Moderate *

WOOD TYPE
Semi-hardwood

BEST MONTHS
Jan – Apr

CUTTING LENGTH
50–75 mm

HORMONE (IBA)
1.5–3 g/L IBA

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Cuttings	Best	Best	Best	Best	OK	–	–	–	–	–	–	OK
Best window												
Workable – see 'Selecting material' in the method												
Avoid / low success												

Plant snapshot

Habit	Small erect or ascending, spreading shrub, 0.1–0.75 m.
Flowering	Jul – Dec (yellow, on long stalks).
Habitat	Grey, white or yellow sand in coastal areas: dunes, plains and limestone.
Local note	Recorded in the City of Joondalup. Grey-leaved and prostrate forms are sold in the trade – keep forms separate and labelled.

Propagation method

Item	Details
Best time to take cuttings	Jan – Apr , after flowering finishes and new growth has begun to firm. Dec and May are workable. Avoid Jun – Nov (cool weather, then flowering).
Wood type	Semi-hardwood – new growth that has started to harden. Firm tip cuttings also work.
Cutting size	50–75 mm long. Stems are fine and leaves small, so keep cuttings short.
Selecting material	Non-flowering side shoots from healthy plants. Snip off any flower stalks or buds.
Pre-treatment	- Strip leaves from the lower half (cut rather than pull on fine stems). - Wounding not needed on fine stems; a light wound is optional on firmer material. - Dip the base in rooting hormone.
Rooting hormone	Semi-hardwood strength (3 g/L IBA, e.g. Clonex Purple). For softer tips use softwood strength (1.5 g/L IBA, e.g. Clonex Green).
Propagation mix	Very free-draining, sand-based mix. Use fresh, clean mix and clean tubes – hibbertias are prone to <i>Phytophthora</i> root rot.
Environment	Humid frame, bright shade, bottom heat. Keep the mix moist but never wet; good airflow.
Expected strike time	Roughly 4–8 weeks in warm weather; may be slower. Record your results.
Post-treatment / aftercare	- Standard hardening off (2 weeks closed by day/open at night, then 2 weeks shade). - Pot on into a free-draining native mix; avoid overwatering. - Keep good hygiene on benches to limit root rot.

Common problems

Problem	Likely cause and fix
Cuttings collapse at the base	Too wet, or root rot. Improve drainage, use clean mix, remove affected cuttings.
Few roots, cuttings flower	Material was taken from flowering shoots. Use non-flowering side shoots.
Slow to strike in winter	Use bottom heat, or wait for the Jan – Apr window.

Notes

Difficulty	Moderate (provisional). Hibbertias are generally easy from cuttings, but some WA species (e.g. <i>H. hypericoides</i>) are known to be difficult. Set larger batches until your own strike rate is known.
Other methods	Seed – generally unreliable in <i>Hibbertia</i> (complex dormancy). Layering – several hibbertias root where stems touch the ground; check low stems on prostrate forms and pot up any rooted pieces.
Confidence	Medium. Little species-specific data published. Based on how the genus behaves and this species' growth cycle.
Sources	See References: 1, 4, 8, 9, 10, 11

Nursery records

Log each batch here to confirm or update the timing and difficulty above.

Date set	Stock source / provenance	No. set	No. struck	Strike %	Notes

Lechenaultia linarioides

Yellow Leschenaultia

Family
Goodeniaceae

DIFFICULTY	WOOD TYPE	BEST MONTHS	CUTTING LENGTH	HORMONE (IBA)
Easy *	Softwood - semi-hardwood	Feb - Apr	50-75 mm	1.5-3 g/L IBA

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Cuttings	OK	Best	Best	Best	OK	-	-	-	-	OK	OK	OK
	Best window	Workable - see 'Selecting material' in the method										

Plant snapshot

Habit	Sprawling, open shrub to 1.5 m, with narrow, linear leaves.
Flowering	Can flower in any month. Flowers yellow with red or pink tones, sometimes cream.
Habitat	Sand (often over limestone) and sandy clay; sandplains, low dunes, ridges and winter-wet flats.
Local note	Recorded in the Wanneroo LGA.

Propagation method

Item	Details
Best time to take cuttings	Feb - Apr (late summer to autumn) while it is warm. One grower reports roots in 4-6 weeks from late-February leschenaultia cuttings. Oct - Jan and May are workable. Avoid Jun - Sep.
Wood type	Softwood to semi-hardwood tip cuttings.
Cutting size	50-75 mm.
Selecting material	Non-flowering tips. This species can flower in any month, so check each cutting and remove buds.
Pre-treatment	- Strip leaves from the lower half (narrow leaves strip easily). - No wounding needed. - Dip the base in rooting hormone.
Rooting hormone	1.5 g/L IBA (Clonex Green) for soft tips; 3 g/L (Clonex Purple) for firmer material.
Propagation mix	Very free-draining - leschenaultias dislike wet roots.
Environment	Humid frame, bottom heat, bright shade. Keep the mix moist, never wet.
Expected strike time	About 4-6 weeks in warm weather.
Post-treatment / aftercare	- Standard hardening off. - Pot on into a free-draining mix; water sparingly. - Tip-prune to counter the sprawling habit.

Common problems

Problem	Likely cause and fix
Base rotting	Overwatering. Water less often; add coarse sand or perlite to the mix.
Wilting in the frame	Material too soft or humidity too low. Use slightly firmer tips; keep the frame closed.
Cuttings flowering	Remove buds; pick vegetative shoots.

Notes

Difficulty	Easy (provisional). <i>Lechenaultia</i> cuttings generally strike very easily; seed is difficult.
Other methods	Seed – slow and unreliable. The Geraldton community nursery soaked seed in smoke water, saw germination at about 46 days, and kept trays for a second year. Division – some <i>Lechenaultias</i> sucker; pot up rooted suckers.
Confidence	Medium. No species-specific cutting data found. Based on the genus, whose cuttings strike very easily.
Sources	See References: 1, 4, 12, 13, 24, 25

Nursery records

Log each batch here to confirm or update the timing and difficulty above.

Date set	Stock source / provenance	No. set	No. struck	Strike %	Notes

Myoporum insulare

Blueberry Tree / Common Boobialla

Family
Scrophulariaceae

DIFFICULTY	WOOD TYPE				BEST MONTHS		CUTTING LENGTH			HORMONE (IBA)		
Easy	Semi-hardwood				Feb – Apr		75–100 mm			3 g/L IBA		
	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Cuttings	OK	Best	Best	Best	OK	–	–	–	–	OK	OK	OK
Best window	Workable – see ‘Selecting material’ in the method							Avoid / low success				

Plant snapshot

Habit	Dense, low spreading to erect shrub or small tree, 0.25–5 m, with thick, fleshy leaves.
Flowering	Jul – Dec, also Jan – Feb (white, purple-spotted), followed by purple berries.
Habitat	White or grey sand on coastal dunes and limestone.
Local note	Recorded in Joondalup and Wanneroo. Forms range from prostrate to tree-like – take cuttings from a plant with the form you want.

Propagation method

Item	Details
Best time to take cuttings	Feb – Apr – warm, and after the main flowering. Oct – Jan and May are workable. Avoid Jun – Sep.
Wood type	Semi-hardwood – firm young growth.
Cutting size	75–100 mm.
Selecting material	Firm, non-flowering young shoots from a stock plant with the growth form you need (prostrate for groundcover, erect for screening).
Pre-treatment	- Strip leaves from the lower half. - Reduce the remaining thick leaves by about one-third. - Dip the base in rooting hormone.
Rooting hormone	3 g/L IBA (Clonex Purple). Strikes readily without it, but hormone gives faster, more even rooting.
Propagation mix	Standard sand/coir cutting mix.
Environment	Standard humid frame with bottom heat. Hairless leaves tolerate normal misting.
Expected strike time	About 4–8 weeks in warm weather.
Post-treatment / aftercare	- Standard hardening off. - Fast-growing – pot on promptly before roots circle. - Tip-prune for a bushier plant.

Common problems

Problem	Likely cause and fix
Leaf drop in the frame	Too dark or too wet. Give brighter shade and water less.
Roots circling in tubes	Pot on sooner – it grows quickly.
Plants not the expected shape	Seed-grown or mixed stock. Propagate from a known-form stock plant.

Notes

Difficulty	Easy. Firm young growth roots readily; even branchlets placed straight into soil can strike.
Other methods	Seed – the Geraldton community nursery ferments ripe (darkest purple) berries for 4–5 days, washes off the pulp and soaks in smoke water; germination took 15–30 days. Seedlings vary in form.
Confidence	High. Published sources agree it strikes readily from firm young growth.
Sources	See References: 1, 4, 12, 13, 23

Nursery records

Log each batch here to confirm or update the timing and difficulty above.

Date set	Stock source / provenance	No. set	No. struck	Strike %	Notes

Phyllanthus calycinus

False Boronia

Current WA Herbarium name: *Lysiandra calycina*

Family
Phyllanthaceae

DIFFICULTY	WOOD TYPE	BEST MONTHS	CUTTING LENGTH	HORMONE (IBA)
Moderate *	Semi-hardwood	Feb – Apr	50–75 mm	3 g/L IBA

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Cuttings	OK	Best	Best	Best	OK	–	–	–	–	–	–	–
	Best window	Workable – see ‘Selecting material’ in the method						Avoid / low success				

Plant snapshot

Habit	Small erect shrub, 0.2–1.2 m, with smooth branchlets.
Flowering	Jun – Dec or Jan (white-cream to pink). Sepals enlarge to enclose the fruit.
Habitat	Usually sandy soils, including coastal alkaline sands.
Name note	FloraBase now lists this plant as <i>Lysiandra calycina</i> ; many nurseries still sell it as <i>Phyllanthus calycinus</i> . Kept under <i>Phyllanthus</i> here to match your plant list.
Local note	Recorded in Joondalup and Wanneroo.

Propagation method

Item	Details
Best time to take cuttings	Feb – Apr – warm, and after the long flowering season ends. Jan and May are workable. Avoid Jun – Dec (flowering, and cool in winter).
Wood type	Semi-hardwood.
Cutting size	50–75 mm.
Selecting material	Non-flowering shoots from healthy plants.
Pre-treatment	- Strip leaves from the lower half. - Light wound on firmer stems. - Dip the base in rooting hormone.
Rooting hormone	3 g/L IBA (Clonex Purple); trial 1.5 g/L (Clonex Green) on softer tips.
Propagation mix	Free-draining sand-based mix.
Environment	Standard humid frame, bottom heat, bright shade.
Expected strike time	Not documented – start checking from about 6 weeks and record the result.
Post-treatment / aftercare	- Standard hardening off. - Pot on into a free-draining native mix.

Common problems

Problem	Likely cause and fix
No roots after 10+ weeks	Try a different hormone strength or timing on the next batch; record what you tried.
Rotting	Mix too wet. Water less often; improve drainage.
Cuttings flowering	Use non-flowering shoots only.

Notes

Difficulty	Moderate (provisional). No cutting data found for this species; the rating is a cautious starting point.
Other methods	Seed – the method most guides mention; use fresh seed. Few published details on pre-treatment.
Confidence	Low. No published cutting data found. Treat as a trial protocol and record results.
Sources	See References: 1, 4, 29, 30

Nursery records

Log each batch here to confirm or update the timing and difficulty above.

Date set	Stock source / provenance	No. set	No. struck	Strike %	Notes

Scaevola canescens

Grey Scaevola

Family
Goodeniaceae

DIFFICULTY	WOOD TYPE	BEST MONTHS	CUTTING LENGTH	HORMONE (IBA)
Easy-Moderate *	Softwood – semi-hardwood	Nov – Jan	50–80 mm	1.5–3 g/L IBA

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Cuttings	Best	OK	OK	OK	OK	–	–	–	–	OK	Best	Best

Best window	Workable – see 'Selecting material' in the method	Avoid / low success
-------------	---	---------------------

Plant snapshot

Habit	Prostrate to ascending shrub, 0.1–0.5 m. Densely hairy grey leaves; stems are brittle.
Flowering	Mar – Oct (white to white-purple), in spikes in the leaf axils.
Habitat	Sand, often with lateritic gravel or limestone.
Local note	Recorded in the northern Perth area (Wanneroo and Stirling LGAs).

Propagation method

Item	Details
Best time to take cuttings	Nov – Jan, on new growth after flowering ends, while it is warm. Oct and Feb – May are workable if you choose shoots without flower spikes. Avoid Jun – Sep (cool, peak flowering).
Wood type	Softwood to semi-hardwood tip cuttings.
Cutting size	50–80 mm tips.
Selecting material	Vigorous non-flowering tips. Flowers form in the leaf axils along the stem, so check the whole cutting and pinch off any spikes.
Pre-treatment	- Handle gently – stems snap easily. Carry in a rigid tray or box. - Strip lower leaves by cutting, not tearing. - Dip the base in rooting hormone.
Rooting hormone	Softwood strength (1.5 g/L IBA, e.g. Clonex Green) for soft tips; semi-hardwood strength (3 g/L, Clonex Purple) for firmer material.
Propagation mix	Extra free-draining: standard sand/coir mix with extra coarse sand or perlite.
Environment	Keep foliage as dry as possible. Dense hairy leaves rot under heavy mist. Water the mix rather than the leaves, give good airflow, and use bottom heat (root zone about 21–24°C).
Expected strike time	About 4–8 weeks in warm weather.
Post-treatment / aftercare	- Standard hardening off. - Pot on into a free-draining mix; keep water off the foliage. - Remove any leaves showing grey mould promptly.

Common problems

Problem	Likely cause and fix
Grey fuzz or blackened leaves	Too humid/wet. Cut back misting, remove affected material, improve airflow.
Snapped stems	Brittle stems – transport in rigid containers and use a dibber when setting.
Cuttings flowering, not rooting	Take shoots without axillary flower spikes.

Notes

Difficulty	Easy-Moderate (provisional). <i>Scaevola</i> cuttings generally strike very readily; the hairy foliage makes rot the main risk.
Other methods	Seed – <i>Scaevola</i> seed is generally difficult. Division – possible if a plant has suckered or rooted along its stems.
Confidence	Medium. Little species-specific data published. Based on the genus (strikes readily) plus this species' hairy foliage.
Sources	See References: 1, 4, 5, 6, 7, 12, 13, 14, 16

Nursery records

Log each batch here to confirm or update the timing and difficulty above.

Date set	Stock source / provenance	No. set	No. struck	Strike %	Notes

Scaevola crassifolia

Thick-leaved Fan-flower

Family
Goodeniaceae

DIFFICULTY	WOOD TYPE				BEST MONTHS		CUTTING LENGTH			HORMONE (IBA)		
Easy	Semi-hardwood				Mar – May		75–100 mm			3 g/L IBA		
	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Cuttings	OK	OK	Best	Best	Best	–	–	–	–	OK	OK	OK
Best window												

Plant snapshot

Habit	Erect or spreading shrub, 0.1–1.5 m and up to 3 m wide. Young growth is sticky; thick, toothed, egg-shaped to round leaves.
Flowering	Jul – Dec, also Jan – Feb (blue, mauve or white fans).
Habitat	Sand on coastal dunes, plains and limestone cliffs.
Local note	Recorded in Joondalup and Wanneroo. Flower colour and habit vary – cuttings keep the parent's form.

Propagation method

Item	Details
Best time to take cuttings	Mar – May – warm, and outside the main flowering period. Oct – Feb is workable with non-flowering shoots (shade well in summer). Avoid Jun – Sep.
Wood type	Semi-hardwood.
Cutting size	75–100 mm.
Selecting material	Non-flowering tips. Flower spikes carry leafy bracts – check closely and pinch them off.
Pre-treatment	- New growth is sticky – wipe the blade often. - Strip lower leaves; cut the large remaining leaves back by a third to a half. - Dip the base in rooting hormone.
Rooting hormone	3 g/L IBA (Clonex Purple).
Propagation mix	Free-draining sand/coir mix.
Environment	Standard humid frame with bottom heat, bright shade.
Expected strike time	About 4–8 weeks in warm weather.
Post-treatment / aftercare	- Standard hardening off. - Pot on into a free-draining mix. - Tip-prune young plants for dense growth.

Common problems

Problem	Likely cause and fix
Wilting in the first week	Too much leaf area. Trim leaves harder; keep the frame closed.
Rotting	Mix too wet. Water less often; improve drainage.
Cuttings flowering	Use non-flowering tips or wait for the Mar – May window.

Notes

Difficulty	Easy. Semi-hardwood cuttings in the warmer months are the standard method.
Other methods	Seed – large seeds are produced and sometimes germinate naturally; cuttings are preferred to keep flower colour and habit.
Confidence	High. Cuttings are the standard method in WA nurseries (e.g. Geraldton community nursery).
Sources	See References: 1, 4, 5, 12, 13, 26, 30

Nursery records

Log each batch here to confirm or update the timing and difficulty above.

Date set	Stock source / provenance	No. set	No. struck	Strike %	Notes

Scaevola globulifera

No widely used common name

Family
Goodeniaceae

DIFFICULTY	WOOD TYPE				BEST MONTHS			CUTTING LENGTH			HORMONE (IBA)	
Easy *	Softwood - semi-hardwood				Jan - Apr			50-80 mm			1.5-3 g/L IBA	

Notes

Difficulty	Easy (provisional). <i>Scaevola</i> cuttings generally strike very readily.
Other methods	Seed – <i>Scaevola</i> seed is generally difficult. Division – check stock plants for rooted stems.
Confidence	Medium. No species-specific data found. Based on the genus, which strikes readily from cuttings.
Sources	See References: 1, 4, 5, 6, 7, 13, 14

Nursery records

Log each batch here to confirm or update the timing and difficulty above.

Date set	Stock source / provenance	No. set	No. struck	Strike %	Notes

Scaevola repens

No widely used common name ('repens' = creeping)

Family
Goodeniaceae

DIFFICULTY	WOOD TYPE	BEST MONTHS	CUTTING LENGTH	HORMONE (IBA)
Easy *	Softwood – semi-hardwood	Mar – May	50–80 mm	1.5–3 g/L IBA

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Cuttings	OK	OK	Best	Best	Best	–	–	–	OK	OK	OK	OK

Best window	Workable – see 'Selecting material' in the method	Avoid / low success
-------------	---	---------------------

Plant snapshot

Habit	Prostrate to ascending or erect shrub, 0.05–1 m; trailing branches to about 50 cm.
Flowering	Jun – Dec, sometimes Jan – Feb (white or cream, occasionally blue-purple; golden hairs on the outside of the flower).
Habitat	Sand dunes and plains on sand, gravel, laterite or limestone.
Local note	Recorded in the City of Joondalup. A variable species with informal subspecies – use local provenance and keep it labelled.

Propagation method

Item	Details
Best time to take cuttings	Mar – May – warm autumn weather, mostly outside the main flowering period. Sep – Feb is workable using non-flowering shoots (shade well in summer). Avoid Jun – Aug.
Wood type	Softwood to semi-hardwood tip cuttings.
Cutting size	50–80 mm tips from the trailing stems.
Selecting material	Non-flowering tips. Flower spikes sit in the leaf axils – pinch off any you find.
Pre-treatment	- Strip lower leaves. - Leaves can be long (up to 90 mm) – trim remaining leaves by about one-third to cut water loss. - Dip the base in rooting hormone.
Rooting hormone	Softwood strength (1.5 g/L IBA, Clonex Green) for soft tips; 3 g/L (Clonex Purple) for firmer material.
Propagation mix	Free-draining sand/coir mix.
Environment	Humid frame with bottom heat, bright shade. Keep the mix moist but on the drier side rather than wet.
Expected strike time	About 4–8 weeks in warm weather.
Post-treatment / aftercare	- Standard hardening off. - Pot on into a free-draining native mix. - Pinch tips once growing to encourage branching.

Common problems

Problem	Likely cause and fix
Wilting in the first week	Too much leaf area or low humidity. Trim leaves harder; keep the frame closed.
Rotting	Mix too wet. Water less often; improve drainage.
Cuttings flowering, not rooting	Take non-flowering tips, or wait for the Mar – May window.

Notes

Difficulty	Easy (provisional). <i>Scaevola</i> cuttings generally strike very readily.
Other methods	Seed – <i>Scaevola</i> seed is generally difficult. Division – some <i>scaevolae</i> sucker; check stock plants for rooted pieces.
Confidence	Medium. Little species-specific data published. Based on the genus, which strikes readily from cuttings.
Sources	See References: 1, 4, 5, 6, 7, 12, 13, 14, 16

Nursery records

Log each batch here to confirm or update the timing and difficulty above.

Date set	Stock source / provenance	No. set	No. struck	Strike %	Notes

Scaevola thesioides

No widely used common name

Family
Goodeniaceae

DIFFICULTY	WOOD TYPE	BEST MONTHS	CUTTING LENGTH	HORMONE (IBA)
Easy-Moderate *	Softwood - semi-hardwood	Apr - May	50-75 mm	1.5-3 g/L IBA

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Cuttings	OK	OK	OK	Best	Best	-	-	-	OK	OK	OK	OK
	Best window	Workable - see 'Selecting material' in the method										

Plant snapshot

Habit	Spreading subshrub or perennial herb, 0.15-1 m. Mostly hairless; thread-like to narrow leaves; ribbed stems.
Flowering	Aug - Dec and Jan - Mar (pale blue to white), in spikes at the ends of branches.
Habitat	Coastal sand and limestone.
Local note	The coastal Perth plant is subsp. <i>thesioides</i> ; recorded in the Wanneroo LGA.

Propagation method

Item	Details
Best time to take cuttings	Apr - May - warm autumn weather between flowering periods. Sep - Mar is workable using non-flowering side shoots. Avoid Jun - Aug.
Wood type	Softwood to semi-hardwood - firm but still green tip or side-shoot cuttings.
Cutting size	50-75 mm.
Selecting material	Non-flowering side shoots. Flowers form at the branch tips, so main tips often carry buds.
Pre-treatment	- Strip leaves from the lower half; leaves are narrow so little trimming is needed. - Stems are slender - always use a dibber hole. - Dip the base in rooting hormone.
Rooting hormone	Softwood strength (1.5 g/L IBA, Clonex Green); 3 g/L (Clonex Purple) for firmer material.
Propagation mix	Free-draining sand/coir mix.
Environment	Hairless leaves tolerate misting better than <i>S. canescens</i> , but fine stems collapse quickly if the mix is waterlogged or dries out. Keep evenly moist, with bottom heat and bright shade.
Expected strike time	About 4-8 weeks in warm weather.
Post-treatment / aftercare	- Standard hardening off. - Pot on gently - fine roots break easily. Single cuttings per tube make this easier. - Pinch tips once growing to encourage branching.

Common problems

Problem	Likely cause and fix
Fine stems wilt or collapse	Mix drying out or waterlogged. Keep evenly moist; check daily.
Cuttings flower instead of rooting	Pinch off terminal flower buds; use side shoots.
Root damage when potting on	Strike one cutting per tube instead of community pots.

Notes

Difficulty	Easy-Moderate (provisional). The genus strikes readily; the fine stems are less forgiving of watering errors.
Other methods	Seed – <i>Scaevola</i> seed is generally difficult. Division – possible if stock plants have suckered.
Confidence	Medium. Little species-specific data published. Based on the genus plus this species' fine, slender growth.
Sources	See References: 1, 4, 5, 6, 7, 12, 13, 14, 16

Nursery records

Log each batch here to confirm or update the timing and difficulty above.

Date set	Stock source / provenance	No. set	No. struck	Strike %	Notes

Spinifex hirsutus

Hairy Spinifex

Family
Poaceae

DIFFICULTY	WOOD TYPE	BEST MONTHS	CUTTING LENGTH	HORMONE (IBA)
Moderate *	Runner (stolon) sections	Mar – May	10–15 cm, 1–2 nodes	Optional (1.5 g/L)

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Cuttings	OK	OK	Best	Best	Best	–	–	–	OK	OK	OK	OK
Best window												
Workable – see 'Selecting material' in the method												
Avoid / low success												

Plant snapshot

Habit	Rhizomatous, tussock-forming grass, 0.3–0.9 m, sending long runners across the sand. Broad, flat, silvery leaves, hairy on both sides.
Flowering	Oct – Jan. Male and female flowers are on separate plants; female heads are large spiky balls that roll in the wind.
Habitat	White sand on coastal dunes; an important stabiliser at the toe of unstable dunes.
Local note	Recorded in the City of Joondalup. Less common than <i>S. longifolius</i> on the Cottesloe foreshore.

Propagation method

Item	Details
Best time to take cuttings	Mar – May, when runners are young, green and actively growing, giving plants ready for winter planting. Sep – Feb is workable. Avoid Jun – Aug (slow growth).
Wood type	Runner (stolon) sections – not woody cuttings. Use young, green runners, not old brown ones.
Cutting size	Nursery tubes: 10–15 cm sections with at least one (ideally two) nodes. Direct planting on site: 40–50 cm runners, buried about halfway in damp sand.
Selecting material	Actively growing runners from the dune edge. Plants are male or female, so collect from several plants if you want seed-producing plantings later.
Pre-treatment	- Keep runners moist and shaded after collecting – they dry out fast. - Cut just below a node; trim leaves back by about half. - Hormone optional – roots form naturally at the nodes.
Rooting hormone	Optional. If used, softwood strength (1.5 g/L IBA, Clonex Green) on the node.
Propagation mix	Coarse washed sand or a very sandy mix. Deep tubes suit the root system.
Environment	Bottom heat and misting gave good rooting at the nodes in <i>S. sericeus</i> trials. The hairy leaves of this species can hold water, so mist lightly.
Expected strike time	Nodes root readily, but plants are slow to reach planting size – in one trial about 80% of cuttings were ready to plant within 18 months.
Post-treatment / aftercare	- Harden off, then grow on in full sun. - Pot on into deep tubes or larger pots. - Plant out in late autumn or winter when rain is reliable.

Common problems

Problem	Likely cause and fix
Runners dry out before setting	Carry in damp bags in an esky; set the same day.
Nodes rot	Mix too wet or too much mist. Use coarser sand; mist less.
Slow growth after rooting	Normal for spinifex – allow time; feed lightly and give full sun.

Notes

Difficulty	Moderate (provisional). Runner sections root well under bottom heat and mist, but it's labour-intensive and slow. Seed is the more common nursery method.
Other methods	Seed – the usual nursery method. The Geraldton community nursery advises collecting from female plants growing near males. <i>S. sericeus</i> research found seed germinates far better in darkness (56%) than light (14%), so cover seed. Division of tussocks also works but survived less well than rooted cuttings in trials (31% vs 40%).
Confidence	Medium. Method based on published trials with the closely related <i>S. sericeus</i> and general spinifex practice.
Sources	See References: 4, 12, 18, 19, 20

Nursery records

Log each batch here to confirm or update the timing and difficulty above.

Date set	Stock source / provenance	No. set	No. struck	Strike %	Notes

Spinifex longifolius

Beach Spinifex

Family
Poaceae

DIFFICULTY	WOOD TYPE	BEST MONTHS	CUTTING LENGTH	HORMONE (IBA)
Moderate *	Runner (stolon) sections	Mar – May	10–15 cm, 1–2 nodes	Optional (1.5 g/L)

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Cuttings	OK	OK	Best	Best	Best	–	–	–	OK	OK	OK	OK
Best window												
Workable – see 'Selecting material' in the method												
Avoid / low success												

Plant snapshot

Habit	Rhizomatous, tussock-forming grass, 0.3–1 m high and up to 2 m wide, with long runners through the sand.
Flowering	Apr – Dec or Jan (green-brown). Male and female plants are separate; female heads up to 250 mm across roll in the wind.
Habitat	White sand on coastal dunes; a key foredune stabiliser.
Local note	Recorded in the Wanneroo LGA; the more common of the two spinifex species on the Cottesloe foreshore.

Propagation method

Item	Details
Best time to take cuttings	Mar – May, when runners are young, green and actively growing, giving plants ready for winter planting. Sep – Feb is workable. Avoid Jun – Aug (slow growth).
Wood type	Runner (stolon) sections – not woody cuttings. Use young, green runners, not old brown ones.
Cutting size	Nursery tubes: 10–15 cm sections with at least one (ideally two) nodes. Direct planting on site: 40–50 cm runners, buried about halfway in damp sand.
Selecting material	Actively growing runners from the dune edge. Plants are male or female, so collect from several plants if you want seed-producing plantings later.
Pre-treatment	- Keep runners moist and shaded after collecting – they dry out fast. - Cut just below a node; trim leaves back by about half. - Hormone optional – roots form naturally at the nodes.
Rooting hormone	Optional. If used, softwood strength (1.5 g/L IBA, Clonex Green) on the node.
Propagation mix	Coarse washed sand or a very sandy mix. Deep tubes suit the root system.
Environment	Bottom heat and misting gave good rooting at the nodes in <i>S. sericeus</i> trials. Keep the sand moist, not wet.
Expected strike time	Nodes root readily, but plants are slow to reach planting size – in one trial about 80% of cuttings were ready to plant within 18 months.
Post-treatment / aftercare	- Harden off, then grow on in full sun. - Pot on into deep tubes or larger pots. - Plant out in late autumn or winter when rain is reliable.

Common problems

Problem	Likely cause and fix
Runners dry out before setting	Carry in damp bags in an esky; set the same day.
Nodes rot	Mix too wet or too much mist. Use coarser sand; mist less.
Slow growth after rooting	Normal for spinifex – allow time; feed lightly and give full sun.

Notes

Difficulty	Moderate (provisional). Runner sections root well under bottom heat and mist, but it's labour-intensive and slow. Seed is the more common nursery method.
Other methods	Seed – the usual nursery method. The Geraldton community nursery collects from female plants growing close to males, separates the seed, cuts off the awns and sows en masse with smoke water (about 10 days to germinate). <i>S. sericeus</i> research found seed germinates far better in darkness, so cover seed. Division of tussocks also works.
Confidence	Medium. Method based on published trials with the closely related <i>S. sericeus</i> and general spinifex practice.
Sources	See References: 4, 12, 18, 19, 20

Nursery records

Log each batch here to confirm or update the timing and difficulty above.

Date set	Stock source / provenance	No. set	No. struck	Strike %	Notes

Threlkeldia diffusa

Coast Bonefruit

Family
Amaranthaceae
(chenopods)

DIFFICULTY Moderate *	WOOD TYPE Softwood tips				BEST MONTHS Dec - Mar			CUTTING LENGTH 50-75 mm			HORMONE (IBA) 1.5 g/L IBA		
	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	
Cuttings	Best	Best	Best	OK	-	-	-	-	-	-	OK	Best	
Best window	Workable – see ‘Selecting material’ in the method						Avoid / low success						

Plant snapshot

Habit	Much-branched, prostrate to erect perennial, 0.1-0.4 m, forming a spreading mat. Semi-succulent leaves.
Flowering	Oct - Nov (tiny, green), followed by small red to black fruits.
Habitat	White or grey sand over limestone, coastal areas and saline flats. Salt- and drought-tolerant.
Local note	Recorded in the City of Joondalup.

Propagation method

Item	Details
Best time to take cuttings	Dec - Mar - warm, with plants actively growing after flowering. Nov and Apr are workable. Avoid May - Oct.
Wood type	Softwood tip cuttings of the semi-succulent stems.
Cutting size	50-75 mm.
Selecting material	Healthy tips without developing fruit.
Pre-treatment	- Strip lower leaves. - No wounding. - Dip the base in rooting hormone.
Rooting hormone	Softwood strength, 1.5 g/L IBA (Clonex Green).
Propagation mix	Very free-draining sandy mix.
Environment	Semi-succulent leaves rot if kept wet - mist very little, keep the mix on the dry side, give good airflow and bright light.
Expected strike time	Not documented - start checking from about 4 weeks and record the result.
Post-treatment / aftercare	- Standard hardening off. - Grow on in full sun; water sparingly.

Common problems

Problem	Likely cause and fix
Cuttings turn soft and rot	Too wet or humid. Water less, mist less, improve airflow.
Weak, stretched growth	Not enough light. Move to a brighter position.
No roots	Try firmer tips or the seed method.

Notes

Difficulty	Moderate (provisional). No cutting data found; seed is the proven method (see Other methods).
Other methods	Seed – the documented method. The Geraldton community nursery soaks seed in smoke water (1:9) for 24 hours; germination takes 10–15 days and happens in cooler weather, so sow in autumn.
Confidence	Low. No published cutting data found. Seed is the documented method; cuttings are a trial.
Sources	See References: 1, 4, 12, 28

Nursery records

Log each batch here to confirm or update the timing and difficulty above.

Date set	Stock source / provenance	No. set	No. struck	Strike %	Notes

References

Numbers match the source list on each species page. Web addresses shortened; accessed September 2026.

#	Source	Address
1	Australian Native Plants Society (ANPSA) – Plant Propagation: cuttings, hormones, mixes, hardening off	anpsa.org.au/plant-propagation/
2	ANPSA – Grevillea vestita plant profile	anpsa.org.au/plant_profiles/grevillea-vestita/
3	Australian Plants Society NSW – Grevillea vestita	resources.austplants.com.au/plant/grevillea-vestita/
4	FloraBase, WA Herbarium (DBCA) – species profiles 5162, 7603, 12585, 12824, 13152	florabase.dbca.wa.gov.au
5	ANPSA – Scaevola ramosissima profile (genus note: cuttings strike very readily, seed difficult)	anpsa.org.au/plant_profiles/scaevola-ramosissima/
6	Sustainable Gardening Australia – Fan Flower: take cuttings during the growing season	sgaonline.org.au/fan-flower-fabulous-native-ground-cover/
7	Greenhouse Product News – Scaevola propagation (IBA rates, flower-free cuttings, bottom heat)	gpnmag.com/article/scaevola-outback-fan-flower/
8	ANPSA – Hibbertia vestita profile (genus note: easy from cuttings, seed unreliable)	anpsa.org.au/plant_profiles/hibbertia-vestita/
9	Australian National Botanic Gardens – Hibbertia pedunculata and H. empetrifolia (cutting timing, root rot)	anbg.gov.au/gnp/gnp11/hibbertia-pedunculata.html
10	Trees and Shrubs Online – Hibbertia (nodal cuttings while in growth; seed dormancy)	treesandshrubsonline.org/articles/hibbertia/
11	DBCA Library – wildflower booklet: H. hypericoides difficult from seed or cuttings	library.dbca.wa.gov.au/FullTextFiles/072542.pdf
12	City of Greater Geraldton Community Nursery – Native Plant Propagation Guide (2019)	cgg.wa.gov.au
13	Growth Technology – Clonex rooting gel strengths (Green 1.5, Purple 3, Red 8 g/L IBA)	growthtechnology.com.au
14	EverydayWits – Propagating Scaevola (small-nursery practice, seasonal success rates)	everydaywits.com/propagate-scaevola/
15	Minimalist Gardener – Propagating Australian natives from cuttings (grevillea strike times, sap irritant)	minimalistgardener.com.au
16	Wikipedia – Scaevola canescens, S. repens, S. thesioides (descriptions)	en.wikipedia.org
17	CompleteGrow – How to propagate Grevillea (low-phosphorus feeding)	completegrow.com.au
18	NZ Department of Conservation – Rehabilitation of coastal foredunes using indigenous sand-binding species (Spinifex sericeus stolon cuttings, seed)	doc.govt.nz/Documents/science-and-technical/sfc122.pdf
19	ABC Gardening Australia – Spinifex (40–50 cm runner cuttings)	abc.net.au/gardening/how-to/spinifex/9432152
20	Cottesloe Coastcare Association – plant pages: Spinifex longifolius, S. hirsutus, Eremophila glabra	cottesloecoastcare.org
21	ANPSA – Eremophila glabra plant profile	anpsa.org.au/plant_profiles/eremophila-glabra/
22	Australian Bushfoods – The Eremophila Page (cutting length, wounding, hormone)	ausbushfoods.com/bushfoodsonline/reports/Plants/EREMO.TXT
23	Australian National Botanic Gardens – Myoporum insulare	anbg.gov.au/gnp/interns-2010/myoporum-insulare.html
24	ANPSA – Lechenaultia biloba profile (cuttings strike very easily; seed difficult)	anpsa.org.au/plant_profiles/lechenaultia-biloba/
25	Better Homes and Gardens – fastest and easiest native plants to propagate (leschenaultia timing)	bhg.com.au/how-to-propagate-natives
26	Nurseries Online – Scaevola crassifolia (semi-hardwood cuttings in warmer months)	nurseriesonline.com.au/plant-index/australian-native-plants/scaevola-crassifolia/

#	Source	Address
27	Natural Area Nursery – WA native plant propagation (H. hypericoides difficult to propagate)	naturalarea.com.au/revegetation-plant-production/wa-native-plant-propagation/
28	APACE WA – Threlkeldia diffusa	apacewa.org.au/plant/threlkeldia-diffusa/
29	Geographe Community Landcare Nursery – Phyllanthus calycinus (Lysiandra calycina)	gcln.org.au/product/phyllanthus-calycinus/
30	Wikipedia – Hibbertia hypericoides, Eremophila glabra subsp. albicans, Phyllanthus calycinus, Scaevola crassifolia	en.wikipedia.org
31	PMC – Sympatric species of Hibbertia vary in dormancy break and germination requirements	pmc.ncbi.nlm.nih.gov/articles/PMC3336950

Revision log

Version	Date	Changes
1.0	Sep 2026	First edition: Grevillea vestita, Hibbertia racemosa, Scaevola canescens, S. repens, S. thesioides.
1.1	Sep 2026	Added 10 species: Eremophila glabra, Hibbertia hypericoides, Lechenaultia linarioides, Myoporum insulare, Phyllanthus calycinus, Scaevola crassifolia, S. globulifera, Spinifex hirsutus, S. longifolius, Threlkeldia diffusa. Added page numbers to the species index.