

ME 202  
LECTURE 31  
TUE 29 MAR 2022

Previously,

Buckling force is the critical force for which

✓ Multiple equilibrium solutions come into existence


OR

✓ System in static equilibrium switches stability i.e. potential energy extremum changes its nature

OR

min (stable)  $\Rightarrow$  max (unstable)

□ Dynamical system changes its time-dependent behavior oscillatory  $\Rightarrow$  exponential



### Dynamical System

Now  $\theta(t)$

$$\underline{I} = \underline{I} \approx \underline{I}$$

$$I \ddot{\theta} = PL \sin \theta - \beta \theta$$

$$I \ddot{\theta} + (\beta - PL) \theta = 0$$

$\theta \approx \sin \theta$  small

$$\beta > PL \Rightarrow P < \frac{\beta}{L}$$

oscillatory motion  $\theta(t)$   
 $\theta(t) = A \cos \omega t + B \sin \omega t$

$$\beta = PL$$

$$\theta(t) = A_0 + A_1 t$$

$$\beta < PL, P > \frac{\beta}{L}$$

$$\theta(t) = A e^{\omega t} + B e^{-\omega t}$$

exponential

### Imperfect/Asymmetric System in


#### Static Equilibrium

Use either  
force/moment or  
energy

As  $P \rightarrow P^*, P_{cr}, \theta \rightarrow \infty$   
uncontrollable

- ① Horiz force  $F$     ② Eccentric Load  $e$     ③ Initial deflection  $\theta_0$

① Horiz Force




$$\Pi(\theta) = \frac{1}{2} \beta \theta^2 - PL(1 - \cos\theta) - FL \sin\theta$$

$$\frac{d\Pi}{d\theta} = 0 \Rightarrow \underbrace{\beta\theta}_{\theta \text{ small}} - \underbrace{PL \sin\theta}_{\theta} - \underbrace{FL \cos\theta}_1 = 0$$

$$\theta = \frac{FL}{\beta - PL}$$

$$\text{As } P \rightarrow \underbrace{\beta/L}_{P^*}, \theta \rightarrow \infty$$

② Eccentric Load



$$\delta = L - L \cos \theta + e \sin \theta$$

$$\Pi = \frac{1}{2} \beta \theta^2 - P(L - L \cos \theta + e \sin \theta)$$


$$\frac{d\Pi}{d\theta} = 0, \quad \theta \text{ small}$$

$$\Rightarrow \beta \theta - PL \sin \theta - Pe \cos \theta = 0$$

$$\theta = \frac{Pe}{\beta - PL}$$

$$\text{As } P \rightarrow \underbrace{\beta/L}_{P^*}, \quad \theta \rightarrow \infty$$

### ③ Initial Angle / Offset




$$\Pi = \frac{1}{2} \beta (\theta - \theta_0)^2 - PL(1 - \cos \theta) + \Pi_0$$

$$\frac{d\Pi}{d\theta} = 0, \quad \beta(\theta - \theta_0) - PL \sin \theta = 0$$

$$\theta = \frac{\beta \theta_0}{\beta - PL} \quad \theta \text{ small}$$

$$P^* = \beta / L$$

## 2 DOF $N=2$ BUCKLING



$\theta=0, \psi=0$  orig eqm config

Find  $P$  at which non-trivial  $\begin{pmatrix} \theta \\ \psi \end{pmatrix}$  exist

$$\Pi(\theta, \psi) = \frac{1}{2} \beta \theta^2 + \frac{1}{2} \beta (\psi - \theta)^2 - PL(2 - \cos \theta - \cos \psi)$$

Eqm

$$\frac{\partial \Pi}{\partial \theta} = 0 \Rightarrow \beta \theta - \beta(\psi - \theta) - \underbrace{PL \sin \theta}_{\theta} = 0$$

$$\frac{\partial \Pi}{\partial \psi} = 0 \Rightarrow \beta(\psi - \theta) - \underbrace{PL \sin \psi}_{\psi} = 0$$

$$2\beta \frac{\theta}{L} - \frac{\beta}{L} \psi - P\theta = 0$$

$$-\frac{\beta}{L} \theta + \frac{\beta}{L} \psi - P\psi = 0$$

$$\begin{pmatrix} 2\beta/L & -\beta/L \\ -\beta/L & \beta/L \end{pmatrix} \begin{pmatrix} \theta \\ \psi \end{pmatrix} = P \begin{pmatrix} \theta \\ \psi \end{pmatrix}$$

eigenvalue problem

$P$

eigenvalues

eigenvectors

$$\begin{pmatrix} \theta \\ \psi \end{pmatrix}$$

$$\begin{pmatrix} 2\frac{\beta}{L} - P & -\frac{\beta}{L} \\ -\frac{\beta}{L} & \frac{\beta}{L} - P \end{pmatrix} \begin{pmatrix} \theta \\ \psi \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

$$\det = 0$$

$$(PL)^2 - 3\beta PL + \beta^2 = 0$$

$$P = \frac{\beta}{2L} (3 \pm \sqrt{5})$$

$$P_1 \approx 0.3820 \frac{\beta}{L}, \quad P_2 \approx 2.621 \frac{\beta}{L}$$

$$\text{for } P_1, \quad \psi = 1.6180 \theta$$

$$\text{for } P_2, \quad \psi = -0.618 \theta$$

Deflections are arbitrarily large/small.



Loads  $P = P_1^* = 0.3820 \frac{\beta}{L}$



$$\psi = 1.618 \theta$$

$P = P_2^* = 2.621 \frac{\beta}{L}$



$$\psi = -0.618 \theta$$

Buckling modes / mode shapes /  
eigenvectors

Hessian  $\tilde{H} = \begin{pmatrix} \frac{\partial^2 \pi}{\partial \theta^2} & \frac{\partial^2 \pi}{\partial \psi \partial \theta} \\ \frac{\partial^2 \pi}{\partial \theta \partial \psi} & \frac{\partial^2 \pi}{\partial \psi^2} \end{pmatrix}$

compute  $\tilde{H}$  at  $\frac{\partial \pi}{\partial \theta} = 0, \frac{\partial \pi}{\partial \psi} = 0$

$H_2$	pos. def	minimum	stable eqm
-------	----------	---------	------------

$H_2$	neg. def	maximum	unstable eqm
-------	----------	---------	--------------

$H_2$	indefinite	extremum	neither min/ max
-------	------------	----------	---------------------

Dnyanesh Pawaskar



10

of 10

