



Contents lists available at ScienceDirect

Environment International

journal homepage: www.elsevier.com/locate/envint



Corrigendum



Corrigendum to “Exposure to air pollutant mixture and gestational diabetes mellitus in Southern California: Results from electronic health record data of a large pregnancy Cohort” [Environ. Int. 158 (2022) 106888]

Yi Sun^a, Xia Li^b, Tarik Benmarhnia^c, Jiu-Chiuan Chen^d, Chantal Avila^b, David A. Sacks^{b,e}, Vicki Chiu^b, Jeff Slezak^b, John Molitor^f, Darios Getahun^{b,g,*}, Jun Wu^{a,*}

^a Department of Environmental and Occupational Health, Program in Public Health, University of California, Irvine, CA, USA

^b Department of Research & Evaluation, Kaiser Permanente Southern California, Pasadena, CA, USA

^c Herbert Wertheim School of Public Health and Scripps Institution of Oceanography, University of California, San Diego, 9500 Gilman Drive #0725, La Jolla, CA 92093, USA

^d Department of Preventive Medicine, University of Southern California, Los Angeles, CA 90033, USA

^e Department of Obstetrics and Gynecology, University of Southern California, Keck School of Medicine, Los Angeles, CA, USA

^f College of Public Health and Human Sciences, Oregon State University, Corvallis, OR 97331, USA

^g Department of Health Systems Science, Kaiser Permanente Bernard J. Tyson School of Medicine, Pasadena, CA, USA

The authors regret the error in Table 3. A revised Table 3 is included below, and the correct statement in the abstract and Results section should read as follows: “In multi-pollutant models, increased ORs for GDM in association with one quartile increase in air pollution mixtures were found for both kriging-based regional air pollutants (NO₂, PM_{2.5},

and PM₁₀, OR = 1.233, 95% CI: 1.199–1.268) and PM_{2.5} constituents (i. e., sulfate, nitrate, ammonium, organic matter and black carbon, OR = 1.694, 95% CI: 1.538–1.875).” The authors would like to apologise for any inconvenience caused.

DOI of original article: <https://doi.org/10.1016/j.envint.2021.106888>.

* Corresponding authors at: 100 S. Los Robles Avenue 2nd Floor, Pasadena, CA 91101, USA (D.T. Getahun). Anteater Instruction & Research Bldg, 2034, 653 East Peltason Drive, University of California, Irvine, CA 92697-3957, USA (J. Wu).

E-mail addresses: Darios.T.Getahun@kp.org (D. Getahun), junwu@hs.uci.edu (J. Wu).

<https://doi.org/10.1016/j.envint.2022.107207>

Table 3

Adjusted odds ratios (ORs) and 95% confidence intervals (CIs) of GDM associated with air pollution during pregnancy in multi-pollutant models.

Pollutants	Coefficient β	Effect of mixtures ψ (log OR)	95% CI	OR	95% CI
Model 1: Krigged air pollutants					
Krigged PM _{2.5}	0.031				
Krigged PM ₁₀	−0.051	0.091	(0.079, 0.103)	1.233	(1.199, 1.268)
Krigged NO ₂	0.111				
Model 2: PM_{2.5} constituents					
PM _{2.5} sulfate	−0.004				
PM _{2.5} nitrate	0.082				
PM _{2.5} ammonium	0.065	0.229	(0.187, 0.273)	1.694	(1.538, 1.875)
PM _{2.5} organic matter	−0.050				
PM _{2.5} black carbon	0.135				

GDM, gestational diabetes mellitus; Model adjusted for maternal age, race/ethnicity, education, block group household income, pre-pregnancy BMI, smoking during pregnancy, insurance type, season of conception, year of birth and county.