

Supplemental Material

Table S1. Polynomial distributed lag (PDL) model posterior probabilities obtained from Bayesian model averaging (BMA) for zero-order through 5th order models using 12 lag periods, for PM_{2.5} and PM₁₀.

Pollutant	PDL model order (for 12 lag periods)	Posterior probability
PM _{2.5}	0	98.4%
	1	1.6%
	2	0.0%
	3	0.0%
	4	0.0%
	5	0.0%
PM ₁₀	0	98.2%
	1	1.8%
	2	0.0%
	3	0.0%
	4	0.0%
	5	0.0%

Table S2. Associations of PM_{2.5} and PM₁₀ exposure metrics and FMD stratified by age (less than 50 years vs. 50 years and older) and sex for increases in normalized PM_{2.5} and PM₁₀ exposure in fully adjusted models.

Age younger than 50 years								
	<i>Men</i>				<i>Women</i>			
Particulate air pollution metrics (normalized)	β	SE	<i>p</i> -value	95% CI	β	SE	<i>p</i> -value	95% CI
PM _{2.5} 12-month moving-average	-0.07	0.05	0.1	-0.17, 0.03	-0.02	0.08	0.7	-0.17, 0.14
PM ₁₀ 12-month moving-average	-0.16	0.05	0.005	-0.25, -0.04	-0.01	0.07	0.8	-0.14, 0.12
Age 50 years and older								
PM _{2.5} 12-month moving-average	-0.13	0.09	0.1	-0.30, 0.04	-0.07	0.07	0.3	-0.20, 0.06
PM ₁₀ 12-month moving-average	-0.17	0.09	0.08	-0.36, 0.02	-0.06	0.07	0.4	-0.18, 0.08


Table S3. Associations of PM_{2.5} and PM₁₀ exposure metrics and FMD across all participants and by sex for a 10 µg m⁻³ increment in PM_{2.5} and PM₁₀ exposure in fully adjusted models.

Particulate air pollution metrics (per 10 µg m ⁻³)	<i>Across all</i>				<i>Men</i>				<i>Women</i>			
	β	SE	<i>p</i> -value	95% CI	β	SE	<i>p</i> -value	95% CI	β	SE	<i>p</i> -value	95% CI
PM _{2.5} 12-month moving-average	-4.0	1.8	0.03	-7.5, -0.5	-4.9	2.4	0.04	-9.6, -0.3	-3.7	3.0	0.2	-9.5, 2.1
PM ₁₀ 12-month moving-average	-4.7	1.7	0.007	-8.2, -1.3	-7.6	2.3	0.001	-12.2, -3.0	-2.8	2.6	0.3	-7.9, 2.4

*All models adjusted for age, sex, age by sex interaction, smoking (except in models for women only because there were no ever smokers), BMI, season, year, hypertension, and base brachial artery diameter.

Figure S1. Effect estimates from the six PDL models (zero-order through 5th order) weighted using BMA as discussed in main text, for: A) PM_{2.5} and B) PM₁₀. The month prior to FMD measurement (lag zero) and eleven months prior are shown, as well as the cumulative effect over all 12 lag periods.

A)



B)

