

Supplemental Material

Table S1. Odds ratio (OR) and 95% confidence intervals (CI) for Short-term associations between takotsubo syndrome (TTS) hospitalization and daily maximum temperature exposure per 1 °C increase in state of California from 2006 to 2019.

Exposure Windows	TTS Hospitalization			TTS hospitalization with angiography but no revascularization		
	OR	Lower 95% CI	Upper 95% CI	OR	Lower 95% CI	Upper 95% CI
Lag 1	0.992	0.984	1.000	0.990	0.981	1.000
Lag 2	0.997	0.989	1.005	0.995	0.986	1.004
Lag 3	1.005	0.997	1.013	1.004	0.995	1.013
Lag 4	1.008	1.000	1.016	1.007	0.998	1.016
Lag 5	1.012	1.004	1.020	1.012	1.003	1.021
Lag 6	1.010	1.002	1.018	1.010	1.001	1.019

Table S2. Odds ratio (OR) and 95% confidence intervals (CI) for takotsubo syndrome (TTS) hospitalization associated with heat wave days compared to non-heat wave days. Heat waves were defined as periods of ≥ 2 consecutive days with daily maximum temperature exceeding the 95th percentile. The first day of each heat wave period was considered as non-heat wave day.

Exposure Windows	TTS Hospitalization			TTS hospitalization with angiography but no revascularization		
	OR	Lower 95% CI	Upper 95% CI	OR	Lower 95% CI	Upper 95% CI
Lag 1	0.935	0.782	1.119	0.839	0.683	1.029
Lag 2	0.936	0.783	1.120	0.881	0.719	1.079
Lag 3	1.129	0.954	1.336	1.117	0.925	1.349
Lag 4	1.248	1.054	1.478	1.181	0.976	1.430
Lag 5	1.382	1.167	1.636	1.288	1.065	1.557
Lag 6	1.266	1.070	1.498	1.201	0.997	1.448
Lag 1-3	1.038	0.909	1.184	0.992	0.855	1.151
Lag 4-6	1.300	1.142	1.479	1.243	1.076	1.437
Lag 1-6	1.152	1.032	1.286	1.110	0.981	1.255

Table S3. Odds ratio (OR) and 95% confidence intervals (CI) for takotsubo syndrome (TTS) hospitalization associated with heat wave days compared to non-heat wave days. Heat waves were defined as periods of ≥ 2 consecutive days with daily maximum temperature exceeding the 98th percentile. The first day of each heat wave period was considered as non-heat wave day.

Exposure Windows	TTS Hospitalization			TTS hospitalization with angiography but no revascularization		
	OR	Lower 95% CI	Upper 95% CI	OR	Lower 95% CI	Upper 95% CI
Lag 1	1.036	0.774	1.388	1.052	0.762	1.453
Lag 2	0.851	0.633	1.144	0.775	0.552	1.088
Lag 3	1.189	0.912	1.545	1.209	0.899	1.629
Lag 4	1.293	0.990	1.688	1.217	0.902	1.643
Lag 5	1.161	0.878	1.536	1.041	0.756	1.434
Lag 6	1.386	1.048	1.833	1.403	1.025	1.921
Lag 1-3	1.052	0.864	1.282	1.056	0.846	1.318
Lag 4-6	1.336	1.099	1.623	1.266	1.018	1.574
Lag 1-6	1.179	1.011	1.376	1.158	1.000	1.342

Table S4. The sensitivity analysis for different degrees of freedom (DF) of day of the year (DoY). Odds ratio (OR) and 95% confidence intervals (CI) for takotsubo syndrome (TTS) hospitalization associated with heat wave days compared to non-heat wave days. Heat waves were defined as periods of ≥ 2 consecutive days with daily maximum temperature exceeding the 95th percentile. The first day of each heat wave period was considered as non-heat wave day.

	DoY DF=5		DoY DF=6	
	TTS Hospitalization	TTS hospitalization with angiography but no revascularization	TTS Hospitalization	TTS hospitalization with angiography but no revascularization
Exposure Windows	OR (95% CI)	OR (95% CI)	OR (95% CI)	OR (95% CI)
Lag 1	0.935 (0.781, 1.119)	0.837 (0.682, 1.027)	0.936 (0.782, 1.120)	0.838 (0.683, 1.028)
Lag 2	0.937 (0.783, 1.120)	0.881 (0.719, 1.079)	0.937 (0.784, 1.121)	0.881 (0.719, 1.080)
Lag 3	1.130 (0.955, 1.337)	1.118 (0.926, 1.350)	1.130 (0.955, 1.338)	1.118 (0.926, 1.350)
Lag 4	1.250 (1.055, 1.480)	1.182 (0.976, 1.430)	1.248 (1.054, 1.477)	1.177 (0.972, 1.425)
Lag 5	1.384 (1.169, 1.638)	1.289 (1.066, 1.558)	1.380 (1.166, 1.634)	1.282 (1.061, 1.550)
Lag 6	1.267 (1.071, 1.499)	1.202 (0.997, 1.448)	1.265 (1.069, 1.496)	1.195 (0.992, 1.440)
Lag 1-3	1.038 (0.910, 1.184)	0.991 (0.854, 1.150)	1.039 (0.910, 1.186)	0.992 (0.855, 1.151)
Lag 4-6	1.302 (1.144, 1.481)	1.244 (1.077, 1.438)	1.299 (1.142, 1.479)	1.239 (1.072, 1.432)
Lag 1-6	1.153 (1.033, 1.288)	1.110 (0.981, 1.256)	1.153 (1.033, 1.287)	1.108 (0.980, 1.254)