

Bathymetry beneath the Amery ice shelf, East Antarctica, revealed by airborne gravity

Junjun Yang^{2,*}, Jingxue Guo^{1,*}, Jamin S. Greenbaum³, Xiangbin Cui¹, Liangcheng Tu^{2,4}, Lin Li¹, Lenneke M. Jong⁵, Xueyuan Tang¹, Bingrui Li¹, Donald D. Blankenship⁶, Jason L. Roberts⁵, Tas van Ommen⁵, and Bo Sun¹

¹Polar Research Institute of China, Shanghai 200136, China

²Institute of Geophysics, PGMF and School of Physics, Huazhong University of Science and Technology, Wuhan 430074, China

³Scripps Institution of Oceanography, University of California, San Diego, La Jolla, CA 92093, USA

⁴TianQin Research Center for Gravitational Physics, School of Physics and Astronomy, Sun Yat-sen University, Zhuhai 519082, China

⁵Australian Antarctic Division, Kingston, Tasmania 7050, Australia

⁶Institute for Geophysics, University of Texas at Austin, Austin, Texas 78758, USA

*J. Yang and J. Guo contributed equally to this work

Contents of this file

Figures S1 to S2




Figure S1. (a) The cost function decreases as the iteration number increases. (b) Differences between the pseudo-observed gravity anomalies and the one forward computed from the gravity-estimated seafloor topography. The white line is the coast/calving front boundary. The thin magenta line is grounding line. The background Landsat 8 image is acquired in January 2018. (c) Differences between the pseudo-observed gravity anomalies and the one forward computed from bed elevations in BedMachine Antarctica v2. (d) Differences between the gravity-estimated seafloor topography and BedMachine Antarctica v2.




Figure S2. Bed elevations (orthometric height in meters referred to the EIGEN-6C4 geoid) around the Amery ice shelf from BedMachine Antarctica v2 model. Thin black lines are contours with 500 m interval.