



Geothermal heat flux from measured temperature profiles in deep ice boreholes in Antarctica

Pavel Talalay, Yazhou Li, Laurent Augustin, Gary Clow, Jialin Hong, Eric Lefebvre, Alexey Markov, Hideaki Motoyama, Catherine Ritz

► To cite this version:

Pavel Talalay, Yazhou Li, Laurent Augustin, Gary Clow, Jialin Hong, et al.. Geothermal heat flux from measured temperature profiles in deep ice boreholes in Antarctica. *The Cryosphere*, Copernicus 2020, 14 (11), pp.4021-4037. 10.5194/tc-14-4021-2020 . hal-03036864

HAL Id: hal-03036864

<https://hal.archives-ouvertes.fr/hal-03036864>

Submitted on 25 Aug 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution | 4.0 International License



Geothermal heat flux from measured temperature profiles in deep ice boreholes in Antarctica

Pavel Talalay¹, Yazhou Li¹, Laurent Augustin², Gary D. Clow³, Jialin Hong¹, Eric Lefebvre⁴, Alexey Markov¹, Hideaki Motoyama⁵, and Catherine Ritz⁴

¹Polar Research Center, Institute for Polar Science and Engineering, Jilin University, 130021 Changchun, China

²Division Technique de l'INSU, CNRS, 83507 La Seyne sur Mer, France

³Institute of Arctic and Alpine Research, University of Colorado Boulder, Boulder, Colorado, USA

⁴Université Grenoble Alpes, CNRS, IRD, IGE, 38000 Grenoble, France

⁵National Institute of Polar Research, Tokyo, Japan

Correspondence: Pavel Talalay (ptalalay@yahoo.com) and Yazhou Li (jlyuzhouli@163.com)

Received: 25 January 2020 – Discussion started: 4 March 2020

Revised: 6 October 2020 – Accepted: 11 October 2020 – Published: 14 November 2020

Abstract. The temperature at the Antarctic Ice Sheet bed and the temperature gradient in subglacial rocks have been directly measured only a few times, although extensive thermodynamic modeling has been used to estimate the geothermal heat flux (GHF) under the ice sheet. During the last 5 decades, deep ice-core drilling projects at six sites – Byrd, WAIS Divide, Dome C, Kohnen, Dome F, and Vostok – have succeeded in reaching or nearly reaching the bed at inland locations in Antarctica. When temperature profiles in these boreholes and steady-state heat flow modeling are combined with estimates of vertical velocity, the heat flow at the ice-sheet base is translated to a geothermal heat flux of $57.9 \pm 6.4 \text{ mW m}^{-2}$ at Dome C, $78.9 \pm 5.0 \text{ mW m}^{-2}$ at Dome F, and $86.9 \pm 16.6 \text{ mW m}^{-2}$ at Kohnen, all higher than the predicted values at these sites. This warm base under the East Antarctic Ice Sheet (EAIS) could be caused by radiogenic heat effects or hydrothermal circulation not accounted for by the models. The GHF at the base of the ice sheet at Vostok has a negative value of $-3.6 \pm 5.3 \text{ mW m}^{-2}$, indicating that water from Lake Vostok is freezing onto the ice-sheet base. Correlation analyses between modeled and measured depth–age scales at the EAIS sites indicate that all of them can be adequately approximated by a steady-state model. Horizontal velocities and their variation over ice-age cycles are much greater for the West Antarctic Ice Sheet than for the interior EAIS sites; a steady-state model cannot precisely describe the temperature distribution here. Even if the correlation factors for the best fitting age–depth

curve are only moderate for the West Antarctic sites, the GHF values estimated here of $88.4 \pm 7.6 \text{ mW m}^{-2}$ at Byrd and $113.3 \pm 16.9 \text{ mW m}^{-2}$ at WAIS Divide can be used as references before more precise estimates are made on the subject.

1 Introduction

The Antarctic geothermal heat flux (GHF), an important boundary condition for ice-sheet behavior, can influence sea-level changes (Golledge et al., 2015) considering its significant influence on the viscosity of basal ice and meltwater content at the ice–base interface. What are the basal ice temperature and mechanical properties? How does GHF control basal melt and affect the internal deformation of the ice sheet? How old is ice at different locations? These questions can be answered only by applying reliable GHF measurements or estimates.

The average global surface GHF is $\sim 86 \text{ mW m}^{-2}$, which varies from 64.7 mW m^{-2} , the mean continental heat flow (including arcs and continental margins), to 95.9 mW m^{-2} , the mean oceanic heat flow (Davies, 2013). However, several geologic factors including heat from the mantle, heat production in the crust by radioactive decay, and tectonic history cause spatially variable GHF in Antarctica (Burton-Johnson et al., 2020).

Most studies of GHF in Antarctica rely on thermal models (Pattyn, 2010; Van Liefferinge and Pattyn, 2013). Modeling studies based on a global seismic survey and the structural similarity of crust and upper mantle showed that the West Antarctic Ice Sheet (WAIS) has a GHF 3 times higher than that of the East Antarctic Ice Sheet (EAIS; Shapiro and Ritzwoller, 2004). For a central point in the WAIS (78° S, 110° W), the average GHF is expected to be 110 mW m^{-2} . The GHF can also be estimated on the basis of geologic information, where uniform values are attributed to large geologically homogeneous areas (An et al., 2015; Goodge, 2018; Llubes et al., 2006; Martos et al., 2017; Pollard et al., 2005).

Some studies use remote methods to estimate the GHF underneath the Antarctic Ice Sheet. For example, satellite magnetic data showed that the GHF underneath the ice sheet varies from 40 to 185 mW m^{-2} and that areas of high GHF coincide with known current volcanism and some areas known to have ice streams (Fox Maule et al., 2005). In the central part of the EAIS, the average GHF was estimated to be in the range of 50 to 60 mW m^{-2} ; however, elevated GHFs were found along the WAIS–EAIS boundary and around the Siple Coast. Similarly, high GHFs were found around Victoria Land, Oates Land, and George V Land. Observations of crustal heat production within the continental crust underneath the Lambert–Amery glacial system in East Antarctica also show high heat flux of at least 120 mW m^{-2} (Pittard et al., 2016).

Direct temperature measurement obviously produces the most reliable GHF estimates and can be used to verify results of preliminary thermal modeling and geological–geophysical studies. While over 10 000 heat flow measurements have been made globally, 90 % are from Europe, North America, and southern Africa. South America, Asia, and Australia have far fewer measurements, while Antarctica has virtually none (Davies, 2013). Drilling through thick ice is extremely complicated, time-consuming, and expensive; therefore, direct temperature measurements in Antarctic subglacial till and bedrock environments have only been conducted twice so far, both under the WAIS: at the subglacial Lake Whillans ($285 \pm 80 \text{ mW m}^{-2}$; Fisher et al., 2015) and near the grounding zone of the Whillans Ice Stream ($88 \pm 7 \text{ mW m}^{-2}$; Beegeman et al., 2017), ~ 100 km apart (Fig. 1). The tremendous difference in the values of GHF between these two adjacent sites suggests high spatial variability in West Antarctica.

More reliable GHF estimates under the Antarctic Ice Sheet can be made from available temperature profiles in ice boreholes. During the last 5 decades, deep ice-core drilling projects at six sites – Byrd (Ueda and Garfield, 1970), WAIS Divide (Slawny et al., 2014), Dome C (Augustin et al., 2007), Kohnen (Wilhelms et al., 2014), Dome F (Motoyama, 2007), and Vostok (Lukin and Vasiliev, 2014; Vasiliev et al., 2011) – have succeeded in reaching or nearly reaching the ice-sheet bed at inland locations in Antarctica. Reported drill site conditions – snow accumulation rate, mean annual surface

air temperature, ice-sheet surface velocity, ice thickness, and drilling depth – are summarized in Table 1.

The Byrd and Kohnen holes encountered water at the base, which welled up into the holes. The borehole at Vostok penetrated the subglacial Lake Vostok at 3769.3 m, and here, water rose from the lake to a height of more than 340 m. Drilling of the other holes was stopped within 10–50 m of the bed. All these holes were temperature-logged and provide a good opportunity to fill the gap in our knowledge of the GHF under the Antarctic ice.

2 Methods

2.1 Temperature and temperature gradient at the base of Antarctic Ice Sheet

Temperatures in the Byrd, WAIS Divide, Vostok, Dome C, Kohnen, and Dome F boreholes were measured using different devices and different methods. All boreholes were mechanically drilled and filled with kerosene-based drilling fluid. Temperature profiles were then obtained by logging with custom-made borehole loggers (Dome C, Kohnen, Dome F, WAIS Divide, and Vostok) or a thermistor embedded in the drill (Byrd).

Measured temperature profiles in four of the boreholes (Vostok, Dome C, Kohnen, and Dome F) increase almost linearly with depth, as expected at locations with minimal annual snow accumulation and hence small vertical velocities (Fig. 2). In contrast, vertical advection is much greater at the Byrd and WAIS Divide sites in West Antarctica; at these locations the upper part of the ice sheet is nearly isothermal, but at depth the temperature gradient is nearly the same as that at the other sites. Temperature gradients at the bed are $2.02\text{--}3.12 \text{ }^{\circ}\text{C}/100 \text{ m}$ at Dome C, Kohnen, Dome F, and Vostok and slightly higher in West Antarctica, $3.70\text{--}3.88 \text{ }^{\circ}\text{C}/100 \text{ m}$ at Byrd and WAIS Divide (Table 2).

Temperature profiles in deep ice-core drilling boreholes are approximated closely by polynomials with correlation factors of > 0.99 (Table 3), indicating a positive relationship between temperature and vertical depth. Ice thicknesses generated by extrapolating the temperature profile to the depth of the pressure melting point assuming a Clausius–Clapeyron slope of 0.0742 K/MPa (Cuffey and Paterson, 2010) are in good agreement with radar data except for WAIS Divide, where the difference is $\sim 30 \text{ m}$. This could be attributed to scintillations on the melted ice–bedrock interface or other effects. However, in-depth temperature extrapolation has limited accuracy and thus often does not provide a correct estimate of GHF. Thus, a steady-state model and genetic algorithm (GA) are applied herein to fit the measured temperatures.




Figure 1. GHF derived in the present study (P.S.) from basal temperature gradients in deep ice boreholes (green bars) compared with modeling. Red circles show locations of deep-ice-drilling sites (Byrd, WAIS Divide, Vostok, Dome C, Kohnen, and Dome F) discussed in the present study. Black squares show locations of boreholes drilled in Antarctic margins in which borehole temperature measurements were carried out and GHF values were estimated (^a Zagorodnov et al., 2012; ^b Nicholls and Paren, 1993; ^c Fisher et al., 2015; ^d Begeman et al., 2017; ^e Engelhardt, 2004; ^f Risk and Hochstein, 1974; ^g Decker and Bucher, 1982; ^h Clow et al., 2011; ⁱ Dahl-Jensen et al., 1999; ^j Zotikov, 1961). Drill sites in ocean and subshelf sediments are not shown. Location of the Antarctic ice divides is shown according to Rignot et al. (2011).


Table 1. Information for Antarctic deep-ice-drilling sites.

| Parameters | WAIS | | EAIS | | | |
|--|------------------------|------------------------|-------------------------------------|---------------------------|------------------------|-------------------------|
| | Byrd | WAIS Divide | Vostok | Dome C | Kohnen | Dome F |
| Coordinates | 80°01' S, 119°31' W | 79°28' S, 112°05' W | 78°28' S, 106°48' E | 75°06' S, 123°24' E | 75° S, 0° E | 77°19' S, 39°40' E |
| Years drilled | 1966–1968 ^a | 2006–2011 ^d | 1990–1998, 2005–2014 ^{f,g} | 1999–2004 ⁱ | 2002–2006 ^k | 2003–2007 ⁿ |
| Surface elevation (m a.s.l.) | 1530 ^a | 1766 ^e | 3488 ^f | 3233 ^j | 2892 ^l | 3810 ^j |
| Drilled depth (m) | 2193 | 3405 ^d | 3769.3 ^g | 3270.2 ⁱ | 2774.2 ^k | 3035.2 ⁿ |
| Ice thickness according to radar or seismic survey (m) | 2300 ^b | 3455 ^e | 3750 ± 20 ^g | 3273 ± 5 ⁱ | 2750 ± 50 ^l | 3028 ± 15 ^j |
| Snow accumulation at surface (mm ice a ⁻¹) | 169.5 ^c | 220 ^e | 24.8 ^h | 28.4 ^j | 70 ^m | 29.9 ^j |
| Ice-sheet surface horizontal velocity (m a ⁻¹) | 12.7 ^o | ~ 3.0 ^{t,u} | 2.00 ± 0.01 ^s | 0.015 ± 0.01 ^p | 0.74 ^r | Negligible ^v |
| Mean surface snow temperature (°C) | -28 ^a | -30 ^e | -57 ^h | -54.6 ^j | -44 ^l | -57.3 ^j |

^a Ueda (2007); ^b Wexler (1961); ^c Gow (1968); ^d Slawny et al. (2014); ^e WAIS Divide Project Members (2013); ^f Vasiliev et al. (2011); ^g Lukin and Vasiliev (2014); ^h Ekaykin et al. (2012); ⁱ Augustin et al. (2007); ^j Parrenin et al. (2007a); ^k Wilhelms et al. (2014); ^l Ueltzhöffer et al. (2010); ^m Huybrechts et al. (2007); ⁿ Motoyama (2007); ^o Whillans (1977); ^p Virtuari et al. (2004); ^r Wesche et al. (2007); ^s Wendt et al. (2006); ^t Conway and Rasmussen (2009); ^u Koutnik et al. (2016); ^v Motoyama et al. (2008).

Table 2. Thermophysical properties at the base of the Antarctic Ice Sheet at sites of deep ice drilling estimated in this study.

| Parameters | WAIS | | EAIS | | | |
|---|----------------|------------------|----------------|-----------------|-----------------|----------------|
| | Byrd | WAIS Divide | Vostok | Dome C | Kohnen | Dome F |
| Temperature ($^{\circ}\text{C}$) | -1.43 | -2.30 | -2.49 | -2.15 | -1.85 | -1.99 |
| Temperature gradient ($^{\circ}\text{C } 100 \text{ m}^{-1}$) | 3.70 | 3.88 | 2.02 | 2.42 | 3.12 | 2.66 |
| Ice thickness according to depth of pressure melting point (m) | 2164 | 3485 | 3759 | 3257 | 2770 | 3016 |
| Basal melt rate (mm a^{-1}) | 1.2 ± 0.8 | 3.7 ± 1.7 | -4.8 ± 0.6 | 1.08 ± 0.27 | 2.8 ± 1.6 | 2.5 ± 0.5 |
| GHF (mW m^{-2}) | 88.4 ± 7.6 | 113.3 ± 16.9 | -3.6 ± 5.3 | 57.9 ± 6.4 | 86.9 ± 16.6 | 78.9 ± 5.0 |

**Figure 2.** Smoothed measured temperature profiles in Antarctic deep ice boreholes. Pressure melting point temperature $T_{\text{melt}}(z)$ is shown under the assumption of a Clausius–Clapeyron slope of 0.0742 K/MPa.

2.2 GHF estimation model

A one-dimensional time-dependent energy-balance equation (Dahl-Jensen et al., 2003; Johnsen et al., 1995) is usually used to model the temperature distribution through the ice as a function of the climate conditions on the surface and the GHF from the bedrock:

$$\rho c \frac{\partial T}{\partial t} = \frac{\partial}{\partial z} \left(k \frac{\partial T}{\partial z} \right) - \rho c w \frac{\partial T}{\partial z} - \rho c u \frac{\partial T}{\partial x}, \quad (1)$$

where T is the temperature as a function of z , x , and t ; z represents the vertical coordinate (at the ice-sheet base, $z=0$, while at ice-sheet surface, $z=H$); H is the ice thickness and is assumed to be constant in time; x is the horizontal coordinate; t is the time; k is the thermal conductivity of ice dependent on T ; ρ is the density of ice; c is the specific heat

capacity of ice dependent on T ; w and u are, respectively, the vertical velocity and horizontal velocity of the ice sheet dependent on z and t .

The temperature measured at the six drill sites can be considered at thermal steady state in their near-base portion. Three drill sites (Dome C, Dome F, Vostok) are in close vicinity to ice divides where horizontal advection and horizontal heat conduction are assumed to be minimal, and the environment approximates a steady state (Cuffey and Paterson, 2010). At first approximation, we also assume WAIS Divide is in a steady state. Byrd and Kohnen are in the slow-moving areas of the interior of the Antarctic Ice Sheet with a relatively smooth bed, where horizontal conduction is much lower than vertical conduction (Hindmarsh, 1999, 2018), and horizontal advection and horizontal heat conduction can be neglected (Robin, 1955; Van Liefferinge et al., 2018). This assumption reduces the non-steady-state heat-transfer equation to

$$\frac{\partial}{\partial z} \left(k \frac{\partial T}{\partial z} \right) - \rho c w \frac{\partial T}{\partial z} = 0, \quad (2)$$

which can be rewritten as

$$\frac{1}{k} \frac{\partial k}{\partial z} \frac{\partial T}{\partial z} + \frac{\partial^2 T}{\partial z^2} - \frac{\rho c}{k} w \frac{\partial T}{\partial z} = 0. \quad (3)$$

Using $\frac{\partial k}{\partial z} = \frac{\partial k}{\partial T} \frac{\partial T}{\partial z}$, Eq. (3) becomes

$$\frac{\partial^2 T}{\partial z^2} + \left(\frac{1}{k} \frac{\partial k}{\partial T} \frac{\partial T}{\partial z} - \frac{\rho c}{k} w \right) \frac{\partial T}{\partial z} = 0. \quad (4)$$

According to Fischer et al. (2013), in the most general terms, the vertical velocity in the ice can be approximated by

$$w(z) = -w_{\text{melt}} - (\text{Acc} - w_{\text{melt}}) \left(\frac{z}{H} \right)^{m+1}, \quad (5)$$

where w_{melt} is the basal melt rate, Acc is the surface accumulation rate dependent on t , and m is an adjustable form factor that accounts for the variation in horizontal velocity.

Substitution of Eq. (5) into Eq. (4) and integrating on the assumption that k is constant gives the following temperature distribution in the ice sheet at steady state:

Table 3. Polynomial approximations of borehole temperature T ($^{\circ}$ C) as a function of true vertical depth z and correlation factors.

| Drill sites | Polynomial | R^2 |
|-------------|--|-------|
| Byrd | $T = -28.343 + 0.8367 \times 10^{-3}z - 6.7651 \times 10^{-6}z^2 + 6.1339 \times 10^{-9}z^3$ | 0.997 |
| WAIS Divide | $T = -31.799 + 8.8595 \times 10^{-3}z - 9.4649 \times 10^{-6}z^2 + 2.657 \times 10^{-9}z^3$ | 0.997 |
| Vostok | $T = -56.034 + 2.9889 \times 10^{-3}z + 3.888 \times 10^{-6}z^2 + 0.2419 \times 10^{-9}z^3$ | 0.999 |
| Dome C | $T = -54.316 + 5.2978 \times 10^{-3}z + 4.4141 \times 10^{-6}z^2 - 0.368 \times 10^{-9}z^3$ | 0.999 |
| Kohnen | $T = -44.428 + 1.7384 \times 10^{-3}z + 4.4124 \times 10^{-6}z^2 + 0.184 \times 10^{-9}z^3$ | 0.999 |
| Dome F | $T = -55.016 + 5.839 \times 10^{-3}z + 5.188 \times 10^{-6}z^2 - 0.446 \times 10^{-9}z^3$ | 0.998 |

$$T = T_s - \left[\frac{\partial T}{\partial z} \right]_B \int_0^z \exp \left(\frac{(w_{melt} - Acc)z^{m+2}}{\alpha_T(m+2)H^{m+1}} - \frac{w_{melt}}{\alpha_T}z \right) dz \\ + \left[\frac{\partial T}{\partial z} \right]_B \int_0^H \exp \left(\frac{(w_{melt} - Acc)z^{m+2}}{\alpha_T(m+2)H^{m+1}} - \frac{w_{melt}}{\alpha_T}z \right) dz, \quad (6)$$

where T_s is the surface temperature, $\left[\frac{\partial T}{\partial z} \right]_B$ is the temperature gradient at the ice-sheet base, and $\alpha_T = k/\rho c$ is the thermal diffusivity of ice.

The least squares method was used to fit measured borehole temperatures with this equation. In fitting, the initial values of the unknown parameters $T_s \left[\frac{\partial T}{\partial z} \right]_B$, Acc, and w_{melt} can only be guessed, and this results in unavoidable uncertainty of fitting. To overcome this large uncertainty, a common genetic algorithm was used to find the optimal global solution of temperature fitting by constraining these unknown parameters to a predetermined range (Reeves and Rowe, 2002). The GA can solve optimization problems by limiting the unknown parameters to a predetermined range with any type of constraints, including integer constraints. In general, the GA generates high-quality solutions for optimization problems and search problems.

In the GA, the crossover fraction is set to be 0.9, while the migration fraction is 0.2 (Reeves and Rowe, 2002). To obtain an accurate solution and save calculation time, we set the population size to be 8000 and the number of generations to be 20. Usually, after 15 generations of iteration, the optimal solution can be found. All the calculations were performed using MATLAB software. The GA provides results for the first generation of the optimal solution in a wide range based on a random combination of the fitting parameters. Thus, for each deep borehole, the fitting experiments were trialed 5 times to avoid random error of the GA caused by the initial random parameter combination. Then, the average value from the five fitting experiments was used as the GHF from bedrock into the ice sheet at the selected site.

Equation (4) can also be re-expressed as follows:

$$w(z) = \left[\alpha_T \left(\frac{\partial^2 T}{\partial z^2} \right) - \alpha_T \left(\frac{1}{k} \frac{\partial k}{\partial T} \right) \left(\frac{\partial T}{\partial z} \right)^2 \right] / \frac{\partial T}{\partial z}. \quad (7)$$

Note that the vertical velocity is markedly affected by $\frac{\partial^2 T}{\partial z^2}(z)$ and $\frac{\partial T}{\partial z}(z)$. At the base of the ice sheet, the melt/freezing rate is $w_{melt} = w(0)$, while the gradient is $\left[\frac{\partial T}{\partial z} \right]_B = \frac{\partial T}{\partial z}(0)$. The geothermal heat flux Q_{geo} from below the ice is balanced by the conductive flux in the ice $q = k \left[\frac{\partial T}{\partial z} \right]_B$ and the rate at which energy is used to melt or freeze ice, $J = \rho L w_{melt}$. Thus, the GHF will be

$$Q_{geo} = \rho L w_{melt} - k \left[\frac{\partial T}{\partial z} \right]_B, \quad (8)$$

where L is the specific latent heat for the melting of ice.

2.3 Uncertainties

In our method, the temperature in the lower portion of the ice sheets is assumed to be in steady state, and the GA is used to fit the measured temperatures in deep ice-core drilling boreholes by varying the four key parameters influencing the temperature distribution: the surface temperature, surface accumulation rate, basal melt, and basal temperature gradient. All these parameters are suggested by the algorithm in order to obtain the best-fitting curve. We assume that the main uncertainties in our fitting model come from temperature measurements, variability of the form factor m , ice-thickness estimation, and the GA itself. It must be noted that the uncertainties we state are lower limits. There are some additional unexamined uncertainties that were missing from our model, including transient effects associated with climate change and ice-sheet dynamics, the horizontal velocity field, the form of the vertical velocity field, the temperature dependence of the thermal conductivity, thermal disturbances due to drilling processes that are unaccounted for, 2D effects, and some other phenomena.

2.3.1 Temperature measurements

Interpretation of temperature measurements in mechanically drilled deep boreholes filled with drilling fluids is complicated by several factors (Clow, 2008). First, the temperature is measured in the borehole fluid, not in the surrounding ice; therefore, an important consideration is the need for thermal equilibration of the ice wall and the borehole fluids following

drilling and prior to measurement. Second, the heat produced during drilling needs to be dissipated from the borehole, or the thermal drilling disturbance needs to be accounted for (Clow, 2015). Third, increasing temperature with depth can cause convective mixing in the borehole. Fourth, the depth of temperature measurements has an inherent uncertainty due to cable slippage in the counting assembly and cable elongation. Thus, all successful temperature measurements in deep boreholes obey a logging protocol in terms of logger tripping speed, measurement direction, borehole settling time, and so on to minimize the effects of these complicating factors. Temperature measurement errors from sensor accuracy and calibration are found to be within the tolerance for large-scale GHF estimates for our six boreholes to interpret ice-sheet basal dynamics.

The temperature in the Byrd borehole was measured with an accuracy of $0.1\text{ }^{\circ}\text{C}$ (Herbert Ueda, personal communication, 2017). Motoyama et al. (2013) reported that temperature measurements at Dome F were carried out with a precision of $0.05\text{ }^{\circ}\text{C}$. The absolute temperature measurement error at Vostok was estimated to be $0.07\text{ }^{\circ}\text{C}$ (Salamatin et al., 1998a). The resolution of the temperature measurements in the Dome C borehole was $0.015\text{ }^{\circ}\text{C}$, while the precision was found to be $0.05\text{ }^{\circ}\text{C}$ (Lefebvre et al., 2002). The logger used in the borehole at Kohnen was calibrated to $0.03\text{ }^{\circ}\text{C}$ (Gundersen et al., 1994). Prior to drilling, a detailed study of the expected temperature measurement uncertainties was made for the WAIS Divide site to optimize the logging system setup (Clow, 2008). The standard uncertainty (accuracy) of the subsequent WAIS Divide temperature logs was $\sim 0.0053\text{ }^{\circ}\text{C}$ (Cuffey et al., 2016). In general, temperature measurement accuracy in the studied boreholes is more than adequate, and the measured drilling depth was recalculated to true vertical depth using available borehole inclinations.

2.3.2 Form factor m

Selection of the appropriate form factor m is a challenging task. Classically, vertical velocity depends linearly on z/H (Cuffey and Paterson, 2010) and $m = 0$. However, at an ice divide, the downward flow of ice is slower for the same depth than at locations away from the divide (Raymond, 1983). This reduces the cooling influence of vertical advection and increases the basal temperature. Therefore, Raymond (1983) suggested the use of $m = 1.0$ for deformation in the vicinity of ice divides.

To set up the vertical velocity profile at Dome C, Fischer et al. (2013) performed three runs with $m = 0.3$, $m = 0.5$, and $m = 0.7$ and found that the temperature profile is only slightly affected by this choice. However, the form factor m exhibited a strong influence on the age profile of the ice. That was the reason why the authors used $m = 0.5$, which is in good agreement with the EDC3 age scale (Parrenin et al., 2007b). Following the method of Fischer et al. (2013), we

modeled the age profile of ice by

$$\hat{A} = \int_z^H \frac{1}{w_{\text{melt}} + (\text{Acc} - w_{\text{melt}}) \left(\frac{z}{H}\right)^{m+1}} dz, \quad (9)$$

where \hat{A} is the modeled ice age. Then, the modeled ice age is used to compare with the published depth–age scales at the studied sites, and the best value for the form factor m is estimated. In order to reduce the run time of multilevel calculations, we examine the form factor m at only five levels: 0, 0.25, 0.50, 0.75, and 1.00. The best value for the form factor m is selected on the basis of the nonlinear correlation analysis between modeled and measured age scales. To calculate the correlation factor R^2 , we first found the average value of the measured age:

$$\bar{A} = \frac{1}{n} \sum_{i=1}^n A_i, \quad (10)$$

where n is the number of measured ice ages A_i . Then the total sum of squares SS_{tot} and the sum of squares of residuals SS_{res} were calculated:

$$SS_{\text{tot}} = \sum_{i=1}^n (A_i - \bar{A})^2, \quad (11)$$

$$SS_{\text{res}} = \sum_{i=1}^n (A_i - \hat{A}_i)^2, \quad (12)$$

where \hat{A}_i is the modeled ice age when $n = i$. The correlation factor R^2 was estimated by

$$R^2 = 1 - \frac{SS_{\text{res}}}{SS_{\text{tot}}}. \quad (13)$$

Finally, the results of the nonlinear correlation analysis were checked by evaluating the root mean square error (RMSE):

$$\text{RMSE} = \sqrt{\frac{1}{n} \sum_{i=1}^n (A_i - \hat{A}_i)^2}. \quad (14)$$

2.3.3 Ice thickness

We assume that the ice-sheet thickness at the studied sites has kept constant at the present-day height; however, it has varied in the past. The 3D thermomechanical model and the simple 1D model showed that the maximum variation in ice-sheet thickness at Dome C and Dome F was less than 250 m in the past (Parrenin et al., 2007a). In general, the typical difference in the ice thickness in the glacial and interglacial periods at Dome C was 150 m (Passalacqua et al., 2017). At the Kohnen site, the local elevation variation is on the order of 100 m (Huybrechts et al., 2007). The ice-thickness variation at Vostok, located in the central part of the East Antarctica

Plateau, exhibits a similar range as at Dome F and Dome C (Ritz et al., 2001).

The best evidence for ice-sheet elevation change in the interior of the West Antarctic Ice Sheet comes from the Ohio Range, to the south of the WAIS Divide site at a height of 1600 m a.s.l., and from Mt. Waesche, to the north of the WAIS Divide site at a height of 2000 m a.s.l. (Ackert et al., 1999, 2007). Moraines at Mt. Waesche were ~ 50 m higher, and trimlines in the Ohio Range were ~ 125 m higher, between 12 and 10 ka. The thinning of ~ 100 m throughout the Holocene occurred as the grounding line retreated by hundreds of kilometres, and the accumulation rates were relatively stable (Anderson et al., 2002; Conway et al., 1999). Cuffey et al. (2016) presented a model which indicates a more likely scenario of 200 m thickening at WAIS Divide when the accumulation rate rose after the last glacial maximum, followed by 300 m of thinning to the mid-Holocene. The elevation change is comparable to the amount of elevation change inferred for interior East Antarctic sites.

Comparison with the modern ice-thickness value indicates that the variation in ice thickness is small, and its influence on ice temperature distribution can be neglected, in particular, on the lower portion of the ice borehole. For example, assuming a 150 m thickness increase from the last glacial maximum (LGM) to 15 ka leads to the change in the reconstructed LGM temperature by less than 0.2°C compared to a constant thickness in the WAIS ice core (Buzert et al., 2015). This is the reason why constant ice thickness is also used by other researchers for GHF estimates (Dahl-Jensen et al., 2003; Engelhardt, 2004; Mony et al., 2020).

2.3.4 Genetic algorithm

For each deep borehole, the fitting experiments were repeated 5 times for the best value of the form factor m , and the average value obtained from the five fitting experiments was used as the representative value of GHF from bedrock into ice. Thus, the uncertainty ranges came from the difference between the maximum and minimum and the average GHF values.

3 Results

3.1 Initial conditions and GHF estimates

GHF estimates were made using the following ice parameters:

- density of 918 kg m^{-3} ;
- specific heat capacity of $c = 152.5 + 7.122 (T + 273.15) \text{ J kg}^{-1} \text{ K}^{-1}$ (Yen, 1981);
- thermal conductivity of $k = 9.828 e^{-0.0057(T+273.15)} \text{ W m}^{-1} \text{ K}^{-1}$ (Yen, 1981);

- specific latent heat of $L = 333.5 \text{ kJ kg}^{-1}$ (Cuffey and Paterson, 2010).

In our model, we assume that k and c are constant and equal to their values at the temperature of the pressure melting point; this can provide a better estimate of the basal melting rate at the base of the ice sheet (Fischer et al., 2013). In this case, $\frac{1}{k} \frac{\partial k}{\partial T} = -5.7 \times 10^{-3} \text{ K}^{-1}$. Figure 3 shows the fitted temperature profiles compared with measured temperatures.

We performed five runs for estimating GHF with $m = 0$, $m = 0.25$, $m = 0.5$, $m = 0.75$, and $m = 1.0$ for each site and compared modeled and measured age scales. As illustrated in Fig. 4, the form factor has a strong influence on the age profile of the ice. The results of our estimates for GHF for different m values are summarized in Table 4. Surprisingly, on three separate occasions the correlation factor is negative. This may occur when SS_{res} is far beyond that of SS_{tot} (see Eqs. 10–12). That is to say that there is no correlation between modeled and measured age scales in these cases. The results of the correlation analyses were confirmed by evaluating how close the modeled lines are to the data points with the aid of the RMSE: the smaller the RMSE, the closer the model is to the data. The GHF and m values with the highest correlation factor and smallest RMSE were selected for further processing and trialed 5 times. The average GHF values for the selected m are added into Table 2. The precision of the GHF estimates and basal melt or freezing rate are also specified here.

The temperature profiles show that the heat flow through the ice at six deep drilling sites in Antarctica must be $> 42.6\text{--}77.1 \text{ mW m}^{-2}$ in order to match the observed temperatures in the boreholes. The basal ice at all sites is at the pressure melting point, and the amount of melt cannot be constrained by the energy-balance equation alone. When the heat flow model is combined with vertical-velocity estimates, the estimated heat flow can be translated to a GHF of $57.9\text{--}113.3 \text{ mW m}^{-2}$, except for Vostok, with GHF of -3.3 mW m^{-2} .

3.2 Data comparison and divergences

3.2.1 Vostok

The surface temperature time curve for the upper bound of the present-day accumulation rate at Vostok corresponds to a GHF of 53 mW m^{-2} (Salamatin et al., 1998b). We calculated that at the base of the ice sheet, the conductive flux is $42.6 \pm 0.4 \text{ mW m}^{-2}$, while the latent heat flux from refreezing of the lake water is $46.3 \pm 5.6 \text{ mW m}^{-2}$. Thus, the GHF heat flux at the base of the ice sheet has a negative value of $-3.6 \pm 5.3 \text{ mW m}^{-2}$. This is in good agreement with the isotope studies that showed that the Vostok ice core consists of ice refrozen from Lake Vostok water, from 3539 m below the surface of the Antarctic Ice Sheet to its bottom (Jouzel et al., 1999). A sufficiently high correlation factor (0.75) between modeled and measured age scales at $m = 1$ indicates that ice




Figure 3. Temperatures measured in Antarctic deep ice boreholes compared with best-fit temperature profiles for the deepest 1500 m.

above Lake Vostok reasonably fits Raymond's (1983) arguments for deformation in the vicinity of ice divides.

At this stage we are not yet able to predict GHF at the bed of 600 m thick subglacial Lake Vostok because the temperature profile in the lake is still indefinite. However, the DNA detection of thermophile bacteria in the near-base accretion ice suggests the existence of near-bottom warm waters with temperatures as high as 50 °C (Bulat et al., 2012). If so, the GHF in the lake sediments can reach 200–240 mW m⁻². These values can be considered as paleo-GHF because microorganisms were picked up thousands of years ago but still actually account for long durations of geological processes.

3.2.2 Dome C

The inverse approach to retrieving GHF from radar-inferred distribution of wet and dry beds at the EPICA drilling site (Passalacqua et al., 2017) gave 54.5 ± 3.5 mW m⁻², slightly

lower than estimates derived from borehole temperature profiling (57.9 ± 6.4 mW m⁻²). The modeled GHF range (43–55 mW m⁻² obtained by An et al., 2015; Fox Maule et al., 2005; Shapiro and Ritzwoller, 2004; and Van Liefferinge and Pattyn, 2013) is also a little less than our estimates. The high value for the correlation factor (0.997) indicates a perfectly strong relationship between modeled and measured depth-age scales, meaning that there is no horizontal advection of heat, and the drill site is located at a perfect dome position. Perhaps that is the reason that the core from Dome C contains the oldest continuous climate record obtained from ice cores so far (Parrenin et al., 2007b). However, a high spatial variation in GHF in the area of Dome C was found from radar-sounding data (Carter et al., 2009). The values of nearly 100 mW m⁻² inferred for the southern shore of Concordia Subglacial Lake, approximately 50 km to the south of the drilling site, are also well outside modeled estimates.




Figure 4. Comparison of the measured age scales (Ahn and Brook, 2008; Bazin et al., 2013; Bereiter et al., 2012; Blunier and Brook, 2001; Kawamura et al., 2007; Neftel et al., 1988; Parrenin et al., 2007b; Sigl et al., 2016; Staffelbach et al., 1991; Veres et al., 2013) and modeled age scales with $m = 0$, $m = 0.25$, $m = 0.5$, $m = 0.75$, and $m = 1.0$ (correlation factors between modeled and measured age scales for each run are stated in Table 4).

Table 4. GHF (mW m^{-2}) calculated for different form factors m in the steady-state model, the correlation factor R^2 between modeled and measured age scales, and RMSE.

| Parameters | WAIS | | EAIS | | | |
|--------------------|--------------|--------------|--------------|--------------|--------------|--------------|
| | Byrd | WAIS Divide | Vostok | Dome C | Kohnen | Dome F |
| GHF for $m = 0$ | 8.72 | 72.9 | -16.16 | 134.6 | 66.04 | 92.78 |
| R^2 | -0.37 | -2.46 | 0.65 | 0.69 | 0.978 | 0.78 |
| RMSE | 28.19 | 36.40 | 58.95 | 107.51 | 4.132 | 40.62 |
| GHF for $m = 0.25$ | 28.44 | 100.0 | -35.28 | 105.3 | 60.95 | 130.9 |
| R^2 | 0.12 | 0.59 | 0.50 | 0.72 | 0.986 | -1.62 |
| RMSE | 22.60 | 12.93 | 70.60 | 102.14 | 2.431 | 140.91 |
| GHF for $m = 0.50$ | 55.05 | 207.4 | -20.33 | 70.06 | 156.4 | 92.96 |
| R^2 | 0.46 | 0.41 | 0.51 | 0.87 | 0.976 | 0.47 |
| RMSE | 17.59 | 15.05 | 69.74 | 69.48 | 4.338 | 62.91 |
| GHF for $m = 0.75$ | 95.84 | 240.3 | -25.39 | 57.40 | 106.0 | 104.8 |
| R^2 | 0.57 | 0.32 | 0.39 | 0.997 | 0.984 | 0.53 |
| RMSE | 14.86 | 16.15 | 78.01 | 22.34 | 3.477 | 59.25 |
| GHF for $m = 1.00$ | 117.8 | 251.3 | -8.90 | 67.3 | 161.5 | 79.20 |
| R^2 | 0.45 | 0.29 | 0.75 | 0.95 | 0.982 | 0.83 |
| RMSE | 17.83 | 16.45 | 49.71 | 43.31 | 3.692 | 35.87 |

GHF values with the highest correlation factor and smallest RMSE are highlighted by bold.

3.2.3 Kohnen

The model with a standard GHF of 54.6 mW m^{-2} predicted a basal temperature 0.3°C below the pressure melting point at Kohnen (Huybrechts et al., 2007). GHF values obtained by nonthermal geophysical models are in the range of $46\text{--}62 \text{ W m}^{-2}$. Our estimate ($86.9 \pm 16.6 \text{ mW m}^{-2}$) is higher than the modeled GHF values suggested by Fox Maule et al. (2005), Martos et al. (2017), and other referenced models. However, subglacial water entering the borehole indicated that the actual GHF should be much higher than that indicated by the regional models. Under these circumstances, our estimate is likely to be closer to the real heat flux. Surprisingly, the depth–age scale is only slightly affected by the choice of the form factor, indicating that the variation in horizontal velocity is low at this site.

3.2.4 Dome F

A previously estimated GHF of 59 mW m^{-2} neglected the bottom ice melt rate (Hondoh et al., 2002) and thus is lower than our estimate ($78.9 \pm 5.0 \text{ mW m}^{-2}$). Mony et al. (2020) estimated the GHF in the Dome F borehole to be even lower, at 50.4 mW m^{-2} . As the drill approached the base (approx. 10 m above), subglacial meltwater leaked into the borehole and froze onto the drill, directly indicating that ice reaches the pressure melting point, placing a lower bound on the GHF. GHF values obtained by nonthermal geophysical methods are in the range of $48\text{--}65 \text{ mW m}^{-2}$, also lower than our estimates. The correlation factor between modeled and mea-

sured depth–age scales is quite high (0.83) at $m = 1$, indicating that ice at the site can be adequately approximated by the steady-state model. Thus, the slightly elevated heat flow at this location appears to represent a regional value.

3.2.5 Byrd

Unfortunately, age scales for the Byrd borehole for all modeled m values are quite far from the measured depth–age data. Tilting measurements (Garfield and Ueda, 1976; Hansen et al., 1989) and modeling (Whillans, 1979) showed that the relative horizontal velocity of ice at this borehole reaches $\sim 3 \text{ m a}^{-1}$ at 1500 m depth. Thus, horizontal conduction at the bottom of the ice sheet is quite high at this site, producing a high divergence from the steady-state model. Even the correlation factor for the best-fitting age–depth curve with $m = 0.75$ is only 0.58; we can use the GHF value of $88.4 \pm 7.6 \text{ mW m}^{-2}$ at this location as a reference until more precise estimates are obtained. This value is higher than the first estimate made immediately after temperature logging (75.4 mW m^{-2} referenced by Ueda, 2007), primarily because the latter one did not account for the basal ice melt. The latest modeling by Martos et al. (2017) revealed a high GHF at the location of Byrd Station (132 mW m^{-2} with an error of $\pm 5 \text{ mW m}^{-2}$) when compared with values obtained from previous models (An et al., 2015; Fox Maule et al., 2005; Van Liefferinge and Pattyn, 2013). Generally, our approximate estimate is in the range of GHF values suggested by previous models.

3.2.6 WAIS Divide

A preliminary estimate of the GHF at this site suggested a high value in the range of $200\text{--}230 \text{ mW m}^{-2}$, depending on the actual ice thickness (Clow et al., 2012). This is more than twice that derived using nonthermal geophysical methods. There is no depression in the local surface topography or drawdown in the subsurface layers detected by ice-penetrating radar, as would be expected over a local hot spot. Using airborne magnetic data, Martos et al. (2017) estimated the highest value for this area to be $\sim 120 \text{ mW m}^{-2}$. Our new estimate is slightly lower, at $113.3 \pm 16.9 \text{ mW m}^{-2}$. Mony et al. (2020) also estimated GHF from the borehole temperature profile at WAIS Divide by combining a heat-transfer equation and the physical properties of the ice sheet in a numerical model. Based on a truncated temperature profile, they estimate a GHF of 90.5 mW m^{-2} , which is less than ours and fairly corresponds to the latest GHF map for Antarctica constructed by empirically relating the upper-mantle seismic structure (Shen et al., 2020). The low correlation factor (0.59) between modeled and measured depth–age scales in our present estimate indicates that there are some important uncertainties that the steady-state model does not account for. Most likely, these are the same unaccounted effects that affect the Byrd borehole temperature profile, i.e., horizontal flow and climate-related transient effects.

The preliminary GHF estimate (Clow et al., 2012) was based on the first temperature log in 2011 in the borehole before it reached its final depth. The reasons why the preliminary GHF estimate may be so high are that (i) temperatures in the borehole were still thermally disturbed in 2011, and (ii) the bottom of the 2011 temperature log was still far from the base of the ice sheet. The borehole was relogged in 2014, and temperature data were obtained much closer to the bed. In addition, Clow et al. (2012) also did not account for horizontal flow effects, and the GHF estimation could have been lower than the one they produced. Further investigations on ice dynamics through WAIS Divide borehole tilt measurements can allow us to determine in-depth stress and velocity distributions and estimate horizontal flow effects on temperature.

3.3 Indirect results

Although a steady-state model is used in the lower portion of the boreholes to describe the temperature distribution, it is worth noting that the measured modern temperature is the cumulative effect of historical climate forcing. Therefore, the best-fitting parameters obtained by the GA are not the real parameters occurring during the ice sheet's history. They can be considered as “equivalent” parameters, which are used for calculating the modern temperature profile by eliminating the historical climate changes. Processing back, the “equivalent” vertical velocity, modern accumulation rate, and temperature can be calculated from the GA results. Estimated




Figure 5. Estimated vertical velocities at drilling sites in West Antarctica (a) and East Antarctica (b). In East Antarctica snow accumulation and thus vertical ice velocities are far less than in West Antarctica.

vertical velocity profiles are shown in Fig. 5. Table 5 lists values of “equivalent” snow accumulation rate and temperature at the ice-sheet surface, which were derived from GA calculations. In all cases, “equivalent” accumulation rates are higher than the modern rates, while the “equivalent” surface temperatures are very close to the modern ones. This can be explained by the fact that the high “equivalent” accumulation rates are used by the GA to eliminate the colder climate effects on the ice temperature profile during the glacial period.

4 Discussion

4.1 Transient model vs. steady-state model

Both transient thermal models (e.g., Dahl-Jensen et al., 2003; Engelhardt, 2004; Martos et al., 2017; Passalacqua et al., 2017; Van Liefferinge et al., 2018) and steady-state models (e.g., Martin and Gudmundsson, 2012; Mony et al., 2020; Parrenin et al., 2017; Price et al., 2002; Zagorodnov et al., 2012) were used intensively in the past and are still used for GHF estimates in Antarctica. Obviously, an exact steady state never occurs in reality, and thus transient models would be expected to give more precise results than steady-state models. However, the answer is not as simple as it is supposed to be.

It is important to recognize that, first, in both cases the models will produce GHF “estimates”, not “measurements”, and second, the thermal gradient can be affected by processes other than GHF, creating local anomalies that may coincide with the point estimate. In order to use a transient model, the accumulation rate and surface temperature in the past should be known. For some of the discussed drill sites these data are available from ice-core studies, while for other sites they are not.


Table 5. Equivalent thermophysical parameters used by the GA in comparison with published data.

| Parameters | Byrd | WAIS Divide | Vostok | Dome C | Kohnen | Dome F |
|--|------|-------------|--------|--------|--------|--------|
| “Equivalent” snow accumulation at surface (cm ice a^{-1}) | 52.8 | 48.8 | 8.95 | 3.87 | 6.92 | 3.00 |
| Modern snow accumulation at surface (cm ice a^{-1}) | 16.9 | 22.0 | 2.48 | 2.84 | 7.00 | 2.99 |
| “Equivalent” surface temperature ($^{\circ}$ C) | -29 | -30.9 | -56.5 | -54.7 | -44.9 | -56.5 |
| Modern surface temperature ($^{\circ}$ C) | -28 | -30 | -57 | -54.6 | -44 | -57.3 |

To evaluate the possibility of using a transient model, the GHF at WAIS Divide was estimated by using the accumulation rate and surface temperature in the past provided by Buizert et al. (2015). In these calculations, we assume that the history of the ice sheet at WAIS Divide is about 68 kyr long. The governing equation for the transient model was solved using the finite-difference method. The equation was discretized by both the central-difference and upwind-difference methods and then solved using MATLAB. To find the best solution, the GA was still used. The central-difference method and upwind-difference method demonstrated the same temperature profile. Therefore, here we present the calculation results obtained via the upwind-difference method.

Unfortunately, the calculation results with the transient model showed the best-fit GHF value of $\sim 500 \text{ mW m}^{-2}$ when $m = 1$, which seems to be unrealistic. Moreover, after running the model, we found that after about 4–8 kyr, the influence of initial temperature on temperature profile can be ignored. Later, we assumed the form factor $m = 0$ and a GHF value of 235 mW m^{-2} , which showed a good fit with the measured temperatures, although the GHF is much higher than our earlier estimate and estimates from regional models. The temperature distribution throughout history was modeled for 61.2, 54.4, … 6.8 kyr ago until modern day (Fig. 6). As expected, the modeled temperature in the upper part of the ice sheet grossly changes with time, but in the lower portion ($\sim 1000 \text{ m}$ above ice-sheet base) these variations are much smaller. This indicates that the heat disturbance related to atmosphere forcing (temperature and snow precipitation) from the ice-sheet surface gradually decays with the depth. From all appearances, the near-basal portion is close to a steady state.

Both models lack additional heat sources (i.e., shear heating, heat advection, and basal frictional heating) that might be generated at the bottom of the ice sheet. Thus, the results of both modeling approaches strongly depend on the selected initial parameters, in particular, from the selected value for the form factor m . Experimental validation of both models for adequacy is extremely difficult, and recently both of them have rights to exist if assumptions are examined analytically.

**Figure 6.** Paleo-temperature profiles at WAIS Divide based on transient model ($m = 0$).

4.2 Implications of elevated heat flux

Most numerical models of the EAIS basal conditions assume the GHF to be $42\text{--}65 \text{ mW m}^{-2}$. However, the presence of basal meltwater beneath most of the Antarctic Ice Sheet requires $\text{GHF} \geq 80 \text{ mW m}^{-2}$ (Budd et al., 1984). In support of this conjecture, processing of available temperature profiles in ice boreholes shows that at sites where subglacial water exists, bedrocks are quite warm. Currently, a warm base beneath the EAIS was also confirmed by tectonic reconstructions (Carson et al., 2014). Additional evidence of high GHF at EAIS locations comes from ice-penetrating radar data that revealed a $\sim 100 \text{ km}$ long and 50 km wide area near the South Pole with GHF of $120 \pm 20 \text{ mW m}^{-2}$, more than double the values expected for this cratonic sector of East Antarctica (Jordan et al., 2018). This warm base could be caused by radiogenic heat effects or hydrothermal circulation, but a coherent explanation for this phenomenon is still required.

Variability of crustal thickness, hydrothermal circulation (Seroussi et al., 2017), magmatic intrusion (Van Wyk De

Vries et al., 2017), and thermal-conductivity variability are the main contributors to the elevated and highly variable values of GHF in West Antarctica (Begeman et al., 2017). One of the first pieces of evidence for an “unreasonably high” GHF ($> 100 \text{ mW m}^{-2}$) under WAIS came from temperature–depth profiles in a 480 m deep borehole drilled at Crary Ice Rise, Antarctica (Bindschadler et al., 1990). Measured GHF in Subglacial Lake Whillans was found to be $285 \pm 80 \text{ mW m}^{-2}$, significantly higher than the continental and regional averages estimated for this site by using geo-physical and glaciological models (Fisher et al., 2015). The GHF at the vents of subglacial volcanoes in West Antarctica can be as high as 25 W m^{-2} , and one such 23 km wide caldera was revealed $\sim 100 \text{ km}$ to the south of the WAIS Divide drill site (Blankenship et al., 1993). Undeniably, more systematic explorations are still required to study how far this heat flow high extends into the interior of the West Antarctic rift system.

5 Conclusions

Prediction of the future behavior of the Antarctic Ice Sheet undeniably requires accurate ice-sheet models. However, GHF models based on seismic tomography, radar data, magnetic field observations, the tectonic age, and geological structure of the bedrock yield mixed results at sites of deep ice-core drilling in Antarctica. We suggested to estimate GHF from ice-borehole temperature profiles using a one-dimensional steady-state energy-balance equation and the genetic algorithm (GA) for determining the optimal solution of temperature fitting. To our knowledge, we used the GA approach for the first time in ice thermodynamics. Comparison of modeled and measured depth–age scales shows that our model is able to assess the variation in GHF estimates from ice-borehole temperature profiles if in-depth horizontal ice velocities are low and can be ignored. The correlation analyses at the EAIS sites indicate that all of them can be adequately approximated by the steady-state model. However, horizontal velocities and their variation over ice-age cycles are much greater at WAIS than at the EAIS sites. Thus, the steady-state model cannot precisely describe temperature distribution here.

At three studied EAIS sites (Dome C, Dome F, and Kohnen), the GHF is higher than that predicted by other models. We assume that this elevated GHF can represent regional value and can be used as a reference point for regional modeling. More precise GHF estimates and explanations for an elevated GHF would be possible after temperature logging and subglacial rock studies from deep boreholes that are required to drill in Antarctica in the distant future. Finally, the proposed method of GHF estimates can be used at other sites in Antarctica and Greenland where the steady-state model is acceptable.

Data availability. The data that support the findings in this study are available from the corresponding authors.

Author contributions. PT and YL developed the concept, wrote the manuscript, and drew all figures. YL also designed and performed all GHF estimations. JH processed all temperature profiles. GDC has made important recommendations and edited the final version of the paper. LA, GDC, EL, AM, HM, and CR provided observed borehole temperature data. All authors contributed to discussion and interpretation of the results.

Competing interests. The authors declare that they have no conflict of interest.

Acknowledgements. We are grateful to Herbert Ueda (retired from USA CRREL) for providing original Byrd borehole temperature log data. We thank Dorthe Dahl-Jensen (Physics of Ice, Climate and Earth group, Niels Bohr Institute, University of Copenhagen, Denmark) for fruitful discussion and useful comments. We also would like to thank the editor Alex Robinson and both anonymous reviewers for fruitful comments and advice.

Financial support. This research has been supported by the National Natural Science Foundation of China (grant nos. 41327804 and 41806220) and the Fundamental Research Funds for the Central Universities (grant no. 2017TD-24).

Review statement. This paper was edited by Alexander Robinson and reviewed by two anonymous referees.

References

- Ackert, R. P., David, J. B., Harold, W. B., Parker, E. C., Mark, D. K., James, L. F., and Eric, J.: Measurements of past ice sheet elevations in interior West Antarctica, *Science*, 286, 276–280, <https://doi.org/10.1126/science.286.5438.276>, 1999.
- Ackert, R. P., Mukhopadhyay, S., Parizek, B. R., and Borns, H. W.: Ice elevation near the West Antarctic Ice Sheet divide during the Last Glaciation, *Geophys. Res. Lett.*, 34, L21506, <https://doi.org/10.1029/2007GL031412>, 2007.
- Ahn, J. and Brook E. J.: Atmospheric CO₂ and Climate on Millennial Time Scales During the Last Glacial Period, *Science*, 322, 83–85, <https://doi.org/10.1126/science.1160832>, 2008.
- Anderson, J. B., Shipp, S. S., Lowe, A. L., Wellner, J. S., and Mosola, A. B.: The Antarctic Ice Sheet during the Last Glacial Maximum and its subsequent retreat history: a review, *Quaternary Sci. Rev.*, 21, 49–70, [https://doi.org/10.1016/S0277-3791\(01\)00083-X](https://doi.org/10.1016/S0277-3791(01)00083-X), 2002.
- An, M., Wiens, D. A., Zhao, Y., Feng, M., Nyblade, A., Kanao, M., Li, Y., Maggi, A., and Lévéque, J.-J.: Temperature, lithosphere–asthenosphere boundary, and heat flux beneath the Antarctic

- Plate inferred from seismic velocities, *J. Geophys. Res.-Sol. Ea.*, 120, 8720–8742, <https://doi.org/10.1002/2015JB011917>, 2015.
- Augustin, L., Panichi, S., and Frascati, F.: EPICA Dome C 2 drilling operations: performances, difficulties, results, *Ann. Glaciol.*, 47, 68–72, <https://doi.org/10.3189/172756407786857767>, 2007.
- Bazin, L., Landais, A., Lemieux-Dudon, B., Toyé Mahamadou Kele, H., Veres, D., Parrenin, F., Martinerie, P., Ritz, C., Capron, E., Lipenkov, V., Loutre, M.-F., Raynaud, D., Vinther, B., Svensson, A., Rasmussen, S. O., Severi, M., Blunier, T., Leuenberger, M., Fischer, H., Masson-Delmotte, V., Chappellaz, J., and Wolff, E.: An optimized multi-proxy, multi-site Antarctic ice and gas orbital chronology (AICC2012): 120–800 ka, *Clim. Past*, 9, 1715–1731, <https://doi.org/10.5194/cp-9-1715-2013>, 2013.
- Begeman, C. B., Tulaczyk, S. M., and Fisher, A. T.: Spatially variable geothermal heat flux in West Antarctica: Evidence and implications, *Geophys. Res. Lett.*, 44, 9823–9832, <https://doi.org/10.1002/2017gl075579>, 2017.
- Bereiter, B., Lüthi, D., Siegrist, M., Schüpbach, S., Stocker, T. F., and Fischer, H.: Mode change of millennial CO₂ variability during the last glacial cycle associated with a bipolar marine carbon seesaw, *P. Natl. Acad. Sci. USA*, 109, 9755–9760, <https://doi.org/10.1073/pnas.1204069109>, 2012.
- Bindschadler, R. A., Roberts, E. P., and Iken, A.: Age of Crary Ice Rise, Antarctica, determined from temperature-depth profiles, *Ann. Glaciol.*, 14, 13–16, <https://doi.org/10.1017/S0260305500008168>, 1990.
- Blankenship, D. D., Bell, R. E., Hodge, S. M., Brozena, J. M., Behrengt, J. C., and Finn, C. A.: Active volcanism beneath the West Antarctic ice sheet and implications for ice-sheet stability, *Nature*, 361, 526–529, <https://doi.org/10.1038/361526a0>, 1993.
- Blunier, T. and Brook, E. J.: Timing of millennial-scale climate change in Antarctica and Greenland during the last glacial period, *Science*, 291, 109–112, <https://doi.org/10.1126/science.291.5501.109>, 2001.
- Budd, W. F., Jenssen, D., and Smith I. N.: A three-dimensional time-dependent model of three West Antarctic ice-sheets, *Ann. Glaciol.*, 5, 29–36, <https://doi.org/10.1017/S026030550000344X>, 1984.
- Buizert, C., Cuffey, K. M., Severinghaus, J. P., Baggensos, D., Fudge, T. J., Steig, E. J., Markle, B. R., Winstrup, M., Rhodes, R. H., Brook, E. J., Sowers, T. A., Clow, G. D., Cheng, H., Edwards, R. L., Sigl, M., McConnell, J. R., and Taylor, K. C.: The WAIS Divide deep ice core WD2014 chronology – Part 1: Methane synchronization (68–31 ka BP) and the gas age–ice age difference, *Clim. Past*, 11, 153–173, <https://doi.org/10.5194/cp-11-153-2015>, 2015.
- Bulat, S. A., Marie, D., and Petit, J.-R.: Prospects for life in the subglacial Lake Vostok, *Ice and Snow*, 52, 92–96, <https://doi.org/10.15356/2076-6734-2012-4-92-96>, 2012.
- Burton-Johnson, A., Dziadek, R., and Martin, C.: Geothermal heat flow in Antarctica: current and future directions, *The Cryosphere Discuss.*, <https://doi.org/10.5194/tc-2020-59>, in review, 2020.
- Carter, S. P., Blankenship, D. D., Young, D. A., and Holt, J. W.: Using radar-sounding data to identify the distribution and sources of subglacial water: application to Dome C, East Antarctica, *J. Glaciol.*, 55, 1025–1040, <https://doi.org/10.3189/002214309790794931>, 2009.
- Carson, C. J., McLaren, S., Roberts, J. L., Boger, S. D., and Blankenship, D. D.: Blankenship, hot rocks in a cold place: High sub-glacial heat flow in East Antarctica, *J. Geol. Soc. London*, 171, 9–12, <https://doi.org/10.1144/jgs2013-030>, 2014.
- Clow, G. D.: USGS Polar temperature logging system, description and measurement uncertainties: U.S. Geological Survey Techniques and Methods 2–E3, 24 pp., 2008.
- Clow, G. D.: A Green's function approach for assessing the thermal disturbance caused by drilling deep boreholes in rock or ice, *Geophys. J. Int.*, 203, 1877–1895, <https://doi.org/10.1093/gji/ggv415>, 2015.
- Clow, G., Waddington, E., Hawley, R., and Dahl-Jensen, D.: Sub-glacial heat flow measurements in Greenland and Antarctica, European Geosciences Union General Assembly, 3–8 April, 2011, Vienna, Austria, *Geophysical Research Abstracts* 13, abstract id EGU2011-6629, 2011.
- Clow, G. D., Cuffey, K. M., and Waddington, E. D.: High heat-flow beneath the central portion of the West Antarctic Ice Sheet, American Geophysical Union, Fall Meeting 2012, 3–7 December, 2012, San-Francisco, USA, abstract id. C31A-0577, 2012.
- Conway, H., Hall, B. L., Denton, G. H., Gades, A. M., and Waddington, E. D.: Past and future grounding-line retreat of the West Antarctic Ice Sheet, *Science*, 286, 280–283, <https://doi.org/10.1126/science.286.5438.280>, 1999.
- Conway, H. and Rasmussen L. A.: Recent thinning and migration of the Western Divide, central West Antarctica, *Geophys. Res. Lett.*, 36, L12502, <https://doi.org/10.1029/2009GL038072>, 2009.
- Cuffey, K. M. and Paterson, W. S. B.: *The physics of glaciers*, 4th edn., Butterworth-Heinemann, Oxford, 2010.
- Cuffey, K. M., Clow, G. D., Steig, E. J., Buizert, C., Fudge, T. J., Koutnik, M., Waddington, E. D., Alley, R. B., and Severinghaus, J. P.: Deglacial temperature history of West Antarctica, *P. Natl. Acad. Sci. USA*, 113, 14249–14254, <https://doi.org/10.1073/pnas.1609132113>, 2016.
- Dahl-Jensen, D., Morgan, V. I., and Elcheikh, A.: Monte Carlo inverse modelling of the Law Dome (Antarctica) temperature profile, *Ann. Glaciol.*, 29, 145–150, <https://doi.org/10.3189/172756499781821102>, 1999.
- Dahl-Jensen, D., Gundestrup, N., Gogineni, S. P., and Miller, H.: Basal melt at NorthGRIP modeled from borehole, ice-core and radio-echo sounder observations, *Ann. Glaciol.*, 37, 217–212, <https://doi.org/10.3189/172756403781815492>, 2003.
- Davis, J. H.: Global map of solid Earth surface heat flow, *Geochem. Geophys. Geosys.*, 14, 4608–4622, <https://doi.org/10.1002/ggge.20271>, 2013.
- Decker, E. R. and Bucher, G. J.: Geothermal studies in the Ross Island-Dry Valley region, *Antarct. Geosci.*, 4, 887–894, 1982.
- Ekaykin, A. A., Lipenkov, V. Y., and Shibaev, Y. A.: Spatial distribution of the snow accumulation rate along the ice flow lines between Ridge B and Lake Vostok, *Ice and Snow*, 4, 122–128, <https://doi.org/10.15356/2076-6734-2012-4-122-128>, 2012.
- Engelhardt, H.: Ice temperature and high geothermal flux at Siple Dome, West Antarctica, from borehole measurements, *J. Glaciol.*, 50, 251–256, <https://doi.org/10.3189/172756504781830105>, 2004.
- Fischer, H., Severinghaus, J., Brook, E., Wolff, E., Albert, M., Alemany, O., Arthern, R., Bentley, C., Blankenship, D., Chappellaz, J., Creyts, T., Dahl-Jensen, D., Dinn, M., Frezzotti, M., Fujita, S., Gallee, H., Hindmarsh, R., Hudspeth, D., Jugie, G., Kawamura, K., Lipenkov, V., Miller, H., Mulvaney, R., Parrenin, F., Pattyn, F., Ritz, C., Schwander, J., Steinhage, D., van Om-

- men, T., and Wilhelms, F.: Where to find 1.5 million yr old ice for the IPICS “Oldest-Ice” ice core, *Clim. Past*, 9, 2489–2505, <https://doi.org/10.5194/cp-9-2489-2013>, 2013.
- Fisher, A. T., Mankoff, K. D., Tulaczyk, S. M., Tyler, S. W., Foley, N., and the WISSARD Science Team: High geothermal heat flux measured below the West Antarctic Ice Sheet, *Sci. Adv.*, 1, e1500093, <https://doi.org/10.1126/sciadv.1500093>, 2015.
- Fox Maule, C. F., Purucker, M. E., Olsen, N., and Mosegaard, K.: Heat flux anomalies in Antarctica revealed by satellite magnetic data, *Science*, 309, 464–467, <https://doi.org/10.1126/science.1106888>, 2005.
- Garfield, D. E. and Ueda, H. T.: Resurvey of the “Byrd” Station, Antarctica, drill hole, *J. Glaciol.*, 17, 29–34, <https://doi.org/10.3189/S0022143000030689>, 1976.
- Golledge, N. R., Kowalewski, D. E., Naish, T. R., Levy R. H., Fogwill, C. J., and Gasson, E. G. W.: The multi-millennial Antarctic commitment to future sea-level rise, *Nature*, 526, 421–425, <https://doi.org/10.1038/nature15706>, 2015.
- Goodge, J. W.: Crustal heat production and estimate of terrestrial heat flow in central East Antarctica, with implications for thermal input to the East Antarctic ice sheet, *The Cryosphere*, 12, 491–504, <https://doi.org/10.5194/tc-12-491-2018>, 2018.
- Gow, A. J.: Deep core studies of the accumulation and densification of snow at Byrd Station and Little America V, Antarctica, CRREL Res. Rep. 197, 45 pp. 1968.
- Gundestrup, N. S., Clausen, H. B., and Hansen, B. L.: The UCPH borehole logger, *Mem. Nat. Inst. Polar Res.*, 49, 224–233, 1994.
- Hansen, B. L., Kelty, J. R., and Gundestrup, N. S.: Resurvey of Byrd Station drill hole, Antarctica, *Cold Reg. Sci. Tech.*, 17, 1–6, [https://doi.org/10.1016/S0165-232X\(89\)80011-4](https://doi.org/10.1016/S0165-232X(89)80011-4), 1989.
- Hindmarsh, R. C. A.: On the numerical computation of temperature in an ice sheet, *J. Glaciol.*, 45, 568–574, <https://doi.org/10.1017/S0022143000001441>, 1999.
- Hindmarsh, R.: Ice-sheet and glacier modelling, in: *Past Glacial Environments*, Elsevier, Amsterdam, Netherlands, 605–661, 2018.
- Hondoh, T., Shoji, H., Watanabe, O., Salamatian, A. N., and Lipenkov, V. Y.: Depth-age and temperature prediction at Dome Fuji station, East Antarctica, *Ann. Glaciol.*, 35, 384–390, <https://doi.org/10.3189/172756402781817013>, 2002.
- Huybrechts, P., Rybak, O., Pattyn, F., Ruth, U., and Steinhage, D.: Ice thinning, upstream advection, and non-climatic biases for the upper 89 % of the EDML ice core from a nested model of the Antarctic ice sheet, *Clim. Past*, 3, 577–589, <https://doi.org/10.5194/cp-3-577-2007>, 2007.
- Johnsen, S., Dahl-Jensen, D., Dansgaard, W., and Gundestrup N.: Greenland palaeotemperatures derived from GRIP bore hole temperature and ice core isotope profiles, *Tellus*, 47B, 624–629, <https://doi.org/10.3402/tellusb.v47i5.16077>, 1995.
- Jordan, T. A., Martin, C., Ferraccioli, F., Matsuoka, K., Corr, H., Forsberg, R., Olesen, A., and Siegert M.: Anomalously high geothermal flux near the South Pole, *Sci. Rep.*, 8, 16785, <https://doi.org/10.1038/s41598-018-35182-0>, 2018.
- Jouzel, J., Petit, J. R., Souchez, R., Barkov, N. I., Lipenkov, V. Y., Raynaud, D., Stievenard, M., Vassiliev, N. I., Verbeke, V., and Vimeux, F.: More than 200 meters of lake ice above subglacial Lake Vostok, Antarctica, *Science*, 286, 2138–2141, <https://doi.org/10.1126/science.286.5447.2138>, 1999.
- Kawamura, K., Parrenin, F., Lisicki, L., Uemura, R., Vimeux, F., Severinghaus, J. P., Hutterli, M. A., Nakazawa, T., Aoki, S., Jouzel, J., Raymo, M. E., Matsumoto, K., Nakata, H., Motoyama, H., Fujita, S., Goto-Azuma, K., Fujii, Y., and Watanabe, O.: Northern Hemisphere forcing of climatic cycles in Antarctica over the past 360,000 years, *Nature*, 448, 912–916, <https://doi.org/10.1038/nature06015>, 2007.
- Koutnik, M. R., Fudge, T. J., Conway, H., Waddington E. D., Neumann T. A., Cuffey K. M., Buijzer C., and Taylor K. C.: Holocene accumulation and ice flow near the West Antarctic Ice Sheet Divide ice core site, *J. Geophys. Res.-Earth*, 121, 907–924, <https://doi.org/10.1002/2015JF003668>, 2016.
- Lefebvre, E., Augustin, L., and Maitre, M.: The EPICA borehole logger, *Mem. Nat. Inst. Polar Res.*, 56, 264–274, 2002.
- Huber, M., Lanseau, C., and Rémy, F.: Relations between basal condition, subglacial hydrological networks and geothermal flux in Antarctica, *Earth Planet. Sc. Lett.*, 241, 655–662, <https://doi.org/10.1016/j.epsl.2005.10.040>, 2006.
- Lukin, V. V. and Vasiliev, N. I.: Technological aspects of the final phase of drilling borehole 5G and unsealing Vostok Subglacial Lake, East Antarctica, *Ann. Glaciol.*, 55, 83–89, <https://doi.org/10.3189/2014AoG65A002>, 2014.
- Martín, C. and Gudmundsson, G. H.: Effects of nonlinear rheology, temperature and anisotropy on the relationship between age and depth at ice divides, *The Cryosphere*, 6, 1221–1229, <https://doi.org/10.5194/tc-6-1221-2012>, 2012.
- Martos, Y. M., Catalán, M., Jordan, T. A., Golynsky, A., Golynsky, D., Eagles, G., and Vaughan, D. G.: Heat flux distribution of Antarctica unveiled, *Geophys. Res. Lett.*, 44, 11417–11426, <https://doi.org/10.1002/2017GL075609>, 2017.
- Mony, L., Roberts, J. L., and Halpin, J. A.: Inferring geothermal heat flux from an ice-borehole temperature profile at Law Dome, East Antarctica, *J. Glaciol.*, 66, 509–519, <https://doi.org/10.1017/jog.2020.27>, 2020.
- Motoyama, H.: The second deep ice coring project at Dome Fuji, Antarctica, *Sci. Drilling*, 5, 41–43, <https://doi.org/10.2204/iodp.sd.5.05.2007>, 2007.
- Motoyama, H., Furukawa, T., and Nishio, F.: Study of ice flow observations in Shirase drainage basin and around Dome Fuji area, East Antarctica by differential GPS method, *Nankyoku Shiryo (Antarctic Record)*, 52, 216–231, 2008.
- Motoyama, H., Furusaki, A., Takahashi, A., Tanaka, Y., Miyahara, M., Takata, M., Sawagaki, T., Matoba, S., Sugiyama, S., Shinbori, K., and Mori, S.: Deep borehole logging at Dome Fuji Station, Antarctica, Abstracts of the Fourth Symposium on Polar Science, 12–15 November, 2013, National Institute of Polar Research, Tachikawa, Tokyo, Japan, 2013.
- Neftel, A., Oeschger, H., Staffelbach, T., and Stauffer, B.: CO₂ record in the Byrd ice core 50000–5000 years BP, *Nature*, 331, 609–611, <https://doi.org/10.1038/331609a0>, 1988.
- Nicholls, K. W. and Paren, J. G.: Extending the Antarctic meteorological record using ice-sheet temperature profiles, *J. Climate*, 6, 141–150, [https://doi.org/10.1175/1520-0442\(1993\)006<0141:ETAMRU>2.0.CO;2](https://doi.org/10.1175/1520-0442(1993)006<0141:ETAMRU>2.0.CO;2), 1993.
- Parrenin, F., Dreyfus, G., Durand, G., Fujita, S., Gagliardini, O., Gillet, F., Jouzel, J., Kawamura, K., Lhomme, N., Masson-Delmotte, V., Ritz, C., Schwander, J., Shoji, H., Uemura, R., Watanabe, O., and Yoshida, N.: 1-D-ice flow modelling at EPICA Dome C and Dome Fuji, East Antarctica, *Clim. Past*, 3, 243–259, <https://doi.org/10.5194/cp-3-243-2007>, 2007a.

- Parrenin, F., Barnola, J.-M., Beer, J., Blunier, T., Castellano, E., Chappellaz, J., Dreyfus, G., Fischer, H., Fujita, S., Jouzel, J., Kawamura, K., Lemieux-Dudon, B., Loulergue, L., Masson-Delmotte, V., Narcisi, B., Petit, J.-R., Raisbeck, G., Raynaud, D., Ruth, U., Schwander, J., Severi, M., Spahni, R., Steffensen, J. P., Svensson, A., Udisti, R., Waelbroeck, C., and Wolff, E.: The EDC3 chronology for the EPICA Dome C ice core, *Clim. Past.*, 3, 485–497, <https://doi.org/10.5194/cp-3-485-2007>, 2007b.
- Parrenin, F., Cavitte, M. G. P., Blankenship, D. D., Chappellaz, J., Fischer, H., Gagliardini, O., Masson-Delmotte, V., Passalacqua, O., Ritz, C., Roberts, J., Siegert, M. J., and Young, D. A.: Is there 1.5-million-year-old ice near Dome C, Antarctica?, *The Cryosphere*, 11, 2427–2437, <https://doi.org/10.5194/tc-11-2427-2017>, 2017.
- Passalacqua, O., Ritz, C., Parrenin, F., Urbini, S., and Frezzotti, M.: Geothermal flux and basal melt rate in the Dome C region inferred from radar reflectivity and heat modelling, *The Cryosphere*, 11, 2231–2246, <https://doi.org/10.5194/tc-11-2231-2017>, 2017.
- Pattyn, F.: Antarctic subglacial conditions inferred from a hybrid ice sheet/ice stream model, *Earth Planet. Sci. Lett.*, 295, 451–461, <https://doi.org/10.1016/j.epsl.2010.04.025>, 2010.
- Pittard, M. L., Galton-Fenzi, B. K., Roberts, J. L., and Watson, C. S.: Organization of ice flow by localized regions of elevated geothermal heat flux, *Geophys. Res. Lett.*, 43, 3342–3350, <https://doi.org/10.1002/2016GL068436>, 2016.
- Pollard, D., DeConto, R. M., and Nyblade, A. A.: Sensitivity of Cenozoic Antarctic ice sheet variations to geothermal heat flux, *Global Planet. Change*, 49, 63–74, <https://doi.org/10.1016/j.gloplacha.2005.05.003>, 2005.
- Price, P. B., Nagornov, O. V., Bay, R., Chirkin, D., He, Y., Miocinovic, P., Richards, A., Woschnagg, K., Koci, B., and Zagorodnov, V.: Temperature profile for glacial ice at the South Pole: Implications for life in a nearby subglacial lake, *P. Natl. Acad. Sci. USA*, 99, 7844–7847, <https://doi.org/10.1073/pnas.082238999>, 2002.
- Raymond, C. F.: Deformation in the vicinity of ice divides, *J. Glaciol.*, 29, 357–373, <https://doi.org/10.1017/S0022143000030288>, 1983.
- Reeves, C. R. and Rowe, J. E.: Genetic Algorithms: Principles and Perspectives. A Guide to GA Theory, Kluwer Academic Publishers, Boston, USA, 2002.
- Rignot, E., Mouginot, J., and Scheuchl, B.: Ice flow of the Antarctic Ice Sheet, *Science*, 333, 1427–1430, <https://doi.org/10.1126/science.1208336>, 2011.
- Risk, G. F. and Hochstein, R.: Heat flow at Arrival Heights, Ross Island, Antarctica, New Zeal. *J. Geol. Geop.*, 17, 629–644, <https://doi.org/10.1080/00288306.1973.10421586>, 1974.
- Ritz, C., Rommelaere, V., and Dumas, C.: Modeling the evolution of Antarctic ice sheet over the last 420,000 years: Implications for altitude changes in the Vostok region, *J. Geophys. Res.-Atmos.*, 106, 31943–31964, <https://doi.org/10.1029/2001JD900232>, 2001.
- Robin, G. de Q.: Ice movement and temperature distribution in glaciers and ice sheets, *J. Glaciol.*, 2, 523–532, <https://doi.org/10.3189/002214355793702028>, 1955.
- Salamatin, A. N., Vostretsov R. N., Petit J. R., Lipenkov, V. Y. and Barkov, N. I.: Geofisicheskiye i paleoklimaticheskie prilozheniya sostavnogo temperaturnogo profilya iz glubokoi skvazhiny na stantsii Vostok (Antarktida) [Geophysical and palaeoclimatic implications of the stacked temperature profile from the deep borehole at Vostok station, Antarctica], *Mater. Glyatsiol. Issled.*, 85, 233–240, 1998a (in Russian with English summary).
- Salamatin, A. N., Lipenkov, V. Y., Barkov, N. I., Jouzel, J., Petit, J. R., and Raynaud, D.: Ice core age dating and paleothermometer calibration based on isotope and temperature profiles from deep boreholes at Vostok Station (East Antarctica), *J. Geophys. Res.*, 103, 8963–8977, <https://doi.org/10.1029/97JD02253>, 1998b.
- Seroussi, H., Ivins, E. R., Wiens D. A., and Bondzio J.: Influence of a West Antarctic mantle plume on ice sheet basal conditions. *J. Geophys. Res.-Sol. Ea.*, 122, 7127–7155, <https://doi.org/10.1002/2017jb014423>, 2017.
- Shapiro, N. M. and Ritzwoller, M. H.: Inferring surface heat flux distributions guided by a global seismic model: particular application to Antarctica, *Earth Planet. Sc. Lett.*, 223, 213–224, <https://doi.org/10.1016/j.epsl.2004.04.011>, 2004.
- Shen, W., Wiens, D., Lloyd, A., and Nyblade, A.: A geothermal heat flux map of Antarctica empirically constrained by seismic structure, *Geophys. Res. Lett.*, 47, e2020GL086955, <https://doi.org/10.1029/2020GL086955>, 2020.
- Sigl, M., Fudge, T. J., Winstrup, M., Cole-Dai, J., Ferris, D., McConnell, J. R., Taylor, K. C., Welten, K. C., Woodruff, T. E., Adolphi, F., Bisiaux, M., Brook, E. J., Buzert, C., Caffee, M. W., Dunbar, N. W., Edwards, R., Geng, L., Iverson, N., Koffman, B., Layman, L., Maselli, O. J., McGwire, K., Muscheler, R., Nishiizumi, K., Pasteris, D. R., Rhodes, R. H., and Sowers, T. A.: The WAIS Divide deep ice core WD2014 chronology – Part 2: Annual-layer counting (0–31 ka BP), *Clim. Past*, 12, 769–786, <https://doi.org/10.5194/cp-12-769-2016>, 2016.
- Slawny, K. R., Johnson, J. A., Mortensen, N. B., Gibson, C. J., Goetz, J. J., Shturmakov, A. J., Lebar, D. A., and Wendricks, A. W.: Production drilling at WAIS Divide, *Ann. Glaciol.*, 55, 147–155, <https://doi.org/10.3189/2014AoG68A018>, 2014.
- Staffelbach, T., Stauffer, B., Sigg, A., and Oeschger, H.: CO₂ measurements from polar ice cores: more data from different sites, *Tellus*, 43B, 91–96, <https://doi.org/10.3402/tellusb.v43i2.15251>, 1991.
- Ueda, H.: Byrd Station drilling 1966–69, *Ann. Glaciol.*, 47, 24–27, <https://doi.org/10.3189/172756407786857631>, 2007.
- Ueda, H. T. and Garfield, D. E.: Deep core drilling at Byrd Station, Antarctica, in: Proceedings of International Symp. on Antarctic Glaciological Exploration (ISAGE), Hanover, New Hampshire, USA, September 3–7, 1968, edited by: Gow, A. J., Association of Scientific Hydrology, Cambridge, UK, 86, 53–62, 1970.
- Ueltzhöffer, K. J., Bendel, V., Freitag, J., Kipfstuhl, S., Wagenbach, D., Faria, S. H., and Garbe, C. S.: Distribution of air bubbles in the EDML and EDC (Antarctica) ice cores, using a new method of automatic image analysis, *J. Glaciol.*, 56, 339–348, <https://doi.org/10.3189/002214310791968511>, 2010.
- Van Wyk De Vries, M., Robert G., Bingham R. G., and Hein A. S.: A new volcanic province: An inventory of subglacial volcanoes in West Antarctica, in: Exploration of Subsurface Antarctica: Uncovering Past Changes and Modern Processes, edited by: Siegert, M. J., Jamieson, S. S. R., and White, D. A., Geol. Soc. London, Special Publications, 461, 231–248, <https://doi.org/10.1144/SP461.7>, 2017.
- Van Liefferinge, B. and Pattyn, F.: Using ice-flow models to evaluate potential sites of million year-old ice in Antarctica,

- Clim. Past, 9, 2335–2345, <https://doi.org/10.5194/cp-9-2335-2013>, 2013.
- Van Liefferinge, B., Pattyn, F., Cavitte, M. G. P., Karlsson, N. B., Young, D. A., Sutter, J., and Eisen, O.: Promising Oldest Ice sites in East Antarctica based on thermodynamical modelling, *The Cryosphere*, 12, 2773–2787, <https://doi.org/10.5194/tc-12-2773-2018>, 2018.
- Vasiliev, N. I., Talalay, P. G., and Vostok Deep Ice Core Drilling Parties: Twenty Years of Drilling the Deepest Hole in Ice, *Sci. Dril.*, 11, 41–45, <https://doi.org/10.2204/iodp.sd.11.05.2011>, 2011.
- Veres, D., Bazin, L., Landais, A., Toyé Mahamadou Kele, H., Lemieux-Dudon, B., Parrenin, F., Martinerie, P., Blayo, E., Blunier, T., Capron, E., Chappellaz, J., Rasmussen, S. O., Severi, M., Svensson, A., Vinther, B., and Wolff, E. W.: The Antarctic ice core chronology (AICC2012): an optimized multi-parameter and multi-site dating approach for the last 120 thousand years, *Clim. Past*, 9, 1733–1748, <https://doi.org/10.5194/cp-9-1733-2013>, 2013.
- Vittuari, L., Vincent, C., Frezzotti, M., Mancini, F., Gandolfi, S., Bitelli, G., and Capra, A.: Space geodesy as a tool for measuring ice surface velocity in the Dome C region and along the ITASE traverse, *Ann. Glaciol.*, 39, 402–408, <https://doi.org/10.3189/172756404781814627>, 2004.
- Wexler, H.: Growth and thermal structure of the deep ice in Byrd Land, Antarctica, *J. Glaciol.*, 3, 1075–1087, <https://doi.org/10.1017/S002214300017482>, 1961.
- WAIS Divide Project Members: Onset of deglacial warming in West Antarctica driven by local orbital forcing, *Nature*, 500, 440–444, <https://doi.org/10.1038/nature12376>, 2013.
- Wendt, J., Dietrich, R., Fritzsche, M., Wendt, A., Yuskevich, A., Kokhanov, A., Senatorov, A. Lukin, V., Shibuya K., and Doi, K.: Geodetic observations of ice flow velocities over the southern part of subglacial Lake Vostok, Antarctica, and their glaciological implications, *Geophys. J. Int.*, 166, 991–998, <https://doi.org/10.1111/j.1365-246X.2006.03061.x>, 2006.
- Wesche, C., Eisen, O., Oerter, H., Schulte, D., and Steinhage, D.: Surface topography and ice flow in the vicinity of the EDML deep-drilling site, Antarctica, *J. Glaciol.*, 53, 442–448, <https://doi.org/10.3189/002214307783258512>, 2007.
- Whillans, I. M.: The equation of continuity and its application to the ice sheet near "Byrd" Station, Antarctica, *J. Glaciol.*, 18, 359–371, <https://doi.org/10.3189/S0022143000021055>, 1977.
- Whillans, I. M.: Ice flow along the Byrd Station strain network, Antarctica, *J. Glaciol.*, 24, 15–28, <https://doi.org/10.3189/S0022143000014611>, 1979.
- Wilhelms, F., Miller, H., Gerasimoff, M. D., Drücker, C., Frenzel, A., Fritzsche, D., Grobe, H., Hansen, S. B., Hilmarsson, S.Æ., Hoffmann, G., Hörnby, K., Jaeschke, A., Jakobsdóttir, S. S., Juckschat, P., Karsten, A., Karsten, L., Kaufmann, P. R., Karlin, T., Kohlberg, E., Kleffel, G., Lambrecht, A., Lambrecht, A., Lawer, G., Schärmeli, I., Schmitt, J., Sheldon, S. G., Takata, M., Trenke, M., Twarloh, B., Valero-Delgado, F., and Wilhelms-Dick, D.: The EPICA Dronning Maud Land deep drilling operation, *Ann. Glaciol.*, 55, 355–366, <https://doi.org/10.3189/2014AoG68A189>, 2014.
- Yen, Y.-C.: Review of thermal properties of snow, ice and sea ice, *CRREL Rep.*, vol. 81, 27 pp., 1981.
- Zagorodnov, V., Nagornov, O., Scambos, T. A., Muto, A., Mosley-Thompson, E., Pettit, E. C., and Tyuflin, S.: Borehole temperatures reveal details of 20th century warming at Bruce Plateau, Antarctic Peninsula, *The Cryosphere*, 6, 675–686, <https://doi.org/10.5194/tc-6-675-2012>, 2012.
- Zotikov, I. A.: Izmerenie geotermicheskogo potoka tepla v Antarktide [Measurement of the geothermal heat flow in Antarcical], Sovetskaia antarkticheskaiia ekspeditsiiia, Informatsionnyi biuletен [Soviet Antarctic Expedition, Information Bulletin] 29, 30–32, 1961 (in Russian).