

## Palaeogeographic evolution of Zealandia: mid-Cretaceous to present

Dominic P. Strogen, Hannu Seebeck, Benjamin R. Hines, Kyle J. Bland & James S. Crampton

To cite this article: Dominic P. Strogen, Hannu Seebeck, Benjamin R. Hines, Kyle J. Bland & James S. Crampton (2022): Palaeogeographic evolution of Zealandia: mid-Cretaceous to present, New Zealand Journal of Geology and Geophysics, DOI: [10.1080/00288306.2022.2115520](https://doi.org/10.1080/00288306.2022.2115520)

To link to this article: <https://doi.org/10.1080/00288306.2022.2115520>



Published online: 12 Sep 2022.



Submit your article to this journal 



Article views: 107



View related articles 



CrossMark

View Crossmark data 



RESEARCH ARTICLE



## Palaeogeographic evolution of Zealandia: mid-Cretaceous to present

Dominic P. Strogen <sup>a</sup>, Hannu Seebeck <sup>a</sup>, Benjamin R. Hines <sup>b</sup>, Kyle J. Bland <sup>a</sup> and James S. Crampton <sup>b</sup>

<sup>a</sup>GNS Science, Lower Hutt, New Zealand; <sup>b</sup>School of Environment, Geography and Earth Sciences, Victoria University of Wellington, Wellington, New Zealand

### ABSTRACT

We present a suite of 15 palaeogeographic maps illustrating the geological evolution of the entirety of Zealandia, from mid-Cretaceous to present, highlighting major tectonic phases, from initial Gondwana rifting through to development of the Neogene plate boundary. They illustrate palaeobathymetric and palaeofacies interpretations along with supporting geological datasets and a synthesis of regional tectonics. The maps are underpinned by a geologically-constrained and structurally-based rigid retro-deformation block model. This model, tied to the global plate circuit, is relatively simple for the main regions of Northern and Southern Zealandia, but breaks central Zealandia into numerous fault-bounded blocks, reflecting complex Neogene deformation associated with the modern plate boundary. Production of maps using GPlates and GIS allows for simple alteration or refinement of the block model and reconstruction of any geological dataset at any time. Reconstructions are within a palaeomagnetic reference frame, allowing assessment of palaeo-latitude, critical for palaeo-climatic and palaeo-biogeographic studies.

### ARTICLE HISTORY

Received 8 September 2021  
Accepted 8 August 2022

### HANDLING EDITOR

Richard Jongens

### KEYWORDS

Palaeogeography; tectonics; Cretaceous; Paleogene; Neogene; palaeobathymetry; palinspastic reconstruction; rifting; subduction


## Introduction

Zealandia, Earth's most recently recognised continent (Mortimer and Campbell 2014; Mortimer et al. 2017), forms a roughly 4.9 million km<sup>2</sup> area of continental crust in the southwest Pacific (Figure 1). Although it includes the landmasses of New Zealand, New Caledonia, and numerous smaller islands, 94% of Zealandia is below sea level. The majority of Zealandia is encompassed by the extended continental shelf (ECS) of New Zealand (Figure 1), with parts of Northern Zealandia falling within the ECS of Australia and France (New Caledonia).

The present-day extent, physiography/bathymetry and tectonic setting of Zealandia have been recently summarised (Mortimer 2020; Mortimer et al. 2020b, 2020c). However, the palaeogeographic evolution of Zealandia as a whole has not previously been considered in detail. Zealandia's present-day architecture is known at a broad scale, and formed from a combination of subduction accretion, magmatism, rifting, and passive-margin sedimentation over the last 100 Myr. Palaeogeographic reconstructions are important for resolving the distribution of the elements that comprise Zealandia, such as its sedimentary basins, volcanic arcs, and fault systems. They provide a valuable means of synthesising large amounts of geological data and help to summarise the long and complex geological history of regions such as Zealandia. Palaeobathymetry, restored to its correct

palaeolatitude, also provides important input data for palaeo-oceanographic, palaeoclimate and biogeographic studies (e.g. Bunce et al. 2009; Trewick and Bland 2012). Palaeogeographic maps also provide a better understanding of our biological heritage, such as the location and size of landmasses and land bridges, and how extensive terrestrial and marine habitats were through time. They also provide an important framework for sediment provenance and dispersal studies (Bull et al. 2019; Sagar et al. 2022).

This paper presents a new set of palaeogeographic maps covering the entirety of Zealandia retro-deformed through time, and represents the culmination of many years of research and compilation. These maps mark a major advance from the tectonic summary maps of King (2000) and the derived schematic palaeogeographic maps (e.g. King et al. 1999; Sutherland et al. 2009; Mortimer and Campbell 2014). Whilst palaeogeographic maps have been produced for parts of Zealandia (see section 2.2), unified maps covering the whole region or covering the entire mid-Cretaceous to present-day time period have not. Nor have maps been produced in a retro-deformed framework, and placed in an accurate latitude/longitude reference frame. It is beyond the scope of this paper to provide a full detailed geological history for all of Zealandia. Instead, a listing of data sources and methods used to build the maps is provided, as well as brief descriptions of the geological evolution as



**Figure 1.** Map of Zealandia showing the distribution of seismic and well datasets upon which palaeogeographic and basin studies are based. This shows the high data density in proximal New Zealand basins, but the very sparse datasets in much of Northern Zealandia and the distal Campbell Plateau. Bathymetry courtesy of GEBCO. 2D seismic dataset from New Zealand Petroleum & Minerals (2015) and Sutherland et al. (2012).

illustrated by each of the palaeogeographic maps, and some discussion of any issues or points of interest.

### Regional geological setting

Zealandia and surrounding regions have experienced a dynamic geological history, reflecting the complex tectonics of the southwest Pacific region over the last 100 Myr. The continent is presently bisected by an active convergent boundary between the Australian and Pacific plates (Figure 1). The area north (Australian Plate) and south (Pacific Plate) of the present day plate boundary are hereafter referred to as Northern and Southern Zealandia, respectively, as defined by Mortimer et al. (2017). Beneath eastern North Island,

the plate boundary manifests as subduction of oceanic Pacific Plate beneath the continental Australian Plate, along the Hikurangi margin (Walcott 1978; Nicol et al. 2007). This margin transitions southwards from subduction of the Pacific Plate along the Tonga-Kermadec Trench (and associated back-arc extension in the Lau Basin-Havre Trough), to continental collision and strike-slip along the Alpine Fault in South Island, New Zealand. To the southwest of South Island, a reversed subduction polarity sees oceanic Australian Plate crust being subducted obliquely beneath Pacific Plate continental crust (Gurnis et al. 2019).

From the Paleozoic to Early Cretaceous, the region that would later become Zealandia formed part of eastern Gondwana, with long-lived subduction along

its palaeo-Pacific eastern margin, and associated terrane accretion and magmatism. Rocks associated with this time period within the New Zealand region form the Austral Supergroup of Mortimer et al. (2014), and are subdivided into a series of basement terranes (Figure 2). Overlying the ‘basement’ rocks of the Austral Superprovince are mid-Cretaceous to Recent sedimentary and igneous ‘cover’ rocks of the Zealandia megasequence (Mortimer et al. 2014), which are the focus of this study. As the early part of the cover sequence straddles the Early to Late Cretaceous boundary, we use the informal term mid-Cretaceous.


Deposition and emplacement of the Zealandia megasequence began immediately following the cessation of subduction along the palaeo-Pacific margin at c. 100 Ma (Crampton et al. 2019; Riefstahl et al. 2020a). Some models infer complex subduction continued for some time (e.g. Cooper and Palin 2018), although a model of early subduction cessation is favoured here (see discussion in Crampton et al. 2019). Widespread rifting occurred throughout much of Zealandia (Laird and Bradshaw 2004; Strogen et al. 2017), which continued until initial separation of Zealandia from the remainder of eastern Gondwana at c. 83 Ma (Gaina et al. 1998b). Following this, sea-floor spreading began in both the southwest Pacific and Tasman Sea, and Zealandia drifted away from both Antarctica and Australia. This was a mostly tectonically quiescent time, although some minor rifting did occur in central Zealandia (Strogen et al. 2017). Tasman Sea spreading ceased by c. 52 Ma (Gaina et al. 1998b), with Zealandia then fully separated from Australia. There are several competing tectonic models for the evolution of northeastern Zealandia and surrounds, especially with regards to Late Eocene ophiolite obduction in New Caledonia (Collot et al. 2020). Onset of Pacific-Australia plate-boundary formation may have affected Northern Zealandia from as early as 50 Ma (Sutherland et al. 2020), and deformation migrated south through Northern Zealandia during the Eocene (Bache et al. 2012; Sutherland et al. 2017), concomitant with complex opening of back-arc basins east of Northern Zealandia through the Eocene–Middle Miocene (Mortimer et al. 2007; Herzer et al. 2011). In the Middle Eocene, intra-plate rifting began in Southern Zealandia (Turnbull et al. 1993) related to Emerald Basin opening (King 2000), which continued through to roughly the end of the Oligocene. From the Early Miocene, deformation related to the modern plate boundary affected much of central Zealandia (Nicol et al. 2007) with establishment of subduction along the full length of the Hikurangi margin. Deformation continued and intensified through the Neogene (Sutherland 1995), with the onset of subduction on the Puysegur subduction zone in Southern Zealandia (Gurnis et al. 2019).

From the Middle Miocene onwards intra-arc rifting occurred (Giba et al. 2013), with opening of the Havre Trough in the Pliocene (Wysoczanski et al. 2019).

## Data sources and methodology

Regional-scale syntheses of many of Zealandia’s sedimentary basins were presented in publications resulting from the Cretaceous-Cenozoic petroleum basins programme (CCP) led by the New Zealand Geological Survey and GNS Science between the mid-1970s and late 1990s (Nathan et al. 1986; Field et al. 1989; Wood et al. 1989; Turnbull et al. 1993; Edbrooke et al. 1994; Isaac et al. 1994; King and Thrasher 1996; Field et al. 1997; Cook et al. 1999). In many cases, the study areas were largely onshore (Nathan et al. 1986; Field et al. 1989), and most of the far-offshore regions were not studied. A number of sedimentary basins, such as Raukumara, Pegasus and Reinga basins, were almost unknown at that time (e.g. Sutherland et al. 2009; Uruski 2010; Bland et al. 2015). Significant advances in mapping and understanding the geological history and architecture of Zealandia were subsequently made during the past two decades, primarily driven by New Zealand’s United Nations Convention for the Law of the Sea (UNCLOS) programme (Uruski 1997; Davy and Uruski 2002; Wood 2002; Wood and Davy 2002; Wright et al. 2002) and by Government-driven seismic data acquisition programmes to encourage petroleum exploration (e.g. Uruski et al. 2005; Stagpoole et al. 2008; Uruski et al. 2008; Stagpoole et al. 2009; Uruski and Bland 2011; Bland et al. 2012; Bland et al. 2014a; Bland et al. 2014b; Crutchley et al. 2016). Recent data compilations and syntheses for the major offshore sedimentary basins, covering the entirety of New Zealand’s extended continental shelf (ECS), have recently been produced by the Atlas of Petroleum Prospectiveity (APP) programme (e.g. Arnot et al. 2016). By far the biggest sources of new geological data have come from regional seismic data (2D and 3D), and deep wells acquired for petroleum exploration. We have not carried out new seismic interpretation as part of this study, rather we have synthesised the open-file observations and the interpretations of others. This phase of discovery extended beyond the main sedimentary basins, and has included numerous dredging campaigns, often in more remote parts of Zealandia (e.g. Herzer et al. 1997; Browne et al. 2016; Mortimer et al. 2018; Lawrence et al. 2020; Mortimer et al. 2020a; Mortimer et al. 2020d) as well as scientific drilling by DSDP/ODP/IODP (e.g. Sutherland et al. 2019).

We compile, evaluate and reconcile this wealth of existing, disparate datasets. A full listing of previous palaeogeographic and geological studies used in this study is given for each region or basin in the



**Figure 2.** The tectonic block model (Seebek et al. 2018) used to retro-deform Zealandia, focusing on central Zealandia, shown at the present day and at c. 40 Ma. Overlain in onshore/nearshore New Zealand are simplified basement terranes, after Mortimer (2004).

supplementary material (<http://doi.org/10.6084/m9.figshare.20500113>). Detailed references to these studies are given where relevant in the detailed description for each map, also in the supplementary material. We have also digitised ‘paper copy’ (non-digital) datasets.

Where no existing palaeogeographic maps were available for a region of Zealandia, we have constructed them using any available, often sparse, datasets. These palaeogeographic interpretations were then placed into a restored tectonic position and a unified palaeogeography for the whole of Zealandia was produced for each time step. The data and maps were all managed and produced in ArcGIS, with tectonic reconstructions using GPlates. These digital compilations then allowed interrogation of data within a common projection and geographic framework.

### Tectonic reconstruction model

Underpinning the palaeogeographic maps is a new rigid block model (58 blocks in total; Figure 2), developed using the open source GPlates plate tectonic reconstruction software (Boyden et al. 2011; Müller et al. 2018b), which we use to sequentially retro-deform Zealandia. An outline of the construction of the block model and methodologies involved in the tectonic reconstructions are given in Seebek et al. (2018) and Hines (2018). Details on the geological justifications for the choice of blocks and their deformation history is discussed in detail in Hines et al. (2022), including in their supplementary material. A forthcoming paper will further describe the GPlates model in more detail, and release the model itself. An advantage of GPlates is that geospatial data of

practically any type can be easily imported into and exported from the associated geographic information system (GIS), and reconstructed for any time step within the model. This leads to a second advantage, that the rigid block model is readily testable, adjustable, and updateable using GPlates. Further iterations of the model will take account of any feedback and new data, and will allow for more complex non-rigid modelling of both extension and compressional deformation (Müller et al. 2018a).

The majority of Northern and Southern Zealandia consists of only a handful of large blocks tied to the global plate circuit. However, the area in central Zealandia associated with the present-day plate boundary, particularly in the East Coast-Marlborough region, is broken into numerous smaller blocks to best approximate the complex deformation that has occurred there. Our model uses more, smaller blocks, particularly in Marlborough, than previous studies (Crampston et al. 2003; Lamb et al. 2016; Lamb and Mortimer 2020; Ghisetti 2021). Block boundaries were chosen to reflect real pre-Miocene structures and basement geology, and block rotations are constrained where possible by palaeomagnetic data (Hines 2018; Hines et al. 2022). Although rigid blocks cannot capture the true ductile deformation that occurred, with gaps between blocks opening up through time (Figure 2), they do allow the relative positions of features to be estimated.

Relative motions between Northern and Southern Zealandia are defined by total reconstruction poles for the Australia-East Antarctica, Australia-Lord Howe Rise, East Antarctica-West Antarctica and Pacific-West Antarctica plate pairs determined from seafloor and magnetic anomaly data along a plate motion chain that links Lord Howe Rise to Australia,

East Antarctica, West Antarctica, and hence the Pacific plate (Gaina et al. 1998b; Cande and Stock 2004; Whittaker et al. 2007; Croon et al. 2008; Granot et al. 2013; Whittaker et al. 2013; Wright et al. 2015; Choi et al. 2017). The block model was reconstructed at selected times (2.58, 4.24, 6.04, 10.95, 20.13, 26.55, 40.13 and 83.5 Ma; corresponding to seafloor magnetic anomalies 2Ay, 3y, 3Ay, 5o, 6o, 15o, 18o and 34y) which define well constrained total reconstruction poles for Australia-Pacific relative motion (e.g. Cande and Stock 2004). These ages also generally correspond to significant tectonic events within the New Zealand plate boundary zone (King 2000) where passive strain markers are progressively restored back through time. The GPlates model then interpolates between these time points. As only relatively minor intraplate deformation is documented within Zealandia between 40 and 83 Ma (King 2000) all the model blocks are fixed in their 40.13 Ma positions (Figure 2b) relative to the main Southern Zealandia (Campbell Plateau) block, and carried back to the onset of seafloor spreading and the break-up of Gondwana at c. 83 Ma (Gaina et al. 1998b). In order to restore the reconstructed plate geometries to their correct latitude/longitude, it is necessary to use an absolute reference frame that describes how the plates have moved relative to a fixed reference system (Torsvik et al. 2008; van Hinsbergen et al. 2015). We utilise the palaeomagnetically-derived true polar wander corrected reference frame of Torsvik et al. (2012), which links to the global plate circuit through Africa, which has moved <500 km over the last 100 Ma (Torsvik et al. 2008; Seton et al. 2012).

Between the cessation of palaeo-Pacific subduction c. 100 Ma (Davy 2014; Reyners et al. 2017b; Crampton et al. 2019; Rieftahl et al. 2020a) and the breakup of eastern Gondwana at c. 83 Ma, there are many uncertainties concerning the disposition of Zealandia. Some models suggest significant strike-slip motion between Northern and Southern Zealandia and East and West Antarctica (e.g. Lamb et al. 2016; Lamb and Mortimer 2020) or at least significant internal deformation of Zealandia (Mortimer 2014) and possible continued complex subduction (Cooper and Palin 2018). Other models also break Northern Zealandia up into a number of smaller blocks to allow better closure of the Tasman Sea (Gaina et al. 1998b). Due to these uncertainties, there are significant issues when producing reconstructions for this time. We have however produced maps for both 85 and 98 Ma, in order to display the distribution of early syn-rift strata (Strogen et al. 2017). For simplicity, we keep the same relative block positions as for 40–83 Ma, with the position of Northern Zealandia, and all the smaller blocks in central Zealandia, dictated by the well-constrained positions of Southern Zealandia and West Antarctica. See further comments in the discussion.

### ***Biostratigraphic, palaeoenvironmental and other map data***

Geological outcrop data for New Zealand, derived from Heron (2018), have been grouped according to age for each map. Onshore measured section data from various New Zealand CCP studies (summarised in King et al. 1999), have been compiled by Galbraith et al. (2015) and provide litho- and biostratigraphic data, which are especially useful for delineating non-to marginal-marine sections. All palaeontological data for the Zealandia megasequence containing both robust age information and foraminiferal assemblages useful for assessing palaeo water depth (Hayward 1986, 2004; Hayward et al. 2013) have been extracted from the New Zealand fossil record file (FRED database; Clowes et al. 2021). Where possible, both age and palaeoenvironmental data have been extracted from DSDP/ODP/IODP wells, which provide the only control points across large parts of Northern Zealandia. These include legacy DSDP leg 21 (sites 206–208; Burns et al. 1973), leg 90 (sites 588–593; Kennett et al. 1986), ODP leg 181 (sites 1120–1125; Carter et al. 1999) and the recent IODP legs 371 (sites U1506–1511; Sutherland et al. 2019) and 372B/375 (sites U1517–1520 & 1526; Wallace et al. 2019). Available updated compilations for DSDP/ODP/IODP wells were also used here (Ayress 1994; Kulhanek and Morgans 2013; Sutherland et al. 2020). These biostratigraphic and palaeoenvironmental data are displayed as point data. For each map, strata are classified as ‘present’ where there is high biostratigraphic confidence, ‘possibly present’ where there is low confidence, or ‘missing’ where there is high confidence that the time interval is absent at that locality.

Dated volcanic samples, both onshore and offshore, were extracted from PETLAB (Strong et al. 2016), and are also shown on the relevant maps. Active faults are taken from existing palaeogeographies, and other studies where relevant (e.g. Giba et al. 2013; Ghisetti et al. 2016; Ghisetti 2021). Additional faults may be present in areas where insufficient data are available to constrain their orientations, so the absence of faults on a map does not necessarily imply an absence of faulting. Faults are omitted from the regional Zealandia maps (Figures 4–18) for clarity, but are shown on the maps centred on New Zealand (see supplementary material). Also shown are folds and areas of uplift where known (e.g. Bache et al. 2013; Sutherland et al. 2020). Faults from the present day map are from the New Zealand active faults database (Langridge et al. 2016).

Palaeobathymetric contours, representing boundaries between the palaeo water depth zones of Hayward (1986), are drawn using existing palaeogeographic mapping or updated based on new

palaeoenvironmental data and to fit the reconstructed geometries. In areas outside of existing mapping, new contours have been drawn based on the compiled palaeoenvironmental data. Palaeofacies boundaries are drawn to mark significant lithological changes, although necessarily simplified at this scale. These boundaries were used to construct both palaeobathymetric and palaeofacies polygons, covering the entirety of Zealandia in a reconstructed geometry for each map time. In some areas, there are sufficient


data to subdivide the shelf into inner, middle and outer shelf zones, elsewhere these are undivided and identified only as shelfal. The palaeobathymetry has been simplified on Figures 4–17 for clarity; all shelfal zones are combined, as are uppermost/upper bathyal, as well as lower深深 lower bathyal. All identified palaeobathymetric zones are shown on the maps in the supplementary data.

Global tectonic features provided with GPlates include; isochrons, spreading ridges and areas of continental crust (Müller et al. 2016), fracture zones (Matthews et al. 2011) and Australian palaeogeography (Totterdell et al. 2001). In some cases, we have replaced these where more detailed or improved data are available. These include seafloor spreading isochrons between the Campbell Plateau and West Antarctica and in the Adare Trough-Emerald Basin (Croon et al. 2008; Wobbe et al. 2012; Granot et al. 2013) and in the South Fiji Basin (Herzer et al. 2011). The extent of Zealandia continental crust is from Mortimer et al. (2020c), and in the Ross Sea its extent is revised using the 1000 m isobath to include Iselin Bank. Sedimentary basins for Zealandia are from Mortimer et al. (2020c), for the Ross Sea from Brancolini et al. (1995), and for Australia from Raymond (2018). The present-day extent of the Hikurangi Plateau is from Mortimer et al. (2020c), after Reyners et al. (2017b), ultimately modified from that of Wood and Davy (1994). Where no palaeogeographical interpretation has been made, the maps are left blank, or coloured grey where continental crust is interpreted. The modern coastlines are also shown for all regions, for reference.

## Palaeogeographic evolution

The series of 15 palaeogeographic maps (see Figure 3 for their ages) presented herein (Figures 4–18) were chosen to illustrate phases of the c. 100 Myr evolution of the Zealandia megasequence. The maps are assigned an age, rounded to the nearest million years (Ma), based on the current New Zealand geological timescale of Raine et al. (2015). However, the majority of biostratigraphic ages upon which maps are constructed are accurate only to the NZ stage level, with stage durations of c. 1–10 Myr. Therefore, the maps are a synthesis of data from an only very approximate snapshot in time. The maps are at c. 10 Myr intervals in the Cretaceous–Eocene and c. 5 Myr intervals from the Middle Eocene to present. They also depict the depositional sequences bounded by regionally mapped seismic horizons (Figure 3) in New Zealand basins (Strogen & King 2014) and are the same ages as those chosen for regional APP studies (Arnot et al. 2016; Sahoo et al. 2017; Arnot et al. 2018), with the addition of new maps for 40 and 30 Ma.

Each of the palaeogeographic maps produced are shown (Figures 4–18) along with a brief description




**Figure 3.** Comparison of the international and New Zealand timescales (Raine et al. 2015), and regional New Zealand seismic reflectors (Strogen & King 2014). The ages of the 15 palaeogeographic maps produced in this study are indicated.

of the tectonics and geological evolution at this time and any major points of interest or issues. We have grouped the map descriptions into four major phases of Zealandia evolution. Significantly more detailed descriptions of each map and its data sources are given in the supplementary material. For the sake of clarity, not all data used can be displayed clearly for the entirety of Zealandia, at a scale of 1:17M. An additional set of maps covering onshore/nearshore New Zealand at a larger scale (1:6M) have been produced, displaying additional data, such as outcrop data, well/outcrop sample data, active faults and more detailed lithofacies. These maps, as well as full-


size regional maps, are provided in the supplementary material along with the relevant map descriptions. For the Zealandia-wide maps (1:17M), the only point data shown are DSDP/ODP/IODP wells, outcrop sections from outlying islands and New Caledonia and dated volcanic samples. Faults are omitted for clarity from these regional maps.

## ***Mid- to Late Cretaceous ‘Zealandia rifting’ (98–85 Ma maps)***

These maps (Figures 4–5) illustrate the ‘Zealandia rifting’ phase (Strogen et al. 2017), with extensional



**Figure 4.** Mid-Cretaceous regional palaeogeographic map of Zealandia (c. 98 Ma). Key applies to all maps. The overlap between Northern Zealandia and Australia in the current reconstruction is also shown, see text for discussion. Australian palaeogeography from Totterdell et al. (2001).



**Figure 5.** Late Cretaceous regional palaeogeographic map of Zealandia (c. 85 Ma). Australian palaeogeography from Totterdell et al. (2001). See Figure 4 for key.

basins forming across much of Zealandia. Further detailed descriptions of the tectonics and geological evolution during this interval, as well as sets of both regional and New Zealand-focused maps, are given in the supplementary material. It follows the end of subduction along the palaeo-Pacific margin of Gondwana (e.g. Crampton et al. 2019), which ran along the present-day eastern margin of Zealandia. This followed the collision and partial subduction of the Hikurangi Plateau (a fragment of a large igneous province) with the Chatham Rise (Davy et al. 2008; Bland et al. 2015; Rieftahl et al. 2020a). The present-day eastern margin of Zealandia, then more north facing, was a passive margin dominated by deep-marine deposition, although some minor tectonic events are recognised within this sequence (Crampton et al. 2019). There was widespread syn-rift deposition of initially generally non-marine facies (Figure 4) in many basins across Zealandia from the Campbell Plateau to New Caledonia (Strogen et al. 2017; Maurizot

et al. 2020a; Barrier et al. 2022; Sahoo et al. 2022). Through this time interval the later part of this syn-rift succession became more widespread and increasingly marine influenced in many basins (Figure 5). The age of the rift succession in many undrilled basins is poorly constrained, especially in western Northern Zealandia (Totterdell et al. 2014; Bland and Strogen 2018). Crustal extension accrued during the ‘Zealandia’ rift phase has not been removed from these maps, and there are a number of refinements that remain poorly constrained with respect to reconstructions of this time (see discussion section and the supplementary material).

#### Late Cretaceous–Early Eocene Tasman Sea opening (77–57 Ma maps)


These maps (Figures 6–8) mark a phase characterised by active seafloor spreading in both the Tasman Sea and between Southern Zealandia and West Antarctica

(Gaina et al. 1998b; Rieftahl et al. 2020b). Detailed discussion of the tectonics and geological evolution, as well as sets of both regional and New Zealand-focused maps, can be found in the supplementary material. Initial spreading in the Tasman Sea propagated northwards, to the west of the Dampier Ridge by the end of the Cretaceous (Figure 7), and finally further northwards until there was full separation of Northern Zealandia and Australia, through the Cato Trough into the Coral Sea, in the Paleocene (Figure 8). Spreading continued into the Early Eocene (c. 52 Ma, Gaina et al. 1998b; Gaina et al. 1999).


Most areas were dominated by post-rift thermal subsidence, with continued marine transgression across most of Zealandia through the Late Cretaceous to Paleocene (Figures 6–8). Non-marine facies remained only in proximal basins, and shelfal facies deepened to bathyal conditions in many areas (Strogen et al. 2017; Maurizot et al. 2020a; Sahoo et al. 2022). Many of the major highs, such as the Lord

Howe Rise-Challenger Plateau, West Norfolk and Norfolk ridges and Campbell Plateau, which separated the former rift basins, were slowly transgressed through this interval and only relatively small areas remained emergent by the end of the Paleocene, although much of central Zealandia remained above sea level (Figure 8). The present-day eastern margin of Zealandia (then north facing), remained in a deep-marine passive margin setting (Field et al. 1997; Crampton et al. 2019).

Extension had ended in most basins, but minor rifting in central Zealandia during the Haumurian–Teurian, was orientated roughly perpendicular to previous rifts and occurred in parts of Taranaki, West Coast, and Ohai (Western Southland) basins ('West Coast-Taranaki' rift phase of Strogen et al. 2017). It is also likely that rifting in basins in the northwest of Zealandia preceded the onset of seafloor spreading, although this is poorly constrained (Uruski 2010; Totterdell et al. 2014).



**Figure 6.** Late Cretaceous regional palaeogeographic map of Zealandia (c. 77 Ma). Australian palaeogeography from Totterdell et al. (2001). See Figure 4 for key.




**Figure 7.** Latest Cretaceous regional palaeogeographic map of Zealandia (c. 67 Ma). Australian palaeogeography from Totterdell et al. (2001). See Figure 4 for key.

### Early Eocene–Oligocene subduction inception in Northern Zealandia (48–24 Ma maps)

This interval marks a period of plate boundary formation, with subduction initiation focused on Northern Zealandia and seafloor spreading to the south of Zealandia (Figures 9–13). Comprehensive descriptions of the tectonics and geological evolution during this interval, as well as sets of both regional and New Zealand-focused maps, are given in the supplementary material. Spreading in the Tasman Sea–Coral Sea was complete by the Early Eocene (c. 52 Ma, Gaina et al. 1998b; Gaina et al. 1999). There is evidence for subduction initiation along the northeast margin of Zealandia from c. 48 Ma (Figure 9) from the oldest arc-related rocks in Tonga (Meffre et al. 2012). This subduction system propagated south along the eastern margin of Northern Zealandia (Sutherland et al. 2020), so that by the end of the Eocene it had extended south to the east of the Three Kings Ridge (Figure 11). Through the Oligocene (Figures 12 and 13), this

subduction system rolled back resulting in the complex opening of the South Fiji Basin and later also the Norfolk Basin from c. 25 Ma (Herzer et al. 2011). We do not attempt to restore the complex tectonics of the South Loyalty Basin, given the conflicting models (see Collot et al. 2020; Sutherland et al. 2020) that exist for its formation, geometry and extent (see discussion section and supplementary material). The closure of the South Loyalty Basin was complete c. 34 Ma and was marked by final ophiolite obduction in New Caledonia (Maurizot et al. 2020b). To the south of Zealandia, a new spreading centre propagated northwards through the Emerald Basin during the Mid to Late Eocene (Figures 10 and 11). Spreading continued during the Oligocene (Figures 12 and 13), although becoming more strike-slip in nature with small disconnected ridge segments (Sutherland 1995).

A period of uplift in the northern Lord Howe Rise and Kenn Plateau (Figure 9) at c. 50 Ma is recognised by Sutherland et al. (2020), and likely related to subduction onset. Facies changes in onshore New




**Figure 8.** Late Paleocene regional palaeogeographic map of Zealandia (c. 57 Ma). Australian palaeogeography from Totterdell et al. (2001). See Figure 4 for key.

Caledonia from c. 46 Ma (Dallanave et al. 2020) also indicate the onset of uplift and imminent tectonism that would intensify through the Mid to Late Eocene with increasingly coarse-grained sedimentation (Maurizot et al. 2020a; Bordenave et al. 2021). Following final ophiolite obduction at c. 34 Ma, New Caledonia would remain a relatively stable high (Sevin et al. 2020; Maurizot et al. 2020b).

There was tectonic quiescence throughout much of central Zealandia during most of the Eocene. Continued, gradual, passive-margin transgression and bathyal conditions occurred across much of Zealandia, with increasingly fine-grained and calcareous facies deposited in distal areas (Figures 9 and 10). There was significant Early Eocene progradation of deltaic to shallow-marine facies in both the Taranaki Basin (King and Thrasher 1996; Higgs et al. 2017) and the Great South Basin (Constable and Crookbain 2011), fed from non-marine topographic highs in central Zealandia.

Compressional deformation would reach the Reinga Basin by the Late Eocene (Figure 11), with intense uplift and folding (Bache et al. 2014; Orr et al. 2020) and uplift of the Norfolk Ridge, Lord Howe Rise and Challenger Plateau (Sutherland et al. 2020). This deformation would reach as far south as northern Taranaki Basin (Figure 11) by the end of the Eocene (Strogen et al. 2014; Strogen et al. 2019). The boundary between compressional tectonics in the north and extension in the south lay within Taranaki Basin during the Late Eocene and Oligocene, with compressional tectonics migrating into southern Taranaki Basin in the Early Oligocene in response to activity along the length of the Taranaki Fault System (Stagpoole and Nicol 2008; Strogen et al. 2014).

A new phase of rifting related to Emerald Basin opening is recognised through much of Western Southland and West Coast basins from the Bortonian (Figure 10). Facies were initially non-marine




**Figure 9.** Early Eocene regional palaeogeographic map of Zealandia (c. 48 Ma). Australian palaeogeography from Totterdell et al. (2001). See Figure 4 for key.

in the Middle Eocene, but deposition would spread and become increasingly marine through the Late Eocene and Oligocene (Nathan et al. 1986; Turnbull et al. 1993). Although poorly constrained, extension in this rift likely reduced during the Oligocene due to sea-floor spreading in the Emerald Basin becoming more strike-slip dominated (Sutherland 1995).

Although active tectonics dominated the plate boundary zone, away from this region, passive subsidence still dominated across much of Zealandia (Field et al. 1997; Morgans 2016). By the Late Oligocene (Figure 13) there was maximum transgression of Zealandia (Landis et al. 2008) with only relatively small land areas preserved, and although there was widespread deposition of carbonate facies, there was still significant clastic input in areas close to the plate boundary zone (e.g. Nathan et al. 1986; Strogen et al. 2014).

#### **Earliest Miocene to present-day plate boundary in central Zealandia (21–0 Ma maps)**

This interval marks the propagation of the present-day plate boundary in the New Zealand region (Figures 14–18). Detailed descriptions of the tectonics and geological evolution during this interval, as well as sets of both regional and New Zealand-focused maps, are located in the supplementary material. In the earliest Miocene, the Hikurangi subduction zone propagated along the entire length of the East Coast Basin, bringing a major change to the tectonics in central Zealandia (Figure 14). The Northland and East Coast allochthons were emplaced in this interval (c. 23–19 Ma). The Northland Allochthon was likely formed due to transpression on the Veining-Meinesz Fracture Zone as the Norfolk Basin opened (Herzer and Mascle 1996). Complex seafloor spreading continued in the South Fiji and Norfolk basins until c. 16 Ma (Herzer et al. 2011),




**Figure 10.** Middle Eocene regional palaeogeographic map of Zealandia (c. 40 Ma). Australian palaeogeography from Totterdell et al. (2001). See Figure 4 for key.

when rollback of the Colville-Lau arc paused. The East Coast Allochthon, and numerous smaller olistostromes through the East Coast were likely related to the propagation of the subduction margin throughout the region, apparently synchronously (Chanier and Ferrière 1991; Rait et al. 1991; Delteil et al. 2006; Bland et al. 2022). There was also a change in depositional style throughout the East Coast Basin with an abrupt increase in clastic sedimentation, and widespread deposition of thick flysch deposits (Field et al. 1997) that indicate the onset of active tectonism all along this part of the margin. There was also an intensification of deformation on the Taranaki Fault System in the Early Miocene (King and Thrasher 1996), and thick flysch packages were deposited within a piggyback basin in the King Country region (Nelson and Hume 1977; Kamp et al. 2004) and the pulldown of the Waitematā Basin (Isaac et al. 1994). Punctuated increases in sedimentation rate, grain size and progradation of shelfal facies (Figures

14–16) during the Miocene in the West Coast through Taranaki basins can be linked to the increasing plate boundary convergence (Bull et al. 2019) as can intensification in compressional deformation (Ghisetti et al. 2016).

Arc volcanism migrated through northern New Zealand (Figures 14–16), initially in Northland (Early Miocene), moving south-eastwards through the Coromandel to Taranaki Basin (Middle to Late Miocene) and to the Tauranga-Kaimai area in the Pliocene (Hayward et al. 2001; Seebeck et al. 2013). Back-arc extension began in northern Taranaki Basin from the Middle Miocene, and gradually propagated into central Taranaki Basin during the Late Miocene to Pliocene (Giba et al. 2010).

Whilst significant sea-floor spreading had finished in the Emerald Basin (Sutherland 1995), minor highly oblique spreading likely continued as late as c. 15 Ma (Shuck et al. 2021). The plate boundary through central Zealandia, linking the Hikurangi




**Figure 11.** Late Eocene regional palaeogeographic map of Zealandia (c. 35 Ma). Australian palaeogeography from Totterdell et al. (2001). See Figure 4 for key.

subduction margin and the Emerald Basin, was initially a zone of distributed deformation (Figure 14). A through-going Alpine Fault (Figure 15) had likely formed by the Middle Miocene (Cox and Sutherland 2007; Reyners 2018). Highly oblique subduction on the Puysegur subduction margin had likely begun during the Middle Miocene (Sutherland et al. 2006) although the precise geometry and the timing of onset of this subduction and associated deformation is somewhat ambiguous (Shuck et al. 2021). There was an increase in coarse sedimentation in many basins (Nathan et al. 1986; Norris and Turnbull 1993), including for example, an intensification of convergence and rapid uplift in the Marlborough Fault System (Little and Roberts 1997; Little and Jones 1998).

By the Pliocene (Figure 17), spreading or crustal extension had begun in the Havre Trough (Caratori Tontini et al. 2019). This linked into spreading in

the Lau Basin to the north (Taylor et al. 1996), and resulted in separation of the Colville-Lau and Tonga-Kermadec arcs. By c. 2 Ma volcanism had migrated into the present-day Taupō Volcanic Zone with extension linked to the Havre Trough (Figure 18). There has been significant strike-slip motion through the North Island Fault System since c. 2 Ma, with mountain building, and intense deformation in the forearc (Beanland et al. 1998; Nicol et al. 2007; Bland et al. 2019).

The 0 Ma map (Figure 18) shows the present-day bathymetry and tectonics in the same style as the previous reconstructions. It is included largely to illustrate that the modern bathymetry is considerably more complex than we can hope to reconstruct in any of the previous palaeogeographic maps, and to emphasise that despite our efforts using the best data available, they still represent a gross simplification of the likely palaeobathymetry.




**Figure 12.** Early Oligocene regional palaeogeographic map of Zealandia (c. 30 Ma). Australian palaeogeography from Totterdell et al. (2001). See Figure 4 for key.

## Discussion

We have produced a set of palinspastically restored palaeogeographic maps for the entirety of Zealandia, from the Cretaceous break-up of eastern Gondwana, through to the present day. Importantly, the tectonic block model (see Hines et al. 2022 for full discussion) that underpins these palinspastic maps is tested and constrained by geological observations, and can be easily updated as required if new data suggest alterations are needed. Although previous studies have provided reconstructed palaeogeographies (e.g. King et al. 1999) and tectonic features (King 2000; Sutherland et al. 2001), these generally cover only the part of Zealandia centred on New Zealand or are focused only on limited time intervals. The majority of these maps were essentially schematic, not located in their restored latitude/longitude as in this study, nor allowing GIS geological datasets to be rigorously restored using a block model tied into the global plate circuit. The study of Cao et al. (2017) did produce palinspastic

maps with limited palaeogeographic data, but these were at a global scale, not focused on Zealandia, and contain much less detail than those in our study. Although previous models have addressed the relative positions of the large blocks of Northern and Southern Zealandia well, they failed to capture the complexity of deformation related to the Neogene plate boundary through central Zealandia, which this and a companion study by Hines et al. (2022) have attempted to address. Further iterations capturing more complexity are likely to be produced.

It is possible to analyse the maps produced in this study (Figures 4–18) to look at changes in the area of land and shelfal environments through time (Figure 19), to provide useful insights for palaeo-biogeographers. This shows a rapid decrease in both land and shelf areas from the Late Cretaceous through Paleocene, slowing somewhat in the Eocene–Oligocene, likely due to diminishing post-rift thermal subsidence, but also due to uplift in Northern Zealandia related to subduction initiation processes from c. 50 Ma onwards



**Figure 13.** Late Oligocene regional palaeogeographic map of Zealandia (c. 24 Ma). Australian palaeogeography from Totterdell et al. (2001). See Figure 4 for key.


(Sutherland et al. 2020). Land areas reach a minimum in the Late Oligocene, at the time of supposed total Zealandia drowning (Landis et al. 2008) but, arguably, still with a significant land area present ( $\sim 150,000 \text{ km}^2$ ). Since the Neogene plate boundary propagated into New Zealand, the land area has approximately doubled, to around  $300,000 \text{ km}^2$ . These maps also illustrate the rapid northward migration of Zealandia since the mid-Cretaceous (Figures 4–18); from  $\sim 80^\circ\text{S}$  to  $\sim 55^\circ\text{S}$  for southernmost Zealandia, and from  $\sim 50^\circ$  to  $\sim 20^\circ\text{S}$  for northernmost Zealandia.

A further advance to these maps would be to introduce estimates of palaeotopography, although this is inherently difficult to estimate, given the lack of preservation of non-marine compared to marine facies. However, it may be possible, based on pollen or other flora from sediments in surrounding areas. Attempts have been made to estimate palaeotopography at least for the last 4 Myr in central Zealandia (Nicol 2011; Trewick and Bland 2012), and similar palaeotopography estimates for older strata would

help to constrain ecosystem and palaeoclimate models.

#### Pre-breakup reconstructions

As previously stated, we have taken a simple approach to Cretaceous reconstructions to produce pre-breakup palaeogeographic maps, by maintaining the blocks in their 40 Ma relative positions and using the position of Southern Zealandia within the global plate circuit to drive block positions. Using this model yields a relatively straight palaeo-Pacific subduction margin from Raukumara Basin through to the Chatham Rise, as well as coherent palaeobathymetry and facies belts from Raukumara to Marlborough (Hines 2018; Hines et al. 2022). Neither of these lines of evidence require the major movements between Northern and Southern Zealandia prior to 83 Ma, as suggested by some workers (Lamb et al. 2016; Lamb and Mortimer 2020). Control on the palaeo-Pacific margin of




**Figure 14.** Early Miocene regional palaeogeographic map of Zealandia (c. 19 Ma). Australian palaeogeography from Totterdell et al. (2001). See Figure 4 for key.

Northern Zealandia from the Three Kings Ridge northwards is very poor.

This simplified model results in a pre-breakup overlap between Northern Zealandia and Australia of roughly 250–350 km, as well as an area of underlap in the southern Tasman Sea (Figure 4), which are obviously in error. One way to reduce this would be the removal of the extension from the basins in Northern Zealandia (present-day width of ~1000–1250 km). Removing this overlap would require extension of ~30–40% across the continent as a whole, which would have been concentrated within the major basins. Retrodeforming this extension may increase gaps present in the southern Tasman Sea although our knowledge of the Tasman Sea facing basins of the middle Lord Howe Rise (LHR) (Totterdell et al. 2014) and Challenger Plateau (CP) (Uruski 2010; Bland and Strogen 2018) is very poor. The effects of removing extension in Southern Zealandia are also

unknown, but should result in a better fit of the Chatham Rise against West Antarctica (Figure 4).

Another method to remove overlap is to break Northern Zealandia into multiple blocks. Gaina et al. (1998b) broke the LHR-CP into several blocks, most notably requiring significant sinistral strike-slip between their middle and northern LHR blocks to close the southern Tasman Sea without overlap to the north, and also closure of the Bellona Trough between the middle LHR and CP. It should be noted that Gaina et al. (1998b) only dealt with the fit of the LHR-CP against Australia and not the remainder of Northern Zealandia (Norfolk Ridge) or Southern Zealandia. Our model agrees with Gaina et al. (1998b) from c. 70 Ma onwards, where they have a contiguous LHR-CP in its present configuration. It is beyond the scope of the present study to address this further, but clearly future iterations of any block models for the period prior to the full separation of



**Figure 15.** Middle Miocene regional palaeogeographic map of Zealandia (c. 14 Ma). Australian palaeogeography from Totterdell et al. (2001). See Figure 4 for key.


Zealandia from eastern Gondwana (at c. 62 Ma) will need to address these issues, whether by removing extension, breaking Northern Zealandia into more blocks—assuming there is geological evidence for suitable structures, or more likely some combination of both approaches.

### Gondwana breakup

The breakup of Southern Zealandia against West Antarctica appears very different in style than that of Northern Zealandia against Australia. Breakup appears to be much sharper in the south, with a very narrow continental slope, possibly reflecting limited upper-crustal extension in the Campbell Plateau. In comparison, in Northern Zealandia, there is much more distributed crustal extension, particularly in the outer basins on the southwestern side of the LHR. Furthermore, many of the rift basins in Southern Zealandia and the eastern Ross Sea are at a high angle to the line of eventual breakup, unlike in the region

adjacent to Tasman Sea opening, where they are generally parallel. An exception is the Bellona Trough (Wood 1991, 1993), which trends at a high angle to the other Tasman Sea basins, and is poorly understood in both its age and mode of origin (Bland and Strogen 2018). The direction of marine flooding into the southern Tasman Sea prior to breakup at c. 83 Ma is unclear. We have suggested a marine connection along the line of subsequent breakup, but this is somewhat tortuous. Other possibilities include through the Bellona Trough and into Aotea Basin, or perhaps to the west of Tasmania, through the Sorrell Basin.

The breakup of Zealandia from eastern Gondwana is often thought of as occurring at c. 83 Ma, and associated with the end of the ‘Zealandia’ rift phase (Strogen et al. 2017). However, final separation of northern Zealandia with Australia did not occur until c. 62 Ma (Gaina et al. 1998a; 1999), almost 20 Myr later, and almost 40 Myr since rifting first began (Strogen et al. 2017). Also, whilst the eventual separation in the southern Tasman Sea and Southern



**Figure 16.** Late Miocene regional palaeogeographic map of Zealandia (c. 10 Ma). Australian palaeogeography from Totterdell et al. (2001). See Figure 4 for key.


Ocean is wide (>1000 km), the final separation of the Kenn Plateau and Australia through the Cato Trough is extremely narrow (~35 km; Mortimer et al. 2020c). The pre-breakup northern termination of Zealandia (and its sedimentary basins) and its relationship with microplates in the region now forming the d'Entrecasteaux to Coral Sea basins is unclear.

The age of extension in basins in northwestern Zealandia is poorly defined, but inferred to be generally c. 100–83 Ma, the same 'Zealandia' rift phase (Strogen et al. 2017) as recognised elsewhere in Zealandia. This appears to be the case in the eastern basins in Northern Zealandia, such as the New Caledonia Basin (Maurizot et al. 2020a). However, as discussed above, breakup in the northern Tasman Sea continued later than elsewhere in Zealandia, and it is possible that extension in the western basins of Northern Zealandia continued later than elsewhere. As yet there are few constraints on this as only seismic data are available for these rift basins (Totterdell et al. 2014).

### Cenozoic plate boundary initiation

Post-breakup, much of the Campbell Plateau deepened rapidly (Sutherland et al. 2010) and was quickly dominated by starved bathyal carbonate sedimentation. Whilst the highs of Northern Zealandia (CP-LHR and Norfolk Ridge) also deepened following the end of rifting, bathymetry in the north is overprinted by the effects of subduction initiation processes from the Early Eocene onwards (Sutherland et al. 2020; Skinner and Sutherland 2022). There was little change in the palaeobathymetry of much of distal Zealandia through the Neogene (Hayward et al. 2004).

An area in central Zealandia close to the present-day plate boundary appears to have supplied coarse sediment to basins to the NW (Taranaki) and SE (Great South) through the Late Cretaceous to Middle Eocene. This supply is cut off immediately prior to Emerald Basin rifting beginning at c. 40 Ma, possibly




**Figure 17.** Pliocene regional palaeogeographic map of Zealandia (c. 4 Ma). Australian palaeogeography from Totterdell et al. (2001). See Figure 4 for key.

reflecting topographic changes affecting sediment transport paths. This results in the reduction of coarse clastic sediments into the basins, and the end of long-lived deltaic sequences in both basins—the Farewell/Kaimiro/Mangahewa system in Taranaki Basin (King and Thrasher 1996) and the Tara-Toroa complex in Great South Basin (Sahoo et al. 2022). From the Middle Eocene onwards, the southern part of the West Coast formed part of the western margin facing into the Emerald Basin, the conjugate margin to the western edge of the Campbell Plateau. This explains the distal deep bathyal nature of deposition in southern Westland, which is quite different to further north on the West Coast (Nathan et al. 1986).

There are uncertainties regarding Eocene reconstructions prior to and during Emerald Basin opening. In the current block model, it is difficult to place Fiordland further south, and this places Western Southland to the east of much of the West Coast Basin, with Fiordland tight against Murchison Basin.

This has implications for provenance studies in the region (Sagar et al. 2022). Previous models (e.g. King et al. 1999) suggest Western Southland was south of, and contiguous with, the West Coast Basin. Whilst the present block model restores the oroclinal bend in Marlborough, as yet, it does not address inferred oroclinal bending in Fiordland (Lamb and Mortimer 2020).

We favour the model for South Fiji Basin opening of Herzer et al. (2011), which infers younger oceanic crust than many previous models (e.g. Sdrolias et al. 2003; Schellart et al. 2006). Whilst the geometry of spreading centres may be complex, overall it describes a simple rollback of the Tonga-Kermadec subduction system which fits well with our palaeogeographic model. Some details of how the subduction system propagated through this region are still unclear, such as the nature of volcanism on the Norfolk and Three Kings ridges (Mortimer et al. 2020d) and the precise timing of subduction propagation.



**Figure 18.** Present day map of Zealandia in the same style as the palaeogeographic maps. Bathymetry courtesy of GEBCO. Australian environments from Totterdell et al. (2001). See Figure 4 for key.


The Northland and East Coast allochthons appear to be similar in age (c. 23–19 Ma). The two allochthons did not form a contiguous single unit and were likely formed by different processes. The Northland Allochthon appears to be related to extension in the South Fiji and Norfolk basins and transpressive deformation along the VMFZ (Herzer and Mascle 1996). The East Coast Allochthon is probably related to transient uplift associated with propagation of the subduction zone south into the area stretching from Raukumara Basin to Marlborough (Chanier and Ferrière 1991; Rait et al. 1991), also reflected in smaller scale olistostromes found further south along this margin (e.g. Delteil et al. 2006; Bland et al. 2022).

It is when rollback in the South Fiji Basin is complete c. 15 Ma (Herzer et al. 2011) that convergence and rotation through central Zealandia intensifies (e.g. Bull et al. 2019), with the formation of a through-going Alpine Fault, and renewed subduction of the Hikurangi Plateau (Reyners et al. 2017a). Neogene

subduction of the Hikurangi Plateau under eastern central Zealandia appears to have had a major impact on deformation, uplift and erosion and is seen in significant Middle–Late Miocene increases in sedimentation rates in surrounding basins, particularly the East Coast (Field et al. 1997; Bland et al. 2019) and Taranaki basins (King and Thrasher 1996).

### Future work

Further work is needed to reconcile the models of New Caledonia ophiolite obduction and South Loyalty Basin tectonics (for discussion see Collot et al. 2020). As yet, we have not attempted to reconstruct the area to the east of New Caledonia prior to Late Eocene obduction. Future iterations of the block model should aim to more accurately capture the tectonic complexity in this northeastern part of Zealandia, as well as addressing the pre-breakup reconstruction issues already discussed. Further



**Figure 19.** Graph showing the variation in the area of land and shelf (0–200 m) in Zealandia since 98 Ma (note log scale). There was an initially rapid drop in the area of both through the Late Cretaceous and Palaeocene, and slower Eocene–Oligocene, drowning through to a minimum at c. 24 Ma, although there was still significant land area ( $\sim 148,000 \text{ km}^2$ ). Since the Neogene propagation of the plate boundary into New Zealand, the land area has roughly doubled from this minimum. Note that Northern Zealandia did not fully separate from Australia until c. 62 Ma (Gaina et al. 1998b). These estimates are approximate with potentially large but difficult to quantify errors.

refinements to the block model in the plate boundary zone are also to be expected, as well as attempts to better capture the ductile deformation occurring there.

Further work is required to fully capture the complex bathymetry and facies in East Coast Neogene accretionary wedge sub-basins as the Hikurangi margin developed (Pettinga 1982; Bailleul et al. 2013), as these have only been mapped at a reconnaissance level in this study. In far-field areas, such as much of Northern Zealandia and the Campbell Plateau, lack of data probably make the construction of more maps untenable at present. However, in areas with better data coverage and higher biostratigraphic resolution, particularly onshore/nearshore New Zealand, the production of further palaeogeographic maps is feasible. This is illustrated by the set of 30 maps produced for the Taranaki Basin by Strogen (2011). It is

anticipated that further maps will be constructed to address intervals of interest, particularly the late Neogene of interest in palaeo-climatic studies.

## Conclusions

A suite of palaeogeographic maps have been produced for Zealandia, using a new tectonic reconstruction model. These maps mark a major advance in the understanding of Zealandia's palaeogeographic evolution, with the main results summarised below.

- (1) The 15 maps (98–0 Ma) illustrate major phases in the tectonic evolution of Zealandia, from initial mid-Cretaceous Gondwana rifting through to the development of the modern Neogene–Quaternary plate boundary. They cover the whole of Zealandia, reconstructed in a rigorous tectonic

- framework. The maps provide palaeobathymetric and palaeofacies interpretations along with reconstructed supporting geological datasets and regional tectonics syntheses.
- (2) The underpinning retro-deformational block model is geologically constrained and structurally based (see companion paper Hines et al. 2022). It breaks central Zealandia into numerous fault-bounded blocks, reflecting complex Neogene–Quaternary plate boundary deformation. Production of maps using GPlates and GIS allows for simple alteration or refinement of the block model and reconstruction of any geological datasets through time.
  - (3) Reconstructions are tied to the global plate circuit within a palaeomagnetic reference frame, allowing assessment of palaeo-latitude, critical for palaeo-climatic and palaeo-biogeographic studies. We have also used these maps to estimate variations in the area of land and shelfal environments through time.
  - (4) Future iterative refinements to the tectonic model are expected. This will include modelling of extensional and contractional deformation, oroclinal bending, particularly to address issues with Cretaceous reconstructions. Palaeogeographic maps will be updated using any improved block model, and also integrating any new regional datasets, as available.

## Acknowledgements

This work has benefitted from discussions with many colleagues both at GNS Science and throughout the wider New Zealand geological community. Special thanks go to our colleagues Peter King and the late Mike Isaac for early inspiration and support for this work, to Nick Mortimer for highly fruitful tectonic discussions and providing volcanic ages from PETLAB, and to Chris Clowes for extracting useful data from the FRED database. Thanks for the use of the open source GPlates software, developed by the EarthByte Group at the University of Sydney. Many thanks to internal reviewers Nick Mortimer and Malcolm Arnot, to the journal reviewers Rupert Sutherland, Catherine Reid and Andy Nicol and to editor Richard Jongens, all of whom have helped to greatly improve this manuscript.

## Disclosure statement

No potential conflict of interest was reported by the author(s).

## Funding

This project was supported by the New Zealand Ministry of Business, Innovation and Employment (MBIE) through the Understanding Zealandia/Te Riu-a-Māui (UZTRM) programme (Strategic Science Investment Fund, contract C05X1702), and previously by the SSIF funded Sedimentary Basins and Petroleum research programmes. It incorporates

many of the results of the Atlas of Petroleum Prospectivity (APP) programme, funded through the MBIE Energy and Minerals targeted research fund. BRH was funded by a Sarah Beanland Memorial Scholarship, awarded by GNS Science.

## Data availability statement

The palaeogeographic reconstructions presented in this study have utilised open-file and published datasets, which are cited through the paper. All biostratigraphic data used are lodged in the New Zealand Fossil Record File (FRF; Clowes et al. 2021), accessible at <https://fred.org.nz/>. Data regarding the ages of onshore and offshore volcanic rock samples were extracted from PETLAB (Strong et al. 2016), accessible at <http://pet.gns.cri.nz/>. Onshore geological map data were derived from the QMAP Seamless 1:250k digital dataset, held by GNS and accessible via <https://www.gns.cri.nz/Home/Products/Maps>. Digital (GIS) palaeogeographic maps, illustrated seismic transects, and digital (GIS) seismic depth-grids relating to the ‘Atlas of Petroleum Prospectivity’ (APP) programme, are accessible for download from <https://data.gns.cri.nz/pbe/>, via digital Petroleum Exploration Data Packs distributed by New Zealand Petroleum & Minerals (MBIE; <https://www.nzpam.govt.nz/>), or by contacting the authors. A supplemental file providing more detailed descriptions of the palaeogeographic reconstructions presented in this paper and all referenced material used in their construction, is available in figshare via <https://doi.org/10.6084/m9.figshare.20500113>.

## ORCID

*Dominic P. Strogen*  <http://orcid.org/0000-0002-1744-1289>  
*Hannu Seebeck*  <http://orcid.org/0000-0002-1749-6626>  
*Benjamin R. Hines*  <http://orcid.org/0000-0001-6177-7137>  
*Kyle J. Bland*  <http://orcid.org/0000-0002-9038-0208>  
*James S. Crampton*  <http://orcid.org/0000-0003-3270-9981>

## References

- Arnot MJ, Bland KJ, Boyes AF, Bull S, Funnell RH, Griffin AG, Hill MG, Kroeger KF, Lukovic B, Scadden PG, et al. 2016. *Atlas of Petroleum Prospectivity, Northwest Province: ArcGIS geodatabase and technical report*. GNS Science data series, 23b. 5 ArcGIS projects, 1 ArcGIS geodatabase, and technical report.
- Arnot MJ, Lawrence MJF, Viskovic GPD, Zhu H, Bland KJ, Strogen DP, Schuurmans R. 2018. *Atlas of petroleum prospectivity, Southwest Province (West Coast, Fiordland, and western Southland basins)*. GNS Science data series, 23e. 1 ArcGIS geodatabase, 3 ArcGIS projects, 23 illustrated seismic transects, and technical report.
- Ayress MA. 1994. Cainozoic palaeoceanographic and subsidence history of the eastern margin of the Tasman Basin based on Ostracoda. In: Van der Lingen GJ, Swanson KM, Muir RJ, editor. *Evolution of the Tasman Sea Basin: Proceedings of the Tasman Sea Conference Christchurch, New Zealand, 27–30 November 1992*. Rotterdam: A.A. Balkema; p. 139–158.
- Bache F, Mortimer N, Sutherland R, Collot J, Rouillard P, Stagpoole V, Nicol A. 2014. Seismic stratigraphic record of transition from Mesozoic subduction to continental

- breakup in the Zealandia sector of eastern Gondwana. *Gondwana Research*. 26(3–4):1060–1078. doi:[10.1016/j.gr.2013.08.012](https://doi.org/10.1016/j.gr.2013.08.012).
- Bache F, Stagpoole VM, Sutherland R. 2013. Seismic stratigraphy of the Reinga Basin, NW New Zealand: tectonic and petroleum implications. In: Rosen NC, Weimer P, dos Anjos SMC, Henrickson S, Marques E, Mayall M, Fillon R, D'Agostino T, Saller A, Campion K and others editors. *New understanding of the petroleum systems of continental margins of the world: SEPM Foundation, Gulf Coast Section, 32nd annual research conference proceedings*. Tulsa, SEPM Society for Sedimentary Geology. p. 221–252. doi:[10.5724/gcs.12.32.0221](https://doi.org/10.5724/gcs.12.32.0221)
- Bache F, Sutherland R, Stagpoole V, Herzer R, Collot J, Rouillard P. 2012. Stratigraphy of the southern Norfolk Ridge and the Reinga Basin: A record of initiation of Tonga-Kermadec-Northland subduction in the southwest Pacific. *Earth and Planetary Science Letters*. 321–322:41–53. doi:[10.1016/j.epsl.2011.12.041](https://doi.org/10.1016/j.epsl.2011.12.041).
- Bailleul J, Chanier F, Ferrière J, Robin C, Nicol A, Mahieux G, Gorini C, Caron V. 2013. Neogene evolution of lower trench-slope basins and wedge development in the central Hikurangi subduction margin, New Zealand. *Tectonophysics*. 591:152–174. doi:[10.1016/j.tecto.2013.01.003](https://doi.org/10.1016/j.tecto.2013.01.003).
- Barrier A, Browne GH, Nicol A, Bassett K. 2022. Sedimentary architecture of a Late Cretaceous under-filled rift basin, Canterbury Basin, New Zealand. *Basin Research*. 34(1):342–365. doi:[10.1111/bre.12622](https://doi.org/10.1111/bre.12622).
- Beanland S, Melhuish A, Nicol A, Ravens J. 1998. Structure and deformational history of the inner forearc region, Hikurangi subduction margin, New Zealand. *New Zealand Journal of Geology and Geophysics*. 41(4):325–342. doi:[10.1080/00288306.1998.9514814](https://doi.org/10.1080/00288306.1998.9514814).
- Bland KJ, Field BD, Griffin AG, Heron DW, Isaac MJ, Juniper Z, Uruski CI. 2012. Play-based assessment of the Pegasus and East Coast Basins. *GNS Science data series*, 9b. 1 CD.
- Bland KJ, Hill MG, Strogen DP, Bache F, Barker DHN, Bull S, Davy BW, Sahoo TR. 2014a. New Zealand national seismic framework: Reinga-Northland, Taranaki, Canterbury-Great South, Bounty Trough, and Pegasus-East Coast-Raukumara basins. In: New Zealand Petroleum & Minerals, editor. 2013 *New Zealand petroleum exploration data pack*. PR4700. p. 55 p. + 51 Enclosures.
- Bland KJ, Lamarche G, Anderson O, Barnes PM, Black J, Bowden D, Chiswell S, Crutchley GJ, Gorman R, Mountjoy JJ, et al. 2014b. Pegasus Basin petroleum prospectivity report. *GNS Science consultancy report*, 2014/103. 139 p. & GIS project.
- Bland KJ, Morgans HEG, Strogen DP, Harvey H. 2022. Litho- and biostratigraphy of a Late Oligocene–Early Miocene succession in the Weber area, southern Hawke's Bay, and implications for early Hikurangi subduction-margin evolution. *New Zealand Journal of Geology and Geophysics*. doi:[10.1080/00288306.2022.2108069](https://doi.org/10.1080/00288306.2022.2108069).
- Bland KJ, Nicol A, Kamp PJJ, Nelson CS. 2019. Stratigraphic constraints on the late Miocene–Pleistocene evolution of the North Island Fault System and axial ranges in the central Hikurangi subduction margin, New Zealand. *New Zealand Journal of Geology and Geophysics*. 62(2):248–272. doi:[10.1080/00288306.2018.1545675](https://doi.org/10.1080/00288306.2018.1545675).
- Bland KJ, Strogen DP. 2018. *Atlas of Petroleum Prospectivity, Tasman frontier (Far Frontiers Province)*: ArcGIS geodatabase, illustrated seismic transects and technical report. *GNS Science data series*, 23f. 1 ArcGIS geodatabase, 1 ArcGIS project, 5 illustrated seismic transects, and technical report.
- Bland KJ, Uruski CI, Isaac MJ. 2015. Pegasus Basin, eastern New Zealand: A stratigraphic record of subsidence and subduction, ancient and modern. *New Zealand Journal of Geology and Geophysics*. 58(4):319–343. doi:[10.1080/00288306.2015.1076862](https://doi.org/10.1080/00288306.2015.1076862).
- Bordenave A, Etienne S, Collot J, Razin P, Patriat M, Grélaud C, Agnini C, Morgans H, Guillemaut F, Moreau A. 2021. Upper Cretaceous to Palaeogene successions of the Gouaro anticline: deepwater sedimentary records of the tectonic events that led to obduction in New Caledonia (SW Pacific). *Sedimentary Geology*. 415:105818. doi:[10.1016/j.sedgeo.2020.105818](https://doi.org/10.1016/j.sedgeo.2020.105818).
- Boyden J, Müller RD, Gurnis M, Torsvik TH, Clark JA, Turner M HI-L, Watson RJ, Cannon JS. 2011. Next-generation plate-tectonic reconstructions using GPlates. In: Keller G, Baru C, editor. *Geoinformatics: cyberinfrastructure for the solid earth sciences*. Cambridge: Cambridge University Press; p. 95–114.
- Brancolini G, Busetti M, Marchetti A, De Santis L, Zanolla C, Cooper AK, Cochrane GR, Zayatz I, Belyaev V, Knyazev M, et al. 1995. Descriptive text for the seismic stratigraphic atlas of the Ross Sea, Antarctica. In: Cooper AK, Barker PF, Brancolini G, editor. *Geology and seismic stratigraphy of the Antarctic margin*. *Geology and seismic stratigraphy of the Antarctic margin*. Vol. 68. Washington: AGU; p. A271–A286.
- Browne GH, Lawrence MJF, Mortimer N, Clowes CD, Morgans HEG, Hollis CJ, Beu AG, Black JA, Sutherland R, Bache F. 2016. Stratigraphy of Reinga and Aotea basins, NW New Zealand: constraints from dredge samples on regional correlations and reservoir character. *New Zealand Journal of Geology and Geophysics*. 59(3):396–415. doi:[10.1080/00288306.2016.1160940](https://doi.org/10.1080/00288306.2016.1160940).
- Bull S, Nicol A, Strogen DP, Kroeger KF, Seebeck HS. 2019. Tectonic controls on Miocene sedimentation in the Southern Taranaki Basin and implications for New Zealand plate boundary deformation. *Basin Research*. 31(2):253–273. doi:[10.1111/bre.12319](https://doi.org/10.1111/bre.12319).
- Bunce M, Worthy TH, Phillips MJ, Holdaway RN, Willerslev E, Haile J, Shapiro B, Scofield RP, Drummond A, Kamp PJJ, et al. 2009. The evolutionary history of the extinct ratite moa and New Zealand Neogene paleogeography. *Proceedings of the National Academy of Sciences*. 106(49):20646–20651. doi:[10.1073/pnas.0906660106](https://doi.org/10.1073/pnas.0906660106).
- Burns RE, Andrews JE, others 1973. Initial reports of the Deep Sea Drilling Project. Washington (DC): US Govt. Printing Office. doi:[10.2973/dsdp.proc.21.1973](https://doi.org/10.2973/dsdp.proc.21.1973)
- Cande S, Stock J. 2004. Pacific-Antarctic-Australian motion and the formation of the Macquarie Plate. *Geophysical Journal International*. 157:399–414.
- Cao W, Zahirovic S, Flament N, Williams S, Golonka J, Müller RD. 2017. Improving global paleogeography since the late Paleozoic using paleobiology. *Biogeosciences*. 14(23):5425–5439. doi:[10.5194/bg-14-5425-2017](https://doi.org/10.5194/bg-14-5425-2017).
- Carter RM, McCave IN, Richter C, Carter L, et al. 1999. Southwest Pacific gateways. *Proceedings of the Ocean Drilling Program, Initial Reports*, 181. College Station, TX, International Ocean Discovery Program. [http://www-odp.tamu.edu/publications/181\\_IR/181ir.htm](http://www-odp.tamu.edu/publications/181_IR/181ir.htm).
- Chanier F, Ferrière J. 1991. From a passive to an active margin; tectonic and sedimentary processes linked to the

- birth of an accretionary prism (Hikurangi Margin, New Zealand). *Bulletin de la Societe Geologique de France*. 162(4):649–660. doi:[10.2113/gssgbull.162.4.649](https://doi.org/10.2113/gssgbull.162.4.649).
- Choi H, Kim S-S, Dymant J, Granot R, Park S-H, Hong JK. 2017. The kinematic evolution of the Macquarie Plate: A case study for the fragmentation of oceanic lithosphere. *Earth and Planetary Science Letters*. 478:132–142. doi:[10.1016/j.epsl.2017.08.035](https://doi.org/10.1016/j.epsl.2017.08.035).
- Clowes CD, Crampton JS, Bland KJ, Collins KS, Prebble JG, Raine JI, Strogen DP, Terezow MG, Womack T. 2021. The New Zealand Fossil Record File: a unique database of biological history. *New Zealand Journal of Geology and Geophysics*. 64(1):62–71. doi:[10.1080/00288306.2020.1799827](https://doi.org/10.1080/00288306.2020.1799827).
- Collot J, Patriat M, Sutherland R, Williams S, Cluzel D, Seton M, Pelletier B, Roest WR, Etienne S, Bordenave A, et al. 2020. Geodynamics of the SW Pacific: a brief review and relations with New Caledonian geology. In: Maurizot P, Mortimer N, editor. *New Caledonia: geology, geodynamic evolution and mineral resources*. New Caledonia: Geology, Geodynamic Evolution and Mineral Resources, Geological Society of London; p. 13–26. doi:[10.1144/M51-2018-5](https://doi.org/10.1144/M51-2018-5)
- Constable RM, Crookbain R. 2011. A sequence stratigraphic study of the Great South Basin. *New Zealand Unpublished Openfile Petroleum Report PR4348*: 223 p.
- Cook RA, Sutherland R, Zhu H, others 1999. Cretaceous-Cenozoic geology and petroleum systems of the Great South Basin, New Zealand. Institute of Geological & Nuclear Sciences Monograph, 20. Lower Hutt, New Zealand. 188 p, 2 enclosures.
- Cooper AF, Palin JM. 2018. Two-sided accretion and poly-phase metamorphism in the Haast Schist belt, New Zealand: constraints from detrital zircon geochronology. *GSA Bulletin*. doi:[10.1130/B31826.1](https://doi.org/10.1130/B31826.1).
- Cox SC, Sutherland R. 2007. Regional geological framework of South Island, New Zealand, and its significance for understanding the active plate boundary. In: Okaya D, Stern TA, Davey FJ, editor. *A continental plate boundary: tectonics at South Island, New Zealand*. Vol. 175. Washington, DC: American Geophysical Union; p. 19–46. doi:[10.1029/175GM03](https://doi.org/10.1029/175GM03)
- Crampton J, Laird M, Nicol A, Townsend D, Van Dissen R. 2003. Palinspastic reconstructions of southeastern Marlborough, New Zealand, for mid-Cretaceous-Eocene times. *New Zealand Journal of Geology and Geophysics*. 46(2):153–175. doi:[10.1080/00288306.2003.9515002](https://doi.org/10.1080/00288306.2003.9515002).
- Crampton JS, Mortimer N, Bland KJ, Strogen DP, Sagar M, Hines BR, King PR, Seebek H. 2019. Cretaceous termination of subduction at the Zealandia margin of Gondwana: the view from the paleo-trench. *Gondwana Research*. 70:222–242. doi:[10.1016/j.gr.2019.01.010](https://doi.org/10.1016/j.gr.2019.01.010).
- Croon MB, Cande SC, Stock JM. 2008. Revised Pacific-Antarctic plate motions and geophysics of the Menard Fracture Zone. *Geochemistry, Geophysics, Geosystems*. 9:7. doi:[10.1029/2008GC002019](https://doi.org/10.1029/2008GC002019).
- Crutchley GJ, Mountjoy JJ, Kellett RL, Black J, Bland KJ, Coffin RB, Field B, Henrys SA, Neil HL, Pecher IA, et al. 2016. Petroleum Prospective Screening Report of the offshore Northern East Coast Basin. GNS Science consultancy report, 2015/225. 150 p.
- Dallanave E, Maurizot P, Agnini C, Sutherland R, Hollis CJ, Collot J, Dickens GR, Bachadse V, Strogen D, Morgans HEG. 2020. Eocene (46–44 Ma) onset of Australia-Pacific Plate motion in the Southwest Pacific inferred from stratigraphy in New Caledonia and New Zealand. *Geochemistry, Geophysics, Geosystems*. 21(7): e2019GC008699. doi:[10.1029/2019GC008699](https://doi.org/10.1029/2019GC008699).
- Davy B. 2014. Rotation and offset of the Gondwana convergent margin in the New Zealand region following Cretaceous jamming of Hikurangi Plateau large igneous province subduction. *Tectonics*. 33(8):1577–1595. doi:[10.1002/2014TC003629](https://doi.org/10.1002/2014TC003629).
- Davy B, Hoernle K, Werner R. 2008. Hikurangi Plateau: crustal structure, rifted formation, and Gondwana subduction history. *Geochemistry, Geophysics, Geosystems*. 9(7):31. doi:[10.1029/2007gc001855](https://doi.org/10.1029/2007gc001855).
- Davy B, Uruski CI. 2002. Results of the 2001 deep seismic survey of the Chatham Rise and Hikurangi Plateau and implications for petroleum exploration. *New Zealand Petroleum Conference Proceedings 2002*. Ministry of Economic Development, Wellington. 10 p.
- Delteil J, de Lepinay BM, Morgans HEG, Field BD. 2006. Olistostromes marking tectonic events, East Coast, New Zealand. *New Zealand Journal of Geology and Geophysics*. 49(4):517–531. doi:[10.1080/00288306.2006.9515185](https://doi.org/10.1080/00288306.2006.9515185).
- Edbrooke SW, Sykes R, Pocknall DT. 1994. *Geology of the Waikato Coal Measures, Waikato Coal Region, New Zealand*. Institute of Geological & Nuclear Sciences Monograph, 6. Lower Hutt. 236 p., 8 maps.
- Field BD, Browne GH, Davy BW, Herzer RH, Hoskins RH, Raine JI, Wilson GJ, Sewell RJ, Smale D, Watters WA. 1989. Cretaceous and Cenozoic sedimentary basins and geological evolution of the Canterbury region, South Island, New Zealand. *New Zealand Geological Survey Basin Studies*, 2. Lower Hutt, New Zealand Geological Survey. 94 p.
- Field BD, Uruski CI, Beu AG, Browne GH, Crampton JS, Funnell RH, Killups SD, Laird M, Mazengarb C, Morgans HEG, et al. 1997. Cretaceous-Cenozoic geology and petroleum systems of the East Coast region, New Zealand. Institute of Geological & Nuclear Sciences Monograph 19: 301 p & 7 enclosures.
- Gaina C, Müller DR, Roest WR, Symonds P. 1998a. The opening of the Tasman Sea: a gravity anomaly animation. *Earth Interactions*. 2(4):1–23. doi:[10.1175/1087-3562\(1998\)002<0001:TOOTTS>2.3.CO;2](https://doi.org/10.1175/1087-3562(1998)002<0001:TOOTTS>2.3.CO;2).
- Gaina C, Müller DR, Royer J-Y, Stock J, Hardebeck J, Symonds P. 1998b. The tectonic history of the Tasman Sea: A puzzle with 13 pieces. *Journal of Geophysical Research*. 103(B6):12413–12433. doi:[10.1029/98jb00386](https://doi.org/10.1029/98jb00386).
- Gaina C, Müller RD, Royer JY, Symonds P. 1999. Evolution of the Louisiade triple junction. *Journal of Geophysical Research: Solid Earth*. 104(B6):12927–12939. doi:[10.1029/1999jb900038](https://doi.org/10.1029/1999jb900038).
- Galbraith EM, Elliott JEG, Field BD, Martin C, Scadden PG, Scowen P, Molina M. 2015. Developing the New Zealand Measured Section Database (MSD). GNS Science internal report, 2015/03. 21 p.
- Ghisetti F, Sibson RH, Hamling I. 2016. Deformed Neogene basins, active faulting and topography in Westland: distributed crustal mobility west of the Alpine Fault transpressive plate boundary (South Island, New Zealand). *Tectonophysics*. 693(Part B):340–362. doi:[10.1016/j.tecto.2016.03.024](https://doi.org/10.1016/j.tecto.2016.03.024).
- Ghisetti FC. 2021. Map-view restorations of the South Island, New Zealand: a reappraisal of the last 10 Myr of evolution of the Alpine and Wairau faults. *New Zealand Journal of Geology and Geophysics*. 1–26. doi:[10.1080/00288306.2021.1878243](https://doi.org/10.1080/00288306.2021.1878243).
- Giba M, Nicol A, Walsh JJ. 2010. Evolution of faulting and volcanism in a back-arc basin and its implications for

- subduction processes. *Tectonics* 29: TC2040 18 p. doi:[10.1029/2009TC002634](https://doi.org/10.1029/2009TC002634).
- Giba M, Walsh JJ, Nicol A, Mouslopoulou V, Seebeck H. 2013. Investigation of the spatio-temporal relationship between normal faulting and arc volcanism on million-year timescales. *Journal of the Geological Society, London*. 170:951–962.
- Granot R, Cande SC, Stock JM, Damaske D. 2013. Revised Eocene-Oligocene kinematics for the West Antarctic rift system. *Geophysical Research Letters*. 40(2):279–284. doi:[10.1029/2012GL054181](https://doi.org/10.1029/2012GL054181).
- Gurnis M, Van Avendonk H, Gulick SPS, Stock J, Sutherland R, Hightower E, Shuck B, Patel J, Williams E, Kardell D, et al. 2019. Incipient subduction at the contact with stretched continental crust: the Puysegur Trench. *Earth and Planetary Science Letters*. 520:212–219. doi:[10.1016/j.epsl.2019.05.044](https://doi.org/10.1016/j.epsl.2019.05.044).
- Hayward BW. 1986. A guide to paleoenvironmental assessment using New Zealand Cenozoic foraminiferal faunas. New Zealand Geological Survey report PAL, 109. Lower Hutt, New Zealand Geological Survey. 73 p.
- Hayward BW. 2004. Foraminifera-based estimates of paleobathymetry using modern analogue technique, and the subsidence history of the Early Miocene Waitemata Basin. *New Zealand Journal of Geology and Geophysics*. 47(4):749–767.
- Hayward BW, Black PM, Smith IEM, Ballance PF, Itaya T, Doi M, Takagi M, Bergman S, Adams CJ, Herzer RH, et al. 2001. K-Ar ages of early Miocene arc-type volcanoes in northern New Zealand. *New Zealand Journal of Geology and Geophysics*. 44(2):285–311. doi:[10.1080/00288306.2001.9514939](https://doi.org/10.1080/00288306.2001.9514939).
- Hayward BW, Grenfell HR, Carter R, Hayward JJ. 2004. Benthic foraminiferal proxy evidence for the Neogene palaeoceanographic history of the Southwest Pacific, east of New Zealand. *Marine Geology*. 205(1):147–184. doi:[10.1016/S0025-3227\(04\)00022-2](https://doi.org/10.1016/S0025-3227(04)00022-2).
- Hayward BW, Sabaa AT, Grenfell HR, Neil H, Bostock H. 2013. Ecological distribution of recent deep-water foraminifera around New Zealand. *Journal of Foraminiferal Research*. 43(4):415–442. doi:[10.2113/gsjfr.43.4.415](https://doi.org/10.2113/gsjfr.43.4.415).
- Heron DW (custodian). 2018. Geological Map of New Zealand 1:250,000. 2nd ed. GNS Science Geological Map, 1. 1 CD.
- Herzer RH, Barker DHN, Roest WR, Mortimer N. 2011. Oligocene-Miocene spreading history of the northern South Fiji Basin and implications for the evolution of the New Zealand plate boundary. *Geochemistry Geophysics Geosystems*. 12(2):Q02004. doi:[10.1029/2010gc003291](https://doi.org/10.1029/2010gc003291).
- Herzer RH, Chaproniere GCH, Edwards AR, Hollis CJ, Pelletier B, Raine JI, Scott GH, Stagpoole V, Strong CP, Symonds P, et al. 1997. Seismic stratigraphy and structural history of the Reinga Basin and its margins, southern Norfolk Ridge system. *New Zealand Journal of Geology and Geophysics*. 40(4):425–451. doi:[10.1080/00288306.1997.9514774](https://doi.org/10.1080/00288306.1997.9514774).
- Herzer RH, Maslce J. 1996. Anatomy of a continent-backarc transform — The Vening Meinesz Fracture Zone northwest of New Zealand. *Marine Geophysical Researches*. 18 (2):401–427. doi:[10.1007/BF00286087](https://doi.org/10.1007/BF00286087).
- Higgs KE, Crouch EM, Raine JI. 2017. An interdisciplinary approach to reservoir characterisation; an example from the early to middle Eocene Kaimiro Formation, Taranaki Basin, New Zealand. *Marine and Petroleum Geology*. 86:111–139. doi:[10.1016/j.marpetgeo.2017.05.018](https://doi.org/10.1016/j.marpetgeo.2017.05.018).
- Hines BR. 2018. Cretaceous to Paleogene palinspastic reconstruction of the East Coast Basin, New Zealand Unpublished Ph.D. thesis, Victoria University of Wellington. 522 p.
- Hines BR, Seebeck HC, Crampton JS, Bland KJ, Strogen DP. 2022. Reconstructing a dismembered basin along the active Hikurangi subduction margin, New Zealand. *GSA Bulletin*. doi:[10.1130/B36308.1](https://doi.org/10.1130/B36308.1).
- Isaac MJ, Herzer RH, Brook FJ, Hayward BW. 1994. Cretaceous and Cenozoic sedimentary basins of Northland, New Zealand. Institute of Geological & Nuclear Sciences Monograph, 8. Lower Hutt. 230 p.
- Kamp PJJ, Vonk AJ, Bland KJ, Hansen RJ, Hendy AJW, McIntyre AP, Ngatai M, Cartwright SJ, Hayton S, Nelson CS. 2004. Neogene stratigraphic architecture and tectonic evolution of Wanganui, King Country, and eastern Taranaki basins, New Zealand. *New Zealand Journal of Geology and Geophysics*. 47(4):625–644. doi:[10.1080/00288306.2004.9515080](https://doi.org/10.1080/00288306.2004.9515080).
- Kennett JP, von der Borch CC, et al. 1986. Initial reports of the Deep Sea Drilling Project, 90. Washington, US Govt. Printing Office. doi:[10.2973/dsdp.proc.90.1986](https://doi.org/10.2973/dsdp.proc.90.1986).
- King PR. 2000. Tectonic reconstructions of New Zealand 40 Ma to the present. *New Zealand Journal of Geology and Geophysics*. 43(4):611–638. doi:[10.1080/00288306.2000.9514913](https://doi.org/10.1080/00288306.2000.9514913).
- King PR, Naish TR, Browne GH, Field BD, Edbrooke SW. (compilers). 1999. Cretaceous to Recent sedimentary patterns in New Zealand. Institute of Geological & Nuclear Sciences Folio Series, 1. Lower Hutt, New Zealand. 35 p, 39 transects, 1 enclosure.
- King PR, Thrasher GP. 1996. Cretaceous-Cenozoic geology and petroleum systems of the Taranaki Basin, New Zealand. Institute of Geological & Nuclear Sciences Monograph, 13. Lower Hutt, New Zealand. 243 p, 6 enclosures.
- Kulhanek DK, Morgans HEG. 2013. Micropaleontological constraints on age and paleodepth across Paleogene unconformities on Lord Howe Rise and in the New Caledonia Basin. GNS Science consultancy report, 2013/201. 67 p.
- Laird MG, Bradshaw JD. 2004. The break-up of a long-term relationship: the Cretaceous separation of New Zealand from Gondwana. *Gondwana Research*. 7(1):273–286. doi:[10.1016/S1342-937X\(05\)70325-7](https://doi.org/10.1016/S1342-937X(05)70325-7).
- Lamb S, Mortimer N. 2020. Taking time to twist a continent — multistage origin of the New Zealand orocline. *Geology*. 49(1):56–60. doi:[10.1130/G47805.1](https://doi.org/10.1130/G47805.1).
- Lamb S, Mortimer N, Smith E, Turner G. 2016. Focusing of relative plate motion at a continental transform fault: Cenozoic dextral displacement >700 km on New Zealand's Alpine Fault, reversing >225 km of Late Cretaceous sinistral motion. *Geochemistry, Geophysics, Geosystems*. 17(3):1197–1213. doi:[10.1002/2015GC006225](https://doi.org/10.1002/2015GC006225).
- Landis CA, Campbell HJ, Begg JG, Mildenhall DC, Paterson AM, Trewick SA. 2008. The Waipounamu erosion surface: questioning the antiquity of the New Zealand land surface and terrestrial fauna and flora. *Geological Magazine*. 145(2):173–197. doi:[10.1017/S0016756807004268](https://doi.org/10.1017/S0016756807004268).
- Langridge RM, Ries WF, Litchfield NJ, Villamor P, Van Dissen RJ, Barrell DJA, Rattenbury MS, Heron DW, Haubrock S, Townsend DB, et al. 2016. The New Zealand Active Faults Database. *New Zealand Journal of Geology and Geophysics*. 59(1):86–96. doi:[10.1080/00288306.2015.1112818](https://doi.org/10.1080/00288306.2015.1112818).

- Lawrence MJF, Morgans HEG, Crundwell MP, Patriat M. **2020**. Carbonate rocks of offshore northern Zealandia. *New Zealand Journal of Geology and Geophysics*. 63(1):66–89. doi:[10.1080/00288306.2019.1626745](https://doi.org/10.1080/00288306.2019.1626745).
- Little TA, Jones A. **1998**. Seven million years of strike-slip and related off-fault deformation, northeastern Marlborough fault system, South Island, New Zealand. *Tectonics*. 17(2):285–302. doi:[10.1029/97tc03148](https://doi.org/10.1029/97tc03148).
- Little TA, Roberts AP. **1997**. Distribution and mechanism of Neogene to present-day vertical axis rotations, Pacific-Australian Plate Boundary Zone, South Island, New Zealand. *Journal of Geophysical Research: Solid Earth*. 102(B9):20447–20468. doi:[10.1029/97JB01279](https://doi.org/10.1029/97JB01279).
- Matthews KJ, Müller RD, Wessel P, Whittaker JM. **2011**. The tectonic fabric of the ocean basins. *Journal of Geophysical Research: Solid Earth*. 116:B12. doi:[10.1029/2011JB008413](https://doi.org/10.1029/2011JB008413).
- Maurizot P, Bordenave A, Cluzel D, Collot J, Etienne S. **2020a**. Late Cretaceous to Eocene cover of New Caledonia: from rifting to convergence. In: Maurizot P, Mortimer N, editor. *New Caledonia: geology, geodynamic evolution and mineral resources*. New Caledonia: geology, geodynamic evolution and mineral resources. 51. London: Geological Society; p. 53–91. doi:[10.1144/m51-2017-18](https://doi.org/10.1144/m51-2017-18)
- Maurizot P, Cluzel D, Patriat M, Collot J, Iseppi M, Lesimple S, Secchiari A, Bosch D, Montanini A, Macera P, et al. **2020b**. The Eocene subduction–obduction complex of New Caledonia. In: Maurizot P, Mortimer N, editor. *New Caledonia: geology, geodynamic evolution and mineral resources*. 51. London: Geological Society of London; p. 93–130. doi:[10.1144/M51-2018-70](https://doi.org/10.1144/M51-2018-70)
- Meffre S, Falloon TJ, Crawford TJ, Hoernle K, Hauff F, Duncan RA, Bloomer SH, Wright DJ. **2012**. Basalts erupted along the Tongan fore arc during subduction initiation: evidence from geochronology of dredged rocks from the Tonga fore arc and trench. *Geochemistry, Geophysics, Geosystems*. 13(12):1–17. doi:[10.1029/2012GC004335](https://doi.org/10.1029/2012GC004335).
- Morgans HEG. **2016**. The Weber Formation at its type location in Southern Hawke's Bay, and other localities on the East Coast, North Island, New Zealand. *GNS Science Report*, 2015/38. 101 p.
- Mortimer N. **2004**. New Zealand's geological foundations. *Gondwana Research*. 7(1):261–272. doi:[10.1016/S1342-937X\(05\)70324-5](https://doi.org/10.1016/S1342-937X(05)70324-5).
- Mortimer N. **2014**. The oroclinal bend in the South Island, New Zealand. *Journal of Structural Geology*. 64:32–38. doi:[10.1016/j.jsg.2013.08.011](https://doi.org/10.1016/j.jsg.2013.08.011).
- Mortimer N. **2020**. Information sources for GNS Science Poster Series 7 and 8: bathymetric and tectonic maps of Te Riu-a-Maui/Zealandia. . *GNS Science Report*, 2020/01. 28 p. doi:[10.21420/10.21420/K8BF-CH36](https://doi.org/10.21420/10.21420/K8BF-CH36).
- Mortimer N, Adams CJ, Wright IC. **2020a**. Early Cretaceous greywacke from Colville Knolls, New Zealand. *New Zealand Journal of Geology and Geophysics*. 63(1):145–150. doi:[10.1080/00288306.2019.1629967](https://doi.org/10.1080/00288306.2019.1629967).
- Mortimer N, Campbell HJ. **2014**. Zealandia: our continent revealed. Auckland, NZ: Penguin; 271 p.
- Mortimer N, Campbell HJ, Tulloch AJ, King PR, Stagpoole VM, Wood RA, Rattenbury MS, Sutherland R, Adams CJ, Collot J, et al. **2017**. Zealandia: Earth's hidden continent. *GSA Today*. 27(3):8 p. doi:[10.1130/GSATG321A.1](https://doi.org/10.1130/GSATG321A.1).
- Mortimer N, Gans PB, Meffre S, Martin CE, Seton M, Williams S, Turnbull RE, Quilty PG, Micklethwaite S, Timm C, et al. **2018**. Regional volcanism of northern Zealandia: post-Gondwana break-up magmatism on an extended, submerged continent. In: Sensarma S, Storey BC editors. *Special Publications* 463(1). London: Geological Society; p. 199–226. doi:[10.1144/sp463.9](https://doi.org/10.1144/sp463.9)
- Mortimer N, Herzer RH, Gans PB, Laporte-Magoni C, Calvert AT, Bosch D. **2007**. Oligocene–Miocene tectonic evolution of the South Fiji Basin and Northland Plateau, SW Pacific Ocean: Evidence from petrology and dating of dredged rocks. *Marine Geology*. 237(1–2):1–24. doi:[10.1016/j.margeo.2006.10.033](https://doi.org/10.1016/j.margeo.2006.10.033).
- Mortimer N, Patriat M, Gans PB, Agranier A, Chazot G, Collot J, Crundwell MP, Durance PMJ, Campbell HJ, Etienne S. **2020d**. The Norfolk Ridge seamounts: Eocene–Miocene volcanoes near Zealandia's rifted continental margin. *Australian Journal of Earth Sciences*. 1–13. doi:[10.1080/08120099.2020.1805007](https://doi.org/10.1080/08120099.2020.1805007).
- Mortimer N, Rattenbury MS, King PR, Bland KJ, Barrell DJA, Bache F, Begg JG, Campbell HJ, Cox SC, Crampton JS, et al. **2014**. High-level stratigraphic scheme for New Zealand rocks. *New Zealand Journal of Geology and Geophysics*. 57(4):402–419. doi:[10.1080/00288306.2014.946062](https://doi.org/10.1080/00288306.2014.946062).
- Mortimer N, Smith Lyttle B, Black J. **2020b**. Bathymetric map of Te Riu-a-Maui/Zealandia. Scale 1:8500000. *GNS Science Poster*, 7. Lower Hutt, Institute of Geological & Nuclear Sciences. 1 map. doi:[10.21420/M90C-5J03](https://doi.org/10.21420/M90C-5J03)
- Mortimer N, Smith Lyttle B, Black J. **2020c**. Tectonic map of Te Riu-a-Maui/Zealandia. Scale 1:8500000. *GNS Science Poster*, 8. Lower Hutt, Institute of Geological & Nuclear Sciences. 1 map. doi:[10.21420/M90C-5J03](https://doi.org/10.21420/M90C-5J03)
- Müller D, Russell S, Zahirovic S, Williams S, Williams C. **2018a**. Modelling and visualising distributed crustal deformation of Australia and Zealandia using GPlates 2.0. *ASEG Extended Abstracts*. 2018(1):1–7.
- Müller RD, Cannon J, Qin X, Watson RJ, Gurnis M, Williams S, Pfaffelmoser T, Seton M, Russell SHJ, Zahirovic S. **2018b**. GPlates: building a virtual Earth through deep time. *Geochemistry, Geophysics, Geosystems*. 19(7):2243–2261. doi:[10.1029/2018GC007584](https://doi.org/10.1029/2018GC007584).
- Müller RD, Seton M, Zahirovic S, Williams SE, Matthews KJ, Wright NM, Shephard GE, Maloney KT, Barnett-Moore N, Hosseinpour M, et al. **2016**. Ocean basin evolution and global-scale plate reorganization events since Pangea breakup. *Annual Review of Earth and Planetary Sciences*. 44(1):107–138. doi:[10.1146/annurev-earth-060115-012211](https://doi.org/10.1146/annurev-earth-060115-012211).
- Nathan S, Anderson HJ, Cook RA, Herzer RH, Hoskins RH, Raine JI, Smale D. **1986**. Cretaceous and Cenozoic sedimentary basins of the West Coast region, South Island, New Zealand. *New Zealand Geological Survey Basin Studies*, 1. Wellington. 89 p.
- Nelson CS, Hume TM. **1977**. Relative intensity of tectonic events revealed by the tertiary sedimentary record in the North Wanganui Basin and adjacent areas, New Zealand. *New Zealand Journal of Geology and Geophysics*. 20(2):369–392. doi:[10.1080/00288306.1977.10420714](https://doi.org/10.1080/00288306.1977.10420714).
- New Zealand Petroleum & Minerals. **2015**. 2015 New Zealand petroleum exploration data pack. 1 hard drive.
- Nicol A. **2011**. Landscape history of the Marlborough Sounds. *New Zealand Journal of Geology and Geophysics*. 54(2):195–208. doi:[10.1080/00288306.2010.523079](https://doi.org/10.1080/00288306.2010.523079).
- Nicol A, Mazengarb C, Chanier F, Rait G, Uruski CI, Wallace LM. **2007**. Tectonic evolution of the active Hikurangi subduction margin, New Zealand, since the

- Oligocene. *Tectonics*. 26(4):TC4002. doi:doi:10.1029/2006TC002090.
- Norris RJ, Turnbull IM. 1993. Cenozoic basins adjacent to an evolving transform plate boundary, southwest New Zealand. In: Ballance PF, editor. *South Pacific sedimentary basins. South Pacific sedimentary basins*, 2. Amsterdam: Elsevier Science Publishers; p. 251–270.
- Orr D, Sutherland R, Stratford WR. 2020. Eocene to Miocene subduction initiation recorded in stratigraphy of Reinga Basin, northwest New Zealand. *Tectonics*. e2019TC005899: 1–21. doi:10.1029/2019TC005899
- Pettinga JR. 1982. Upper Cenozoic structural history, coastal Southern Hawke's Bay, New Zealand. *New Zealand Journal of Geology and Geophysics*. 25(2):149–191. doi:10.1080/00288306.1982.10421407.
- Raine JI, Beu AG, Boyes AF, Campbell HJ, Cooper RA, Crampton JS, Crundwell MP, Hollis CJ, Morgans HEG, Mortimer N. 2015. New Zealand geological timescale NZGT 2015/1. *New Zealand Journal of Geology and Geophysics*. 58(4):398–403. doi:10.1080/00288306.2015.1086391.
- Rait G, Chanier F, Waters DW. 1991. Landward- and seaward-directed thrusting accompanying the onset of subduction beneath New Zealand. *Geology*. 19(3):230–233. doi:10.1130/0091-7613(1991)019<0230:LASDTA>2.3.CO;2.
- Raymond OL. 2018. Australian geological provinces 2018, 1st edition. Canberra: Geoscience Australia. <http://pid.geoscience.gov.au/dataset/ga/116823>.
- Reyners M. 2018. Impacts of Hikurangi Plateau subduction on the origin and evolution of the Alpine Fault. *New Zealand Journal of Geology and Geophysics*. 61(3):260–271. doi:10.1080/00288306.2018.1454481.
- Reyners M, Eberhart-Phillips D, Bannister S. 2017a. Subducting an old subduction zone sideways provides insights into what controls plate coupling. *Earth and Planetary Science Letters*. 466:53–61. doi:10.1016/j.epsl.2017.03.004.
- Reyners M, Eberhart-Phillips D, Upton P, Gubbins D. 2017b. Three-dimensional imaging of impact of a large igneous province with a subduction zone. *Earth and Planetary Science Letters*. 460:143–151. doi:10.1016/j.epsl.2016.12.025.
- Rieftahl F, Gohl K, Davy B, Barrett R. 2020a. Extent and cessation of the mid-Cretaceous Hikurangi Plateau underthrusting: impact on global plate tectonics and the submarine Chatham Rise. *Journal of Geophysical Research: Solid Earth*. 125(8):e2020JB019681. doi:10.1029/2020JB019681.
- Rieftahl F, Gohl K, Davy B, Hoernle K, Mortimer N, Timm C, Werner R, Hochmuth K. 2020b. Cretaceous intracontinental rifting at the southern Chatham Rise margin and initialisation of seafloor spreading between Zealandia and Antarctica. *Tectonophysics*. 776:228298. doi:10.1016/j.tecto.2019.228298.
- Sagar MW, Higgs KE, Strogen DP, Bland KJ, Browne GH. 2022. Provenance of sandstones in the Murchison Basin, New Zealand: a record of Neogene strike-slip, convergence and basement exhumation along the Australian-Pacific plate boundary. *New Zealand Journal of Geology and Geophysics*. doi:10.1080/00288306.2022.2108067.
- Sahoo TR, Bland KJ, Arnot MJ, Boyes AF, Funnell RH, Griffin AG, Kroeger K, Lawrence MJF, O'Brien G, Scadden PG, et al. 2017. *Atlas of Petroleum Prospectivity, Southeast Province: ArcGIS geodatabase and technical report*. GNS Science data series, 23c. 1 ArcGIS geodatabase, 5 ArcGIS projects, 1 technical report.
- Sahoo TR, Strogen DP, Browne GH, Nicol A. 2022. Evolution of syn- to early post-rift facies in rift basins: insights from the Cretaceous–Paleocene of the Great South Basin, New Zealand. *Basin Research*. 34(3):1113–1142. doi:10.1111/bre.12652.
- Schellart W, Lister GS, Toy VG. 2006. A Late Cretaceous and Cenozoic reconstruction of the Southwest Pacific region: tectonics controlled by subduction and slab rollback processes. *Earth Science Reviews*. 76:191–233.
- Sdrolias M, Müller RD, Gaina C. 2003. Tectonic evolution of the southwest Pacific using constraints from backarc basins. In: Hillis RR, Müller RD, editor. *Evolution and dynamics of the Australian Plate*. Geological Society of America; p. 343–359.
- Seebeck H, Nicol A, Giba M, Pettinga J, Walsh J. 2013. Geometry of the subducting Pacific plate since 20 Ma, Hikurangi margin, New Zealand. *Journal of the Geological Society*. 171:131–143. doi:10.1144/jgs2012-145.
- Seebeck H, Strogen D, King P, Nicol A, Hines B, O'Brien G. 2018. Cretaceous to present-day tectonic reconstructions of Zealandia. *The APPEA Journal*. 58(2):852–857. doi:10.1071/AJ17117.
- Seton M, Müller RD, Zahirovic S, Gaina C, Torsvik T, Shephard G, Talsma A, Gurnis M, Turner M, Maus S, et al. 2012. Global continental and ocean basin reconstructions since 200Ma. *Earth-Science Reviews*. 113(3):212–270. doi:10.1016/j.earscirev.2012.03.002.
- Sevin B, Maurizot P, Cluzel D, Tournadour E, Etienne S, Folcher N, Jeanpert J, Collot J, Iseppi M, Meffre S, et al. 2020. Post-obduction evolution of New Caledonia. In: Maurizot P, Mortimer N, editor. *New Caledonia: geology, geodynamic evolution and mineral resources*. London: Geological Society of London; p. 147–188. doi:10.1144/M51-2018-74.
- Shuck B, Van Avendonk H, Gulick SPS, Gurnis M, Sutherland R, Stock J, Patel J, Hightower E, Sastrup S, Hess T. 2021. Strike-Slip enables subduction initiation beneath a failed rift: New seismic constraints from Puysegur Margin, New Zealand. *Tectonics*. 40(5): e2020TC006436. doi:10.1029/2020TC006436.
- Skinner C, Sutherland R. 2022. Cretaceous rift-drift tectonics then Paleogene prearc subsidence related to subduction initiation: Aotea Basin, Zealandia, Southwest Pacific. *Tectonics*. 41(1):e2021TC006820. doi:10.1029/2021TC006820.
- Stagpoole VM, Browne GH, Ilg B, Herzer RH, Reid E, Bland KJ, Griffin AG, Uruski CI. 2009. Petroleum prospectivity of the Reinga Basin, New Zealand. Unpublished Openfile Petroleum Report 4037: 112 p.
- Stagpoole VM, Field BD, Sutherland R, Uruski CI, Zink K. 2008. Petroleum prospectivity of the Raukumara Basin, New Zealand. Unpublished Openfile Petroleum Report PR3921: 111 p.
- Stagpoole VM, Nicol A. 2008. Regional structure and kinematic history of a large subduction back thrust: Taranaki Fault, New Zealand. *Journal of Geophysical Research*. 113:B01403. doi:10.1029/2007JB005170.
- Strogen DP. 2011. Updated paleogeographic maps for the Taranaki Basin and surrounds. GNS Science Report, 2010/53. 83 p.
- Strogen DP, Bland KJ, Nicol A, King PR. 2014. Paleogeography of the Taranaki Basin region during the latest Eocene–Early Miocene and implications for the

- 'total drowning' of Zealandia. *New Zealand Journal of Geology and Geophysics*. 57(2):110–127. doi:[10.1080/00288306.2014.901231](https://doi.org/10.1080/00288306.2014.901231).
- Strogen DP, Higgs KE, Griffin AG, Morgans HEG. 2019. Late Eocene – Early Miocene facies and stratigraphic development, Taranaki Basin, New Zealand: the transition to plate boundary tectonics during regional transgression. *Geological Magazine*. 156(10):1751–1770. doi:[10.1017/S0016756818000997](https://doi.org/10.1017/S0016756818000997).
- Strogen DP, King PR 2014. A new Zealandia-wide seismic horizon naming scheme. GNS Science Report, 2014/34. 20 p.
- Strogen DP, Seebeck H, Nicol A, King PR. 2017. Two-phase Cretaceous–Paleocene rifting in the Taranaki Basin region, New Zealand; implications for Gondwana break-up. *Journal of the Geological Society*. 174(5):929–946. doi:[10.1144/jgs2016-160](https://doi.org/10.1144/jgs2016-160).
- Strong DT, Turnbull RE, Haubrock S, Mortimer N. 2016. Petlab: New Zealand's national rock catalogue and geoanalytical database. *New Zealand Journal of Geology and Geophysics*. 59(3):475–481. doi:[10.1080/00288306.2016.1157086](https://doi.org/10.1080/00288306.2016.1157086).
- Sutherland R. 1995. The Australia-Pacific boundary and Cenozoic plate motions in the SW Pacific: some constraints from Geosat data. *Tectonics*. 14(4):819–831. doi:[10.1029/95tc00930](https://doi.org/10.1029/95tc00930).
- Sutherland R, Barnes PM, Uruski CI. 2006. Miocene–Recent deformation, surface elevation, and volcanic intrusion of the overriding plate during subduction initiation, offshore southern Fiordland, Puysegur margin, southwest New Zealand. *New Zealand Journal of Geology and Geophysics*. 49:131–149.
- Sutherland R, Collot J, Bache F, Henrys S, Barker D, Browne GH, Lawrence MJF, Morgans HEG, Hollis CJ, Clowes C, et al. 2017. Widespread compression associated with Eocene Tonga-Kermadec subduction initiation. *Geology*. 45(4):355–358. doi:[10.1130/g38617.1](https://doi.org/10.1130/g38617.1).
- Sutherland R, Dickens GR, Blum P, Agnini C, Alegret L, Asatryan G, Bhattacharya J, Bordenave A, Chang L, Collot J, et al. 2020. Continental-scale geographic change across Zealandia during Paleogene subduction initiation. *Geology*. 1–6. doi:[10.1130/G47008.1](https://doi.org/10.1130/G47008.1).
- Sutherland R, Dickens GR, Blum P, the Expedition 371 Scientists. 2019. Tasman Frontier Subduction Initiation and Paleogene Climate. *Proceedings of the International Ocean Discovery Program*, 371. College Station, TX, International Ocean Discovery Program. <https://doi.org/10.14379/iodp.proc.371.2019>.
- Sutherland R, King PR, Wood RA. 2001. Tectonic evolution of cretaceous rift basins in south-eastern Australia and New Zealand: implications for exploration risk assessment. In: Hill KC, Bernecker T, editor. *Eastern Australasian Basins Symposium 2001*. Eastern Australasian Basins Symposium 2001. Melbourne: PESA; p. 3–13.
- Sutherland R, Spasojevic S, Gurnis M. 2010. Mantle upwelling after Gondwana subduction death explains anomalous topography and subsidence histories of eastern New Zealand and West Antarctica. *Geology*. 38(2):155–158. doi:[10.1130/g30613.1](https://doi.org/10.1130/g30613.1).
- Sutherland R, Stagpoole V, Uruski C, Kennedy C, Bassett D, Henrys S, Scherwath M, Kopp H, Field B, Toulmin S, et al. 2009. Reactivation of tectonics, crustal underplating, and uplift after 60 Myr of passive subsidence, Raukumara Basin, Hikurangi-Kermadec fore arc, New Zealand: Implications for global growth and recycling of continents. *Tectonics*. 28(5):TC5017. doi:[10.1029/2008tc002356](https://doi.org/10.1029/2008tc002356).
- Sutherland R DVGP, Bache F, Stagpoole VM, Collot J, Rouillard P, Hashimoto T, Hackney R, Higgins KL, Rollet N, et al. 2012. Compilation of seismic reflection data from the Tasman Frontier region, southwest Pacific. GNS Science Report, 2012/01. 72 p.
- Taylor B, Zellmer K, Martinez F, Goodliffe A. 1996. Seafloor spreading in the Lau back-arc basin. *Earth and Planetary Science Letters*. 144(1):35–40. doi:[10.1016/0012-821X\(96\)00148-3](https://doi.org/10.1016/0012-821X(96)00148-3).
- Tontini FC, Bassett D, de Ronde CEJ, Timm C, Wysoczanski R. 2019. Early evolution of a young back-arc basin in the Havre Trough. *Nature Geoscience*. 12(10):856–862. doi:[10.1038/s41561-019-0439-y](https://doi.org/10.1038/s41561-019-0439-y).
- Torsvik TH, Müller RD, Van der Voo R, Steinberger B, Gaina C. 2008. Global plate motion frames: toward a unified model. *Reviews of Geophysics*. 46:3. doi:[10.1029/2007RG000227](https://doi.org/10.1029/2007RG000227).
- Torsvik TH, Van der Voo R, Preeden U, Mac Niocaill C, Steinberger B, Doubrovine PV, van Hinsbergen DJJ, Domeier M, Gaina C, Tohver E, et al. 2012. Phanerozoic polar wander, palaeogeography and dynamics. *Earth-Science Reviews*. 114(3):325–368. doi:[10.1016/j.earscirev.2012.06.007](https://doi.org/10.1016/j.earscirev.2012.06.007).
- Totterdell J, Hall L, Hashimoto T, Owen K, Bradshaw M. 2014. Petroleum geology inventory of Australia's offshore frontier basins. *Geoscience Australia Record*, 2014/009. Geoscience Australia, Canberra. <http://doi.org/10.11636/Record.2014.009>.
- Totterdell JM, Cook PJ, Bradshaw MT, Wilford GE, Yeates AN, Yeung M, Truswell EM, Brakel AT, Isem AR, Ollisoff S, et al. 2001. Palaeogeographic Atlas of Australia (a set of ten volumes). Record 1 to 10. Canberra, Geoscience Australia. <http://pid.geoscience.gov.au/dataset/ga/35880>.
- Trewick SA, Bland KJ. 2012. Fire and slice: palaeogeography for biogeography at New Zealand's North Island/South Island juncture. *Journal of the Royal Society of New Zealand*. 42(3):153–183. doi:[10.1080/03036758.2010.549493](https://doi.org/10.1080/03036758.2010.549493).
- Turnbull IM, Uruski CI, Anderson HJ, Lindqvist JK, Scott GH, Morgans HEG, Hoskins RH, Raine JI, Mildenhall DC, Pocknall DT, et al. 1993. Cretaceous and Cenozoic sedimentary basins of western Southland, South Island, New Zealand. Institute of Geological & Nuclear Sciences Monograph, 1. Lower Hutt, Institute of Geological & Nuclear Sciences. 86 p.
- Uruski CI. 1997. New Zealand UNCLOS report: preliminary interpretation report Line TL-01, New Caledonia Basin. GNS Science Client Report 52655A. 11 p.
- Uruski CI. 2010. New Zealand's deepwater frontier. *Marine and Petroleum Geology*. 27(9):2005–2026.
- Uruski CI, Bland KJ. 2011. Pegasus Basin and the prospects for oil and gas. New Zealand Unpublished Openfile Petroleum Report PR4326: 117 p.
- Uruski CI, Browne GH, Stagpoole VM, Herzer RH, Bland KJ, Griffin AG, Sykes R, Roncaglia L, Schiøler P, Funnell RH, et al. 2008. Petroleum prospectivity of the Northland Basin, New Zealand. New Zealand Unpublished Openfile Petroleum Report PR3933: 89 p.
- Uruski CI, Field BD, Funnell RH. 2005. The East Coast Basin of New Zealand, an emerging petroleum province. *APPEA Journal*. 45(1):563–580.
- van Hinsbergen DJJ, de Groot LV, van Schaik SJ, Spakman W, Bijl PK, Sluijs A, Langereis CG, Brinkhuis H. 2015. A paleolatitude calculator for paleoclimate studies. *PLOS ONE*. 10(6):e0126946. doi:[10.1371/journal.pone.0126946](https://doi.org/10.1371/journal.pone.0126946).

- Walcott RI. 1978. Present tectonics and Late Cenozoic evolution of New Zealand. *Geophysical Journal of the Royal Astronomical Society*. 52(1):137–164.
- Wallace LM, Saffer DM, Barnes PM, Pecher IA, Petronotis KE, LeVay LJ, the Expedition 372B/375 Scientists. 2019. Hikurangi Subduction Margin Coring, Logging, and Observatories. *Proceedings of the International Ocean Discovery Program*, 372B/375. College Station, TX, International Ocean Discovery Program. doi:[10.14379/iodp.proc.372B375.2019](https://doi.org/10.14379/iodp.proc.372B375.2019)
- Whittaker JM, Müller RD, Leitchenkov G, Stagg H, Sdrolias M, Gaina C, Goncharov A. 2007. Major Australian-Antarctic Plate reorganization at Hawaiian-Emperor bend time. *Science*. 318(5847):83. doi:[10.1126/science.1143769](https://doi.org/10.1126/science.1143769).
- Whittaker JM, Williams SE, Müller RD. 2013. Revised tectonic evolution of the Eastern Indian Ocean. *Geochemistry, Geophysics, Geosystems*. 14(6):1891–1909. doi:[10.1002/ggge.20120](https://doi.org/10.1002/ggge.20120).
- Wobbe F, Gohl K, Chambord A, Sutherland R. 2012. Structure and breakup history of the rifted margin of West Antarctica in relation to Cretaceous separation from Zealandia and Bellingshausen plate motion. *Geochemistry, Geophysics, Geosystems*. 13:Q04W12. doi:[10.1029/2011GC003742](https://doi.org/10.1029/2011GC003742).
- Wood R, Davy B. 1994. The Hikurangi Plateau. *Marine Geology*. 118(1):153–173. doi:[10.1016/0025-3227\(94\)90118-X](https://doi.org/10.1016/0025-3227(94)90118-X).
- Wood RA. 1991. Structure and seismic stratigraphy of the western Challenger Plateau. *New Zealand Journal of Geology and Geophysics*. 34(1):1–9. doi:[10.1080/00288306.1991.9514433](https://doi.org/10.1080/00288306.1991.9514433).
- Wood RA. 1993. The Challenger Plateau. In: Ballance PF, editor. *South Pacific sedimentary basins*, 2. Amsterdam: Elsevier Science Publishers; p. 351–364.
- Wood RA. 2002. Finding the continental shelf: integration of geology and geophysics. *New Zealand Petroleum Conference Proceedings* 2002. Ministry of Economic Development, Wellington. p. 391–401.
- Wood RA, Andrews PB, Herzer RH, others. 1989. Cretaceous and Cenozoic geology of the Chatham Rise region, South Island, New Zealand. *New Zealand Geological Survey Basin Studies*, 3. Lower Hutt, New Zealand Geological Survey. 75 p.
- Wood RA, Davy BW. 2002. New Zealand's UNCLOS project: defining the continental margin: a summary and way forward. *New Zealand Science Review*. 64(3/4):59–64.
- Wright I, Sheppard J, Turner R, Barnes PM, Davy BW, Herzer RH, Lamarche G, Lewis K, Nodder S, Stagpoole VM, et al. 2002. New Zealand Continental Shelf Project: a status report of the survey programme. *New Zealand Petroleum Conference Proceedings* 2002. Ministry of Economic Development, Wellington. p. 373–380.
- Wright NM, Müller RD, Seton M, Williams SE. 2015. Revision of Paleogene plate motions in the Pacific and implications for the Hawaiian-Emperor bend. *Geology*. 43(5):455–458. doi:[10.1130/G36303.1](https://doi.org/10.1130/G36303.1).
- Wysoczanski R, Leonard G, Gill J, Wright I, Calvert A, McIntosh W, Jicha B, Gamble J, Timm C, Handler M, et al. 2019. Ar-Ar age constraints on the timing of Havre Trough opening and magmatism. *New Zealand Journal of Geology and Geophysics*. 62(3):371–377. doi:[10.1080/00288306.2019.1602059](https://doi.org/10.1080/00288306.2019.1602059).