


Reduced Order Models Using a Non-Local Calculus on Finite Weighted Graphs



Matthew Duschenes, Siddhartha Srivastava & Krishna Garikipati

Applied Physics, Mechanical Engineering, and
Michigan Institute for Computational Discovery & Engineering
University of Michigan, Ann Arbor


July 26, 2021

arXiv:2105.01740

USNCCM #309 Data-Driven Science



Outline



1. Systems of Interest
2. Graph Theoretic Approaches
3. Applications




1. Systems of Interest

2. Graph Theoretic Approaches

3. Applications



Diffusion Processes



Consider a local diffusion process for a scalar field $\phi(x, t)$, driven by gradient-flow:

$$\frac{\partial \phi}{\partial t} = - M_\phi \frac{\delta \psi}{\delta \phi} \quad (1)$$

which can be described by a free energy density


$$\psi = f(\phi) + \lambda |\nabla \phi|^2 \quad (2)$$

where the Landau f is algebraic in ϕ , and the total free energies are

$$\Psi = \int_{\Omega} d\Omega \psi , \quad F = \int_{\Omega} d\Omega f , \quad G = \int_{\Omega} d\Omega |\nabla \phi|^2 , \quad F' = \int_{\Omega} d\Omega f' . \quad (3)$$



Reduced Order Modelling



Can we use global observables, such as the field's moments:

$$\varphi_k = \frac{1}{\Omega} \int_{\Omega} d\Omega \underbrace{I(\phi)}_{\text{indicator function}} \phi^k \quad (4)$$

or its derivatives

$$\varphi_{\nabla^k} = \frac{1}{\Omega} \int_{\Omega} d\Omega I(\phi) \nabla^k \phi, \quad \varphi_{\nabla_k} = \frac{1}{\Omega} \int_{\Omega} d\Omega I(\phi) |\nabla \phi|^k \quad (5)$$


or functional derivatives (and how may we precisely define these?)

$$\frac{\delta \Psi}{\delta \varphi_k}, \frac{\delta \Psi}{\delta \varphi_{\nabla^k}}, \frac{\delta \Psi}{\delta \varphi_{\nabla_k}} \dots \quad (6)$$

to develop a reduced order model?



Models of Interest



We aim to find a model for a global gradient-flow (Banerjee *et al.*, *Comp. Meth. App. Mech. Eng.*, **351**, (2019)):

$$\frac{\partial \varphi}{\partial t} = -M_\varphi \frac{\delta \Psi}{\delta \varphi}(\varphi_k, \varphi_{\nabla^k}, \varphi_{\nabla_k}, \frac{\delta \Psi}{\delta \varphi_k}, \dots) \quad (7)$$

and a free energy functional representation

$$\Psi = \Psi(\varphi_k, \varphi_{\nabla^k}, \varphi_{\nabla_k}, \frac{\delta \Psi}{\delta \varphi_k}, \dots). \quad (8)$$

Taken's Theorem: (Stark, *J. Nonlinear Sci.*, **351**, (1999))

For $\phi(x, t) \in \mathcal{M} \subset \mathbb{R}^d$, a mapping $P : \phi(x, (k+1)\tau) = P_\tau(\phi(x, k\tau))$, and an observable, $\varphi[\phi] \in \mathbb{R}$, \exists an embedding

$$\mathcal{T} : \mathcal{M} \rightarrow \mathbb{R}^n, \quad \mathcal{T}[\phi(x, t)] = \{\varphi^{(1)}, \varphi^{(2)}, \dots, \varphi^{(n)}\} \quad (9)$$

for (P, φ) dense and open in $D^r \times C^r(\mathcal{M}, \mathbb{R})$ for $n \geq 2d + 1$ observations.





1. Systems of Interest

2. Graph Theoretic Approaches

3. Applications




Graph Theory Formalism






Graph Theory Formalism




- State Vertices and State Dimensions: $V = \{1, 2, \dots, n\}$, $\chi = \{1, 2, \dots, p\}$
- State Vector: $X = \{x = \{x^\mu\}_{\mu \in \chi} : x \in V\}$
i.e) $X = \{\varphi_k, \varphi_{\nabla^k}, \varphi_{\nabla_k}, \Psi, \dots\} : x \in V\}$
- State Transition Edges: $E = \{(x, y) : x, y \in V\}$
- Edge Weights: $w = \{w(x, y) : (x, y) \in E\}$
- Neighborhoods of Adjacent Vertices:
 $\delta = \{\delta(x) = \{y \in V : (x, y) \in E\} : x \in V\}$
- Derived Quantities (Gilboa and Osher, *Multiscale Model. Simul.*, 7 (2008)):
i.e) $D = \{\{\varphi(x), \frac{\delta \Psi}{\delta \varphi_k}(x), \frac{\delta \Psi}{\delta \varphi_{\nabla^k}}(x), \frac{\delta \Psi}{\delta \varphi_{\nabla_k}}(x), \dots\} : x \in V\}$






Graph Theory Formalism




- State Vertices and State Dimensions: $V = \{1, 2, \dots, n\}$, $\chi = \{1, 2, \dots, p\}$
- State Vector: $X = \{x = \{x^\mu\}_{\mu \in \chi} : x \in V\}$
i.e) $X = \{\varphi_k, \varphi_{\nabla^k}, \varphi_{\nabla_k}, \Psi, \dots\} : x \in V\}$
- State Transition Edges: $E = \{(x, y) : x, y \in V\}$
- Edge Weights: $w = \{w(x, y) : (x, y) \in E\}$
- Neighborhoods of Adjacent Vertices:
 $\delta = \{\delta(x) = \{y \in V : (x, y) \in E\} : x \in V\}$
- Derived Quantities (Gilboa and Osher, *Multiscale Model. Simul.*, 7 (2008)):
i.e) $D = \{\{\varphi(x), \frac{\delta \Psi}{\delta \varphi_k}(x), \frac{\delta \Psi}{\delta \varphi_{\nabla^k}}(x), \frac{\delta \Psi}{\delta \varphi_{\nabla_k}}(x), \dots\} : x \in V\}$






Graph Theory Formalism




- State Vertices and State Dimensions: $V = \{1, 2, \dots, n\}$, $\chi = \{1, 2, \dots, p\}$
- State Vector: $X = \{x = \{x^\mu\}_{\mu \in \chi} : x \in V\}$
i.e) $X = \{\varphi_k, \varphi_{\nabla^k}, \varphi_{\nabla_k}, \Psi, \dots\} : x \in V\}$
- State Transition Edges: $E = \{(x, y) : x, y \in V\}$
- Edge Weights: $w = \{w(x, y) : (x, y) \in E\}$
- Neighborhoods of Adjacent Vertices:
 $\delta = \{\delta(x) = \{y \in V : (x, y) \in E\} : x \in V\}$
- Derived Quantities (Gilboa and Osher, *Multiscale Model. Simul.*, 7 (2008)):
i.e) $D = \{\{\varphi(x), \frac{\delta \Psi}{\delta \varphi_k}(x), \frac{\delta \Psi}{\delta \varphi_{\nabla^k}}(x), \frac{\delta \Psi}{\delta \varphi_{\nabla_k}}(x), \dots\} : x \in V\}$






Graph Theory Formalism




- State Vertices and State Dimensions: $V = \{1, 2, \dots, n\}$, $\chi = \{1, 2, \dots, p\}$
- State Vector: $X = \{x = \{x^\mu\}_{\mu \in \chi} : x \in V\}$
i.e) $X = \{\varphi_k, \varphi_{\nabla^k}, \varphi_{\nabla_k}, \Psi, \dots\} : x \in V\}$
- State Transition Edges: $E = \{(x, y) : x, y \in V\}$
- Edge Weights: $w = \{w(x, y) : (x, y) \in E\}$
- Neighborhoods of Adjacent Vertices:
 $\delta = \{\delta(x) = \{y \in V : (x, y) \in E\} : x \in V\}$
- Derived Quantities (Gilboa and Osher, *Multiscale Model. Simul.*, 7 (2008)):
i.e) $D = \{\{\varphi(x), \frac{\delta \Psi}{\delta \varphi_k}(x), \frac{\delta \Psi}{\delta \varphi_{\nabla^k}}(x), \frac{\delta \Psi}{\delta \varphi_{\nabla_k}}(x), \dots\} : x \in V\}$






Graph Theory Formalism




- State Vertices and State Dimensions: $V = \{1, 2, \dots, n\}$, $\chi = \{1, 2, \dots, p\}$
- State Vector: $X = \{x = \{x^\mu\}_{\mu \in \chi} : x \in V\}$
i.e) $X = \{\varphi_k, \varphi_{\nabla^k}, \varphi_{\nabla_k}, \Psi, \dots\} : x \in V\}$
- State Transition Edges: $E = \{(x, y) : x, y \in V\}$
- Edge Weights: $w = \{w(x, y) : (x, y) \in E\}$
- Neighborhoods of Adjacent Vertices:
 $\delta = \{\delta(x) = \{y \in V : (x, y) \in E\} : x \in V\}$
- Derived Quantities (Gilboa and Osher, *Multiscale Model. Simul.*, 7 (2008)):
i.e) $D = \{\{\varphi(x), \frac{\delta \Psi}{\delta \varphi_k}(x), \frac{\delta \Psi}{\delta \varphi_{\nabla^k}}(x), \frac{\delta \Psi}{\delta \varphi_{\nabla_k}}(x), \dots\} : x \in V\}$






Graph Theory Formalism




- State Vertices and State Dimensions: $V = \{1, 2, \dots, n\}$, $\chi = \{1, 2, \dots, p\}$
- State Vector: $X = \{x = \{x^\mu\}_{\mu \in \chi} : x \in V\}$
i.e) $X = \{\varphi_k, \varphi_{\nabla^k}, \varphi_{\nabla_k}, \Psi, \dots\} : x \in V\}$
- State Transition Edges: $E = \{(x, y) : x, y \in V\}$
- Edge Weights: $w = \{w(x, y) : (x, y) \in E\}$
- Neighborhoods of Adjacent Vertices:
 $\delta = \{\delta(x) = \{y \in V : (x, y) \in E\} : x \in V\}$
- Derived Quantities (Gilboa and Osher, *Multiscale Model. Simul.*, 7 (2008)):
i.e) $D = \{\{\varphi(x), \frac{\delta\Psi}{\delta\varphi_k}(x), \frac{\delta\Psi}{\delta\varphi_{\nabla^k}}(x), \frac{\delta\Psi}{\delta\varphi_{\nabla_k}}(x), \dots\} : x \in V\}$





Non-Local Calculus




Partial derivatives, where $z = x - \tilde{x} \sim O(h)$:

$$\frac{\delta \varphi(\tilde{x})}{\delta x^\mu} \equiv \sum_{x \in \delta(\tilde{x})} \frac{\varphi(x) - \varphi(\tilde{x})}{z^\mu} w(\tilde{x}, x) \quad (10)$$




Non-Local Calculus




Partial derivatives, where $z = x - \tilde{x} \sim O(h)$:

$$\frac{\delta \varphi(\tilde{x})}{\delta x^\mu} \equiv \sum_{x \in \delta(\tilde{x})} \frac{\varphi(x) - \varphi(\tilde{x})}{z^\mu} w(\tilde{x}, x) \quad (10)$$





Non-Local Calculus




Partial derivatives, where $z = x - \tilde{x} \sim O(h)$:

$$\frac{\delta \varphi(\tilde{x})}{\delta x^\mu} \equiv \sum_{x \in \delta(\tilde{x})} \frac{\varphi(x) - \varphi(\tilde{x})}{z^\mu} w(\tilde{x}, x) \quad (10)$$


Modified k -order Taylor series based at \tilde{x} :

$$\begin{aligned} \varphi^{(k)}(x|\tilde{x}) &= \varphi(\tilde{x}) + \sum_{\mu \in \chi} \gamma^\mu \frac{\delta \varphi(\tilde{x})}{\delta x^\mu} z^\mu \\ &\quad + \frac{1}{2!} \sum_{\mu, \nu \in \chi} \gamma^{\mu\nu} \frac{\delta^2 \varphi(\tilde{x})}{\delta x^\mu \delta x^\nu} z^{\mu\nu} \\ &\quad + \dots \end{aligned} \quad (11)$$





Consistency of Modified Taylor Series



Given a k -order model, an r -order accurate *stencil* yields as $h \rightarrow 0$:
The *non-local derivatives* scale as

$$\frac{\delta^l \varphi(\tilde{x})}{\delta x^l} \rightarrow \frac{\partial^l \varphi(\tilde{x})}{\partial x^l} + O(h^{r+1-l}) , \quad (12)$$

the *linear coefficients* scale as

$$\gamma_l \rightarrow 1 + O(h^{\min(k,r)+1-l}) , \quad (13)$$

and so the *model error* scales as

$$\varphi^{(k)}(x|\tilde{x}) \rightarrow \varphi(x) + O(h^{\min(k,r)+1}) , \quad (14)$$

which corresponds with Sobolev interpolation error.




1. Systems of Interest

2. Graph Theoretic Approaches


3. Applications



Evolution of Diffusing Fields




$$\text{Local: } \frac{\partial \phi}{\partial t} = -M_\phi \frac{\delta \psi}{\delta \phi} \quad \rightarrow \quad \text{Global: } \frac{\partial \varphi}{\partial t} = -M_\varphi \frac{\delta \Psi}{\delta \varphi} - \tilde{\mu}_\varphi$$





Evolution of Diffusing Fields



$$\text{Local: } \frac{\partial \phi}{\partial t} = -M_\phi \frac{\delta \psi}{\delta \phi} \quad \rightarrow \quad \text{Global: } \frac{\partial \varphi}{\partial t} = -M_\varphi \frac{\delta \Psi}{\delta \varphi} - \tilde{\mu}_\varphi$$


- Specify Landau potential:
 $f(\phi) = -\phi^2(\xi^2 - \phi^2) \rightarrow \psi = f(\phi) + \lambda|\nabla\phi|^2$
- Specify local material parameters:
 $M_\phi, \lambda, \xi, \text{BCs, ICs, ...}$

- Solve system for N trajectories:
 $\mathcal{D}_j = \{\varphi_k, \varphi_{\nabla^2}, \Psi, \frac{\delta \Psi}{\delta \varphi}, \dots\}^{(j)}$

- Choose global model basis:
 $v = \{\varphi_k, \varphi_{\nabla^2}, F, F', \bar{\varphi}_k, \bar{\varphi}_{\nabla^2}, \bar{F}, \bar{F}'\}$
 $M_\varphi = \sum_\alpha \gamma^\alpha v_\alpha, \quad \tilde{\mu}_\varphi = \sum_\beta \gamma^\beta v_\beta$



Evolution of Diffusing Fields



$$\text{Local: } \frac{\partial \phi}{\partial t} = -M_\phi \frac{\delta \psi}{\delta \phi} \quad \rightarrow \quad \text{Global: } \frac{\partial \varphi}{\partial t} = -M_\varphi \frac{\delta \Psi}{\delta \varphi} - \tilde{\mu}_\varphi$$

- Specify Landau potential:

$$f(\phi) = -\phi^2(\xi^2 - \phi^2) \rightarrow \psi = f(\phi) + \lambda |\nabla \phi|^2$$

- Specify local material parameters:

$$M_\phi, \lambda, \xi, \text{BCs, ICs, \dots}$$

- Solve system for N trajectories:

$$\mathcal{D}_j = \{\varphi_k, \varphi_{\nabla^2}, \Psi, \frac{\delta \Psi}{\delta \varphi}, \dots\}^{(j)}$$


- Choose global model basis:

$$v = \{\varphi_k, \varphi_{\nabla^2}, F, F', \bar{\varphi}_k, \bar{\varphi}_{\nabla^2}, \bar{F}, \bar{F}'\}$$

$$M_\varphi = \sum_\alpha \gamma^\alpha v_\alpha, \quad \tilde{\mu}_\varphi = \sum_\beta \gamma^\beta v_\beta$$



Evolution of Diffusing Fields



$$\text{Local: } \frac{\partial \phi}{\partial t} = -M_\phi \frac{\delta \psi}{\delta \phi} \quad \rightarrow \quad \text{Global: } \frac{\partial \varphi}{\partial t} = -M_\varphi \frac{\delta \Psi}{\delta \varphi} - \tilde{\mu}_\varphi$$


- Specify Landau potential:
 $f(\phi) = -\phi^2(\xi^2 - \phi^2) \rightarrow \psi = f(\phi) + \lambda|\nabla\phi|^2$
- Specify local material parameters:
 $M_\phi, \lambda, \xi, \text{BCs, ICs, ...}$

- Solve system for N trajectories:
 $\mathcal{D}_j = \{\varphi_k, \varphi_{\nabla^2}, \Psi, \frac{\delta \Psi}{\delta \varphi}, \dots\}^{(j)}$

- Choose global model basis:
 $v = \{\varphi_k, \varphi_{\nabla^2}, F, F', \bar{\varphi}_k, \bar{\varphi}_{\nabla^2}, \bar{F}, \bar{F}'\}$
 $M_\varphi = \sum_\alpha \gamma^\alpha v_\alpha, \quad \tilde{\mu}_\varphi = \sum_\beta \gamma^\beta v_\beta$



Evolution of Diffusing Fields



$$\text{Local: } \frac{\partial \phi}{\partial t} = -M_\phi \frac{\delta \psi}{\delta \phi} \quad \rightarrow \quad \text{Global: } \frac{\partial \varphi}{\partial t} = -M_\varphi \frac{\delta \Psi}{\delta \varphi} - \tilde{\mu}_\varphi$$

- Specify Landau potential:
 $f(\phi) = -\phi^2(\xi^2 - \phi^2) \rightarrow \psi = f(\phi) + \lambda|\nabla\phi|^2$
- Specify local material parameters:
 $M_\phi, \lambda, \xi, \text{BCs, ICs, ...}$

- Solve system for N trajectories:

$$\mathcal{D}_j = \{\varphi_k, \varphi_{\nabla^2}, \Psi, \frac{\delta \Psi}{\delta \varphi}, \dots\}^{(j)}$$

- Choose global model basis:


$$v = \{\varphi_k, \varphi_{\nabla^2}, F, F', \bar{\varphi}_k, \bar{\varphi}_{\nabla^2}, \bar{F}, \bar{F}'\}$$

$$M_\varphi = \sum_\alpha \gamma^\alpha v_\alpha, \quad \tilde{\mu}_\varphi = \sum_\beta \gamma^\beta v_\beta$$

Figure 1: Field evolution for specific material parameters.



Evolution of Diffusing Fields



$$\text{Local: } \frac{\partial \phi}{\partial t} = -M_\phi \frac{\delta \psi}{\delta \phi} \quad \rightarrow \quad \text{Global: } \frac{\partial \varphi}{\partial t} = -\textcolor{orange}{M}_\varphi \frac{\delta \Psi}{\delta \varphi} - \tilde{\mu}_\varphi$$

- Specify Landau potential:
 $f(\phi) = -\phi^2(\xi^2 - \phi^2) \rightarrow \psi = f(\phi) + \lambda|\nabla\phi|^2$
- Specify local material parameters:
 $M_\phi, \lambda, \xi, \text{BCs, ICs, ...}$
- Solve system for N trajectories:
 $\mathcal{D}_j = \{\varphi_k, \varphi_{\nabla^2}, \Psi, \frac{\delta \Psi}{\delta \varphi}, \dots\}^{(j)}$
- Choose global model basis:
 $v = \{\varphi_k, \varphi_{\nabla^2}, F, F', \bar{\varphi}_k, \bar{\varphi}_{\nabla^2}, \bar{F}, \bar{F}'\}$
 $\textcolor{orange}{M}_\varphi = \sum_\alpha \gamma^\alpha v_\alpha, \quad \tilde{\mu}_\varphi = \sum_\beta \gamma^\beta v_\beta$

Figure 1: Field evolution for specific material parameters.

Stepwise Regression




Figure 2: Loss curves as model becomes more parsimonious.

Stepwise Regression





Figure 3: Fitted dynamics for \mathcal{D}_5 as model becomes more parsimonious.

Stepwise Regression




Figure 3: Fitted dynamics for \mathcal{D}_5 as model becomes more parsimonious.

Stepwise Regression





Figure 2: Loss curves.

$$\frac{\partial \varphi}{\partial t} = -M_\varphi \frac{\delta \Psi}{\delta \varphi} - \tilde{\mu}_\varphi$$




Figure 3: Fitted dynamics for \mathcal{D}_5 .



Stepwise Regression





Figure 2: Loss curves.

$$\textcolor{brown}{M}_\varphi = \gamma^{\frac{\delta\Psi}{\delta\varphi}\varphi_3} \varphi_3 + \gamma^{\frac{\delta\Psi}{\delta\varphi}\varphi} \varphi + \gamma^{\frac{\delta\Psi}{\delta\varphi}\bar{\varphi}_{\nabla^2}} \bar{\varphi}_{\nabla^2} + \gamma^{\frac{\delta\Psi}{\delta\varphi}\varphi_{\nabla^2}} \varphi_{\nabla^2}$$

$$\frac{\partial \varphi}{\partial t} = -\textcolor{brown}{M}_\varphi \frac{\delta\Psi}{\delta\varphi} - \tilde{\mu}_\varphi$$




Figure 3: Fitted dynamics for \mathcal{D}_5 .

$$\tilde{\mu}_\varphi = \gamma^{F'F'} + \gamma^{\varphi_{\nabla^2}} \varphi_{\nabla^2} + \gamma^{\varphi_3} \varphi_3 + \gamma^{\varphi_4} \varphi_4 + \gamma^{\bar{\varphi}_5} \bar{\varphi}_5 + \gamma^{\bar{\varphi}_2} \bar{\varphi}_2 + \gamma^{\bar{\varphi}_4} \bar{\varphi}_4$$



Stepwise Regression





Figure 2: Loss curves.

$$M_\varphi = \gamma^{\frac{\delta\Psi}{\delta\varphi}} \varphi_3 + \gamma^{\frac{\delta\Psi}{\delta\varphi}} \varphi + \gamma^{\frac{\delta\Psi}{\delta\varphi}} \bar{\varphi}_{\nabla^2} + \gamma^{\frac{\delta\Psi}{\delta\varphi}} \varphi_{\nabla^2}$$

$$\tilde{M}_\varphi = \gamma^{\frac{\delta\Psi}{\delta\varphi}} \varphi_3 + \gamma^{\frac{\delta\Psi}{\delta\varphi}} \varphi + \gamma^{\frac{\delta\Psi}{\delta\varphi}} \bar{\varphi}_{\nabla^2}$$

$$\frac{\partial \varphi}{\partial t} = -M_\varphi \frac{\delta \Psi}{\delta \varphi} - \tilde{\mu}_\varphi$$




Figure 3: Fitted dynamics for \mathcal{D}_5 .

$$\tilde{\mu}_\varphi = \gamma^{F'} F' + \gamma^{\varphi_{\nabla^2}} \varphi_{\nabla^2} + \gamma^{\varphi_3} \varphi_3 + \gamma^{\varphi_4} \varphi_4 + \gamma^{\bar{\varphi}_5} \bar{\varphi}_5 + \gamma^{\bar{\varphi}_2} \bar{\varphi}_2 + \gamma^{\bar{\varphi}_4} \bar{\varphi}_4$$

$$\tilde{\mu}_\varphi = \gamma^{F'} F' + \gamma^{\varphi_{\nabla^2}} \varphi_{\nabla^2} + \gamma^{\varphi_3} \varphi_3 + \gamma^{\varphi_4} \varphi_4 + \gamma^{\bar{\varphi}_5} \bar{\varphi}_5 + \gamma^{\bar{\varphi}_2} \bar{\varphi}_2 + \gamma^{\bar{\varphi}_4} \bar{\varphi}_4$$



Stepwise Regression





Figure 2: Loss curves.

$$M_\varphi = \gamma^{\frac{\delta\Psi}{\delta\varphi}}\varphi_3 + \gamma^{\frac{\delta\Psi}{\delta\varphi}}\varphi + \gamma^{\frac{\delta\Psi}{\delta\varphi}}\bar{\varphi}_{\nabla^2} + \gamma^{\frac{\delta\Psi}{\delta\varphi}}\varphi_{\nabla^2}\varphi_{\nabla^2}$$

$$M_\varphi = \gamma^{\frac{\delta\Psi}{\delta\varphi}}\varphi_3 + \gamma^{\frac{\delta\Psi}{\delta\varphi}}\varphi + \gamma^{\frac{\delta\Psi}{\delta\varphi}}\bar{\varphi}_{\nabla^2}\bar{\varphi}_{\nabla^2}$$

$$\textcolor{orange}{M}_\varphi = \gamma^{\frac{\delta\Psi}{\delta\varphi}}\varphi_3 + \gamma^{\frac{\delta\Psi}{\delta\varphi}}\varphi$$

$$\frac{\partial \varphi}{\partial t} = -\textcolor{orange}{M}_\varphi \frac{\delta \Psi}{\delta \varphi} - \tilde{\mu}_\varphi$$




Figure 3: Fitted dynamics for \mathcal{D}_5 .

$$\tilde{\mu}_\varphi = \gamma^{F'} F' + \gamma^{\varphi_{\nabla^2}}\varphi_{\nabla^2} + \gamma^{\varphi_3}\varphi_3 + \gamma^{\varphi_4}\varphi_4 + \gamma^{\bar{\varphi}_5}\bar{\varphi}_5 + \gamma^{\bar{\varphi}_2}\bar{\varphi}_2 + \gamma^{\bar{\varphi}_4}\bar{\varphi}_4$$

$$\bar{\mu}_\varphi = \gamma^{F'} F' + \gamma^{\varphi_{\nabla^2}}\varphi_{\nabla^2} + \gamma^{\varphi_3}\varphi_3 + \gamma^{\varphi_4}\varphi_4 + \gamma^{\bar{\varphi}_5}\bar{\varphi}_5 + \gamma^{\bar{\varphi}_2}\bar{\varphi}_2 + \gamma^{\bar{\varphi}_4}\bar{\varphi}_4$$

$$\tilde{\mu}_\varphi = \gamma^{F'} F' + \gamma^{\varphi_{\nabla^2}}\varphi_{\nabla^2} + \gamma^{\varphi_3}\varphi_3 + \gamma^{\varphi_4}\varphi_4 + \gamma^{\bar{\varphi}_5}\bar{\varphi}_5 + \gamma^{\bar{\varphi}_2}\bar{\varphi}_2 + \gamma^{\bar{\varphi}_4}\bar{\varphi}_4$$



Stepwise Regression





Figure 2: Loss curves.

$$M_\varphi = \gamma^{\frac{\delta\Psi}{\delta\varphi}} \varphi_3 + \gamma^{\frac{\delta\Psi}{\delta\varphi}} \varphi + \gamma^{\frac{\delta\Psi}{\delta\varphi}} \bar{\varphi}_{\nabla^2} + \gamma^{\frac{\delta\Psi}{\delta\varphi}} \varphi_{\nabla^2} \varphi_{\nabla^2}$$

$$M_\varphi = \gamma^{\frac{\delta\Psi}{\delta\varphi}} \varphi_3 + \gamma^{\frac{\delta\Psi}{\delta\varphi}} \varphi + \gamma^{\frac{\delta\Psi}{\delta\varphi}} \bar{\varphi}_{\nabla^2} \bar{\varphi}_{\nabla^2}$$

$$M_\varphi = \gamma^{\frac{\delta\Psi}{\delta\varphi}} \varphi_3 + \gamma^{\frac{\delta\Psi}{\delta\varphi}} \varphi$$

$$\textcolor{orange}{M}_\varphi = \gamma^{\frac{\delta\Psi}{\delta\varphi}} \varphi_3$$

$$\frac{\partial \varphi}{\partial t} = -\textcolor{orange}{M}_\varphi \frac{\delta \Psi}{\delta \varphi} - \tilde{\mu}_\varphi$$




Figure 3: Fitted dynamics for \mathcal{D}_5 .

$$\tilde{\mu}_\varphi = \gamma^{F'} F' + \gamma^{\varphi \nabla^2} \varphi_{\nabla^2} + \gamma^{\varphi_3} \varphi_3 + \gamma^{\varphi_4} \varphi_4 + \gamma^{\bar{\varphi}_5} \bar{\varphi}_5 + \gamma^{\bar{\varphi}_2} \bar{\varphi}_2 + \gamma^{\bar{\varphi}_4} \bar{\varphi}_4$$

$$\bar{\mu}_\varphi = \gamma^{F'} F' + \gamma^{\varphi \nabla^2} \varphi_{\nabla^2} + \gamma^{\varphi_3} \varphi_3 + \gamma^{\varphi_4} \varphi_4 + \gamma^{\bar{\varphi}_5} \bar{\varphi}_5 + \gamma^{\bar{\varphi}_2} \bar{\varphi}_2 + \gamma^{\bar{\varphi}_4} \bar{\varphi}_4$$

$$\tilde{\mu}_\varphi = \gamma^{F'} F' + \gamma^{\varphi \nabla^2} \varphi_{\nabla^2} + \gamma^{\varphi_3} \varphi_3 + \gamma^{\varphi_4} \varphi_4 + \gamma^{\bar{\varphi}_5} \bar{\varphi}_5 + \gamma^{\bar{\varphi}_2} \bar{\varphi}_2 + \gamma^{\bar{\varphi}_4} \bar{\varphi}_4$$

$$\bar{\mu}_\varphi = \gamma^{F'} F' + \gamma^{\varphi \nabla^2} \varphi_{\nabla^2} + \gamma^{\varphi_3} \varphi_3 + \gamma^{\varphi_4} \varphi_4 + \gamma^{\bar{\varphi}_5} \bar{\varphi}_5 + \gamma^{\bar{\varphi}_2} \bar{\varphi}_2 + \gamma^{\bar{\varphi}_4} \bar{\varphi}_4$$



Stepwise Regression





Figure 2: Loss curves.

$$M_\varphi = \gamma^{\frac{\delta\Psi}{\delta\varphi}} \varphi_3 + \gamma^{\frac{\delta\Psi}{\delta\varphi}} \varphi + \gamma^{\frac{\delta\Psi}{\delta\varphi}} \bar{\varphi}_{\nabla^2} \bar{\varphi}_{\nabla^2} + \gamma^{\frac{\delta\Psi}{\delta\varphi}} \varphi_{\nabla^2} \varphi_{\nabla^2}$$

$$M_\varphi = \gamma^{\frac{\delta\Psi}{\delta\varphi}} \varphi_3 + \gamma^{\frac{\delta\Psi}{\delta\varphi}} \varphi + \gamma^{\frac{\delta\Psi}{\delta\varphi}} \bar{\varphi}_{\nabla^2} \bar{\varphi}_{\nabla^2}$$

$$M_\varphi = \gamma^{\frac{\delta\Psi}{\delta\varphi}} \varphi_3 + \gamma^{\frac{\delta\Psi}{\delta\varphi}} \varphi$$

$$M_\varphi = \gamma^{\frac{\delta\Psi}{\delta\varphi}} \varphi_3$$

$$\textcolor{orange}{M}_\varphi = 0$$

$$\frac{\partial \varphi}{\partial t} = -\textcolor{orange}{M}_\varphi \frac{\delta \Psi}{\delta \varphi} - \tilde{\mu}_\varphi$$




Figure 3: Fitted dynamics for \mathcal{D}_5 .

$$\tilde{\mu}_\varphi = \gamma^{F'} F' + \gamma^{\varphi_{\nabla^2}} \varphi_{\nabla^2} + \gamma^{\varphi_3} \varphi_3 + \gamma^{\varphi_4} \varphi_4 + \gamma^{\bar{\varphi}_5} \bar{\varphi}_5 + \gamma^{\bar{\varphi}_2} \bar{\varphi}_2 + \gamma^{\bar{\varphi}_4} \bar{\varphi}_4$$

$$\bar{\mu}_\varphi = \gamma^{F'} F' + \gamma^{\varphi_{\nabla^2}} \varphi_{\nabla^2} + \gamma^{\varphi_3} \varphi_3 + \gamma^{\varphi_4} \varphi_4 + \gamma^{\bar{\varphi}_5} \bar{\varphi}_5 + \gamma^{\bar{\varphi}_2} \bar{\varphi}_2 + \gamma^{\bar{\varphi}_4} \bar{\varphi}_4$$


$$\tilde{\mu}_\varphi = \gamma^{F'} F' + \gamma^{\varphi_{\nabla^2}} \varphi_{\nabla^2} + \gamma^{\varphi_3} \varphi_3 + \gamma^{\varphi_4} \varphi_4 + \gamma^{\bar{\varphi}_5} \bar{\varphi}_5 + \gamma^{\bar{\varphi}_2} \bar{\varphi}_2 + \gamma^{\bar{\varphi}_4} \bar{\varphi}_4$$

$$\bar{\mu}_\varphi = \gamma^{F'} F' + \gamma^{\varphi_{\nabla^2}} \varphi_{\nabla^2} + \gamma^{\varphi_3} \varphi_3 + \gamma^{\varphi_4} \varphi_4 + \gamma^{\bar{\varphi}_5} \bar{\varphi}_5 + \gamma^{\bar{\varphi}_2} \bar{\varphi}_2 + \gamma^{\bar{\varphi}_4} \bar{\varphi}_4$$

$$\tilde{\mu}_\varphi = \gamma^{F'} F' + \gamma^{\varphi_{\nabla^2}} \varphi_{\nabla^2} + \gamma^{\varphi_3} \varphi_3 + \gamma^{\varphi_4} \varphi_4 + \gamma^{\bar{\varphi}_5} \bar{\varphi}_5 + \gamma^{\bar{\varphi}_2} \bar{\varphi}_2 + \gamma^{\bar{\varphi}_4} \bar{\varphi}_4$$




Summary



- Taken's theorem shows existence of *embeddings* of high dimensional dynamical systems
- Constraining non-locality of operators ensures *consistency* of model
- Reduced order models can be computed using physical ansatz and a *basis* of operators
- Interpretation of reduced order model is very *basis-dependent*
- Very *general* framework for many physical systems



Thank you!



Computational Physics Group:

Dr. Krishna Garikipati

Dr. Xiaoxuan Zhang

Dr. Zhenlin Wang

Dr. Greg Teichert

Liz Livingston

Siddhartha Srivastava

Collaborators:

Dr. Brendan Kochunas

Dean Price

Thomas Folk

