

*"One picture is worth more than
Ten thousand words"*

*"Now you can see why I say
A - nePmarion
GRANH THEORY*

Chapter 1

Directed Graphs and Graphs

The first four chapters of this book are concerned with *Graph Theory*. In this chapter we first introduce and illustrate the concepts of *directed graphs* and *graphs*. Then we present some basic material concerned with graphs and related concepts*.

1.1 Directed Graphs

Look at the diagram shown below. This diagram consists of four vertices A, B, C, D and three edges AB, CD, CA with *directions attached to them*, the directions being indicated by arrows. Because of attaching directions to the edges, the edge AB has to be interpreted as an edge *from the vertex A to the vertex B* and it cannot be written as BA . Similarly, the edge CD is from C to D and cannot be written as DC , and the edge CA is from C to A and cannot be written as AC . Thus, here, the edges AB, CD, CA are *directed edges***.




Figure 1.1

The directed edge AB is determined by the vertices A and B in that order and may therefore be represented by the *ordered pair* (A, B) . Similarly, the directed edges CD and CA may be represented by the ordered pairs (C, D) and (C, A) respectively. Thus, the diagram in Figure 1.1 consists of a nonempty set of vertices, namely $\{A, B, C, D\}$, and a set of directed edges represented by *ordered pairs* of vertices taken from this set, namely $\{(A, B), (C, D), (C, A)\}$. Such a diagram is called a diagram of a *directed graph* (or a *diagraph* for brevity).

The formal definition of a directed graph is given below.

*In graph theory, the definitions, notation and terminology are not yet standardized; they generally vary from one author to the other. The reader has to keep this in mind while using different books.

**If one wishes, these edges may also be denoted by $\overrightarrow{AB}, \overrightarrow{CD}, \overrightarrow{CA}$ to emphasize that these are actually directed edges.

Definition of a Directed graph

A directed graph (or a digraph) is a pair (V, E) , where V is a nonempty set and E is a set of ordered pairs of elements taken from the set V .

For a directed graph (V, E) , the elements of V are called *vertices* (points or nodes) and the elements of E are called *directed edges*. The set V is called the *vertex set* and the set E is called the *directed edge set*.

For brevity in terminology, a directed edge is often referred to as just an "edge". Similarly, the directed edge set is referred to as the "edge set".

The directed graph (V, E) is also denoted $D = (V, E)$, or $D = D(V, E)$, or just D when there is no ambiguity.

A geometrical figure that depicts a directed graph is called a *diagram of the directed graph*.

Thus, Figure 1.1 is a diagram of the directed graph for which the vertex set is

$$V = \{A, B, C, D\}$$

and the edge set is

$$E = \{AB, CD, CA\} = \{(A, B), (C, D), (C, A)\}.$$

Figure 1.2 depicts the directed graph for which the vertex set is $V = \{A, B, C, D\}$ and the edge set is $E = \{AB, CD, AC\} = \{(A, B), (C, D), (A, C)\}$.




Figure 1.2

It has to be noted that the directed graph depicted in Figure 1.2 is *not* the same as the directed graph depicted in Figure 1.1. Although both of these two directed graphs have the same vertex set, their directed edge sets are different; whereas the directed graph in Figure 1.1 has CA as a directed edge, the directed graph in Figure 1.2 does not have CA as a directed edge – it has AC as a directed edge, and the directed edges AC and CA are not one and the same. It has to be mentioned that in a diagram of a directed graph the directed edges need not be straight line segments; they can be curved lines (arcs) also. For example, a directed edge AB of a directed graph can be represented by an arbitrary arc drawn from the vertex A to the vertex B as shown in Figure 1.3.




Figure 1.3

Thus, a directed graph can be depicted in a diagram in more than one way. For brevity in terminology, a diagram of a directed graph itself is often referred to as a directed graph.


Vertices of directed graphs are denoted by upper or lower case letters, like $A, B, C, \dots, u, v, \dots$, or, by letters with suffixes appended to them, like v_1, v_2, v_3, \dots

As illustrated in Figure 1.1, every directed edge of a digraph (directed graph) is determined by two vertices of the digraph – a vertex from which it begins and a vertex at which it ends. Thus, if AB is a directed edge of a digraph D , then it is understood that this directed edge begins at the vertex A of D and terminates at the vertex B of D . Here, we say that A is the *initial vertex* and B is the *terminal vertex* of AB . Equivalently, we say that AB is incident *out of* A and incident *into* B .

It should be mentioned that for a directed edge (in a digraph) the initial vertex and the terminal vertex need not be different. A directed edge beginning and ending at the same vertex A is denoted by AA or (A, A) and is called a *directed loop*. The directed edge shown in Figure 1.4 is a directed loop which begins and ends at the vertex A .



(a)



(b)

Figure 1.4

Figure 1.5

A digraph can have more than one directed edge having the same initial vertex and the same terminal vertex. Two directed edges having the same initial vertex and the same terminal vertex are called *parallel directed edges*. Two parallel directed edges are shown in Figure 1.5(a). Two or more directed edges having the same initial vertex and the same terminal vertex are called *multiple directed edges**. Three multiple edges are shown in Figure 1.5(b).

*When a digraph contains multiple edges, the edge set becomes a *multiple set*; by a multiple set we mean a set in which repetition of elements is also taken into account.

A vertex of a digraph which is neither an initial vertex nor a terminal vertex of any directed edge is called an *isolated vertex* of the digraph. A non-isolated vertex happens to be an initial vertex or a terminal vertex for a (some) directed edge. A non-isolated vertex which is not a terminal vertex for any directed edge is called a *source* and a non-isolated vertex which is not an initial vertex for any directed edge is called a *sink*.




Figure 1.6

In the digraph shown in Figure 1.6, the vertex v_4 is an isolated vertex, the vertices v_3 and v_1 are sources and the vertex v_2 is a sink. The vertices v_5 and v_6 do not belong to any of the categories.

In-degree and Out-degree

If v is a vertex of a digraph D , the number of edges for which v is the initial vertex is called the *out-going degree* or the *out-degree* of v and the number of edges for which v is the terminal vertex is called the *incoming degree* or the *in-degree* of v . The out-degree of v is denoted by $d^+(v)$ or $od(v)$ and the in-degree of v is denoted by $d^-(v)$ or $id(v)$.

It follows that (i) $d^+(v) = 0$ if v is a sink, (ii) $d^-(v) = 0$ if v is a source, and (iii) $d^+(v) = d^-(v) = 0$ if v is an isolated vertex.




Figure 1.7

1.1. Directed Graphs

For the digraph shown in Figure 1.7, the out-degrees and the in-degrees of the vertices are as given below:

$$\begin{array}{ll} d^+(v_1) = 2, & d^-(v_1) = 1, \\ d^+(v_2) = 1, & d^-(v_2) = 3, \\ d^+(v_3) = 1, & d^-(v_3) = 2, \\ d^+(v_4) = 0, & d^-(v_4) = 0, \\ d^+(v_5) = 2, & d^-(v_5) = 1, \\ d^+(v_6) = 2, & d^-(v_6) = 1. \end{array}$$

We note that, in the above digraph, there is a directed loop at the vertex v_3 and this loop contributes a count 1 to each of $d^+(v_3)$ and $d^-(v_3)$.

We further observe that the above digraph has 6 vertices and 8 edges and that the sums of the out-degrees and in-degrees of its vertices are

$$\sum_{i=1}^6 d^+(v_i) = 8, \quad \sum_{i=1}^6 d^-(v_i) = 8.$$

This illustrates the following property common to all digraphs. This property is referred to as the *First Theorem of the Digraph Theory*.

Property : In every digraph D , the sum of the out-degrees of all vertices is equal to the sum of the in-degrees of all vertices, each sum being equal to the number of edges in D .

Proof: Suppose D has n vertices v_1, v_2, \dots, v_n and m edges. Let r_1 be the number of edges going out of v_1 , r_2 be the number of edges going out of v_2 , and so on. Then

$$d^+(v_1) = r_1, \quad d^+(v_2) = r_2, \dots, \quad d^+(v_n) = r_n.$$

Since every edge terminates at some vertex and since there are m edges, we should have

$$r_1 + r_2 + \dots + r_n = m.$$

Accordingly,

$$d^+(v_1) + d^+(v_2) + \dots + d^+(v_n) = r_1 + r_2 + \dots + r_n = m.$$

Similarly, if s_1 is the number of edges coming into v_1 , s_2 is the number of edges coming into v_2 , and so on, we get

$$d^-(v_1) + d^-(v_2) + \dots + d^-(v_n) = s_1 + s_2 + \dots + s_n = m.$$

Thus,

$$\sum_{i=1}^n d^+(v_i) = \sum_{i=1}^n d^-(v_i) = m.$$

This completes the proof.

Example 1 Find the in-degrees and the out-degrees of the vertices of the digraph shown in Figure 1.8.




Figure 1.8

► The given digraph has 7 vertices and 12 directed edges. The out-degree of a vertex is got by counting the number of edges that go out of the vertex and the in-degree of a vertex is got by counting the number of edges that end at the vertex. Thus, we obtain the following data:

Vertex:	v_1	v_2	v_3	v_4	v_5	v_6	v_7
Out-degree:	4	2	2	1	3	0	0
In-degree:	0	1	2	2	1	2	4

This table gives the out-degrees and in-degrees of all vertices. We note that v_1 is a source and v_6 and v_7 are sinks.

We also check that

$$\text{sum of out-degrees} = \text{sum of in-degrees} = 12 = \text{No. of edges.}$$

Example 2 Let D be a digraph with an odd number of vertices. If each vertex of D has an odd out-degree, prove that D has an odd number of vertices with odd in-degrees.

► Let v_1, v_2, \dots, v_n be the n vertices of D , where n is odd. Also, let m be the number of edges in D . Then, we have

$$d^+(v_1) + d^+(v_2) + \dots + d^+(v_n) = m \quad (i)$$

$$\text{and} \quad d^-(v_1) + d^-(v_2) + \dots + d^-(v_n) = m \quad (ii)$$

If each vertex has odd out-degree, then the left hand side of (i) is a sum of n odd numbers. Since n is odd, this sum must also be odd. Thus, m is odd.

1.1. Directed Graphs

Let k be the number of vertices with odd in-degrees. Then $n - k$ number of vertices have even in-degrees. Without loss of generality, let us take v_1, v_2, \dots, v_k to be the vertices with odd in-degrees and $v_{k+1}, v_{k+2}, \dots, v_n$ to be the vertices with even in-degrees. Then, expression (ii) may be rewritten as

$$\sum_{i=1}^k d^-(v_i) + \sum_{i=k+1}^n d^-(v_i) = m \quad (\text{iii})$$


Now, the second sum on the left hand side of this expression is even. Also, m is odd. Therefore, the first sum must be odd. That is

$$d^-(v_1) + d^-(v_2) + \dots + d^-(v_k) = \text{odd} \quad (\text{iv})$$


But, each of $d^-(v_1), d^-(v_2), \dots, d^-(v_k)$ is odd. Therefore, the number of terms in the left hand side of (iv) must be odd. That is, k is odd. This proves the required result. ■

Exercises

1. Write down the vertex set and the directed edge set of each of the following digraphs.



(i)



(ii)

Figure 1.9

2. For the digraph shown in Figure 1.6, determine the out-degrees and in-degrees of all the vertices.
3. For the digraphs of Exercise 1 above, determine the out-degrees and in-degrees of all the vertices.
4. Let D be the digraph whose vertex set is $V = \{v_1, v_2, v_3, v_4, v_5\}$ and the directed edge set is

$$E = \{(v_1, v_4), (v_2, v_3), (v_3, v_5), (v_4, v_2), (v_4, v_4), (v_4, v_5), (v_5, v_1)\}.$$

Write down a diagram of D and indicate the out-degrees and in-degrees of all the vertices.

5. Verify the First theorem of Digraph theory for (i) the digraphs shown in Figures 1.6 and 1.9, and
(ii) the digraphs shown below:




Figure 1.10




Figure 1.11

Answers

1. (i) Vertex set is $V = \{A, B, C\}$ and the directed edge set is

$$E = \{(B, A), (C, A), (C, B), (C, B)\}.$$

- (ii) Vertex set is $V = \{v_1, v_2, v_3, v_4\}$ and the directed edge set is

$$E = \{(v_1, v_2), (v_1, v_3), (v_1, v_3), (v_2, v_3), (v_3, v_2), (v_3, v_4), (v_4, v_4)\}.$$

2. $d^-(v_1) = 0, d^-(v_2) = 3, d^-(v_3) = 0, d^-(v_4) = 0, d^-(v_5) = 1, d^-(v_6) = 1.$

$$d^+(v_1) = 2, d^+(v_2) = 0, d^+(v_3) = 1, d^+(v_4) = 0, d^+(v_5) = 1, d^+(v_6) = 1.$$

3. (i) $d^+(A) = 0, d^+(B) = 1, d^+(C) = 4, d^-(A) = 3, d^-(B) = 2, d^-(C) = 0.$

$$(ii) d^+(v_1) = 3, d^+(v_2) = 1, d^+(v_3) = 2, d^+(v_4) = 1, d^-(v_1) = 0, d^-(v_2) = 2, d^-(v_3) = 3, d^-(v_4) = 2.$$

- 4.





Figure 1.12

Vertices	v_1	v_2	v_3	v_4	v_5
d^+	1	1	1	3	1
d^-	1	1	1	2	2

1.2 Graphs

Consider again the diagram in Figure 1.1. Let us redraw this diagram by *dropping* the arrows present in the edges. The rewritten diagram appears as shown below.



511.5
CHA

Figure 1.13

We say that this (rewritten) diagram represents the diagram of an *underlying graph* of the directed graph shown in Figure 1.1. This underlying graph has the same vertices and the same edges as the original directed graph; but its edges are all *undirected*.

We observe that Figure 1.13 depicts a diagram of the underlying graph of the directed graph shown in Figure 1.2 as well.

The underlying graph of a directed graph is an example of what is called an *undirected graph*. Like a directed graph, an undirected graph also consists of a nonempty set of vertices and a set of edges, but the edges are all “undirected”. An edge in a undirected graph is determined and represented by an *unordered pair* of vertices. For example, in the undirected graph shown in Figure 1.13, the edge AB is determined by the vertices A and B and is represented by the unordered pair $\{A, B\} = \{B, A\}$.* For brevity in terminology, an undirected graph is referred to as just a *graph* (– the adjective “undirected” being understood).†

The formal definition of a graph is given below.

Definition of a Graph

A graph is a pair (V, E) , where V is a nonempty set and E is a set of unordered pairs of elements taken from the set V .

For a graph (V, E) , the elements of V are called *vertices* (or *points* or *nodes*) and the elements of E are called *undirected edges* or just *edges*. The set V is called the *vertex set* and the set E is called the *edge set*.

The graph (V, E) is also denoted by $G = (V, E)$ or $G = G(V, E)$ or just G when there is no ambiguity.

*We use the notation (A, B) to denote the *ordered pair* of A and B in that order and the notation $\{A, B\} = \{B, A\}$ for the *unordered pair* of A and B .

†The terms *directed graph* and *digraph* mean one and the same. The terms *undirected graph* and *graph* mean one and the same.

511.5 CHA




67711

A geometrical figure that depicts a graph is called a *diagram of the graph*. Thus, Figure 1.13 is a diagram of the graph for which the vertex set is $V = \{A, B, C, D\}$ and the edge set is $E = \{AB, AC, CD\} = \{\{A, B\}, \{A, C\}, \{C, D\}\}$.

According to the definition of a graph/digraph, the vertex set in a graph/digraph has to be non-empty. Thus, a *graph/digraph must contain at least one vertex*. But, the edge set can be empty. This means that a graph/digraph need not contain any edge.

A graph/digraph containing no edges is called a *null graph*. A null graph with only one vertex is called a *trivial graph*. Figure 1.14 depicts a null graph with three vertices.



In diagrams depicting non-null graphs, an edge is represented by a line. The line may be *straight* or *curved*, *long* or *short*. The important thing is that the diagram should represent the vertices and edges correctly, irrespective of whether the edges are drawn as straight lines or as curves.

The way one draws a diagram of a graph is basically immaterial. There can be more than one diagram for the same graph. For instance, the two diagrams in Figure 1.15(a) and 1.15(b) *look* different, yet they represent the same graph since each conveys the same information, namely that the graph has four vertices A, B, C, D with AB, AC, AD, BC and CD as edges. For brevity in terminology, a diagram of a graph itself is often referred to as a *graph*.




Figure 1.15

The definition of a graph/digraph does not impose any upper limits for the number of vertices and the number of edges. Thus, a graph/digraph can have infinitely many vertices and/or infinitely many edges. A graph/digraph with only a finite number of vertices as well as only a finite number of edges is called a *finite graph/digraph*; otherwise, it is called an *infinite graph/digraph*. In what follows, we will be concerned only with finite graphs/digraphs. Accordingly, by a graph/digraph we will mean only a finite graph/digraph.

In the following discussions (in this Chapter and in Chapters 2 – 4), our major interest will be towards graphs. Only when it is needed, a digraph will be brought in. All the definitions and results that hold for graphs can be extended to digraphs, but with appropriate modifications in terminology.

Order and Size

The number of vertices in a (finite) graph is called the *order* of the graph and the number of edges in it is called its *size*. In other words, for a graph $G = (V, E)$, the cardinality of the set V , namely $|V|$, is called the order of G and the cardinality of the set E , namely $|E|$, is called the size of G . A graph of order n and size m is called a (n, m) graph. Thus, the graph depicted in Figure 1.15(a) is a $(4, 5)$ graph. A null graph with n vertices is a $(n, 0)$ graph.

End vertices, loop, multiple edges

Generally, the vertices of a graph are denoted by A, B, C , etc., or v_1, v_2, v_3 , etc. When it is convenient, we denote the edges of a graph by e_1, e_2, e_3 , and so on. If v_i and v_j denote two vertices of a graph and if e_k denotes an edge joining v_i and v_j , then v_i and v_j are called the *end vertices* (or *end points*) of e_k . This is symbolically written as $e_k = \{v_i, v_j\} = v_i v_j$. For example, in the graph shown in Figure 1.15(a), suppose we denote the edges AB, BC, AC, AD and DC by e_1, e_2, e_3, e_4 and e_5 respectively. Then e_1 joins A and B ; that is, $e_1 = \{A, B\} = AB$ so that A and B are the end vertices of e_1 . Similarly, e_2 joins B and C ; that is, $e_2 = \{B, C\} = BC$, so that B and C are the end vertices of e_2 , and so on.

Now, consider the graph shown in Figure 1.16. We note that this graph consists of four vertices v_1, v_2, v_3, v_4 , and six edges $e_1, e_2, e_3, e_4, e_5, e_6$. Although the edges e_2 and e_3 seem to intersect (cross over) in the figure,* their point of intersection (even when it exists) is *not* a vertex of the graph. We observe that the edges e_1, e_2, e_3 have distinct end vertices, but the edge e_4 has the same vertex v_3 as both of its end vertices; that is, $e_4 = \{v_3, v_3\}$. An edge such as e_4 is called a *loop*. We also observe that both of the edges e_5 and e_6 have the same end vertices v_1, v_4 ; that is, $e_5 = \{v_1, v_4\}$ and $e_6 = \{v_1, v_4\}$. Edges such as these are called *parallel edges*. If in a graph there are two or more edges with the same end vertices, the edges are called *multiple edges*.

*If the vertices are in different planes, e_2 and e_3 may not intersect.




Figure 1.16

Simple graph, Multigraph, General graph

A graph which does not contain loops and multiple edges is called a *simple graph*. A graph which does not contain a loop is called a *loop-free* graph. A graph which contains multiple edges but no loops is called a *multigraph*[†]. A graph which contains multiple edges and loops (or both) is called a *general graph*.

Figure 1.13 represents a simple graph and Figure 1.16 represents a general graph. Figure 1.17 represents a multigraph.




Figure 1.17

In our discussions here, by a graph we will mean a general graph, unless otherwise mentioned.

In many situations, the names assigned to vertices are inconsequential. In such situations we do not assign any names to the vertices. Such graphs are called *unlabeled graphs*. On the other hand, if names are assigned to vertices of a graph (for some specific purpose or upon certain grounds), the graph is called a *labeled graph*. For example, the graphs shown in Figures 1.13 to 1.16 are labeled graphs. The graph shown in Figure 1.17 is unlabeled.

[†]In a multigraph, the edge set is a multiset.

Incidence

When a vertex v of a graph G is an end vertex of an edge e of the graph G , we say that the edge e is *incident on* (or *to*) the vertex v . Since every edge has two end vertices, every edge is incident on two vertices, one at each end. The two end vertices are coincident if the edge is a loop.

When an edge e is incident on a vertex v , we also say that v is *incident with* e . Note that whereas an edge is incident only on two vertices (namely its end vertices), a vertex may be incident with any number of edges.

Two *non-parallel* edges are said to be *adjacent edges* if they are incident on a common vertex (that is, if they have a vertex in common). Two vertices are said to be *adjacent vertices* (or *neighbours*) if there is an edge joining them.

In the graph shown in Figure 1.18, A and B are adjacent vertices and e_1 and e_2 are adjacent edges. But, A and C are not adjacent vertices, and e_1 and e_3 are not adjacent edges.




Figure 1.18

Complete graph

A *simple graph* of order ≥ 2 in which there is an edge between *every pair* of vertices is called a *complete graph* (or a *full graph*).

In other words, a *complete graph* is a *simple graph* of order ≥ 2 in which *every pair of distinct vertices are adjacent*.

A complete graph with n (≥ 2) vertices is denoted by K_n .

Complete graphs with two, three, four and five vertices are shown in Figures 1.19(a) to 1.19(d) respectively. Of these complete graphs, the complete graph with five vertices, namely K_5 (shown in Figure 1.19(d)), is of great importance. This graph is called the *Kuratowski's first graph*.*

*Named after the Polish mathematician Kasimir Kuratowski.




Figure 1.19

Bipartite graph

Suppose a *simple graph* G is such that its vertex set V is the union of two of its mutually disjoint nonempty subsets V_1 and V_2 which are such that each edge in G joins a vertex in V_1 and a vertex in V_2 . Then G is called a **bipartite graph**. If E is the edge set of this graph, the graph is denoted by $G = (V_1, V_2; E)$, or $G = G(V_1, V_2; E)$. The sets V_1 and V_2 are called **bipartites** (or *partitions*) of the vertex set V .

For example, consider the graph G shown in Figure 1.20 for which the vertex set is $V = \{A, B, C, P, Q, R, S\}$ and the edge set is $E = \{AP, AQ, AR, BR, CQ, CS\}$. Note that the set V is the union of two of its subsets $V_1 = \{A, B, C\}$ and $V_2 = \{P, Q, R, S\}$ which are such that (i) V_1 and V_2 are disjoint, (ii) every edge in G joins a vertex in V_1 and a vertex in V_2 , (iii) G contains no edge that joins two vertices both of which are in V_1 or V_2 . This graph is a bipartite graph with $V_1 = \{A, B, C\}$ and $V_2 = \{P, Q, R, S\}$ as bipartites.




Figure 1.20

Complete Bipartite graph

A bipartite graph $G = (V_1, V_2; E)$ is called a **complete bipartite graph** if there is an edge between every vertex in V_1 and every vertex in V_2 .

The bipartite graph shown in Figure 1.20 is *not* a complete bipartite graph. Observe, for example, that the graph does not contain an edge joining A and S .

A complete bipartite graph $G = (V_1, V_2; E)$ in which the bipartites V_1 and V_2 contain r and s vertices respectively, with $r \leq s$, is denoted by $K_{r,s}$. In this graph, each of r vertices in V_1 is joined to each of s vertices in V_2 . Thus, $K_{r,s}$ has $r + s$ vertices and rs edges; that is $K_{r,s}$ is of order $r + s$ and of size rs ; it is therefore a $(r + s, rs)$ graph.

Figures 1.21(a) to 1.21(d) depict some complete bipartite graphs. Observe that in Figure 1.21(a), the bipartites are $V_1 = \{A\}$ and $V_2 = \{P, Q, R\}$; the vertex A is joined to each of the vertices P, Q, R by an edge. In Figure 1.21(b), the bipartites are $V_1 = \{A\}$ and $V_2 = \{M, N, P, Q, R\}$; the vertex A is joined to each of the vertices M, N, P, Q, R by an edge.




Figure 1.21

In Figure 1.21(c), the bipartites are $V_1 = \{A, B\}$ and $V_2 = \{P, Q, R\}$; each of the vertices A and B is joined to each of the vertices P, Q, R by an edge. In Figure 1.21(d), the bipartites are $V_1 = \{A, B, C\}$ and $V_2 = \{P, Q, R\}$; each of the vertices A, B, C is joined to each of the vertices P, Q, R .

Of these complete bipartite graphs, the graph $K_{3,3}$ shown in Figure 1.21(d) is of great importance. This is known as the *Kuratowski's second graph*.

It is to be noted that a bipartite graph G is not a complete graph even if G is a complete bipartite graph. Because, in such a graph, there exists no edge between two vertices if they belong to the same bipartite.

Example 1 Draw a diagram of the graph $G = (V, E)$ in each of the following cases:

- (i) $V = \{A, B, C, D\}, E = \{AB, AC, AD, CD\}$
- (ii) $V = \{v_1, v_2, v_3, v_4, v_5\}, E = \{v_1v_2, v_1v_3, v_2v_3, v_4v_5\}$
- (iii) $V = \{P, Q, R, S, T\}, E = \{PS, QR, QS\}$
- (iv) $V = \{v_1, v_2, v_3, v_4, v_5, v_6\}, E = \{v_1v_4, v_1v_6, v_4v_6, v_3v_2, v_3v_5, v_2v_5\}$

- The required diagrams are shown below:




Figure 1.22

Example 2 Let P, Q, R, S, T represent five cricket teams. Suppose that the teams P, Q, R have played one game with each other, and the teams P, S, T have played one game with each other. Represent this situation in a graph.

Hence determine (i) the teams that have not played with each other, and (ii) the number of games played by each team.

► Let the teams be represented by vertices and an edge represent the playing. Then the graph representing the given situation is as shown in Figure 1.23:




Figure 1.23

We observe that there is no edge between Q and S , between Q and T , between R and S , and between R and T . Therefore, the teams Q and S , Q and T , R and S , and R and T have not played with each other.

From the graph, we note that two edges are incident on each of the vertices Q, R, S, T and four edges are incident on P . Thus, the teams Q, R, S, T have played two games each and the team P has played four games.

Example 3 Which of the following is a complete graph?




Figure 1.24

- The first of the graphs is *not* complete. It is not simple on the one hand and there is no edge between A and C on the other hand. The second of the graphs is complete. It is a simple graph and there is an edge between every pair of vertices.

Example 4 If $G = G(V, E)$ is a simple graph, prove that $2|E| \leq |V|^2 - |V|$.

- Each edge of a graph is determined by a pair of vertices. In a simple graph there occur no multiple edges. As such, in a simple graph, the number of edges *cannot exceed* the number of pairs of vertices. The number of pairs of vertices that can be chosen from n vertices is (from the theory of combinations)^f

$${}^nC_2 = \frac{n!}{(n-2)!2!} = \frac{1}{2}n(n-1).$$

Thus, for a simple graph with $n (\geq 2)$ vertices, the number of edges cannot exceed $\frac{1}{2}n(n-1)$. Accordingly, if a simple graph $G = G(V, E)$ has n vertices and m edges, then $m \leq \frac{1}{2}n(n-1)$, or, $2m \leq n^2 - n$; that is : $2|E| = |V|^2 - |V|$.

Remark: If a graph is not simple, it can have any number of edges (because, in such a graph multiple edges and loops are allowed).

Example 5 Show that a complete graph with n vertices, namely K_n , has $\frac{1}{2}n(n-1)$ edges.

- In a complete graph, there exists exactly one edge between every pair of vertices. As such, the number of edges in a complete graph is equal to the number of pairs of vertices. If the number of vertices is n , then the number of pairs of vertices is

$${}^nC_2 = \frac{n!}{(n-2)!2!} = \frac{1}{2}n(n-1)$$

^fSee Chapter 5. The symbol nC_r is also denoted by ${}_nC_r$, $C(n, r)$ and $\binom{n}{r}$.

Thus, the number of edges in a complete graph with n vertices is $\frac{1}{2}n(n - 1)$.

Note: For another proof of this result, see Example 13, Section 1.2.1.

Example 6 Show that a simple graph of order $n = 4$ and size $m = 7$ and a complete graph of order $n = 4$ and size $m = 5$ do not exist.


► For $n = 4$, we have

$$\frac{1}{2}n(n - 1) = \frac{1}{2} \times 4 \times 3 = 6.$$


Since $m = 7$ exceeds this number, a simple graph of order $n = 4$ and size $m = 7$ does not exist.

Similarly, since $m = 5$ is not equal to $\frac{1}{2}n(n - 1) = 6$, a complete graph of order 4 and size $m = 5$ does not exist.

Example 7 Which of the following is a bipartite graph?



(a)



(b)

Figure 1.25

► The first of the graphs is a bipartite graph, with $V_1 = \{P, S\}$ and $V_2 = \{Q, R\}$ as the bipartites. The second graph is *not* a bipartite graph. (Why?)

Example 8 (a) How many vertices and how many edges are there in the complete bipartite graphs $K_{4,7}$ and $K_{7,11}$?

(b) If the graph $K_{r,12}$ has 72 edges, what is r ?

► Recall that the complete bipartite graph $K_{r,s}$ has $r + s$ vertices and rs edges. Accordingly:

(a) The graph $K_{4,7}$ has $4 + 7 = 11$ vertices and $4 \times 7 = 28$ edges, and the graph $K_{7,11}$ has 18 vertices and 77 edges.

(b) If the graph $K_{r,12}$ has 72 edges, we have $12r = 72$ so that $r = 6$.

Example 9 Let $G = (V, E)$ be a simple graph of order $|V| = n$ and size $|E| = m$. If G is a bipartite graph, prove that $4m \leq n^2$.

► Let V_1 and V_2 be the bipartites of G , with $|V_1| = r$ and $|V_2| = s$. Since $|V| = n$, we should have $r + s = n$, so that $r = n - s$ and $s = n - r$.

The graph G has the maximum number of edges when each of the r vertices in V_1 is joined by an edge to each of the s vertices in V_2 , and this maximum is equal to rs . This means that $|E| = m \leq rs$.

When n is even, we may express rs as

$$rs = r(n - r) = rn - r^2 = \left(\frac{n}{2}\right)^2 - \left(r - \frac{n}{2}\right)^2$$

Evidently, rs is maximum when $r = (n/2)$, the maximum being $(n/2)^2$. Thus, when n is even, we have

$$m \leq rs \leq \left(\frac{n}{2}\right)^2.$$

When n is odd, we may express rs as

$$rs = r(n - r) = \frac{(n-1)(n+1)}{4} - \left(r - \frac{n-1}{2}\right)\left(r - \frac{n+1}{2}\right)$$

Evidently, rs is maximum when $r = (n-1)/2$ or $r = (n+1)/2$, the maximum being $(n-1)(n+1)/4$. Thus, when n is odd, we have

$$m \leq rs \leq \frac{(n-1)(n+1)}{4} = \frac{n^2-1}{4} \leq \frac{n^2}{4} = \left(\frac{n}{2}\right)^2$$

Accordingly, in both of the possible cases (n is even or odd), we have

$$m \leq \left(\frac{n}{2}\right)^2, \quad \text{or} \quad 4m \leq n^2.$$

Example 10 Show that a simple graph of order $n = 4$ and size $m = 5$ cannot be a bipartite graph.

► Here, $4m = 20$ and $n^2 = 16$, so that $4m > n^2$. Therefore, the given simple graph cannot be a bipartite graph.

Exercises

1. Indicate the order and size of each of the graphs shown below.




Figure 1.26

2. Identify the adjacent vertices in the graphs of the preceding exercise.
3. Identify the adjacent vertices and adjacent edges in the graph shown in Figure 1.16.
4. Which of the following graphs is a simple graph? a multigraph? a general graph?




Figure 1.27

5. Which of the following are complete graphs?





Figure 1.28

6. Identify the graph shown below:




Figure 1.29

7. Verify that the following are bipartite graphs. What are their bipartites?




(a)




(b)

Figure 1.30


8. Which of the graphs shown below are bipartite graphs?



(a)



(b)



(c)

Figure 1.31

9. Company X has offices in cities B , D and K ; company Y in cities B and M ; company Z in cities C and M . Represent this situation by a bipartite graph. Is this a complete bipartite graph?

10. State whether the following graphs can exist or cannot exist.

- (1) Simple graph of order 3 and size 2.
- (2) Simple graph of order 5 and size 12.
- (3) Complete graph of order 5 and size 10.
- (4) Bipartite graph of order 4 and size 3.
- (5) Bipartite graph of order 3 and size 4.
- (6) Complete bipartite graph of order 4 and size 4.

Answers

1. (i) There are 6 vertices and 5 edges; therefore, the order is 6 and size is 5.
 (ii) There are 5 vertices and 7 edges; therefore, the order is 5 and size is 7.
2. (i) $\{A, P\}, \{P, B\}, \{P, C\}, \{C, Q\}, \{C, R\}$ are pairs of adjacent vertices.
 (ii) $\{P, Q\}, \{P, S\}, \{Q, S\}, \{Q, R\}, \{Q, T\}, \{R, T\}, \{S, T\}$ are pairs of adjacent vertices.
3. Adjacent vertices: v_1 and v_2 , v_1 and v_3 , v_1 and v_4 , v_2 and v_4 .
 Adjacent edges: e_1 and e_2 , e_1 and e_3 , e_1 and e_5 , e_1 and e_6 ,
 e_2 and e_4 , e_2 and e_5 , e_2 and e_6 , e_3 and e_5 ,
 e_3 and e_6 .
4. (i) general graph (ii) simple graph, (iii) multigraph.
5. None is a complete graph.
6. The complete graph K_6 .
7. (a) $V_1 = \{A, C, Q\}$, $V_2 = \{B, P, R\}$ (b) $V_1 = \{A, B, C, D\}$, $V_2 = \{P, Q, R, S\}$
8. (a) and (c) are bipartite graphs with $\{A, B, C\}$ and $\{P, Q, R\}$ as bipartites. (b) is not a bipartite graph.
- 9.




Figure 1.32

This is a not a complete bipartite graph.

10. (1), (3), (4), (6) : can exist

(2), (5) : cannot exist

1.2.1 Vertex degree and Handshaking property

Let $G = (V, E)$ be a graph and v be a vertex of G . Then, the number of edges of G that are incident on v (that is, the number of edges that join v to other vertices of G) *with the loops counted twice* is called the *degree* of the vertex v and is denoted by $\deg(v)$, or $d(v)$.*

The degrees of all vertices of a graph arranged in non-decreasing order is called the *degree sequence* of the graph. Also, the *minimum* of the degrees of vertices of a graph is called the *degree of the graph*.

For example, the degrees of vertices of the graph shown in Figure 1.33 are as given below.

$$d(v_1) = 3, \quad d(v_2) = 4, \quad d(v_3) = 4, \quad d(v_4) = 3.$$

Therefore, the degree sequence of this graph is 3, 3, 4, 4, and the degree of the graph is 3.

Note that the loop at v_3 is counted twice for determining the degree of v_3 .




Figure 1.33

Isolated Vertex, Pendant Vertex

A vertex in a graph which is not an end vertex of any edge of the graph is called an *isolated vertex*. Obviously, a vertex is an isolated vertex if and only if its degree is zero.

A vertex of degree 1 is called a *pendant vertex*. An edge incident on a pendant vertex is called a *pendant edge*.

In the graph shown in Figure 1.34, the vertices v_4 and v_6 are isolated vertices, v_5 and v_7 are pendant vertices and the edges e_4 and e_5 are pendant edges.

*Since, in a graph, all edges are undirected, no distinction is made between the in-degree and the out-degree. If G is an underlying graph of a directed graph D , then G and D have the same vertex set V and for any $v \in V$, we have $d(v) = id(v) + od(v)$.




Figure 1.34

As mentioned before, a null graph contains no edges. It therefore follows that *in a null graph every vertex is an isolated vertex*.

Regular graph

A graph in which all the vertices are of the *same degree k* is called a **regular graph of degree k**, or a **k-regular graph**.

In particular, a 3-regular graph is called a **cubic graph**.

The graphs shown in Figures 1.35 and 1.36 are 2-regular and 4-regular graphs respectively.



Figure 1.35




Figure 1.36

The graph shown in Figure 1.37 is a 3-regular graph (cubic graph). This particular cubic graph, which contains 10 vertices and 15 edges, is called the **Petersen graph**.




Figure 1.37

The graph shown in Figure 1.38 is a cubic graph with $8 = 2^3$ vertices. This particular graph is called the *three-dimensional hypercube* and is denoted by Q_3 .




Figure 1.38

In general, for any positive integer k , a loop-free k -regular graph with 2^k vertices is called the *k -dimensional hypercube* (or *k -cube*) and is denoted by Q_k .* (Note that the graph shown in Figure 1.35 is Q_2).

Handshaking Property

Let us refer back to the graph shown in Figure 1.33. As noted earlier, we have, in this graph,

$$\deg(v_1) = 3, \quad \deg(v_2) = 4, \quad \deg(v_3) = 4, \quad \deg(v_4) = 3.$$

Also, the graph has 7 edges. We observe that

$$\deg(v_1) + \deg(v_2) + \deg(v_3) + \deg(v_4) = 14 = 2 \times 7.$$

This observation illustrates the following important property common to all (finite) graphs.

Property : *The sum of the degrees of all the vertices in a graph is an even number; and this number is equal to twice the number of edges in the graph.*

In an alternative form, this property reads as follows:

For a graph $G = (V, E)$,

$$\sum_{v \in V} \deg(v) = 2|E|.$$

This property is obvious from the fact that while counting the degrees of vertices, each edge is counted twice (once at each end).

The aforesaid property is popularly called the *handshaking property*.† Because, it essentially states that if several people shake hands, then the total number of hands shaken must be even, because just two hands are involved in each handshake.

*The graph Q_k arises in the study of *Computer Architecture*.

†Handshaking property was the first result noted in Graph Theory. This property is also known as the *First Theorem of Graph Theory*.

The following theorem is a direct consequence of the handshaking property.

Theorem : In every graph, the number of vertices of odd degrees is even.

Proof: Consider a graph with n vertices. Suppose k of these vertices are of odd degree so that the remaining $n - k$ vertices are of even degree. Denote the vertices with odd degree by $v_1, v_2, v_3, \dots, v_k$ and the vertices with even degree by $v_{k+1}, v_{k+2}, \dots, v_n$. Then the sum of the degrees of the vertices is

$$\sum_{i=1}^n \deg(v_i) = \sum_{i=1}^k \deg(v_i) + \sum_{i=k+1}^n \deg(v_i) \quad (1)$$

In view of the hand shaking property, the sum on the left hand side of the above expression is equal to twice the number of edges in the graph. As such, this sum is even. Further, the second sum in the right hand side is the sum of the degrees of vertices with even degrees. As such, this sum is also even. Therefore, the first sum in the right hand side must also be even, that is,

$$\deg(v_1) + \deg(v_2) + \dots + \deg(v_k) = \text{even.} \quad (2)$$

But, each of $\deg(v_1), \deg(v_2), \dots, \deg(v_k)$ is odd. Therefore, the number of terms in the left hand side of (2) must be even, that is, k is even.

This completes the proof of the theorem.

Note: According to the above theorem, in any graph, there is an even number of vertices of odd degrees. But, it is *not true* in general that a graph must have an odd number of vertices of even degrees. Observe that the graph shown in Figure 1.33 has an even number of vertices of even degrees.

Example 1 For the graph shown in Figure 1.39, indicate the degree of each vertex and verify the handshaking property:




Figure 1.39

- By examining the graph, we find that the degrees of its vertices are as given below:

$$\begin{aligned}\deg(a) &= 3, \quad \deg(b) = 2, \quad \deg(c) = 4, \\ \deg(d) &= 2, \quad \deg(e) = 0, \quad \deg(f) = 2, \\ \deg(g) &= 2, \quad \deg(h) = 1.\end{aligned}$$

We note that e is an isolated vertex and h is a pendant vertex.

Further, we observe that the sum of the degrees of vertices is equal to 16. Also, the graph has 8 edges. Thus, the sum of the degrees of vertices is equal to twice the number of edges. This verifies the handshaking property for the given graph. ■

Example 2 Can there be a graph consisting of the vertices A, B, C, D with $\deg(A) = 2$, $\deg(B) = 3$, $\deg(C) = 2$, $\deg(D) = 2$?

- In every graph, the sum of the degrees of the vertices has to be an even number. Here, this sum is 9 which is not even. Therefore, there does not exist a graph of the given kind. ■

Example 3 Can there be a graph with 12 vertices such that two of the vertices have degree 3 each and the remaining 10 vertices have degree 4 each?

- Here, the sum of the degrees of vertices is $(3 \times 2) + (4 \times 10) = 46$. Therefore, if $m = 23$, we have $2m = 46$ and the handshaking property holds. Hence there can be a graph of the desired type (whose size is 23). ■

Example 4 For a graph $G = (V, E)$, what is the largest possible value for $|V|$ if $|E| = 19$ and $\deg(v) \geq 4$ for all $v \in V$?

- Since all vertices are of degree greater than or equal to 4, the sum of the degrees of vertices is greater than or equal to $4n$, where $n = |V|$ is the number of vertices. This sum is equal to twice $|E|$, by handshaking property. Therefore, $2|E| \geq 4n$, that is, $2 \times 19 \geq 4n$, or $n \leq (38/4) < 10$. Thus, the largest possible value of $|V|$ is 9. ■

Example 5 Show that the hypercube Q_3 is a bipartite graph which is not a complete bipartite graph.

- The hypercube Q_3 is shown in Figure 1.40, with the vertices labeled as A, B, C, D, P, Q, R, S so that, for this graph, the vertex set is $V = \{A, B, C, D, P, Q, R, S\}$.

Let

$$V_1 = \{A, C, Q, S\} \quad \text{and} \quad V_2 = \{B, D, P, R\}$$

Then, we note that $V_1 \cup V_2 = V$, $V_1 \cap V_2 = \emptyset$ and that every edge of the graph has one end vertex in the set V_1 and the other end vertex in the set V_2 ; further, no edge of the graph has both of its end vertices in V_1 or V_2 . Hence, this graph is a bipartite graph.




Figure 1.40

We observe that this graph is *not* a complete bipartite graph. Because, there is no edge between every vertex in V_1 and every vertex in V_2 ; for example, there is no edge joining A and R .

Example 6 Prove that the k -dimensional hypercube Q_k has $k2^{k-1}$ edges.

Determine the number of edges in Q_8 .

- In the hypercube Q_k , the number of vertices is 2^k and each vertex is of degree k . Therefore, the sum of degrees of vertices of Q_k is $k \times 2^k$. By handshaking property, we should have $k \times 2^k = 2|E|$, where $|E|$ is the size of Q_k . Thus, $|E| = \frac{1}{2}(k \times 2^k) = k \times 2^{k-1}$. This means that Q_k has $k2^{k-1}$ edges.

It follows that the number of edges in Q_8 is $8 \times 2^7 = 1024$.

Example 7 (a) What is the dimension of the hypercube with 524288 edges?

(b) How many vertices are there in a hypercube with 4980736 edges?

- For the k -dimensional hypercube Q_k , the number of vertices is 2^k and the number of edges is $k2^{k-1}$.

(a) If Q_k has 524288 edges, we have $k2^{k-1} = 524288$. We check that

$$524288 = 2^{19} = 2^4 \times 2^{15} = 16 \times 2^{15}.$$

Accordingly, $k2^{k-1} = 524288$ holds if $k = 16$. Thus, the dimension of the hypercube with 524288 edges is $k = 16$.

(b) We check that $4980736 = 19 \times 2^{18}$, which indicates that Q_k has 4980736 edges when $k = 19$. The number of vertices in this hypercube is $2^k = 2^{19} = 524288$.

Example 8 Determine the order $|V|$ of the graph $G = (V, E)$ in the following cases :

- (1) G is a cubic graph with 9 edges.
- (2) G is regular with 15 edges.
- (3) G has 10 edges with 2 vertices of degree 4 and all other vertices of degree 3.

- (1) Suppose the order of G is n . Since G is a cubic graph, all vertices of G have degree 3, and therefore the sum of the degrees of vertices is $3n$. Since G has 9 edges, we should have $3n = 2 \times 9$ (by the handshaking property) so that $n = 6$. Thus, the order of G is 6.
- (2) Since G is regular, all vertices of G must be of the same degree, say k . If G is of order n , then the sum of the degrees of vertices is kn . Since G has 15 edges, we should have $kn = 2 \times 15$ so that $k = 30/n$. Since k has to be a positive integer, it follows that n must be a divisor of 30. Thus, the possible orders of G are 1, 2, 3, 5, 6, 10, 15 and 30.
- (3) Suppose the order of G is n . Since two vertices of G are of degree 4 and all others are of degree 3, the sum of the degrees of vertices of G is $2 \times 4 + (n - 2) \times 3$. Since G has 10 edges, we should have $2 \times 4 + (n - 2) \times 3 = 2 \times 10$. This gives $n = 6$. Thus, the order of G is 6. ■

Example 9 Let G be a graph of order 9 such that each vertex has degree 5 or 6. Prove that at least 5 vertices have degree 6 or atleast 6 vertices have degree 5.

- Let p be the number of vertices of G which have degree 5. Then the number of vertices of G which have degree 6 is $9 - p = q$ (say). Evidently, $0 \leq p \leq 9$, $0 \leq q \leq 9$.

We note that the sum of the degrees of vertices of G is

$$(5 \times p) + (6 \times q) = 5p + 6(9 - p) = 54 - p.$$

Since this sum has to be an even number, p cannot be odd. Thus, $p = 0, 2, 4, 6$, or 8. Consequently, the following possible cases arise :

$$\begin{aligned} p &= 0, & q &= 9 \\ p &= 2, & q &= 7 \\ p &= 4, & q &= 5 \\ p &= 6, & q &= 3 \\ p &= 8, & q &= 1 \end{aligned}$$

We observe that in all the above possible cases either $q \geq 5$ or $p \geq 6$. This means that atleast 5 vertices have degree 6 or atleast 6 vertices have degree 5. ■

Example 10 Show that there is no graph with 28 edges and 12 vertices in the following cases:

(i) The degree of a vertex is either 3 or 4.

(ii) The degree of a vertex is either 3 or 6.

► Suppose there is a graph with 28 edges and 12 vertices, of which k vertices are of degree 3 (each). Then:

(i) If all of the remaining $(12 - k)$ vertices have degree 4, then we should have (by the handshaking property) $3k + 4(12 - k) = 2 \times 28 = 56$, or $k = -8$ which is not possible (because k has to be nonnegative).

(ii) If all of the remaining $(12 - k)$ vertices have degree 6, then we should have $3k + 6(12 - k) = 56$, or $k = 16/3$. This is not possible (because k has to be a nonnegative integer).

Hence, in both of the two given cases, the graph of the desired type cannot exist. ■

Example 11 (a) If a graph with n vertices and m edges is k -regular, show that $m = kn/2$.

(b) Does there exist a cubic graph with 11 vertices?

(c) Does there exist a 4-regular graph with (i) 15 edges? (ii) 10 edges?

►(a) If a graph G is k -regular, then the degree of every vertex is k . Therefore, if G has n vertices, then the sum of the degrees of vertices is nk . By handshaking property, this sum must be equal to $2m$ if G has m edges. Thus, $nk = 2m$, or $m = nk/2$.

(b) If there is a cubic (3-regular) graph with $n = 11$ vertices, the number of edges it should have is $m = \frac{1}{2}(3n) = \frac{1}{2}(11 \times 3)$. This is not possible, because $(33/2)$ is not a whole number. Thus, the graph of the desired type does not exist.

(c) (i) If there is a 4-regular graph with $m = 15$ edges, the number of vertices n it should have is given by $4n = 2m = 30$. This is not possible, because $30/4$ is not a whole number. Thus, the graph of the desired type does not exist.

(ii) If there is a 4-regular graph with $m = 10$ edges, the number of vertices n it should have is given by $4n = 2m = 20$. This is possible (with $n = 5$). Thus, such a graph does exist. We note that this graph is the complete graph K_5 . See Figure 1.19(d) which is reproduced below,




Figure 1.41

Example 12 (a) If k is odd, show that the number of vertices in a k -regular graph is even.

(b) Show that it is not possible to have a set of seven persons such that each person in the set knows exactly three other persons in the set.

►(a) In a k -regular graph, the degree of each vertex is k . Therefore, if such a graph has n vertices, then the sum of the degrees is nk , and this has to be an even number (by handshaking property). If k is odd, this is possible only if n is even.

(b) Let G be a graph with seven vertices, each vertex representing a person in the given set and each edge representing an acquaintance. If each person in the set knows exactly three other persons in the set, then there will be exactly three edges incident on each vertex, and the graph G will be 3-regular. This is not possible, because G has an odd number of vertices. Hence, the graph G of the desired type does not exist. This means that it is not possible to have a set of seven persons such that each person in the set knows exactly three other persons in the set.

Example 13 (a) Show that in a complete graph of n vertices (namely K_n) the degree of every vertex is $(n - 1)$ and that the total number of edges is $\frac{1}{2}n(n - 1)$.

(b) If K_n has m edges, show that $n(n + 1) = 2(n + m)$.

►(a) Recall that a complete graph is a simple graph in which every vertex is joined with every other vertex through exactly one edge. Therefore, if there are n vertices, each vertex is joined to $(n - 1)$ vertices through exactly one edge. Thus, there occur $n - 1$ edges at every vertex. This means that the degree of every vertex is $n - 1$.

Consequently, the sum of the degrees of vertices is $n(n - 1)$. By handshaking property, this sum must be equal to $2m$, where m is the number of edges. Thus, $n(n - 1) = 2m$, or $m = \frac{1}{2}n(n - 1)$. Thus, K_n has $\frac{1}{2}n(n - 1)$ edges.

- (b) If K_n has m edges, then we have, from what has been just proved, $m = \frac{1}{2}n(n - 1)$. This gives $n + m = n + \frac{1}{2}n(n - 1)$, or $2(n + m) = 2n + n^2 - n = n(n + 1)$.

Remark: Since the degree of every vertex in K_n is $n - 1$, it follows that K_n is a $(n - 1)$ -regular graph.

Example 14 Show that if a bipartite graph $G(V_1, V_2, E)$ is regular, then $|V_1| = |V_2|$.

- Since G is simple and every edge of G has one end in V_1 and the other end in V_2 ,

$$\sum_{v \in V_1} \deg(v) = \sum_{v \in V_2} \deg(v) \quad (\text{i})$$

Since G is regular, all vertices of G have the same degree, say $k > 0$. Consequently, if V_1 has r elements and V_2 has s elements,

$$\sum_{v \in V_1} \deg(v) = kr \quad \text{and} \quad \sum_{v \in V_2} \deg(v) = ks \quad (\text{ii})$$

Using this result in (i), we get $kr = ks$, so that $r = s$. Thus, $|V_1| = |V_2|$.

Example 15 Show that every simple graph of order ≥ 2 must have at least two vertices of the same degree.

- Let G be a simple graph with n vertices. Suppose all the vertices have different degrees.
- Then, since every vertex must have a degree and since all such degrees must be between 0 and $n - 1$, the degrees must be $0, 1, 2, 3, \dots, n - 1$. Let A be the vertex whose degree is 0 and B be the vertex whose degree is $n - 1$. Then $n - 1$ edges are incident on B . This means that B is joined to all other vertices by an edge and in particular to A also. Hence the degree of A is not zero. This is a contradiction. Hence all vertices of G cannot have different degrees; at least two of them must have the same degree.

Example 16 Is there a simple graph with 1, 1, 3, 3, 3, 4, 6, 7 as the degrees of its vertices?

- Assume that there is such a graph. Since the degrees of vertices are 8 in number, the graph should have 8 vertices, say P, Q, R, S, T, U, V, W , arranged in the order of degrees as given.

Then, since there are 8 vertices and the vertex W is of degree 7, W should have an edge to all other vertices. In particular, W must have an edge to both of the vertices P and Q which are of degree 1. Then P, Q are not joined to any other vertex and in particular to the vertex V which is of degree 6. Since the graph is simple, there cannot be an edge joining V to itself. Therefore, V can be joined only to five vertices W, R, S, T, U . Then V cannot have the degree 6. This is a contradiction.

Hence there is no simple graph for which the degrees of vertices are as given.

Example 17 For a graph with n vertices and m edges, if δ is the minimum and Δ is the maximum of the degrees of vertices, show that

$$\delta \leq \frac{2m}{n} < \Delta.$$

► Let $d_1, d_2, d_3, \dots, d_n$ be the degrees of the vertices. Then, by handshaking property, we have

$$d_1 + d_2 + d_3 + \dots + d_n = 2m. \quad (\text{i})$$

Since $\delta = \min(d_1, d_2, d_3, \dots, d_n)$, we have

$$d_1 \geq \delta, \quad d_2 \geq \delta, \dots, d_n \geq \delta.$$

Adding these n inequalities, we get

$$d_1 + d_2 + \dots + d_n \geq n\delta. \quad (\text{ii})$$

Similarly, since $\Delta = \max(d_1, d_2, \dots, d_n)$, we get

$$d_1 + d_2 + \dots + d_n \leq n\Delta. \quad (\text{iii})$$

From (i), (ii) and (iii), we get $2m \geq n\delta$ and $2m \leq n\Delta$, so that $n\delta \leq 2m \leq n\Delta$,

or

$$\delta \leq \frac{2m}{n} \leq \Delta$$

Example 18 Let D be a directed graph with n vertices. If the underlying graph of D is K_n , prove that

$$\sum_{v \in V} [od(v)]^2 = \sum_{v \in V} [id(v)]^2.$$

► By the First theorem of Digraph Theory, we have

$$\sum_{v \in V} od(v) = \sum_{v \in V} id(v) \quad (*)$$

This gives

$$\sum_{v \in V} \{od(v) - id(v)\} = 0 \quad (\text{i})$$

*Here, the following notation is used:

$$\sum_{v \in V} od(v) = \sum_{k=1}^n d^+(v_k), \quad \sum_{v \in V} id(v) = \sum_{k=1}^n d^-(v_k)$$

where $v_k, k = 1, 2, \dots, n$, are the vertices of D so that $V = \{v_1, v_2, \dots, v_n\}$.

For an underlying graph of a digraph, we have $d(v) = id(v) + od(v)$ for any $v \in V$. If the underlying graph is K_n , we have $d(v) = n - 1$ for every $v \in V$. Therefore, if K_n is the underlying graph of the given digraph D , we have

$$id(v) + od(v) = n - 1 \quad \text{for every } v \in D. \quad (\text{ii})$$

Now, we find that

$$\begin{aligned} \sum_{v \in V} [od(v)]^2 - [id(v)]^2 &= \sum_{v \in V} [od(v) + id(v)][od(v) - id(v)] \\ &= (n - 1) \sum_{v \in V} [od(v) - id(v)], \quad \text{using (ii)} \\ &= (n - 1) \cdot 0, \quad \text{using (i)} \\ &= 0 \end{aligned}$$


This gives

$$\sum_{v \in V} [od(v)]^2 = \sum_{v \in V} [id(v)]^2$$

as required.

Exercises

- Find the degrees of all the vertices of the graph shown in Figure 1.16. Also, verify the handshaking property for this graph.
- Verify the handshaking property for the graph shown in Figure 1.34.
- Are the following graphs regular?



(i)

(ii)

(iii)

Figure 1.42

4. Draw a diagram of a graph where the degrees of the vertices are 1, 1, 1, 2, 3, 4, 5, 7.
5. Draw a diagram for the four-dimensional hypercube Q_4 .
6. Show that, in a simple graph having n vertices, the degree d of every vertex satisfies the inequality $0 \leq d \leq n - 1$.
7. Consider a graph having n vertices and m edges. If p number of vertices are of degree k and the remaining vertices are of degree $k + 1$, prove that $p = (k + 1)n - 2m$.
8. For a graph $G = (V, E)$, what is the largest possible value of $|V|$ if $|E| = 35$ and $\deg v \geq 3$ for all $v \in V$?
9. How many vertices will the following graphs have, if they contain
 - (i) 16 edges and all vertices of degree 4?
 - (ii) 21 edges, 3 vertices of degree 4, and other vertices of degree 3?
 - (iii) 12 edges, 6 vertices of degree 3, and other vertices of degree less than 3.
10. Find the fewest vertices needed to construct a complete graph with at least 1000 edges.
11. Show that there is no simple graph with four vertices such that three vertices have degree 3 and one vertex has degree 1.
12. Prove that there is no simple graph with seven vertices, one of which has degree 2, two have degree 3, three have degree 4 and the remaining vertex has degree 5.
13. Let G be a graph with exactly $n - 1$ edges. Prove that G has either a pendant vertex or an isolated vertex.
14. Prove that every cubic graph has an even number of vertices.
15. Prove that there is no simple graph for which the degree sequence is 1, 1, 1, 2, 3, 4, 5, 7.
16. Show that the maximum number of edges in a bipartite graph of order $2n$ is n^2 .
17. Let G be a graph with n vertices and m edges and let k be the smallest positive integer such that $k \geq 2m/n$. Prove that G has a vertex of degree at least k .
18. Show that in any set of at least two persons, there are always two persons with exactly the same number of friends in the set.
19. Prove that it is impossible to have a set of nine people at a party such that each one knows exactly five of the others in the party.
20. Prove that there can be a gathering of five persons in which there are no three persons who all know each other and no three persons none of whom knows either of the other two.

Answers

1. $\deg(v_1) = 4, \deg(v_2) = 2, \deg(v_3) = 2, \deg(v_4) = 3.$ 3. Yes, all are 3-regular

4. See Figure below:




Figure 1.43

5.




Figure 1.44

8. 23 9. (i) 8 (ii) 13 (iii) at least 9 10. 46.

1.3 Isomorphism

Consider two graphs $G = (V, E)$ and $G' = (V', E')$. Suppose there exists a function $f : V \rightarrow V'$ such that (i) f is a one-to-one correspondence* and (ii) for all vertices A, B of G , $\{A, B\}$ is an edge of G if and only if $\{f(A), f(B)\}$ is an edge of G' . Then f is called an **isomorphism** between G and G' , and we say that G and G' are **isomorphic graphs**.

*Recall the definition of one-to-one correspondence; see D.S.C. : *Discrete Mathematical Structures* (4rded), Section 5.3.1.

In other words, two graphs G and G' are said to be isomorphic (to each other) if there is a one-to-one correspondence between their vertices and between their edges such that the adjacency of vertices is preserved[†]. Such graphs will have the same structure; they differ only in the way their vertices and edges are labeled or only in the way they are represented geometrically. For many purposes, we regard them as essentially the same graphs.

When G and G' are isomorphic, we write $G \cong G'$.

When a vertex A of G corresponds to the vertex $A' = f(A)$ of G' under a one-to-one correspondence $f : G \rightarrow G'$, we write $A \leftrightarrow A'$. Similarly, we write $\{A, B\} \leftrightarrow \{A', B'\}$ to mean that the edge AB of G and the edge $A'B'$ of G' correspond to each other, under f .

For example, look at the two graphs shown in Figure 1.45.




Figure 1.45

Consider the following one-to-one correspondence between the vertices of these two graphs:

$$A \leftrightarrow P, \quad B \leftrightarrow Q, \quad C \leftrightarrow R, \quad D \leftrightarrow S.$$

Under this correspondence, the edges in the two graphs correspond with each other, as indicated below:

$$\begin{aligned} \{A, B\} &\leftrightarrow \{P, Q\}, & \{A, C\} &\leftrightarrow \{P, R\}, & \{A, D\} &\leftrightarrow \{P, S\}, \\ \{B, C\} &\leftrightarrow \{Q, R\}, & \{B, D\} &\leftrightarrow \{Q, S\}, & \{C, D\} &\leftrightarrow \{R, S\} \end{aligned}$$

We check that the above-indicated one-to-one correspondence between the vertices/edges of the two graphs preserves the adjacency of the vertices. The existence of this correspondence proves that the two graphs are isomorphic. (Note that both the graphs represent the complete graph K_4)

Next, consider the two graphs shown in Figure 1.46.

[†]"adjacency of vertices is preserved" means that, for any two vertices u and v in G , if u, v are adjacent in G then the corresponding vertices u', v' in G' are also adjacent in G' .



Figure 1.46

We observe that both of these two graphs have the same number of vertices but different number of edges. Therefore, although there can exist one-to-one correspondence between the vertices, there cannot be a one-to-one correspondence between the edges. The two graphs are therefore *not isomorphic*.

From the definition of isomorphism of graphs, it follows that if two graphs are isomorphic, then they must have:

1. The same number of vertices
2. The same number of edges
3. An equal number of vertices with a given degree

These conditions are necessary but not sufficient. This means that two graphs for which these conditions hold need not be isomorphic. (See Example 8 below)

In particular, *two graphs of the same order and the same size need not be isomorphic*. To see this, consider the two graphs shown in Figure 1.47.



Figure 1.47

We note that both of these graphs are of order 4 and size 3. But the two graphs are *not isomorphic*. Observe that there are *two* pendant vertices in the first graph whereas there are *three* pendant vertices in the second graph. As such, under any one-to-one correspondence between the vertices and the edges of the two graphs, the adjacency of vertices is not preserved.

It is not hard to realize that every two complete graphs with the same number of vertices, n , are isomorphic. For this reason, we speak of *the complete graph of n vertices*, and all complete graphs with n vertices are denoted by K_n .

Similarly, any two complete bipartite graphs with bipartites containing r and s vertices are isomorphic. For this reason, all complete bipartite graphs with bipartites containing r and s vertices are denoted by $K_{r,s}$.

Given two graphs G and G' , there is no set procedure for proving or disproving that they are isomorphic. It is only by carefully examining the nature of vertices and edges of both G and G' that one can find whether or not they are isomorphic. If G and G' are not isomorphic, it is relatively easy to find it out. If G and G' are isomorphic, the work involved in proving it is quite hard – it gets harder as the orders and sizes of G and G' get larger.

Isomorphism of Digraphs

The definition of isomorphism of graphs can be extended to digraphs in a natural way. Two digraphs D_1 and D_2 are said to be isomorphic if there is a one-to-one correspondence between their vertices and between their edges such that adjacency of vertices *along with directions* is preserved.

Example 1 Prove that the two graphs shown below are isomorphic.




Figure 1.48

► We first observe that both graphs have four vertices and four edges. Consider the following one-to-one correspondence between the vertices of the graphs:

$$u_1 \leftrightarrow v_1, \quad u_2 \leftrightarrow v_4, \quad u_3 \leftrightarrow v_3, \quad u_4 \leftrightarrow v_2.$$

This correspondence gives the following correspondence between the edges:

$$\begin{aligned} \{u_1, u_2\} &\leftrightarrow \{v_1, v_4\}, & \{u_1, u_3\} &\leftrightarrow \{v_1, v_3\}, \\ \{u_2, u_4\} &\leftrightarrow \{v_4, v_2\}, & \{u_3, u_4\} &\leftrightarrow \{v_3, v_2\} \end{aligned}$$

These represent one-to-one correspondence between the edges of the two graphs under which the adjacent vertices in the first graph correspond to adjacent vertices in the second graph and vice-versa.

Accordingly, the two graphs are isomorphic.

Example 2 Verify that the two graphs shown below are isomorphic.




Figure 1.49

► Let us consider the one-to-one correspondence between the vertices of the two graphs under which the vertices A, B, C, D, P, Q, R, S of the first graph correspond to the vertices $A', B', C', D', P', Q', R', S'$ respectively of the second graph, and vice-versa. In this correspondence, the edges determined by the corresponding vertices correspond so that the adjacency of vertices is retained. As such, the two graphs are isomorphic.

We note that the first graph is the hypercube Q_3 ; see Figure 1.40. The second graph is just another drawing of Q_3 .

Example 3 Show that the following two graphs are isomorphic:




Figure 1.50

► We first note that both the graphs have six vertices each of degree three, and nine edges.

Bearing the edges in the two graphs in mind, consider the correspondence between the edges as shown below.

$$\{u_1, u_4\} \leftrightarrow \{v_1, v_2\}, \quad \{u_1, u_5\} \leftrightarrow \{v_1, v_3\}, \quad \{u_1, u_6\} \leftrightarrow \{v_1, v_6\},$$

$$\begin{aligned} \{u_2, u_5\} &\leftrightarrow \{v_4, v_3\}, & \{u_2, u_4\} &\leftrightarrow \{v_4, v_2\}, & \{u_2, u_6\} &\leftrightarrow \{v_4, v_6\}, \\ \{u_3, u_6\} &\leftrightarrow \{v_5, v_6\}, & \{u_3, u_4\} &\leftrightarrow \{v_5, v_2\}, & \{u_3, u_5\} &\leftrightarrow \{v_5, v_3\}, \end{aligned}$$

These yield the following correspondence between the vertices:

$$\begin{aligned} u_1 &\leftrightarrow v_1, & u_2 &\leftrightarrow v_4, & u_3 &\leftrightarrow v_5, \\ u_4 &\leftrightarrow v_2, & u_5 &\leftrightarrow v_3, & u_6 &\leftrightarrow v_6. \end{aligned}$$

We observe that the above correspondences between the edges and the vertices are one-to-one correspondences and that these preserve the adjacency of vertices. In view of the existence of these correspondences, we infer that the two graphs are isomorphic.

We note that the first graph is the complete bipartite graph $K_{3,3}$; see Figure 1.21(d). The second graph is just another drawing of $K_{3,3}$. ■

Example 4 Show that the following two graphs are isomorphic.




Figure 1.51

► We first note that each of the two graphs is 3-regular (cubic) and has 10 vertices. Consider the one-to-one correspondence between the vertices as shown below:


$$v_i \leftrightarrow u_i, \quad \text{for } i = 1, 2, 3, \dots, 10.$$

This correspondence has been arrived at after closely examining the structures of the two graphs.


We check that the above mentioned correspondence yields one-to-one correspondence between the edges in the two graphs with the property that adjacent vertices in the first graph correspond to the adjacent vertices in the second graph and vice-versa. The two graphs are therefore isomorphic.

We note that the first graph is the Petersen graph, see Figure 1.37. The second graph is just another drawing of the Petersen graph. ■

Example 5 Show that the following graphs are not isomorphic.



(a)




(b)


Figure 1.52

► We observe that the first graph has 4 vertices and 6 edges and the second graph has 4 vertices and 7 edges. As such, one-to-one correspondence between the edges is not possible. Hence the two graphs are *not* isomorphic. ■

Example 6 Show that the following graphs are not isomorphic



(a)




(b)


Figure 1.53

► We note that each of the two graphs has 6 vertices and 9 edges. But, the first graph has 2 vertices of degree 4 whereas the second graph has 3 vertices of degree 4. Therefore, there cannot be any one-to-one correspondence between the vertices and between the edges of the two graphs which preserves the adjacency of vertices. As such, the two graphs are *not* isomorphic. ■

Example 7 Show that the following graphs are not isomorphic.



(a)



(b)

Figure 1.54

► We note that the first graph has a pair of vertices of degree 4 which are *not* adjacent whereas the second graph has a pair of vertices of degree 4 which are adjacent. (The reader is required to identify them!). Therefore the two graphs are *not* isomorphic.

Example 8 Show that two graphs need not be isomorphic even if they have the same number of vertices, the same number of edges and equal number of vertices with the same degree.

► Consider the two graphs shown in Figure 1.55:




Figure 1.55

We observe that both graphs have the same (6) number of vertices and the same (5) number of edges. Further, in each of them there are 3 vertices of degree 1 (namely, v_1, v_5, v_6 in the first graph and A, P, R in the second graph), there are 2 vertices of degree 2 (namely, v_2, v_3 in the first graph and B, Q in the second graph), and there is 1 vertex of degree 3 (namely, v_4 in the first graph and C in the second graph). Thus, the two graphs have equal number of vertices with the same degree.

But, the two graphs are *not* isomorphic. Because there are 2 pendant vertices adjacent to the vertex v_4 (which is of degree 3) in the first graph but there is only one pendant vertex adjacent to the vertex C (which is of degree 3) in the second graph. As such, the adjacency of vertices cannot be preserved under a one-to-one correspondence between the vertices of the graphs.

Example 9 Let $G = G(V, E)$ and $G' = G'(V', E')$ be two graphs and $f : G \rightarrow G'$ be an isomorphism. Prove the following:

(i) $f^{-1} : G' \rightarrow G$ is also an isomorphism.

(ii) For any vertex v in G , $\deg(v)$ in $G = \deg(f(v))$ in G' .

► (i) Since $f : G \rightarrow G'$ is an isomorphism, f is a one-to-one correspondence (that is : f is one-to-one and onto). Therefore, f is invertible, and $f^{-1} : G' \rightarrow G$ is also a one-to-one correspondence.

Further, since f is an isomorphism, for all vertices a, b in G , $\{a, b\}$ is an edge in G if and only if $\{f(a), f(b)\}$ is an edge in G' . Since $a = f^{-1}\{f(a)\}$ and $b = f^{-1}\{f(b)\}$, this is equivalent to saying that $\{f(a), f(b)\}$ is an edge in G' if and only if $\{a, b\} = \{f^{-1}\{f(a)\}, f^{-1}\{f(b)\}\}$ is an edge in G .

It now follows that $f^{-1} : G' \rightarrow G$ is an isomorphism.

(ii) For any $v \in V$, let k be its degree in G . Then there exist k distinct vertices v_1, v_2, \dots, v_k such that $\{v, v_1\}, \{v, v_2\}, \dots, \{v, v_k\}$ are edges in G .

Consequently, $\{f(v), f(v_1)\}, \{f(v), f(v_2)\}, \dots, \{f(v), f(v_k)\}$ are edges in G' .

Since v_1, v_2, \dots, v_k are distinct and f is a one-to-one correspondence, $f(v_1), f(v_2), \dots, f(v_k)$ are also distinct and their number is k . Thus, there are k edges in G' which are incident on $f(v)$. Hence $\deg\{f(v)\}$ in G' is k . This proves the required result. ■

Example 10 Show that the following digraphs are isomorphic.




Figure 1.56

► Consider the following one-to-one correspondence between the vertices of the given digraphs:

$$A \leftrightarrow Q, \quad B \leftrightarrow P, \quad C \leftrightarrow S, \quad D \leftrightarrow R.$$

Under this correspondence, the directed edges of the two graphs correspond with each other as shown below.

$$(A, B) \leftrightarrow (Q, P), \quad (B, C) \leftrightarrow (P, S), \quad (C, D) \leftrightarrow (S, R), \quad (D, A) \leftrightarrow (R, Q)$$

Evidently, under this correspondence, the adjacency of vertices including directions of the edges is preserved.

Hence the given digraphs are isomorphic. ■

Example 11 Show that the following digraphs are not isomorphic.




Figure 1.57

- The two digraphs have the same number of vertices (5) and the same number of directed edges (7). We observe that the vertex A of the first digraph has 1 as its out-degree and 2 as its in-degree. There is no such vertex in the second digraph. Therefore, there cannot be any one-to-one correspondence between the vertices of the two digraphs which preserves the direction of edges. The two digraphs are therefore *not* isomorphic. ■

Exercises

- Show that the following graphs are isomorphic.




Figure 1.58

- Show that the following graphs are isomorphic.




Figure 1.59

3. Show that the following graphs are isomorphic.




Figure 1.60

4. Show that the following graphs are not isomorphic.




Figure 1.61

5. Show that the following graphs are not isomorphic.




Figure 1.62

6. Show that the following graphs are not isomorphic.




Figure 1.63

7. Show that the following graphs are isomorphic.




Figure 1.64

8. Verify that the following graphs are isomorphic.




Figure 1.65

9. Show that the following digraphs are isomorphic.




Figure 1.66

10. Show that the following digraphs are not isomorphic.




Figure 1.67

1.4 Subgraphs

Given two graphs G and G_1 , we say that G_1 is a *subgraph* of G if the following conditions hold.

- (i) All the vertices and all the edges of G_1 are in G .
- (ii) Each edge of G_1 has the same end vertices in G as in G_1 .

Essentially, a subgraph is a graph which is a part of another graph. Any graph isomorphic to a subgraph of a graph G is also referred to as a subgraph of G .

Consider the two graphs G_1 and G shown in Figures 1.68(a) and 1.68(b) respectively. We observe that all vertices and all edges of the graph G_1 are in the graph G and that every edge in G_1 has the same end vertices in G as in G_1 . Therefore, G_1 is a subgraph of G . In the diagram of G , the part G_1 is shown in thicker lines.




Figure 1.68

The following results are immediate consequences of the definition of a subgraph.

- (1) Every graph is a subgraph of itself.
- (2) Every simple graph of n vertices is a subgraph of the complete graph K_n .
- (3) If G_1 is a subgraph of a graph G_2 and G_2 is a subgraph of a graph G , then G_1 is a subgraph of G .
- (4) A single vertex in a graph G is a subgraph of G .
- (5) A single edge in a graph G , together with its end vertices, is a subgraph of G .

Spanning Subgraph

Given a graph $G = (V, E)$, if there is a subgraph $G_1 = (V_1, E_1)$ of G such that $V_1 = V$, then G_1 is called a *spanning subgraph* of G .

In other words, a subgraph G_1 of a graph G is a spanning subgraph of G whenever G_1 contains all vertices of G . Thus, a graph and all its spanning subgraphs have the same vertex set. Obviously, every graph is its own spanning subgraph.

For example, for the graph shown in Figure 1.69(a), the graph shown in Figure 1.69(b) is a spanning subgraph whereas the graph shown in Figure 1.69(c) is a subgraph but not a spanning subgraph.




Figure 1.69

Induced Subgraph

Given a graph $G = (V, E)$, suppose there is a subgraph $G_1 = (V_1, E_1)$ of G such that every edge $\{A, B\}$ of G , where $A, B \in V_1$ is an edge of G_1 also. Then G_1 is called an *induced subgraph* of G (*induced by* V_1) and is denoted by $\langle V_1 \rangle$.

If follows that a subgraph $G_1 = (V_1, E_1)$ of a graph $G = (V, E)$ is *not* an induced subgraph of G if for some $A, B \in V_1$, there is an edge $\{A, B\}$ which is in G but not in G_1 .

For example, for the graph shown in the Figure 1.70(a), the graph shown in Figure 1.70(b) is an induced subgraph – induced by the set of vertices $V_1 = \{v_1, v_2, v_3, v_5\}$, whereas the graph shown in Figure 1.70(c) is not an induced subgraph.




Figure 1.70

Edge-disjoint and Vertex-disjoint Subgraphs

Let G be a graph and G_1 and G_2 be two subgraphs of G . Then:

- (1) G_1 and G_2 are said to be *edge-disjoint* if they do not have any edge in common.
- (2) G_1 and G_2 are said to be *vertex-disjoint* if they do not have any common edge *and* any common vertex.

It is to be noted that edge-disjoint subgraphs may have common vertices. Subgraphs that have no vertices in common cannot possibly have edges in common. That is, vertex-disjoint subgraphs must be edge-disjoint also but the converse is not necessarily true.

For example, for the graph shown in Figure 1.71(a), the graphs shown in Figures 1.71(b) and 1.71(c) are edge-disjoint but not vertex-disjoint subgraphs.




Figure 1.71

Example 1 Given a graph G_1 , can there exist a graph G_2 such that G_1 is a subgraph of G_2 but not a spanning subgraph of G_2 and yet G_1 and G_2 have the same size?

► Yes. Consider a graph G_1 which contains all the vertices and all the edges of G and at least one extra isolated vertex. See Figures 1.72(a), (b) for instance.




Figure 1.72

Example 2 Consider the graph G shown in Figure 1.73(a).

- Verify that the graph G_1 shown in Figure 1.73(b) is an induced subgraph of G . Is this a spanning subgraph of G ?
- Draw the subgraph G_2 of G induced by the set $V_2 = \{v_3, v_4, v_6, v_8, v_9\}$.




Figure 1.73

► (a) The vertex set of the graph G_1 , namely $V_1 = \{v_1, v_3, v_4, v_6, v_7\}$, is a subset of the vertex set $V = \{v_1, v_2, \dots, v_9\}$ of G .

Also, all the edges of G_1 are in G . Further, each edge in G_1 has the same end vertices in G as in G_1 . Therefore, G_1 is a subgraph of G .

We further check that every edge $\{v_i, v_j\}$ of G where $v_i, v_j \in V_1$ is an edge of G_1 . Therefore, G_1 is an induced subgraph of G . Since $V_1 \neq V$, G_1 is not a spanning subgraph of G .

- (b) The subgraph $G_2 = \langle V_2 \rangle$ is shown in Figure 1.73(c):




Figure 1.73 (c)

Example 3 Consider the graph G shown in Figure 1.74(a). Verify that the graphs G_1 and G_2 shown in Figures 1.74(b) and 1.74(c) are induced subgraphs of G whereas the graph G_3 shown in Figure 1.74(d) is not an induced subgraph of G .




Figure 1.74

► We note that the vertex sets of G_1 , G_2 and G_3 are all subsets of the vertex set of G . Further, all edges in each of G_1 , G_2 , G_3 have the same end vertices in G as in these. Therefore, all of G_1 , G_2 , G_3 are subgraphs of G .

We further check that every edge in G whose end vertices belong to G_1 is an edge in G_1 . Therefore, G_1 is an induced subgraph of G . In fact, G_1 is induced by the set $\{v_1, v_2, v_4, v_5\}$.

Similarly, G_2 is an induced subgraph of G , induced by the set $\{v_2, v_3, v_5, v_6\}$.

The subgraph G_3 of G is not an induced subgraph of G . Because, for example, $\{v_1, v_5\}$ is an edge in G whose end vertices belong to G_3 , but $\{v_1, v_5\}$ is not an edge in G_3 . ■

Example 4 For the graph shown in Figure 1.75, find two edge-disjoint subgraphs and two vertex-disjoint subgraphs.




Figure 1.75

► For the given graph, two edge-disjoint subgraphs are shown in Figure 1.76(a) and two vertex-disjoint subgraphs are shown in Figure 1.76(b).




Figure 1.76

Remark: The given graph has other pairs of edge-disjoint subgraphs and vertex-disjoint subgraphs. The reader can identify these. ■

Exercises

1. Let G be the graph shown in Figure 1.77. Verify whether $G_1 = (V_1, E_1)$ is a subgraph of G in the following cases:

- (i) $V_1 = \{P, Q, S\}, E_1 = \{(PQ, PS)\}$
- (ii) $V_1 = \{Q\}, E_1 = \Phi$, the null set
- (iii) $V_1 = \{P, Q, R\}, E_1 = \{PQ, QR, QS\}$,




Figure 1.77

2. Three graphs G_1 , G_2 , G_3 are shown in Figures 1.78(a), (b), (c) respectively. Are G_2 and G_3 induced subgraphs of G_1 ? Are they spanning subgraphs?


(a) : G_1 (b) : G_2 (b) : G_3

Figure 1.78

3. Can a finite graph be isomorphic to one of its subgraphs (other than itself)?

Answers

1. (i) No. (ii) Yes. (iii) No.
2. G_2 is an induced subgraph of G_1 , it is not a spanning subgraph.
 G_3 is not an induced subgraph of G_1 , it is a spanning subgraph.
3. No.

1.5 Operations on Graphs

Consider two graphs $G_1 = (V_1, E_1)$ and $G_2 = (V_2, E_2)$.

Then the graph whose vertex set is $V_1 \cup V_2$ and the edge set is $E_1 \cup E_2$ is called the *union* of G_1 and G_2 , it is denoted by $G_1 \cup G_2$. Thus,

$$G_1 \cup G_2 = (V_1 \cup V_2, E_1 \cup E_2).$$

Similarly, if $V_1 \cap V_2 \neq \emptyset$, the graph whose vertex set is $V_1 \cap V_2$ and the edge set is $E_1 \cap E_2$ is called the *intersection* of G_1 and G_2 ; it is denoted by $G_1 \cap G_2$. Thus,

$$G_1 \cap G_2 = (V_1 \cap V_2, E_1 \cap E_2). \quad \text{if } V_1 \cap V_2 \neq \emptyset.$$

Next, suppose we consider the graph whose vertex set is $V_1 \cup V_2$ and the edge set is $E_1 \Delta E_2$, where $E_1 \Delta E_2$ is the *symmetric difference* of E_1 and E_2 ^{*}. This graph is called the *ring sum* of

*The symmetric difference $E_1 \Delta E_2$ denotes the set of all those elements (–here, the edges) which are in E_1 or E_2 but not in both. That is:

$$E_1 \Delta E_2 = (E_1 \cup E_2) - (E_1 \cap E_2).$$

See, for example, D.S.C: *Discrete Mathematical Structures* (4th ed), Section 1.2.

G_1 and G_2 , it is denoted by $G_1 \Delta G_2$. Thus,

$$G_1 \Delta G_2 = (V_1 \cup V_2, E_1 \Delta E_2).$$

For the two graphs G_1 and G_2 shown in Figures 1.79(a) and (b), their union, intersection and ring sum are shown in Figures 1.80(a), (b) and (c) respectively.




Figure 1.79




Figure 1.80

Decomposition

We say that a graph G is *decomposed* (or *partitioned*) into two subgraphs G_1 and G_2 if

$$G_1 \cup G_2 = G \quad \text{and} \quad G_1 \cap G_2 = \text{Null graph.}$$

Deletion

If v is a vertex in a graph G , then $G - v$ denotes the subgraph of G obtained by deleting v and all edges incident on v , from G . This subgraph, $G - v$, is referred to as *vertex-deleted subgraph* of G .

It should be noted that the deletion of a vertex always includes the deletion of all edges incident on that vertex.