

# Learning L<sup>A</sup>T<sub>E</sub>X

EE24BTECH11053 - S A Aravind Eswar\*

14. Let  $S$  be set of all column matrix  $\begin{bmatrix} b_1 \\ b_2 \\ b_3 \end{bmatrix}$  such that  $b_1, b_2, b_3 \in \mathbb{R}$  and the system of equations (in real variables)

$$-x + 2y + 5z = b_1, 2x - 4y + 3z = b_2, x - 2y + 2z = b_3$$

has at least one solution. Then, which of the following system(s) (in real variables) has (have) at least one solution for each  $\begin{bmatrix} b_1 \\ b_2 \\ b_3 \end{bmatrix} \in S$  (JEE Adv. 2018)

- 1)  $x + 2y + 3z = b_1, 4y + 5z = b_2$  and  $x + 2y + 6z = b_3$
- 2)  $x + y + 3z = b_1, 5x + 2y + 6z = b_2$  and  $-2x - y - 3z = b_3$
- 3)  $-x + 2y - 5z = b_1, 2x - 4y + 10z = b_2$  and  $x - 2y + 5z = b_3$
- 4)  $5x + 2y + 5z = b_1, 2x + 3z = b_2, x + 4y - 5z = b_3$

15. Let  $M = \begin{bmatrix} 0 & 1 & a \\ 1 & 2 & 3 \\ 3 & b & 1 \end{bmatrix}$  and  $(adj M) = \begin{bmatrix} -1 & 1 & -1 \\ 8 & -6 & 2 \\ -5 & 3 & 1 \end{bmatrix}$  where  $a$  and  $b$  are real numbers. Which of the following options is/are correct? (JEE Adv. 2019)

- 1)  $a + b = 3$
- 2)  $\det(adj M^2) = 81$
- 3)  $(adj M)^{-1} + adj M^{-1} = -M$
- 4) If  $M \begin{bmatrix} \alpha \\ \beta \\ \gamma \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}$  then  $\alpha - \beta + \gamma = 3$

16. Let

$$P_1 = I = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}, P_2 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 1 \end{bmatrix}, P_3 = \begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix}, P_4 = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{bmatrix}, P_5 = \begin{bmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix}, P_6 = \begin{bmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 1 \end{bmatrix}$$