

CSE-420

Final

Ishtiaq Ahmed

Sec: 08

ID: 20301429

$\frac{2 \cdot e}{3} \cdot \rho = \frac{1 - \rho}{3} - \rho$

shirwan not belong to a

$\rho = 1 + 8 = 9$


$\rho = 9$

$\rho = \frac{e \cdot \rho}{3} + \rho = \frac{1 - \rho}{3} + 1$

can give 25 in 100000

①

Ans: to the Q. NO-3)



expression is ~~(a+a)-a~~

$$(a+a)+(a+a)-(a+a)-a$$

$$\cancel{(a+a) + (a+a) - a}$$

$t_1 = \text{unipus } a$

$$t_1 = (a+a)$$

$$t_2 = \cancel{(a+a)} + t_1$$

$$t_2 = t_1 + \cancel{(a+a)} t_1$$

$$t_3 =$$

$$t_3 = t_2 - \cancel{(a+a)} t_1$$

$$t_4 = t_3 - a$$

②

Quadruples

	Op	arg1	arg2	result
0	+	a	a	t ₁
1	+	t ₁	t ₁	t ₂
2	-	t ₂	t ₁	t ₃
3	-	t ₃	a	t ₄

Triple


	Op	arg1	arg2
0	+	a	a
1	+	(0)	(0)
2	-	(1)	(0)
3	-	(2)	a

Ans: to the Q: NO-1

Pose Tree


085, 2, 48
52 56

$T = \text{float, int}$
 $B = 8 \text{ } 4$



T: Type array (2, array, 3, float)
T: width = 48

B: type = float
B: width = 8



Symbol tables

<u>variable</u>	<u>type</u>	<u>offset</u>
a	array (2, array (3; float))	0
b	int	48
p	int	52

Ans: to the Q: NO-2

Using list of hash tables.

