


Name: Adinur Rahman Abir

ID: 19101055

Sec: 07

CSE: 420

Answer to the question No-2



Answer to the question No-1

$P \rightarrow MD$ [offset=0;]

$M \rightarrow \epsilon$

$D \rightarrow Tid;$

$D,$

$D \rightarrow \epsilon$

$T \rightarrow \text{record } \{$

$D \}$

$T \rightarrow B$
 c



$B \rightarrow \text{int.}$

$B \rightarrow \text{float}$

$C \rightarrow \epsilon$

$C \rightarrow [\text{num}]C,$

input: record{ float[2][3]a; int b; }p;



05

~~T.type = int, w = 4~~

offset = 0
24

~~T.type = int, c.width = 4~~

~~Stack.push(0); offset = 0~~

~~T.type = float; T.width = 8~~

~~top.put(~~int~~ [2], int, 0)~~

~~offset =~~

~~C.type = array(2,~~

~~c.width = 2 × 8 = 16~~

type	width	offset
int	16	24

type	offset	width
float	0	8
array	8	16

Answer to the question No - 3



~~a + a + a~~
 $x = (a + a) - (a - (a + a))$

0 $t_1 = a + a$

1 $t_2 = a - t_1$

2 $t_3 = t_1$

3 $t_4 = t_1 - t_2$

4 ~~$x = t_4$~~ $x = t_4$

quadruple:

	opp	arg 1	arg 2	result
0	+	a	a	t_1
1	-	a	t_1	t_2
2	=	t_1		t_3
3	-	t_1	t_2	t_4
4	=	t_4		x

triple

i	op	a_1	a_2
0	+	a	a
1	-	a	(0)
2	=	(0)	(0)
3	-	(0)	(1)
4	=	(3)	

07