

Ans to the Question No-4

record { float [3] [3] a, int b; } P;

a	float	0
b	int	48
P		52

top record = 52

float = 6, 8, 10

int = 10

float = 52 + 32 = 84


int = 52 + 32 = 84

float = 52 + 32 = 84


int = 52 + 32 = 84

float = 52 + 32 = 84

int = 52 + 32 = 84




Ans to the Question No 2



Ans to the Question NO-3

(a)

$$(a+a+a) - (a+a-a)$$



So the expression will be: ~~$((a+a) + (a+a)) - ((a+a) - a)$~~
 $\Rightarrow a+a+a+a-a-a-a$

(b)

Now,

$$\begin{aligned} t_1 &= a+a \\ t_2 &= t_1+a \\ t_3 &= t_1+t_1 \\ t_4 &= t_3-t_2 \\ t_5 &= t_4-t_3 \end{aligned}$$

~~triples~~

(b)

triples

	OP	arg1	arg2
(0)	unary	a	a
(1)	+	(0)	a
(2)	+	(0)	(1)
(3)	+	(2)	(1)
(4)	-	(3)	(2)

7

quadriples

	OP	arg1	arg2	result
(0)	unary	a	a	t ₁
(1)	-	t ₁	a	t ₂
(2)	+	t ₁	t ₁	t ₃
(3)	+	t ₃	t ₂	t ₄
(4)	-	t ₄	t ₃	s