MedBrains Sans Accessible: one change per column (light)

Change	**v0 (shipped)** IBM Plex Sans, slashed zero Latin WOFF2 36.8 KB (+0.0 KB vs v0) baseline for every column	**1 Spacing +20 units** +2% em side bearings, all glyphs Latin WOFF2 36.9 KB (+0.1 KB vs v0) line length +4.5% ships: variable WOFF2, all weights	**2 Regular at 440** body text at wght 440, not 400 Latin WOFF2 36.8 KB (+0.0 KB vs v0) (no change) web: CSS font-weight 440 ships: variable WOFF2; statics re-cut	**3 Tail on q** breaks p/q mirror (b/d already spurred) Latin WOFF2 36.9 KB (+0.1 KB vs v0) ships: variable WOFF2, all weights	**4 Open apertures** c e s a terminals trimmed statics only: a boolean cut breaks interpolation; needs a designer for the variable font	**5 Dotted zero** Plex ss04 instead of the slash Latin WOFF2 36.9 KB (+0.1 KB vs v0) ships: variable WOFF2, all weights	**6 Atkinson b d p q l 1** borrowed outlines, scaled to Plex Latin WOFF2 37.1 KB (+0.2 KB vs v0) ships: variable WOFF2 (advance variations stay Plex's: check bold)	**Recommended mix** spacing +12, Regular 440, tail on q Latin WOFF2 36.7 KB (-0.1 KB vs v0) slashed zero kept ships: variable WOFF2, all weights
Mirror and confusable 24 px	b d p q rn m cl d Il1| O0o 5S 2Z 6b 8B ceas case sea	b d p q rn m cl d Il1| O0o 5S 2Z 6b 8B ceas case sea	b d p q rn m cl d Il1| O0o 5S 2Z 6b 8B ceas case sea	b d p q rn m cl d Il1| O0o 5S 2Z 6b 8B ceas case sea	b d p q rn m cl d Il1| O0o 5S 2Z 6b 8B ceas case sea	b d p q rn m cl d Il1| O0o 5S 2Z 6b 8B ceas case sea	b d p q rn m cl d Il1| O0o 5S 2Z 6b 8B ceas case sea	b d p q rn m cl d Il1| O0o 5S 2Z 6b 8B ceas case sea
Low vision 14 px, blur 1.5 px just-AA grey	b d p q rn m cl d Il1| O0o 5S 2Z 6b 8B 10 mg 0.5 mL 1.25 mg/kg INR 1.10	b d p q rn m cl d Il1| O0o 5S 2Z 6b 8B 10 mg 0.5 mL 1.25 mg/kg INR 1.10	b d p q rn m cl d Il1| O0o 5S 2Z 6b 8B 10 mg 0.5 mL 1.25 mg/kg INR 1.10	b d p q rn m cl d Il1| O0o 5S 2Z 6b 8B 10 mg 0.5 mL 1.25 mg/kg INR 1.10	b d p q rn m cl d Il1| O0o 5S 2Z 6b 8B 10 mg 0.5 mL 1.25 mg/kg INR 1.10	b d p q rn m cl d Il1| O0o 5S 2Z 6b 8B 10 mg 0.5 mL 1.25 mg/kg INR 1.10	b d p q rn m cl d Il1| O0o 5S 2Z 6b 8B 10 mg 0.5 mL 1.25 mg/kg INR 1.10	b d p q rn m cl d Il1| O0o 5S 2Z 6b 8B 10 mg 0.5 mL 1.25 mg/kg INR 1.10
Patient instruction 16 px, line 1.5	Take one 500 mg tablet twice a day after food, for 5 days. Do not take it with milk. If you get a rash, swelling of the face or trouble breathing, stop the tablets and come to the hospital at once.	Take one 500 mg tablet twice a day after food, for 5 days. Do not take it with milk. If you get a rash, swelling of the face or trouble breathing, stop the tablets and come to the hospital at once.	Take one 500 mg tablet twice a day after food, for 5 days. Do not take it with milk. If you get a rash, swelling of the face or trouble breathing, stop the tablets and come to the hospital at once.	Take one 500 mg tablet twice a day after food, for 5 days. Do not take it with milk. If you get a rash, swelling of the face or trouble breathing, stop the tablets and come to the hospital at once.	Take one 500 mg tablet twice a day after food, for 5 days. Do not take it with milk. If you get a rash, swelling of the face or trouble breathing, stop the tablets and come to the hospital at once.	Take one 500 mg tablet twice a day after food, for 5 days. Do not take it with milk. If you get a rash, swelling of the face or trouble breathing, stop the tablets and come to the hospital at once.	Take one 500 mg tablet twice a day after food, for 5 days. Do not take it with milk. If you get a rash, swelling of the face or trouble breathing, stop the tablets and come to the hospital at once.	Take one 500 mg tablet twice a day after food, for 5 days. Do not take it with milk. If you get a rash, swelling of the face or trouble breathing, stop the tablets and come to the hospital at once.
Clinical 14 px	10 mg 0.5 mL 1.25 mg/kg 100 IU 5 mL BD UHID MB-2026-0010 / OPD-11071 lot B10I-O01l Hb 10.1 g/dL K+ 5.1 INR 1.10 hydrOXYzine / hydrALAZINE CeleBREX / CeleXA खुराक 10 mg, दिन में दो बार (BD) · रक्तचाप 120/80	10 mg 0.5 mL 1.25 mg/kg 100 IU 5 mL BD UHID MB-2026-0010 / OPD-11071 lot B10I-O01l Hb 10.1 g/dL K+ 5.1 INR 1.10 hydrOXYzine / hydrALAZINE CeleBREX / CeleXA खुराक 10 mg, दिन में दो बार (BD) · रक्तचाप 120/80	10 mg 0.5 mL 1.25 mg/kg 100 IU 5 mL BD UHID MB-2026-0010 / OPD-11071 lot B10I-O01l Hb 10.1 g/dL K+ 5.1 INR 1.10 hydrOXYzine / hydrALAZINE CeleBREX / CeleXA खुराक 10 mg, दिन में दो बार (BD) · रक्तचाप 120/80	10 mg 0.5 mL 1.25 mg/kg 100 IU 5 mL BD UHID MB-2026-0010 / OPD-11071 lot B10I-O01l Hb 10.1 g/dL K+ 5.1 INR 1.10 hydrOXYzine / hydrALAZINE CeleBREX / CeleXA खुराक 10 mg, दिन में दो बार (BD) · रक्तचाप 120/80	10 mg 0.5 mL 1.25 mg/kg 100 IU 5 mL BD UHID MB-2026-0010 / OPD-11071 lot B10I-O01l Hb 10.1 g/dL K+ 5.1 INR 1.10 hydrOXYzine / hydrALAZINE CeleBREX / CeleXA खुराक 10 mg, दिन में दो बार (BD) · रक्तचाप 120/80	10 mg 0.5 mL 1.25 mg/kg 100 IU 5 mL BD UHID MB-2026-0010 / OPD-11071 lot B10I-O01l Hb 10.1 g/dL K+ 5.1 INR 1.10 hydrOXYzine / hydrALAZINE CeleBREX / CeleXA खुराक 10 mg, दिन में दो बार (BD) · रक्तचाप 120/80	10 mg 0.5 mL 1.25 mg/kg 100 IU 5 mL BD UHID MB-2026-0010 / OPD-11071 lot B10I-O01l Hb 10.1 g/dL K+ 5.1 INR 1.10 hydrOXYzine / hydrALAZINE CeleBREX / CeleXA खुराक 10 mg, दिन में दो बार (BD) · रक्तचाप 120/80	10 mg 0.5 mL 1.25 mg/kg 100 IU 5 mL BD UHID MB-2026-0010 / OPD-11071 lot B10I-O01l Hb 10.1 g/dL K+ 5.1 INR 1.10 hydrOXYzine / hydrALAZINE CeleBREX / CeleXA खुराक 10 mg, दिन में दो बार (BD) · रक्तचाप 120/80

MedBrains Sans Accessible: one change per column (dark)

Change	**v0 (shipped)** IBM Plex Sans, slashed zero Latin WOFF2 36.8 KB (+0.0 KB vs v0) baseline for every column	**1 Spacing +20 units** +2% em side bearings, all glyphs Latin WOFF2 36.9 KB (+0.1 KB vs v0) line length +4.5% ships: variable WOFF2, all weights	**2 Regular at 440** body text at wght 440, not 400 Latin WOFF2 36.8 KB (+0.0 KB vs v0) (no change) web: CSS font-weight 440 ships: variable WOFF2; statics re-cut	**3 Tail on q** breaks p/q mirror (b/d already spurred) Latin WOFF2 36.9 KB (+0.1 KB vs v0) ships: variable WOFF2, all weights	**4 Open apertures** c e s a terminals trimmed statics only: a boolean cut breaks interpolation; needs a designer for the variable font	**5 Dotted zero** Plex ss04 instead of the slash Latin WOFF2 36.9 KB (+0.1 KB vs v0) ships: variable WOFF2, all weights	**6 Atkinson b d p q l 1** borrowed outlines, scaled to Plex Latin WOFF2 37.1 KB (+0.2 KB vs v0) ships: variable WOFF2 (advance variations stay Plex's: check bold)	**Recommended mix** spacing +12, Regular 440, tail on q Latin WOFF2 36.7 KB (-0.1 KB vs v0) slashed zero kept ships: variable WOFF2, all weights
Mirror and confusable 24 px	b d p q rn m cld Il1| O0o 5S 2Z 6b 8B ceas case sea	b d p q rn m cld Il1| O0o 5S 2Z 6b 8B ceas case sea	b d p q rn m cld Il1| O0o 5S 2Z 6b 8B ceas case sea	b d p q rn m cld Il1| O0o 5S 2Z 6b 8B ceas case sea	b d p q rn m cld Il1| O0o 5S 2Z 6b 8B ceas case sea	b d p q rn m cld Il1| O0o 5S 2Z 6b 8B ceas case sea	b d p q rn m cld Il1| O0o 5S 2Z 6b 8B ceas case sea	b d p q rn m cld Il1| O0o 5S 2Z 6b 8B ceas case sea
Low vision 14 px, blur 1.5 px just-AA grey	b d p q rn m cld Il1| O0o 5S 2Z 6b 8B 10 mg 0.5 mL 1.25 mg/kg INR 1.10	b d p q rn m cld Il1| O0o 5S 2Z 6b 8B 10 mg 0.5 mL 1.25 mg/kg INR 1.10	b d p q rn m cld Il1| O0o 5S 2Z 6b 8B 10 mg 0.5 mL 1.25 mg/kg INR 1.10	b d p q rn m cld Il1| O0o 5S 2Z 6b 8B 10 mg 0.5 mL 1.25 mg/kg INR 1.10	b d p q rn m cld Il1| O0o 5S 2Z 6b 8B 10 mg 0.5 mL 1.25 mg/kg INR 1.10	b d p q rn m cld Il1| O0o 5S 2Z 6b 8B 10 mg 0.5 mL 1.25 mg/kg INR 1.10	b d p q rn m cld Il1| O0o 5S 2Z 6b 8B 10 mg 0.5 mL 1.25 mg/kg INR 1.10	b d p q rn m cld Il1| O0o 5S 2Z 6b 8B 10 mg 0.5 mL 1.25 mg/kg INR 1.10
Patient instruction 16 px, line 1.5	Take one 500 mg tablet twice a day after food, for 5 days. Do not take it with milk. If you get a rash, swelling of the face or trouble breathing, stop the tablets and come to the hospital at once.	Take one 500 mg tablet twice a day after food, for 5 days. Do not take it with milk. If you get a rash, swelling of the face or trouble breathing, stop the tablets and come to the hospital at once.	Take one 500 mg tablet twice a day after food, for 5 days. Do not take it with milk. If you get a rash, swelling of the face or trouble breathing, stop the tablets and come to the hospital at once.	Take one 500 mg tablet twice a day after food, for 5 days. Do not take it with milk. If you get a rash, swelling of the face or trouble breathing, stop the tablets and come to the hospital at once.	Take one 500 mg tablet twice a day after food, for 5 days. Do not take it with milk. If you get a rash, swelling of the face or trouble breathing, stop the tablets and come to the hospital at once.	Take one 500 mg tablet twice a day after food, for 5 days. Do not take it with milk. If you get a rash, swelling of the face or trouble breathing, stop the tablets and come to the hospital at once.	Take one 500 mg tablet twice a day after food, for 5 days. Do not take it with milk. If you get a rash, swelling of the face or trouble breathing, stop the tablets and come to the hospital at once.	Take one 500 mg tablet twice a day after food, for 5 days. Do not take it with milk. If you get a rash, swelling of the face or trouble breathing, stop the tablets and come to the hospital at once.
Clinical 14 px	10 mg 0.5 mL 1.25 mg/kg 100 IU 5 mL BD UHID MB-2026-0010 / OPD-11071 lot B10I-O01l Hb 10.1 g/dL K+ 5.1 INR 1.10 hydrOXYzine / hydrALAZINE CeleBREX / CeleXA खुराक 10 mg, दिन में दो बार (BD) · रक्तचाप 120/80	10 mg 0.5 mL 1.25 mg/kg 100 IU 5 mL BD UHID MB-2026-0010 / OPD-11071 lot B10I-O01l Hb 10.1 g/dL K+ 5.1 INR 1.10 hydrOXYzine / hydrALAZINE CeleBREX / CeleXA खुराक 10 mg, दिन में दो बार (BD) · रक्तचाप 120/80	10 mg 0.5 mL 1.25 mg/kg 100 IU 5 mL BD UHID MB-2026-0010 / OPD-11071 lot B10I-O01l Hb 10.1 g/dL K+ 5.1 INR 1.10 hydrOXYzine / hydrALAZINE CeleBREX / CeleXA खुराक 10 mg, दिन में दो बार (BD) · रक्तचाप 120/80	10 mg 0.5 mL 1.25 mg/kg 100 IU 5 mL BD UHID MB-2026-0010 / OPD-11071 lot B10I-O01l Hb 10.1 g/dL K+ 5.1 INR 1.10 hydrOXYzine / hydrALAZINE CeleBREX / CeleXA खुराक 10 mg, दिन में दो बार (BD) · रक्तचाप 120/80	10 mg 0.5 mL 1.25 mg/kg 100 IU 5 mL BD UHID MB-2026-0010 / OPD-11071 lot B10I-O01l Hb 10.1 g/dL K+ 5.1 INR 1.10 hydrOXYzine / hydrALAZINE CeleBREX / CeleXA खुराक 10 mg, दिन में दो बार (BD) · रक्तचाप 120/80	10 mg 0.5 mL 1.25 mg/kg 100 IU 5 mL BD UHID MB-2026-0010 / OPD-11071 lot B10I-O01l Hb 10.1 g/dL K+ 5.1 INR 1.10 hydrOXYzine / hydrALAZINE CeleBREX / CeleXA खुराक 10 mg, दिन में दो बार (BD) · रक्तचाप 120/80	10 mg 0.5 mL 1.25 mg/kg 100 IU 5 mL BD UHID MB-2026-0010 / OPD-11071 lot B10I-O01l Hb 10.1 g/dL K+ 5.1 INR 1.10 hydrOXYzine / hydrALAZINE CeleBREX / CeleXA खुराक 10 mg, दिन में दो बार (BD) · रक्तचाप 120/80	10 mg 0.5 mL 1.25 mg/kg 100 IU 5 mL BD UHID MB-2026-0010 / OPD-11071 lot B10I-O01l Hb 10.1 g/dL K+ 5.1 INR 1.10 hydrOXYzine / hydrALAZINE CeleBREX / CeleXA खुराक 10 mg, दिन में दो बार (BD) · रक्तचाप 120/80