Article Navigation

JOURNAL ARTICLE

The effect of severe psoriasis on the quality of life of 369 patients Get access

A.Y.

Finlay , E.C. Coles

British Journal of Dermatology, Volume 132, Issue 2, 1 February 1995, Pages 236-244, https://doi.org/10.1111/j.1365-2133.1995.tb05019.x

Article

Published: 01 February 1995 history 2

? Views 2

? Cite

? Permissions

? Share ?

Summary

The aim of this survey was to quantify the level of handicap experienced by patients with severe psoriasis, and to assess the value that patients place on their disease using various questionnaire techniques. Dermatologists throughout the U.K. each gave a questionnaire to up to five psoriasis patients, who were either being admitted for in-patient care or were starting systemic therapy. Three hundred and sixtynine questionnaires were completed. Of the 150 patients currently working, 59·3% had lost a mean of 26 days (SD 21·9) from work during the preceding year because of their psoriasis, and of the 180 not working 33.9% attributed not working to their psoriasis. The mean Psoriasis Disability Index (PDI) score was 38·2% (SD 23·3, n=248), with the mean sub-scores of the 'daily activities' and 'treatment' sections being greater than those of the other three sections. Despite having severe psoriasis, the majority of patients felt that it would be worse to have diabetes, asthma or bronchitis than to have psoriasis. Forty-six, 42 and 32% considered it would be either 'better' or 'the same' to have diabetes, asthma or bronchitis, respectively. However, in those patients who also had the comparative disease, 87, 80 and 77% considered it would be 'better', or 'the same' to have the comparative disease. Forty-nine per cent of patients (n=362) stated they would be prepared to spend 2 or 3 h each day on treatment if this might result in normal skin for the rest of the day. Three hundred and sixty-four (98.9%) of patients stated they would prefer to have a complete cure of their psoriasis rather than be given £1000 cash. Seventyone per cent (n=336) of patients said they would be prepared to pay £1000 or more, and 38% (n=336) said they would pay £10,000, for a cure for their psoriasis. There was a substantial correlation (0.48) between the PDI score and the amount a patient indicated they would pay for a cure.

This survey of patients with severe psoriasis reveals major handicap caused by the disease, and illustrates the extent to which patients would value effective therapy.

This article is published and distributed under the terms of the Oxford University Press, Standard Journals Publication Model (https://ac ademic.oup.com/journals/pages/open_access/funder_policies/chorus/standard_publication_model)

Issue Section: Original Articles

You do not currently have access to this article.

Sign in

• Get help with access

British Association of Dermatologists members

Sign in through society site

All other society members

Sign in through society site

Personal account

- Sign in with email/username & password
- Get email alerts
- Save searches
- Purchase content
- Activate your purchase/trial code
- Add your ORCID iD

Sign in

Register

Institutional access

Sign in through your institution

Sign in through your institution

Sign in with a library card

Sign in with username/password

Recommend to your librarian

Institutional account management

Sign in as administrator

Purchase

Subscription prices and ordering for this journal

Purchasing options for books and journals across Oxford Academic

Short-term Access

To purchase short-term access, please sign in to your personal account above.

Don't already have a personal account? Register

The effect of severe psoriasis on the quality of life of 369 patients - 24 Hours access

EUR €58.00

GBP £50.00

USD \$62.00

Rental



This article is also available for rental through DeepDyve.



Email alerts

Article activity alert

Advance article alerts

New issue alert

Receive exclusive offers and updates from Oxford Academic

Citing articles via

Google Scholar

Latest Most Read Most Cited

Publications in the *BJD*: the impact of being read (and cited)

"When you do not feel comfortable in your skin, you cannot get out of it" – A Qualitative Interview Study exploring Psychosocial Impact and Coping Strategies among Patients with Prurigo Nodularis A loss-of-function variant in *KLF4* affecting zinc finger motifs causes progressive symmetric erythrokeratodermia

Measurement properties and interpretability of the Patient-Reported Impact of Dermatological Diseases (PRIDD) measure

Urticaria and the risk of cancer: a Danish population-based cohort study

More from Oxford Academic

Clinical Medicine Dermatology

Medicine and Health Books Journals

About British Journal of Dermatology

Editorial Board

Author Guidelines

Recommend to Your Librarian

Advertising and Corporate Services

Online ISSN 1365-2133 Print ISSN 0007-0963

Copyright © 2024 British Association of Dermatologists

Journals Career Network

About Oxford Academic

Publish journals with us

University press partners

What we publish

New features

Get help with access

Accessibility

Contact us

Advertising

Media enquiries

Authoring

Open access

Purchasing

Institutional account management

Rights and permissions

Oxford University Press

News

Oxford Languages

University of Oxford

Oxford University Press is a department of the University of Oxford. It furthers the University's objective of excellence in research, scholarship, and education by publishing worldwide

Copyright © 2024 Oxford University Press Cookie settings Cookie policy Privacy policy Legal notice