
Response	Percentage
Yes	10%
No	90%

<ul style="list-style-type: none">

<ul style="list-style-type: none">

<ul style="list-style-type: none">

<ul style="list-style-type: none">	

	<ul style="list-style-type: none">

<ul style="list-style-type: none">	
<ul style="list-style-type: none">	
<ul style="list-style-type: none">	

<ul style="list-style-type: none">

<ul style="list-style-type: none">

Table 8: Methotrexate

- Effective, but potential for life-threatening side effects
- Use of the NPF brochure on methotrexate helps educate patients about potential risks and can help physicians easily document that they have provided information on the risks of treatment
- Start with 5 mg test dose, increase in 5 mg/wk intervals until clinical response or maximal dose of 20-25 mg/week is achieved (reduce the initial and incremental doses to 2.5 mg in the elderly or others with suspected renal impairment)
- Check complete blood count and liver function tests one week after initial dose and after each increase in dose
- Trimethoprim-sulfamethoxazole also affects folic acid and should not be taken during methotrexate therapy. Non-steroidal anti-inflammatories can reduce renal function; increased monitoring for methotrexate toxicity is indicated when these agents are started during methotrexate therapy
- Be prepared to use folinic acid (not folic acid!) if needed to treat acute toxicity
- To reduce nausea, a divided dose may be taken one day a week. Folic acid (1-5 mg/day) may also be helpful.
- The cost of oral therapy may be reduced by use of the intravenous solution in place of the pill formulation

Laboratory studies should be repeated one week after every increase in dose of the medication. After a stable dose is reached, the frequency of the laboratory testing may be reduced. There are published recommendations for continued repeat blood tests 4 to 6 weeks, although such frequent testing is probably not necessary in patients on stable doses of methotrexate. A number of factors may increase methotrexate blood levels including concomitant administration of another inhibitor of folate metabolism (the commonly used antibiotic trimethoprim-sulfamethoxazole [Septra, Bactrim] and reduction of renal clearance, which may occur with nonsteroidal anti-inflammatories).[43] It is essential that patients be aware of these potential interactions, however it is not clear that the frequency of laboratory testing alone will prevent these events. Folinic acid (Leucovorin) may be given if acute hematologic toxicity occurs; it should be instituted as early as possible. Folic acid is not a replacement for folinic acid in this situation.

In a review of the literature most physicians did not document the history/physical exam findings on which they based giving methotrexate, nor did they carefully document informing their patients of the risks of this treatment.[44] Very helpful in this regard is the NPF brochure on methotrexate. Written in language for the lay person and reviewed by dermatologists, the brochure provides patients the necessary information on the risks of treatment. It is good medicine both to provide this information and to document that it was provided.

Liver biopsies are done to monitor for chronic hepatotoxicity that may not be revealed by blood tests. Pretreatment biopsy is probably unnecessary in patients without a history suggestive of preexisting hepatotoxicity.[42] If a baseline biopsy is to be done, it may be best to wait until it is clear that the patient will tolerate the drug (some do not due to severe nausea), that the drug is effective in controlling the patient's psoriasis (sometimes it is not), and that the drug will be needed long-term. Repeat liver biopsies may be done after approximately every 1.5 g cumulative dose. There is question whether liver biopsy needs to be done at all.[45] If in doubt, it may be wise to defer the decision to a gastroenterologist who specializes in liver disease and liver biopsy. There are certainly some scenarios in which the risk of liver biopsy is less than the risk of chronic methotrexate hepatotoxicity. Intolerance to methotrexate due to nausea can be extremely severe and sometimes may be managed by switching to intramuscular methotrexate or by administration of folic acid 1-5 mg/day.[46]

The cost of methotrexate therapy may be reduced by switching from pills to oral administration of the intravenous solution in appropriate patients. The cost is approximately \$100/100 mg of methotrexate pills versus \$15/100 mg for the solution (50mg methotrexate in 2cc vial). Accurate dosing is essential. Patients should consider purchasing insulin syringes when purchasing methotrexate and returning to the office for instructions on how much should be drawn up. The methotrexate solution can be put into a glass of water or flavored beverage and consumed.

Cyclosporine

Short-term use of methotrexate may be effective for some patients who have acute exacerbation of otherwise stable psoriasis

[47,48,49] Short-term use of cyclosporine allows very rapid control of psoriasis and is safer (though more expensive) than two tapers of oral corticosteroids. Administration of cyclosporine for as little as 5 days has been reported effective for the treatment of guttate psoriasis. Thus, cyclosporine may be best used in acute situations, followed by a switch to safer long-term treatments. Such an approach might include the initial use of cyclosporine to achieve rapid control followed by a transition to either acitretin or UVB phototherapy to maintain control.

Table 9: Cyclosporine

- Very useful for rapid, short-term control of psoriasis followed by transition to another treatment
- Blood pressure and creatinine measurements should be assessed twice before initiating treatment to establish accurate baseline values
- Elevation of blood pressure during cyclosporine therapy may be treated with antihypertensive medications
- Reduction in cyclosporine dose is recommended if blood creatinine increases by 30%
- Numerous potential drug interactions

Although cyclosporine is quite safe and effective in short-term use, it is associated with long-term toxicity including hypertension and decreased renal function[50,51] The prescribing dermatologist should be familiar with the 1998 consensus conference recommendations for monitoring cyclosporine therapy and delineates the numerous potential drug interactions[52] Among other interactions, increased renal toxicity may occur with nonsteroidal anti-inflammatories; increased cyclosporine levels may be seen with ketoconazole, cimetidine and macrolide antibiotics; and rhabdomyolysis may occur with HMG CoA reductase inhibitors.

Of all the oral agents for severe psoriasis, cyclosporine may be the safest for women who may become pregnant. Surprisingly the FDA lists cyclosporine as a pregnancy category B drug.

Other Treatments

Other immune inhibitors/modulators such as hydroxyurea may be used. The advantage of hydroxyurea is that it may be used in place of methotrexate in patients with cirrhosis. Monitoring for bone marrow toxicity is necessary.

Mycophenolate mofetil (CellCept) may be a helpful adjunctive immune modulator in psoriasis. The initial dose is 1 g by mouth daily. Doses of up to 3 to 4 g/day may be needed. Neutrophil counts should be followed closely, as severe neutropenia may occur in 2% of patients.

Summary

Psoriasis is a multifaceted disease and its treatment generally requires the expertise of dermatologists. The National Psoriasis Foundation is a tremendous resource for patient education and advocacy in psoriasis care. To plan for acute and long-term control of localized psoriasis, combinations of topical corticosteroids and either calcipotriene or tazarotene are the most effective approaches. For generalized disease, UVB treatment provides the safest means of achieving long-term control of disease. Acitretin is a very helpful adjunct for improving the efficacy of phototherapy. For patients with severe, refractory disease, methotrexate may be most effective, while cyclosporine may be most valuable for patients needing rapid, short-term improvement.

References

1. E. M. Farber and M. L. Nall. The natural history of psoriasis in 5,600 patients. *Dermatologica* 148 (1):1-18, 1974.
2. S. R. Rapp, M. L. Exum, D. M. Reboussin, S. R. Feldman, A. Fleischer, and A. Clark. The physical, psychological impact of psoriasis. *J Health Psychol* 2:525-537, 1997.
3. S. R. Rapp, S. R. Feldman, M. L. Exum, A. B. Fleischer, Jr., and D. M. Reboussin. Psoriasis causes as much disability as medical diseases. *J Am Acad Dermatol* 41 (3 Pt 1):401-407, 1999.
4. A. B. Fleischer, Jr., S. R. Feldman, S. R. Rapp, D. M. Reboussin, Exum, ML, A. R. Clark, and V. Rajashekhar. Disease severity measures in a population of psoriasis patients: the symptoms of psoriasis correlate with self-administered psoriasis severity index scores. *J Invest Dermatol* 107 (1):26-29, 1996.
5. A. B. Fleischer, Jr., S. R. Feldman, S. R. Rapp, D. M. Reboussin, M. L. Exum, and A. R. Clark. Alternative therapies used within a population of patients with psoriasis. *Cutis* 58 (3):216-220, 1996.

6. S. R. Feldman, A. B. Fleischer, Jr., D. M. Reboussin, S. R. Rapp, D. D. Bradham, M. L. Exum, and A. R. Clark. The association of psoriasis increases with psoriasis severity. *J Am Acad Dermatol* 37 (4):564-569, 1997.
7. S. R. Rapp, S. R. Feldman, A. B. Fleischer, Jr., D. M. Reboussin, and M. L. Exum. Health related quality of life in psoriasis: biopsychosocial model and measures. In: *Care Management of Skin Diseases: Life Quality and Economic Impact*, edited by F Rajagopalan, E. F. Sherertz, and R. Anderson, New York, New York:Marcel Dekker, Inc., 1998, p. 125-145.
8. S. R. Feldman and S. Al-Suwaidan. Clearance is not a realistic goal of psoriasis treatment. *J Am Acad Dermatol* 2000 in press.
9. I. E. Veronikis, A. O. Malabanan, and M. F. Holick. Comparison of calcipotriene (Dovonex) with a coal tar emulsion (Exorex) in treating psoriasis in adults: a pilot study [letter]. *Arch Dermatol* 135 (4):474-475, 1999.
10. O. Sanmartin. Skin Cap analysis in Spain [letter]. *Dermatol Online*.J 3 (2):11g, 1997.
11. J. K. Tan. Pustular psoriasis and hepatotoxicity associated with use of Skin Cap spray [letter]. *Dermatol Online*.J 3 (2):11d, 1997.
12. K. Smith. Skin Cap: what have we learned, and when did we learn it? [editorial]. *Dermatol Online*.J 3 (2):11c, 1997.
13. E. A. Olsen, D. L. Cram, C. N. Ellis, J. G. Hickman, C. Jacobson, E. E. Jenkins, A. E. Lasser, M. Lebwohl, R. S. Leisner, and et al. A double-blind, vehicle-controlled study of clobetasol propionate 0.05% (Temovate) scalp application in the treatment of moderate to severe scalp psoriasis. *J Am Acad Dermatol* 24 (3):443-447, 1991.
14. T. J. Franz, D. A. Parsell, R. M. Halualani, J. F. Hannigan, J. P. Kalbach, and W. S. Harkonen. Betamethasone foam 0.12%: a novel vehicle with enhanced delivery and efficacy. *Int J Dermatol* 38 (8):628-632, 1999.
15. S. R. Feldman, N. D. Sangha, and V. Setaluri. Topical corticosteroid in foam vehicle offers comparable coverage compared with traditional vehicles. *J Am Acad Dermatol* in press, 2000.
16. M. Lebwohl, S. B. Siskin, W. Epinette, D. Breneman, T. Funicella, R. Kalb, and J. Moore. A multicenter trial of calcipotriene ointment and halobetasol ointment compared with either agent alone for the treatment of psoriasis. *J Am Acad Dermatol* 35 (2 Pt 1):268-269, 1996.
17. M. Lebwohl. Topical application of calcipotriene and corticosteroids: combination regimens. *J Am Acad Dermatol* 37 (3 Pt 2):S55-8, 1997.
18. M. Lebwohl, A. Yoles, K. Lombardi, and W. Lou. Calcipotriene ointment and halobetasol ointment in the long-term treatment of psoriasis: Effects on the duration of improvement. *J Am Acad Dermatol* 39:447-450, 1998.
19. B. Patel, S. Siskin, R. Krazmien, and M. Lebwohl. Compatibility of calcipotriene with other topical medications. *Arch Dermatol* 38 (6 Pt 1):1010-1011, 1998.
20. D. Hecker and M. Lebwohl. Topical calcipotriene in combination with UVB phototherapy for psoriasis. *Int J Dermatol* 37 (3):300-303, 1997.
21. M. Lebwohl, D. Hecker, J. Martinez, A. Sapadin, and B. Patel. Interactions between calcipotriene and ultraviolet light. *J Am Acad Dermatol* 37 (1):93-95, 1997.
22. G. D. Weinstein, G. G. Krueger, N. J. Lowe, M. Duvic, D. J. Friedman, B. V. Jegasothy, J. L. Jorizzo, E. Shih, D. A. Tischenko, D. A. Lew-Kaya, J. C. Lue, J. Sefton, J. R. Gibson, and R. A. Chandraratna. Tazarotene gel, a new retinoid, for the treatment of psoriasis: vehicle-controlled study of safety, efficacy, and duration of therapeutic effect. *J Am Acad Dermatol* 37 (1):85-92, 1997.
23. G. G. Krueger, L. A. Drake, P. M. Elias, N. J. Lowe, C. Guzzo, G. D. Weinstein, D. A. Lew-Kaya, J. C. Lue, J. Sefton, and R. A. Chandraratna. The safety and efficacy of tazarotene gel, a topical acetylenic retinoid, in the treatment of psoriasis. *J Am Acad Dermatol* 34 (1):57-60, 1998.
24. M. Lebwohl, E. Ast, J. P. Callen, S. I. Cullen, S. R. Hong, C. L. Kulp-Shorten, N. J. Lowe, T. J. Phillips, B. T. Wofford, M. Quell, J. Sefton, J. C. Lue, J. R. Gibson, and R. A. Chandraratna. Once-daily tazarotene gel versus twice-daily triamcinolone cream in the treatment of plaque psoriasis. *J Am Acad Dermatol* 38 (5 Pt 1):705-711, 1998.
25. M. G. Lebwohl, D. L. Breneman, B. S. Goffe, J. R. Grossman, M. R. Ling, J. Milbauer, and et al. Tazarotene 0.1% gel plus corticosteroid cream in the treatment of plaque psoriasis. *J Am Acad Dermatol* 39:590-596, 1998.
26. J. Koo. Tazarotene in combination with phototherapy. *J Am Acad Dermatol* 39:S144-148, 1998.
27. W. Torinuki and H. Tagami. Incidence of skin cancer in Japanese psoriatic patients treated with either methoxsalen

phototherapy, Goeckerman regimen, or both therapies. A 10-year follow-up study. *J Am Acad Dermatol* 18 (6):1278-1281, 1988.

28. S. A. Muller and H. O. Perry. The Goeckerman treatment in psoriasis: six decades of experience at the Mayo Clinic. *Cutis* (3):265-268, 1927.

29. S. N. Horwitz, R. A. Johnson, J. Sefton, and P. Frost. Addition of a topically applied corticosteroid to a modified Goeckerman regimen for treatment of psoriasis: effect on duration of remission. *J Am Acad Dermatol* 13 (5 Pt 1):784-791, 1985.

30. D. H. Cort, N. R. Schleider, R. S. Moskowitz, S. N. Horwitz, and P. Frost. Retrospective analysis of a modified Goeckerman for the treatment of psoriasis. *Cutis* 25 (2):201-203, 1978.

31. S. R. Feldman, A. Clark, D. M. Reboussin, and A. B. Fleischer, Jr. An assessment of potential problems of home phototherapy of psoriasis. *Cutis* 58 (1):71-73, 1996.

32. A. B. Fleischer, Jr., A. R. Clark, S. R. Rapp, D. M. Reboussin, and S. R. Feldman. Commercial tanning bed treatment of psoriasis: results from an uncontrolled clinical trial. *J Invest Dermatol* 109 (2):170-174, 1997.

33. R. S. Stern and N. Laird. The carcinogenic risk of treatments for severe psoriasis. *Photochemotherapy Follow-up Study*. *Cutis* 73 (11):2759-2764, 1994.

34. R. S. Stern, K. T. Nichols, and L. H. Vakeva. Malignant melanoma in patients treated for psoriasis with methoxsalen (psoralen) and ultraviolet A radiation (PUVA). *New Engl J Med* 336:1041-1045, 1997.

35. C. E. Orfanos, C. C. Zouboulis, B. Almond-Roesler, and C. C. Geilen. Current use and future potential role of retinoids in dermatology. [Review] [262 refs]. *Drugs* 53 (3):358-388, 1997.

36. H. E. Murray, A. W. Anhalt, R. Lessard, R. K. Schacter, J. B. Ross, W. D. Stewart, and J. M. Geiger. A 12-month study of severe psoriasis with acitretin: results of a Canadian open multicenter study. *J Am Acad Dermatol* 24 (4):598-602, 1991.

37. R. S. Stern, E. Fitzgerald, C. N. Ellis, N. Lowe, M. T. Goldfarb, and R. D. Baughman. The safety of etretinate therapy for psoriasis: results of the etretinate follow-up study. *J Am Acad Dermatol* 33 (1):44-52, 1995.

38. R. C. Chou, R. Wyss, C. A. Huselton, and U. W. Wiengand. A potentially new metabolic pathway: ethylesterification of acil. *Xenobiotica* 22:993-996, 1992.

39. A. Anstey and J. L. Hawk. Isotretinoin-PUVA in women with psoriasis [letter]. *Br J Dermatol* 136 (5):798-799, 1997.

40. N. J. Lowe, J. H. Prystowsky, T. Bourget, J. Edelstein, S. Nychay, Armstrong, and R. Acitretin plus UVB therapy for psoriasis: Comparisons with placebo plus UVB and acitretin alone. *J Am Acad Dermatol* 24 (4):591-594, 1991.

41. A. Tanew, A. Guggenbichler, H. Honigsmann, J. M. Geiger, and P. Fritsch. Photochemotherapy for severe psoriasis with combination with acitretin: a randomized, double-blind comparison study. *J Am Acad Dermatol* 25 (4):682-684, 1991.

42. H. H. Roenigk, Jr., R. Auerbach, H. Maibach, G. Weinstein, and M. Lebwohl. Methotrexate in psoriasis: consensus conference. *Am Acad Dermatol* 38 (3):478-485, 1998.

43. B. Mayall, G. Poggi, and J. D. Parkin. Neutropenia due to low-dose methotrexate therapy for psoriasis and rheumatoid arthritis may be fatal. *Medical Journal of Australia* 155 (7):480-484, 1991.

44. D. J. Bilsland, L. E. Rhodes, I. Zaki, S. M. Wilkinson, K. E. McKenna, S. E. Handfield-Jones, and R. E. Williams. PUVA and methotrexate therapy of psoriasis: how closely do dermatology departments follow treatment guidelines? Psoriasis Audit Workgroup of the British Association of Dermatologists. *Br J Dermatol* 131 (2):220-225, 1994.

45. M. J. Boffa, R. J. Chalmers, N. Y. Haboubi, M. Shomaf, and D. M. Mitchell. Sequential liver biopsies during long-term methotrexate treatment for psoriasis: a reappraisal. *Br J Dermatol* 133 (5):774-778, 1995.

46. P. Duhra. Treatment of gastrointestinal symptoms associated with methotrexate therapy for psoriasis. *J Am Acad Dermatol* 28 (3):466-469, 1993.

47. E. Christophers, U. Mrowietz, H. H. Henneicke, L. Farber, and D. Welzel. Cyclosporine in psoriasis: a multicenter dose-finding study in severe plaque psoriasis. The German Multicenter Study. *J Am Acad Dermatol* 26 (1):86-90, 1992.

48. C. N. Ellis, M. S. Fradin, J. M. Messana, M. D. Brown, M. T. Siegel, A. H. Hartley, L. L. Rocher, S. Wheeler, T. A. Hamilton, Parish, and et al. Cyclosporine for plaque-type psoriasis. Results of a multidose, double-blind trial [see comments]. *New Engl J Med* 324 (5):277-284, 1991.

49. J. Koo and J. Lee. Cyclosporine: what clinicians need to know. [Review] [69 refs]. Dermatologic Clinics 13 (4):897-907, 1997.
50. A. L. Brown, R. Wilkinson, T. H. Thomas, N. Levell, C. Munro, J. Marks, and T. H. Goodship. The effect of short-term low-dose cyclosporin on renal function and blood pressure in patients with psoriasis. Br J Dermatol 128 (5):550-555, 1993.
51. H. Zachariae, K. Kragballe, H. E. Hansen, N. Marcussen, Olsen, and S. Renal biopsy findings in long-term cyclosporin treatment of psoriasis [see comments]. Br J Dermatol 136 (4):531-535, 1997.
52. H. I. Katz. Dermatologist's Guide to Adverse Therapeutic Interactions, Philadelphia, PA:Lippincott-Raven, 1997.

© 2000 Dermatology Online Journal

Jump To

Article

└ Main Content

Metrics

Author & Article Info

Related Items

[A case of melanoma in situ of the central forehead repair by V-Y advancement flap and Burow graft](#)

Leigh,
Diamond, Carrie
Rebekah Rundle, Chandler
Flynn, M Seth
Stamey, Christopher

[Improving pediatric atopic dermatitis care in the emergency department and primary care setting: a collaborative quality improvement pilot project](#)

Haller, Courtney
Paladichuk, Han
Shivaprakash, Nanditha
N; Diaz, Lucia Z
Wilkinson, Matthew
Levy, Moise J
Iyer, Sujit

[A patient with acute-type adult T-cell leukemia/lymphoma that rapidly progressed after Helicobacter pylori eradication therapy](#)

Matsudate,
Yoshihiro
Saeki, Kyosuke

[Extragenital lichen sclerosus et atrophicus-morphea overlap as an initial presentation of genital lichen sclerosus](#)

Trager, Megan
Gordon, Emily R
Adeuyan, Oluwaseyi
Fahmy, Lauren M
Schreidah, Celine M;
Hussain, Sameem
Goskin, Larissa I
Lapolla, Brigit A

[Corrigendum: Spindle cell lipoma: a rare case report on the hallux](#)

Dahle, Sara
Liu, Caroline C
Schulman, Joshua M;
Alley, Shelby M

Top

[Home](#)

[About eScholarship](#)

[Campus Sites](#)

[UC Open Access Policy](#)

[eScholarship Publishing](#)

[Accessibility](#)

[Privacy Statement](#)

[Site Policies](#)

[Terms of Use](#)

[Admin Login](#)

[Help](#)

Powered by the
California Digital Library
Copyright © 2017
The Regents of the University of California