

Pediatric P-08 Pediatric Seizure

Priorities

Primary Survey
Prevent trauma
Early notification of receiving hospital
Treat fever if necessary

Stabilize airway using appropriate BLS adjuncts

Oxygen – High flow if tolerated
Pulse Oximetry
Cardiac Monitor
IV/IO Access
Check Blood Glucose

If patient is actively seizing

Midazolam 0.1mg/kg IV/IO/IM/IN to a max single dose of 4mg
May repeat x1 if seizure persists or returns after 2 minutes from initial dose.

Base Hospital Physician Order needed for any additional doses.

For blood glucose <50mg/dl in patients <1 year old

Dextrose 10% 0.5g/kg (5mL/kg) IV/IO
to max dose of 25g (250mL)

For blood glucose <60m/dl in patients 1-14 years old

Dextrose 10% 0.5g/kg (5mL/kg) IV/IO
to max dose of 25g (250mL)

For fever >100.4°F/38°C and patient is alert and has gag reflex:

Consider **Acetaminophen** 15mg/kg PO
Max dose 650mg PO

Contraindications to Acetaminophen

- Liver disease
- Taken acetaminophen-containing products within last 4 hours
- Infant: Taken >45mg/kg in past 24 hours
- Child <40kg: Taken >60mg/kg in past 24 hours
- Child >40kg: Taken >2,400mg in past 24 hours

Dextrose Dosage Calculations

Dextrose 10% is the preferred concentration. If Dextrose 10% is not available, please use the doses below for other concentrations.

D10% 5mL/kg (max 250mL)
0.5g/kg (max 25g) = D25% 2mL/kg (max 100mL)
D50% 1mL/kg (max 50mL)

DISRUPTED COMMUNICATIONS

In the event of a “disrupted communications” situation, Solano County Paramedics may NOT utilize all portions of this treatment protocol without Base Hospital Contact as needed to stabilize an immediate patient.