





We go from right to left






we have this initial drawing






v_0








$\forall x \in]x(m), x(M)] \exists y \text{ of } B_1 \text{ at } x \text{ such that}$
 $y < y\text{-value of } x$

$\Rightarrow$ L.P. logic with the counter-example






Ψ of $S(\Gamma)$

ab	ac	ad	ae	bc	bd	be	cd	ce	de
$\uparrow$	$\uparrow$	$\checkmark$	$\checkmark$	$\uparrow$	$\checkmark$	$\checkmark$	$\times$	$\times$	A
fb	fh	fi	fj	gl	gi	gj	hi	hj	ij
$\uparrow$	$\uparrow$	$\uparrow$	$\checkmark$	$\uparrow$	$\uparrow$	$\checkmark$	$\uparrow$	$\checkmark$	$\times$
kel									
$\uparrow$									

no of the solution



1	1	0	0	1	1	1	0	0	0
1	1	1	0	1	1	1	1	1	0
1									






1 1 0 0 1 1 1 0 0 0
 1 1 1 0 1 1 1 1 0
 1

ab ac ad ae bc bd be cf ce de
 f g h i j k l m n o p q r s t u v w x y z
 A
 f g h i j k l m n o p q r s t u v w x y z
 A



The stretched edges represent the transitivity
 (we can't have the last comp of h -an (transitive))

because there is always a stretch edge
after interval edges;



$\Rightarrow$ we can draw
an equivalent
 k -monotone
embedding.
(no edge changes
whether it passes from
above or below a point)

I have to think
about it more