

BPROAD Trial Critical Appraisal

Intensive systolic BP target <120 mm Hg versus standard target <140 mm Hg in type 2 diabetes

Journal club appraisal

RCT

The NEW ENGLAND JOURNAL *of* MEDICINE

ESTABLISHED IN 1812

MARCH 27, 2025

VOL. 392 NO. 12

Intensive Blood-Pressure Control in Patients with Type 2 Diabetes

Y. Bi,^{1,2} M. Li,^{1,2} Y. Liu,³ T. Li,⁴ J. Lu,^{1,2} P. Duan,⁵ F. Xu,⁶ Q. Dong,⁷ Ailiang Wang,⁸ T. Wang,^{1,2} R. Zheng,^{1,2} Y. Chen,^{1,2} M. Xu,^{1,2} X. Wang,⁹ Xinhuan Zhang,¹⁰ Y. Niu,¹¹ Z. Kang,¹² C. Lu,¹³ Jing Wang,¹⁴ X. Qiu,¹⁵ An Wang,¹⁶ S. Wu,^{1,2,17} J. Niu,^{1,2,18} Jingya Wang,¹⁹ Z. Zhao,^{1,2} H. Pan,²⁰ X. Yang,²¹ X. Niu,²² S. Pang,²³ Xiaoliang Zhang,²⁴ Y. Dai,²⁵ Q. Wan,²⁶ S. Chen,²⁷ Q. Zheng,²⁸ S. Dai,²⁹ J. Deng,^{1,2} L. Liu,³⁰ G. Wang,³¹ H. Zhu,³² W. Tang,³³ H. Liu,³⁴ Z. Guo,³⁵ G. Ning,^{1,2} J. He,³⁶ Y. Xu,^{1,2} and W. Wang,^{1,2} for the BPROAD Research Group*

Trial Identity and Citation

TRIAL NAME & CITATION

BPROAD Trial: Intensive Blood-Pressure Control in Patients with Type 2 Diabetes

Citation: Bi Y et al. Intensive Blood-Pressure Control in Patients with Type 2 Diabetes. N Engl J Med 2025;392:1155-1167.

FUNDING & RESEARCH GROUPS

- Funded by the National Key Research and Development Program of the Ministry of Science and Technology of China and others.
- Conducted by the BPROAD Research Group across 145 clinical sites in China.
- Independent international data and safety monitoring board monitored design, data quality, and patient safety.

TRIAL REGISTRATION & SPONSORS

- ClinicalTrials.gov: NCT03808311.
- Parallel-design randomized clinical trial; article published online November 16, 2024 and in NEJM March 27, 2025.
- Authors state that trial and data coordinating center personnel performed analyses.

Executive Summary / Practice Bottom Line

12,821

randomized adults with T2D

4.2 yr

median follow-up

HR 0.79

primary composite, 95% CI 0.69-0.90

NNT ~65

event-count ARR 1.55%

- Clinical question: In older adults with type 2 diabetes, elevated SBP, and increased cardiovascular risk, does targeting SBP <120 mm Hg reduce major cardiovascular events compared with <140 mm Hg?
- Primary endpoint: first nonfatal stroke, nonfatal MI, treatment or hospitalization for heart failure, or cardiovascular death.
- Primary result: 393/6414 versus 492/6407 events; HR 0.79 (95% CI 0.69-0.90; P<0.001).
- Harms: serious adverse events similar; symptomatic hypotension and serum potassium >5.5 mmol/L were more frequent with intensive treatment.
- Practice bottom line: consider in selected high-risk, closely monitored T2D patients; do not apply blindly to frail older adults, low-risk patients, advanced CKD, orthostatic symptoms, or poor monitoring settings.

Why This Paper Matters

Clinical problem: Clinical problem: Hypertension is common in type 2 diabetes and strongly drives stroke, heart failure, MI, CKD, and cardiovascular death.

Evidence gap: Evidence gap: ACCORD did not show significant overall cardiovascular benefit for SBP <120 mm Hg in diabetes, while SPRINT excluded diabetes.

Research question: Research question: BPROAD directly tests whether the SPRINT-like SBP target helps patients with type 2 diabetes and elevated cardiovascular risk.

The NEW ENGLAND JOURNAL *of* MEDICINE

ESTABLISHED IN 1812

MARCH 27, 2025

VOL. 392 NO. 12

Intensive Blood-Pressure Control in Patients with Type 2 Diabetes

Y. Bi,^{1,2} M. Li,^{1,2} Y. Liu,³ T. Li,⁴ J. Lu,^{1,2} P. Duan,⁵ F. Xu,⁶ Q. Dong,⁷ Ailiang Wang,⁸ T. Wang,^{1,2} R. Zheng,^{1,2} Y. Chen,^{1,2} M. Xu,^{1,2} X. Wang,⁹ Xinhuan Zhang,¹⁰ Y. Niu,¹¹ Z. Kang,¹² C. Lu,¹³ Jing Wang,¹⁴ X. Qiu,¹⁵ An Wang,¹⁶ S. Wu,^{1,2,17} J. Niu,^{1,2,18} Jingya Wang,¹⁹ Z. Zhao,^{1,2} H. Pan,²⁰ X. Yang,²¹ X. Niu,²² S. Pang,²³ Xiaoliang Zhang,²⁴ Y. Dai,²⁵ Q. Wan,²⁶ S. Chen,²⁷ Q. Zheng,²⁸ S. Dai,²⁹ J. Deng,^{1,2} L. Liu,³⁰ G. Wang,³¹ H. Zhu,³² W. Tang,³³ H. Liu,³⁴ Z. Guo,³⁵ G. Ning,^{1,2} J. He,³⁶ Y. Xu,^{1,2} and W. Wang,^{1,2} for the BPROAD Research Group*

Clinical Question & PICO Framework

P — Adults age ≥ 50 years with type 2 diabetes, elevated systolic BP, and increased cardiovascular risk.

I — Intensive antihypertensive strategy targeting SBP < 120 mm Hg.

C — Standard antihypertensive strategy targeting SBP < 140 mm Hg.

O — Primary composite major cardiovascular outcome over up to 5 years.

Design and Trial Schema

145

clinical sites in 7 Chinese regions

1:1

central web randomization

<120

intensive SBP target

<140

standard SBP target

- Superiority, parallel-group randomized clinical trial.
- Randomization stratified by clinical site with randomly selected block sizes of 2, 4, and 6.
- Patients and treating physicians were not blinded; outcome assessors, adjudicators, and statisticians were blinded.
- Blood-pressure treatment algorithms were similar to SPRINT; glucose and lipid care followed current clinical guidelines.

INTRODUCTION

Clinical need and rationale

Introduction Summary

- Elevated SBP is common in diabetes and is a major modifiable cardiovascular risk factor.
- Guidelines recommend BP reduction in diabetes, but the best SBP target has been uncertain.
- ACCORD was underpowered for BP intervention and complicated by factorial glycemic-control interactions.
- SPRINT supported intensive BP control but excluded diabetes; BPROAD fills this evidence gap.

Introduction Critical Appraisal

Strengths

- Clear clinical uncertainty after ACCORD and SPRINT.
- Outcome is cardiovascular morbidity/mortality, not a surrogate.
- Large event-driven design makes a clinically useful answer plausible.

Concerns

- Prior evidence largely from non-diabetes or underpowered diabetes subgroups.
- Expected benefit may vary by ethnicity, baseline stroke risk, monitoring quality, and medication access.

The rationale is strong: a common clinical decision with prior uncertainty and patient-important outcomes.

METHODS

Design, eligibility, endpoints, and analysis

Methods Overview

90%

planned power

0.045

nominal alpha after interim look

50

imputed datasets for missing outcomes

ITT

all randomized patients analyzed

- Sample size assumed a standard-arm primary-event rate of 2.0% per year and 20% effect size.
- Cox proportional-hazards model adjusted for geographic region.
- Proportional-hazards assumption was tested using Schoenfeld residuals.
- One interim analysis occurred when 54.5% of adjudicated primary events had accrued.

Eligibility and Setting

Included

- Age ≥ 50 years with type 2 diabetes.
- Elevated SBP: 130-180 mm Hg on antihypertensives or ≥ 140 mm Hg without antihypertensives.
- Increased cardiovascular risk: clinical/subclinical CVD, multiple risk factors, or CKD with eGFR 30 to <60 .

Implications

- Mean age 63.8 years; 45.3% women; 22.5% had clinical CVD.
- Only 7.6% had eGFR <60 ml/min/1.73 m² at baseline.
- Generalizability is strongest for monitored, high-risk outpatient diabetes care.

This is a high-risk diabetes population, not a trial of all adults with diabetes and mild hypertension.

Randomisation / Comparator / Blinding

Internal validity supports

- Central web-based randomization with site stratification.
- Outcome assessment, adjudication, and statistical analysis were blinded.
- Background glucose and lipid care were similar during follow-up.

Residual concerns

- Patients and treating clinicians knew treatment assignment.
- More visits and medication changes may alter detection, adherence, and co-interventions.
- Allocation concealment appears adequate, but individual block details by site are not deeply visible in the main paper.

Open-label treatment is unavoidable for BP targets but still leaves room for performance bias.

Outcomes and Endpoint Hierarchy

Primary Composite

- Nonfatal stroke
- Nonfatal MI
- Treatment or hospitalization for heart failure
- Cardiovascular death

Secondary Clinical

- Individual primary components
- Death from any cause
- Expanded composite with all-cause death
- CKD progression, CKD development, incident albuminuria

Safety

- Serious adverse events
- Hypotension, syncope, falls
- Renal failure
- Electrolyte alerts

Endpoint Integrity

- Primary endpoint is patient-important and clinically coherent, although the components differ in seriousness and frequency.
- The benefit appears driven mainly by stroke reduction; MI, heart failure, and cardiovascular death point in the same direction but are less precise.
- Outcome documents were centrally and blindly adjudicated.
- Secondary outcomes and subgroups were not adjusted for multiplicity and should be interpreted as supportive or hypothesis-generating.
- Composite endpoint is acceptable, but practice interpretation should focus on absolute stroke/CV event prevention.

Primary endpoint drives the trial claim; secondary/subgroup results should not be over-sold.

Analysis Sets, Missing Data, Estimand

1.2%

discontinued intervention

4.7%

lost to follow-up

3.3%

withdrew consent

MAR

primary missing-data assumption

- Primary analysis followed intention-to-treat using all randomized patients.
- Multiple imputation assumed missing outcomes were missing at random.
- Sensitivity analyses included censoring missing data, not-missing-at-random assumptions, and Fine-Gray competing-risk modelling.
- Clinical estimand: effect of assignment to an intensive BP-treatment strategy, not the pure biologic effect of SBP alone.

Statistics: Multiplicity, Interim, Time-to-Event Caveats

- Primary outcome was time-to-first event analyzed by Cox regression; HR should not be read as a simple risk ratio.
- Nominal alpha was updated to 0.045 after one interim analysis using Lan-DeMets alpha spending.
- Secondary outcome CIs were not adjusted for multiplicity and cannot support confirmatory claims.
- The Kaplan-Meier curves separate after about 1 year; number at risk becomes very small near year 5.
- Proportional-hazards assumption was reported as tested and confirmed, but RMST was not reported in the main article.

Methods Critical Appraisal

What strengthens trust

- Large sample and event-driven planning.
- Central randomization with blinded adjudication.
- Protocol, statistical analysis plan, and prespecified subgroups reported.

What tempers trust

- Open-label treatment can influence care intensity and symptom reporting.
- COVID-era telephone follow-up and home BP measurements may affect measurement uniformity.
- Only about 60% in intensive arm reached SBP target after 1 year.

Methods are robust for a pragmatic BP-target RCT; main concerns are open-label care and reliance on missing-at-random imputation.

RESULTS

Population, outcomes, figures, and harms

Table 1A. Baseline Characteristics

Table 1. Baseline Characteristics of the Patients.*		
Characteristic	Intensive Treatment (N = 6414)	Standard Treatment (N = 6407)
Female sex — no. (%)	2923 (45.6)	2880 (45.0)
Age — yr	63.7±7.4	63.9±7.5
Age group — no. (%)		
<65 yr	3607 (56.2)	3500 (54.6)
≥65 yr	2807 (43.8)	2907 (45.4)
Systolic blood pressure — mm Hg	140.0±10.2	140.4±10.2
Diastolic blood pressure — mm Hg	76.3±9.2	76.3±9.1
Duration of hypertension — yr	11.8±9.4	11.6±9.4
Duration of diabetes — yr	10.3±7.6	10.3±7.7
History of clinical cardiovascular disease — no. (%)	1480 (23.1)	1408 (22.0)
History of subclinical cardiovascular disease — no. (%)	2173 (33.9)	2251 (35.1)
Educational level: high school diploma or greater — no./total no. (%)	3090/6401 (48.3)	3045/6396 (47.6)
Current smoker — no./total no. (%)	1580/6401 (24.7)	1636/6392 (25.6)
Body-mass index†	26.7±3.2	26.7±3.3
Waist circumference — cm	95.2±8.2	95.2±8.1
Fasting plasma glucose — mg/dl	148.7±48.1	148.4±48.0
Glycated hemoglobin — %	7.6±1.4	7.6±1.4
Total cholesterol — mg/dl	157.8±48.1	157.2±47.8
LDL cholesterol — mg/dl	84.3±34.6	83.7±34.1
HDL cholesterol — mg/dl	40.2±12.7	40.1±11.8
Median triglycerides (IQR) — mg/dl	130.2 (92.1–192.2)	131.1 (92.1–194.0)
eGFR — ml/min/1.73 m²‡	88.6±17.8	88.7±18.0
eGFR of <60 ml/min/1.73 m² — no. (%)‡	501 (7.8)	469 (7.3)
Median urinary albumin-to-creatinine ratio (IQR)§	20.5 (9.6–61.4)	19.3 (9.5–56.9)
Urinary albumin-to-creatinine ratio of ≥30 — no./total no. (%)§	2528/6378 (39.6)	2452/6378 (38.4)
Medications at baseline		
Any antihypertensive drug — no. (%)	6356 (99.1)	6338 (98.9)
No. of antihypertensive drugs per participant	1.4±0.6	1.4±0.6
Antihypertensive drugs — no./total no. (%)		
ACE inhibitor	904/6395 (14.1)	885/6386 (13.9)
Angiotensin-receptor blocker	2790/6395 (43.6)	2814/6386 (44.1)
Calcium-channel blocker	3774/6395 (59.0)	3780/6386 (59.2)
Diuretic	471/6395 (7.4)	447/6386 (7.0)
α-Receptor blocker	45/6395 (0.7)	46/6386 (0.7)
β-Receptor blocker	960/6395 (15.0)	906/6386 (14.2)
Other	206/6395 (3.2)	198/6386 (3.1)
Any hypoglycemic drug — no. (%)	6321 (98.6)	6294 (98.2)
Hypoglycemic drugs — no./total no. (%)		
Insulin	3114/6394 (48.7)	3069/6385 (48.1)

Baseline characteristics of the patients, first part.

Bi Y et al. Intensive Blood-Pressure Control in Patients with Type 2 Diabetes. N Engl J Med 2025;392:1155-1167.

Table 1B. Baseline Characteristics Continued

Table 1. (Continued.)		
Characteristic	Intensive Treatment (N = 6414)	Standard Treatment (N = 6407)
Metformin	4228/6395 (66.1)	4287/6385 (67.1)
Thiazolidinedione	164/6395 (2.6)	178/6385 (2.8)
Sulfonylurea	961/6395 (15.0)	949/6385 (14.9)
α-Glucosidase inhibitor	2170/6395 (33.9)	1976/6385 (30.9)
GLP-1 receptor agonist	283/6395 (4.4)	277/6385 (4.3)
DPP-4 inhibitor	642/6395 (10.0)	641/6385 (10.0)
SGLT2 inhibitor	674/6395 (10.5)	658/6385 (10.3)
Statin — no. (%)	4192 (65.4)	4159 (64.9)
Aspirin — no. (%)	3225 (50.3)	3200 (49.9)

* Plus-minus values are means ±SD. To convert plasma glucose values to millimoles per liter, multiply by 0.0555. To convert total, low-density lipoprotein (LDL), and high-density lipoprotein (HDL) cholesterol values to millimoles per liter, multiply by 0.0259. To convert triglyceride values to millimoles per liter, multiply by 0.0113. ACE denotes angiotensin-converting enzyme, DPP-4 dipeptidyl peptidase 4, GLP-1 glucagon-like peptide-1, IQR interquartile range, and SGLT2 sodium–glucose cotransporter 2.

† The body-mass index is the weight in kilograms divided by the square of the height in meters.

‡ The estimated glomerular filtration rate (eGFR) was based on the serum creatinine level and was calculated with the use of the Chronic Kidney Disease Epidemiology Collaboration equation.

§ The albumin-to-creatinine ratio was calculated as the ratio of urinary albumin (in milligrams) to urinary creatinine (in grams).

Baseline Comparability Appraisal

- Groups were well balanced for age, sex, baseline SBP/DBP, diabetes duration, antihypertensive use, kidney function, albuminuria, statin use, and aspirin use.
- Small numerical differences in prior clinical/subclinical CVD and smoking are unlikely to explain the primary result.
- No major clinically meaningful baseline imbalance is apparent in the main table.
- Baseline balance does not prove randomization quality, but it supports the absence of obvious allocation failure.

Do not over-read baseline p values; the practical question is whether any imbalance can plausibly account for the effect.

Figure 1. Systolic Blood Pressure Separation




Figure 1. Systolic Blood Pressure throughout the Trial.
Shown are the mean systolic blood-pressure values among patients who received intensive treatment that targeted a systolic blood pressure of less than 120 mm Hg and among patients who received standard treatment that targeted a systolic blood pressure of less than 140 mm Hg. I bars represent 95% confidence intervals.

BP Separation and Treatment Fidelity

121.6

mean SBP at 1 year, intensive

133.2

mean SBP at 1 year, standard

60%

intensive arm at target after 1 year

2.2

mean BP meds later in intensive arm

- The trial achieved a sustained between-group SBP separation of roughly 11-12 mm Hg.
- Intensive treatment required more antihypertensive medication.
- Incomplete target attainment makes the result more pragmatic: benefit occurred despite imperfect achievement of <120 mm Hg.
- In practice, the intervention is a monitored treatment strategy, not simply writing a lower target in the chart.

Table 2. Primary Outcome and Main Secondary Outcomes

Table 2. Primary Outcome and Main Secondary Outcomes.*						
Outcome	Intensive Treatment (N = 6414)		Standard Treatment (N = 6407)		Hazard Ratio (95% CI)†	P Value‡
	No. of Events	Incidence Rate	No. of Events	Incidence Rate		
		<i>no. of events/100 person-yr</i>		<i>no. of events/100 person-yr</i>		
Primary outcome: nonfatal stroke, nonfatal MI, treatment or hospitalization for heart failure, or death from cardiovascular causes	393	1.65 (1.50–1.82)	492	2.09 (1.91–2.28)	0.79 (0.69–0.90)	<0.001
Secondary outcomes						
Fatal or nonfatal MI	68	0.28 (0.22–0.35)	81	0.33 (0.27–0.41)	0.84 (0.60–1.16)	—
Fatal or nonfatal stroke	284	1.19 (1.06–1.33)	356	1.50 (1.35–1.66)	0.79 (0.67–0.92)	—
Treatment or hospitalization for heart failure	31	0.13 (0.09–0.18)	46	0.19 (0.14–0.25)	0.66 (0.41–1.04)	—
Death from cardiovascular causes	60	0.24 (0.19–0.31)	79	0.32 (0.26–0.40)	0.76 (0.55–1.06)	—
Death from any cause	169	0.69 (0.59–0.80)	179	0.73 (0.63–0.84)	0.95 (0.77–1.17)	—
Primary-outcome event or death from any cause	493	2.07 (1.90–2.26)	584	2.48 (2.28–2.69)	0.83 (0.74–0.94)	—
CKD outcomes						
CKD progression	24	1.61 (1.08–2.41)	16	1.11 (0.68–1.80)	1.36 (0.71–2.59)	—
CKD development	232	1.14 (1.00–1.29)	214	1.05 (0.92–1.20)	1.11 (0.92–1.34)	—
Incident albuminuria	554	11.29 (10.39–12.27)	648	13.84 (12.81–14.95)	0.87 (0.77–0.97)	—


* Patients were counted only once for each outcome. Confidence intervals for outcomes other than the primary outcome have not been adjusted for multiplicity and may not be used for hypothesis testing. CKD denotes chronic kidney disease, and MI myocardial infarction.

Primary outcome and main secondary outcomes; secondary-outcome CIs not adjusted for multiplicity.

Figure 2. Kaplan-Meier Curves for the Primary Outcome

nts
or
CI,
ps.⁵
out-
the
ate
vs.
cal
eat-
jor
ear
d a
sta-
nci-
rd-
rs)

on-
red
ysis
as-
ing
ce-
red
ing



Effect Size Interpretation

6.13%

primary events, intensive

7.68%

primary events, standard

1.55%

event-count absolute risk
reduction

0.44

fewer events/100 person-years

- Prespecified estimate: HR 0.79 (95% CI 0.69-0.90) for the primary composite.
- Approximate event-count NNT: $1 / 0.0155 = 65$ over median 4.2 years.
- Stroke event-count ARR: about 1.13%; approximate NNT 89.
- All-cause mortality was not clearly reduced: HR 0.95 (95% CI 0.77-1.17).
- Because the primary analysis is time-to-event, event-count NNT is a clinical approximation, not the trial estimand.

Precision and Uncertainty

- Primary outcome CI excludes no effect and is clinically meaningful: HR 0.69-0.90.
- Stroke reduction is directionally consistent and statistically precise, but remains a secondary outcome.
- MI, heart failure, cardiovascular death, CKD progression, and CKD development are imprecise individually.
- Incident albuminuria was lower with intensive therapy, but kidney clinical outcomes did not clearly improve.
- The near-tail Kaplan-Meier numbers at risk are small; avoid interpreting late-curve details.

The most defensible claim is fewer major cardiovascular composite events, probably with a substantial stroke component.

Figure 3. Prespecified Subgroup Analysis




Figure 3. Prespecified Subgroup Analysis. Shown are the effects of the interventions on the primary composite outcome across prespecified subgroups. Data on the glycated hemoglobin level were missing for 94 patients in the intensive-treatment group and for 80 patients in the standard-treatment group; on the duration of diabetes for 30 and 37, respectively; and on the duration of high blood pressure for 36 and 44, respectively. Confidence intervals for estimates in the prespecified subgroups have not been adjusted for multiplicity and may not be used for hypothesis testing.

Subgroup estimates are prespecified but not adjusted for multiplicity.

Subgroups: 4 Ps Discipline

Credibility supports

- Subgroups were prespecified.
- Most point estimates favor intensive treatment.

Why not practice-changing

- Subgroup confidence intervals were not adjusted for multiplicity.
- Some subgroups are underpowered, especially age ≥ 80 and CKD.
- The main article does not foreground clinically decisive interaction tests.

Subgroups support consistency; they should not define narrow target populations without stronger interaction evidence.

Table 3. Adverse Events

Table 3. Adverse Events.*						
Outcome	Intensive Treatment (N = 6414)		Standard Treatment (N = 6407)		Hazard Ratio (95% CI)	P Value
	No. of Events	Percentage of Participants	No. of Events	Percentage of Participants		
Serious adverse event†	2340	36.5	2328	36.3	1.00 (0.94–1.06)	0.96
Conditions of interest‡						
Arrhythmia	69	1.1	68	1.1	1.01 (0.72–1.41)	0.95
Electrolyte abnormality	36	0.6	35	0.6	1.03 (0.65–1.64)	0.91
Injurious fall	65	1.0	61	1.0	1.06 (0.75–1.51)	0.74
Symptomatic hypotension	8	0.1	1	<0.1	7.92 (0.99–63.34)	0.05
Syncope	10	0.2	10	0.2	1.00 (0.41–2.39)	0.99
Acute renal failure	4	0.1	5	0.1	0.79 (0.21–2.95)	0.73
Clinical safety alerts§						
Serum sodium <130 mmol/liter	46	0.7	47	0.8	0.97 (0.65–1.46)	0.89
Serum sodium >150 mmol/liter	22	0.4	25	0.4	0.88 (0.49–1.56)	0.65
Serum potassium <3.0 mmol/liter	32	0.5	33	0.5	0.97 (0.60–1.58)	0.90
Serum potassium >5.5 mmol/liter	177	2.8	125	2.0	1.41 (1.12–1.77)	0.003

* Patients were counted only once for each adverse event.
† Serious adverse events were events that were fatal or life-threatening, resulted in substantial or persistent disability, resulted in or prolonged hospitalization, or were important medical events that investigators judged to represent substantial hazards or harm to research participants.
‡ Conditions of interest were a selected list of events that were serious adverse events or led to an emergency department visit.
§ Data were missing for 184 patients in the intensive-treatment group and 187 patients in the standard-treatment group.

Harms and Benefit-Risk

36.5%

serious AE, intensive

36.3%

serious AE, standard

NNH ~120

K >5.5 mmol/L event-count ARI
0.83%

NNH ~916

symptomatic hypotension ARI
0.11%

- Serious adverse events were nearly identical between arms: HR 1.00 (95% CI 0.94-1.06).
- Symptomatic hypotension was rare but more frequent with intensive therapy: 8 versus 1 patients.
- Hyperkalemia >5.5 mmol/L was more frequent with intensive therapy: 177/6230 versus 125/6220.
- No clear excess of injurious falls, syncope, or acute renal failure in the main table.
- Benefit-risk is favorable only if clinics can monitor BP, symptoms, creatinine, potassium, adherence, and orthostatic risk.

Results Critical Appraisal

Persuasive

- Large sample with adequate event rate close to assumptions.
- Primary endpoint statistically and clinically positive.
- Sensitivity analyses reportedly supported the primary result.

Limits

- Open-label management may influence nonfatal or care-dependent outcomes.
- Only main article available here; detailed protocol/supplemental flow was not fully incorporated into this deck.
- Ethnic and health-system generalizability beyond China is uncertain.

The primary result is convincing for a monitored intensive-treatment strategy; translation requires patient selection and monitoring infrastructure.

APPRAISAL

Risk of bias, GRADE, applicability, and takeaways

Risk of Bias 2 (RoB 2) Summary

Domain 1: Randomisation process

Central web randomization stratified by site; baseline balance acceptable.

LOW RISK

Domain 2: Deviations from intended interventions

Open-label target and treatment adjustment; background cardiometabolic care appeared similar.

SOME CONCERNS

Domain 3: Missing outcome data

Lost to follow-up/withdrawal present; primary analysis used missing-at-random imputation with sensitivity analyses.

SOME CONCERNS

Domain 4: Measurement of outcome

Outcome assessors and adjudicators were blinded; objective cardiovascular outcomes.

LOW RISK

Domain 5: Selection of reported result

Registered trial and protocol/SAP referenced; alpha spending reported.

LOW RISK

OVERALL RISK OF BIAS JUDGEMENT

SOME CONCERNS

GRADE Certainty of Evidence

GRADE FINAL CERTAINTY

**MODERATE
CERTAINTY**

Large positive RCT, downgraded conservatively for open-label care, missing-data dependence, and local indirectness.

Primary composite major cardiovascular outcome

Risk of Bias

Some concerns, mainly open-label treatment strategy and imputation assumptions.

Inconsistency / Publication Bias

Single direct BPROAD trial but direction aligns with intensive BP-control evidence from SPRINT/ESPRIT contexts.

Indirectness

Chinese high-risk T2D population; applicability to frail, very elderly, advanced CKD, and low-resource Indian clinics is limited.

Imprecision

Primary outcome precise; several component, kidney, and harm outcomes are less precise.

Chronology of Key Trials and Guidance

Year	Study / guidance	Population	Main relevance
2010	ACCORD BP	Type 2 diabetes	SBP <120 vs <140 did not significantly lower composite CV outcome; underpowered for BP question.
2015/2021	SPRINT	High-risk adults without diabetes	Intensive SBP target reduced major CV events; diabetes excluded.
2024	ESPRIT	High CV-risk patients with/without diabetes	SBP <120 strategy reduced major vascular events overall; diabetes subgroup less definitive.
2025	BPROAD	Type 2 diabetes, high CV risk	Direct large RCT in diabetes: HR 0.79 for primary composite.
2025-2026	AHA/ACC and ADA guidance	Hypertension / diabetes care	Trend toward tighter targets for high-risk patients, with individualization and monitoring.

Comparison uses article references plus official 2025-2026 guideline context.

Current Guideline Consensus on Targets

- AHA/ACC 2025: overarching BP treatment goal <130/80 mm Hg for adults, with medication recommended at lower thresholds for diabetes, CKD, prior stroke/CVD, or elevated predicted risk.
- ADA 2026: announced updated BP goals for people with high cardiovascular or kidney health risk, with tighter goals for those at risk and more relaxed goals for many older adults.
- ESC 2024: updated European hypertension guidance emphasizes simplified classification and contemporary evidence-based management.
- BPROAD provides diabetes-specific RCT support for a lower SBP target strategy, but guideline translation remains individualized rather than automatic <120 for everyone.
- Practical consensus: confirm BP measurement quality, intensify gradually, monitor potassium/renal function/orthostasis, and relax targets when frailty, falls, limited life expectancy, or adverse effects dominate.

For local practice: use <120 SBP as a selected-patient strategy, not a default prescription for all T2D patients.

Local Applicability in North Indian / Ghaziabad Patients

Applicable

- High diabetes and hypertension burden; stroke prevention is locally relevant.
- Most first-line antihypertensives are available and inexpensive as generics.
- Protocol-like clinic follow-up can be implemented in medicine OPD for selected patients.

Needs caution

- Home BP devices may be inaccurate or unaffordable; clinic measurement quality varies.
- Monitoring potassium/creatinine is essential when RAAS blockers, diuretics, and CKD coexist.
- Frailty, orthostasis, dehydration, heat exposure, and polypharmacy may increase harm.

Best candidates are high-CV-risk, ambulatory, adherent patients who can return for BP, creatinine, potassium, and symptom checks.

Clinical Takeaways for OPD, Wards, & ICU/HDU

OPD

- Consider intensive SBP strategy in high-risk T2D if monitoring is reliable.
- Use validated BP technique and review home logs when available.
- Check creatinine and potassium after intensification.

Wards

- Do not chase <120 during acute illness, sepsis, dehydration, AKI, or poor oral intake.
- Reconcile chronic regimen and plan outpatient titration.
- Watch orthostatic symptoms and drug interactions.

ICU/HDU

- Trial does not answer acute inpatient BP lowering.
- Avoid extrapolation to shock, ACS, stroke thrombolysis windows, or perioperative instability.
- Restart/intensify only after hemodynamics and renal function stabilize.

What Would Change My Mind?

- A large independent trial outside China with similar diabetes-specific benefit and detailed adverse-event capture.
- Longer follow-up showing renal and mortality benefit rather than only composite/stroke-driven benefit.
- Better data in age ≥ 80 years, advanced CKD, frailty, orthostatic hypotension, and multimorbidity.
- Implementation studies showing that Indian public-sector OPDs can achieve safe monitoring at scale.
- Cost-effectiveness and adherence data comparing protocolized <120 strategy versus pragmatic $<130/80$ care.

Journal Club Discussion Questions

Q1. Target or strategy?

Should we interpret BPROAD as proof for SBP <120 itself, or for a closely monitored intensive-treatment strategy?

Q2. Composite endpoint

Is the primary composite clinically coherent enough, or is the result mainly a stroke-prevention story?

Q3. Missing data

How much confidence should we place in missing-at-random imputation given loss to follow-up and withdrawal?

Q4. Local practice

Which Ghaziabad/North Indian patients should be offered intensive BP lowering, and which should be excluded?

References

- Bi Y et al. Intensive Blood-Pressure Control in Patients with Type 2 Diabetes. N Engl J Med 2025;392:1155-1167. DOI: 10.1056/NEJMoa2412006.
- The ACCORD Study Group. Effects of intensive blood-pressure control in type 2 diabetes mellitus. N Engl J Med 2010;362:1575-1585.
- The SPRINT Research Group. A randomized trial of intensive versus standard blood-pressure control. N Engl J Med 2015;373:2103-2116; final report 2021.
- Liu J et al. Lowering systolic blood pressure to less than 120 mm Hg versus less than 140 mm Hg in patients with high cardiovascular risk with and without diabetes or previous stroke. Lancet 2024;404:245-255.
- AHA/ACC et al. 2025 Guideline for the prevention, detection, evaluation, and management of high blood pressure in adults. Circulation. Published online Aug 14, 2025.
- American Diabetes Association. Standards of Care in Diabetes - 2026. Diabetes Care 2026;49(Suppl 1).