

Unit-1: Counting Principles.

1) Fundamental Counting principle

• Addition Principle:-

• Formula:- $m + n$

- eg:- Student wants to buy a drink & can choose from 5 drinks or 4 sodas. Total no. of ways to choose one drink is

$$5 + 4 = 9 \text{ ways.}$$

• Multiplication Principle:-

• Formula: $n_1 \times n_2 \times \dots \times n_k$

- eg:- Person is choosing an outfit consisting of one shirt and one pair of pants. If they have 10 shirts and 7 pants, the total no. of different outfits possible is

$$10 \times 7 = 70 \text{ outfits.}$$

2. Linear Permutation :-

• Permutation of n distinct objects:-

• Formula : $n!$

- eg:- Calculating the number of ways to arrange the letters in the word "FLOWERS". Since there are 7 distinct letters, the total arrangements are:

$$\begin{aligned}\rightarrow 7! &= 7 \times 6 \times 5 \times 4 \times 3 \times 2 \times 1 \\ &= 5,040.\end{aligned}$$

• Permutations of n objects taken r at a time:

• Formula: $P(n, r) = \frac{n!}{(n-r)!}$

- eg:- in a race with 10 sprinters, how many ways can the gold, silver and bronze medals be awarded? This is a permutation of 10 object taken 3 at a time:

$$P(10, 3) = \frac{10!}{(10-3)!} = \frac{10!}{7!}$$

$$= \frac{10 \times 9 \times 8 \times \cancel{7!}}{\cancel{7!}} = \underline{\underline{720}}$$

3.

Circular Permutation :-

- clockwise and anticlockwise arrangements are different:

• Formula: $(n-1)!$

- eg:- How many ways can 5 friends be seated around a campfire? Since the relative position matters, no. of arrangements is

$$(5-1)! = 4! = \underline{\underline{24}}$$

- Clockwise and Anticlockwise arrangements are identical:-

• Formula: $\frac{(n-1)!}{2}$

- eg:- A jeweler is making a bracelet using 8 different colored gemstone. Because the bracelet can be flipped over, the clockwise & anticlockwise orders are the same. No. of ways to arrange a stone is:

$$\frac{(n-1)!}{2} = \frac{7!}{2} = \underline{\underline{2520}}$$

4) Permutation with Repetitions

- objects of the same kind:-

- Formula :-
$$\frac{n!}{n_1 \times n_2 \times \dots \times n_k}$$

- eg:- Find the number of distinct ways to arrange the letters in the word "MISSISSIPPI".

- $(n) = 11$

$$I = 4, S = 4, P = 2$$

$$\frac{11!}{4! \times 4! \times 2!} = \frac{11 \times 10 \times 9 \times 8 \times 7 \times 6 \times 5 \times 4!}{4! \times 4! \times 2!}$$

$$= \frac{11 \times 10 \times 9 \times \overset{4}{\cancel{8}} \times 7 \times \overset{5}{\cancel{6}} \times 5}{\cancel{4} \times \cancel{3} \times \cancel{2} \times \cancel{2}}$$

$$= 11 \times 10 \times 9 \times 7 \times 5$$

$$= 110 \times 9 \times 35$$

$$= 990 \times 35 = \underline{\underline{34650}}$$

Unit 2 -Principle of
Inclusion & Exclusion1.) Principle of Inclusion & Exclusion:- (PIE):-• Formula for 2 Sets (A & B):-

$$\rightarrow |A \cup B| = |A| + |B| - |A \cap B|$$

eg:- in a class, 20 students know Java (A) and 15 know Python (B). if 10 know both ($A \cap B$), the no. of students who know at least one is

$$20 + 15 - 10 = \underline{25}$$

• Formula for 3 Sets (A, B & C)

$$\rightarrow |A \cup B \cup C| = |A| + |B| + |C| - (|A \cap B| + |A \cap C| + |B \cap C|) + |A \cap B \cap C|$$

eg:- in a survey of 200 people, 120 like pizza (A), 100 like burger (B), and 80 like tacos (C), intersections are $A \cap B = 60$, $A \cap C = 30$, $B \cap C = 40$ and all three ($A \cap B \cap C$) = 20

$$120 + 100 + 80 - (60 + 30 + 40) + 20 \\ = \underline{190}$$

2) Set Difference or Set Minus

- Formula for 2 sets.

$$\rightarrow |A \setminus B| = |A| - |A \cap B|$$

→ eg:- if 40 employees know C (A) and 15 of them also know Java ($A \cap B$), those who know C++ but not Java.

$$|A \setminus B| = 40 - 15 = 25$$

- Formula for 3 sets:-

$$\rightarrow |A \setminus (B \cap C)| = |A| - (|A \cap B| + |A \cap C| - |A \cap B \cap C|)$$

→ in a study of 1500 rivers, 520 were polluted by sulphur (A), 335 by phosphate (B), and 425 by oil (C). To find rivers polluted only by sulphur:

$$520 - (180 + 100 - 28) = \underline{268}.$$

3) Other set difference:-

- Symmetric difference \oplus :-

$$\rightarrow |A \oplus B| = |A \cup B| - |A \cap B| = |A| + |B| - 2|A \cap B|$$

→ eg:- if 18 like cricket, 12 like football, and 6 like both, those who like only one sport:

$$18 + 12 - 2(6) = \underline{18}$$

• Complement (A^c):

$$\rightarrow |A^c| = |U| - |A|$$

\rightarrow eg:- in a class of 60 students (U), if 40 classes Maths (A), then (20) did not pass ($60-40$)

• Cartesian Product:

$$\rightarrow |A \times B| = |A| \times |B|$$

\rightarrow eg:- 4 starter choice (A) and 3 drink choices (B) result in $4 \times 3 = \underline{\underline{12}}$

4) Derangement :-

$$\rightarrow !n = n! \left(1 - \frac{1}{1!} + \frac{1}{2!} - \frac{1}{3!} + \dots + (-1)^n \frac{1}{n!} \right)$$

\rightarrow eg:- if 8 distinct coats are returned randomly, there are 14,833 ways for nobody to get their own coat back.

5) Non-negative integer value solution

• Permutation:-

$$\rightarrow P(n, r) = \frac{n!}{(n-r)!}$$

• Combination:-

$$\rightarrow C(n, r) = \binom{n}{r} = \frac{n!}{r!(n-r)!}$$

eg:- from a group of 10 students, a committee of 3 students is to be formed. In how many way can this be done?

$$\begin{aligned} {}^nC_r &= \binom{10}{3} = \frac{10!}{3!(10-3)!} \\ &= \frac{10 \times 9 \times 8}{3 \times 2 \times 1} = \underline{\underline{120}} \end{aligned}$$

• nC_r (with repetition):-

$$\rightarrow \binom{n+r-1}{r}$$

→ eg:-

in how many ways can 6 icecream be chosen from 10 flavours

$$\begin{aligned} &\binom{10+6-1}{6} = \binom{15}{6} \\ &= \frac{15 \times 14 \times 13 \times 12 \times 11 \times 10}{6 \times 5 \times 4 \times 3 \times 2 \times 1} \\ &= 5 \times 7 \times 13 \times 11 = \underline{\underline{5005}} \end{aligned}$$

Unit 3 -Probability1) How to calculate Probability:-• Probability

$$\rightarrow \frac{(\text{No. of favourable outcomes})}{(\text{Total no. of possible outcomes})}$$

→ eg:- what is the probability of getting atleast one head?

$$\rightarrow S = \{HH, HT, TT, TH\}$$

$$A = \{HH, HT, TH\}$$

$$P = A/S = 3/4 = 0.75 = \underline{\underline{75\%}}$$

→ eg:- if 3 coins are flipped, what is the probability of getting exactly 1 tail.

$$\rightarrow S = \{HHH, HHT, HTH, THH, HTT, THT, TTH, TTT\}$$

$$A = \{HHT, HTH, THH\}$$

$$P = A/S = 3/8 = \underline{\underline{0.375}}$$

2) Axioms of Probability:-• Non-Negativity:-

→ Probability of an event is never negative, meaning it can't be less than zero

→ for any event A,

$$P(A) \geq 0$$

- Normalization:

→ probability that something in the sample space happen is 1

→ for the sample space S ,
$$P(S) = 1$$

- Additivity:

→ if A and B are mutually exclusive events.

→
$$P(A \cup B) = P(A) + P(B)$$

→ eg:- Toss 2 coins.

Sample space = $\{HH, HT, TT, TH\}$

$A = \{HH\}$ $B = \{HT\}$

$$P(A \cup B) = P(A) + P(B) = 1/4 + 1/4 = 1/2$$

- if A, B, C are disjoint, then;

- Formula for probability of union of 3 disjoint sets.

→
$$P(A \cup B \cup C) = P(A) + P(B) + P(C)$$

- if A & B are not disjoint sets:-

→
$$P(A \cup B) = P(A) + P(B) - P(A \cap B)$$

- formula for probability of union of 3 sets

→
$$P(A \cup B \cup C) = P(A) + P(B) + P(C) - P(A \cap B) - P(B \cap C) - P(C \cap A) + P(A \cap B \cap C)$$

→ eg:-

Rolling a Dice = $SS = \{1, 2, 3, 4, 5, 6\}$ 1st event $A = \{1, 2, 5\}$ 2nd event $B = \{2, 5, 6\}$ 3rd event $C = \{3, 5, 6\}$

$$P(A) = 3/6 \quad P(B) = 3/6 \quad P(C) = 3/6$$

$$P(A \cap B) = 2/6 \quad P(B \cap C) = 2/6 \quad P(A \cap C) = 1/6$$

$$P(A \cap B \cap C) = 1/6$$

→ For these axioms, we derive useful properties:

$$(i) \quad 0 \leq P(A) \leq 1$$

$$(ii) \quad P(\emptyset) = 0$$

$$(iii) \quad P(A^c) = 1 - P(A)$$

↓

$$P(A) + P(A^c) = 1$$

3) Addition Theorem :-

$$\rightarrow P(A \text{ or } B) = P(A) + P(B) \quad (2 \text{ events})$$

$$\rightarrow P(A \text{ or } B \text{ or } C) = P(A) + P(B) + P(C)$$

$$\rightarrow P(A \cup B) = P(A) + P(B) - P(A \cap B) \quad (2 \text{ events})$$

→ eg:- 4 king and 4 queen in pack of 52 cards

$$\text{king} = 4/52$$

$$\text{queen} = 4/52$$

Probability that it is either a king or

$$\text{queen} = 4/52 + 4/52$$

$$= 8/52 = 2/13$$

→ eg: Probability that query is slow due to heavy load = 0.30.

Probability that query is slow due to indexing issue = 0.25

Probability that query is slow due to both load & indexing issue = 0.10

$$P(A) + P(B) - P(A \cap B) = 0.30 + 0.25 - 0.10 \\ = \underline{0.45}$$

• for 3 events:-

$$\rightarrow P(A \cup B \cup C) = P(A) + P(B) + P(C) - P(A \cap B) - P(A \cap C) - P(B \cap C) + P(A \cap B \cap C)$$

4) Multiplication Theorem:-

• events are independent:-

$$\rightarrow P(A \text{ and } B) = P(A) * P(B)$$

• events are dependent:-

$$\rightarrow P(A \text{ and } B) = P(A) * P(B|A)$$

• union:-

$$\rightarrow (A \cup B) \rightarrow \text{Read as } A \text{ or } B$$

• intersection:-

$$\rightarrow (A \cap B) \rightarrow \text{Read as } A \text{ and } B$$

→ eg:- 3 independent checks.

password correct : 0.90

OTP correct : 0.85

Biometric correct : 0.95

$$P(A \text{ and } B \text{ and } C) = P(A) \cdot P(B) \cdot P(C) \\ = \underline{0.72}$$

- neither A or B, when both try:-

$$\rightarrow P(\text{NO SALE}) = P(A' \cap B') = P(A') \cdot P(B')$$

5) Conditional Probability:-

$$\rightarrow P(A|B) = \frac{P(A \cap B)}{P(B)}, \text{ where } P(B) > 0$$

- conditional probability of A given B:

$$\rightarrow P(A|B) = \frac{P(A \cap B)}{P(B)}, \text{ provided } P(B) > 0$$

- conditional probability of B given A:-

$$\rightarrow P(B|A) = \frac{P(A \cap B)}{P(A)}, \text{ provided } P(A) > 0$$

- Complement of an Event:-

$$\rightarrow P(A') = 1 - P(A)$$

- if $A_1, A_2, A_3 \dots A_n$ are mutually exclusive and exhaustive events:-

$$\rightarrow P(A_i|B) = \frac{P(A_i) \cdot P(B|A_i)}{\sum_{j=1}^n P(A_j) \cdot P(B|A_j)}$$

- Baye's Theorem Formula :-

$$\rightarrow P(A|B) = \frac{P(B|A) \cdot P(A)}{P(B)}$$

- 6) Law of Total Probability :-

$$\rightarrow P(A) = P(B_1) \cdot P(A|B_1) + P(B_2) \cdot P(A|B_2) + \dots + P(B_n) \cdot P(A|B_n)$$

Unit 4:Random variable & Mathematical Expectation.• Discrete Random Variable :-• expected value :-

$$\rightarrow E(x) = \sum_{i=1}^n x_i p(x_i)$$

i.e

$$E(x) = x_1 p_1 + x_2 p_2 + x_3 p_3 + \dots + x_n p_n$$

• Variance :-

$$\rightarrow \text{var}(x) = E(x^2) - [E(x)]^2$$

• Standard deviation :-

$$\rightarrow \sqrt{\text{Var}(x)}$$

Unit 5:-Discrete and Continuous
Probability Distribution• Discrete probability distribution:-• Binomial distribution:-

'x' success out of 'n' trials

→

$$P(X=x) = \binom{n}{x} p^x q^{n-x}$$

Mean:

→

$$E(x) = np$$

Variance :

→

$$\text{var}(x) = npq$$

Failure :

→

$$q = 1 - p$$

Standard Deviation:

→

$$\sigma = \sqrt{npq}$$

• Poisson Distribution:-

PMF of PD is:

→

$$P(X=r) = \frac{e^{-\lambda} \lambda^r}{r!}$$

→

Mean = λ , variance = λ

2) Continuous Probability Distribution:-

• Normal distribution:-

→

• z-score is

$$z = \frac{X - \mu}{\sigma}$$

where X = score

μ = mean

σ = standard deviation

• Uniform distribution:-

→

• Mean:-

$$\mu = \frac{a+b}{2}$$

→

• Variance:-

$$\sigma^2 = \frac{(b-a)^2}{12}$$

→

• Standard

Deviation:-

$$\sigma = \sqrt{\sigma^2}$$