

# NON-ZEEMAN CIRCULAR POLARIZATION OF MOLECULAR SPECTRAL LINES PRESENT IN SEVERAL STAR-FORMING REGIONS

M. A. CHAMMA<sup>1</sup>, M. HOUDE<sup>1</sup>, J. M. GIRART<sup>2</sup> AND R. RAO<sup>3</sup>

<sup>1</sup>Department of Physics and Astronomy, The University of Western Ontario, London, ON, N6A 3K7, Canada

<sup>2</sup>Institut de Ciències de l'Espai (IEEC-CSIC), Can Magrans, S/N, E-08193 Cerdanyola del Vallès, Catalonia, Spain  
and

<sup>3</sup>Submillimeter Array, Academia Sinica Institute of Astronomy and Astrophysics, 645 N. Aohoku Place, Hilo, HI 96720, USA

## Abstract

Understanding the magnetic field in star-forming regions allows us to test ideas about free-fall collapse and support mechanisms in molecular clouds, filling in details about the star formation process. These magnetic fields are usually probed through linear polarization observations from dust continuum and molecular spectral transitions and analysed through the Davis-Chandrasekhar-Fermi method. For molecular lines circular polarization is usually ignored, largely because of difficulty in measurement and its assumed irrelevance. We find in archival data of the Submillimeter Array several examples of circular polarization in common molecular tracers, most notably CO. This circular polarization possibly arises from anisotropic resonant scattering implying that some background linearly polarized flux is being converted and lost to circular polarization. We find circular polarization in NGC7538, IRC+10216 and Orion KL to sufficient degrees that we believe the presence of circular polarization in these spectral lines is widespread for such objects, implying that an important piece of information has been missed when studying magnetic fields through linear polarization from molecular spectral lines in star-forming regions.

## 1. OBSERVATIONS

We collected radio interferometric archival polarimetry observations from the Submillimeter Array (SMA) archive that had been measured using the circular feeds, a similar setup to that used by [Muñoz et al. \(2012\)](#) to measure circular polarization in Sgr A\*. Because the archival observations were not taken with measurements of circular polarization in mind the SNR is often very low, and we had to average velocity channels to increase the SNR at the cost of spectral resolution. This generally increases the SNR from 3-4 to 6-10.

The four objects we present here are Orion KL, NGC7538, NGC1333 and IRC+10216. The first three are well-known star-formation regions while IRC+10216 is an evolved carbon star. We find significant Stokes V signals in all objects except for NGC1333. The visibility data are corrected for beam squint, an instrumental artifact that gives rise to spurious Stokes V signals. Squint gives rise to distinct pairs of positive and negative peaks of Stokes V throughout the inverted image. Maps are then made of the continuum and lines for each object. The squint correction is confirmed visually by inspecting the Stokes V maps, where we see the pairs of peaks largely disappear. More details on the squint correction will be given in the next section.

Figure 1 shows corrected Stokes I and Stokes V spectra obtained at the peak of Stokes V on the corresponding maps. A comparison of Stokes V spectra before and after correction are shown in Figure 4. Notice in all cases the Stokes V

Object	Coordinates (J2000)	Array Configuration	Date Observed
<b>Orion KL</b>	RA 05 <sup>h</sup> 35 <sup>m</sup> 14.501 <sup>s</sup> Dec -05°22'30.40"	Compact	2008-01-06
<b>NGC7538</b>	RA 23 <sup>h</sup> 13 <sup>m</sup> 44.771 <sup>s</sup> Dec +61°26'48.85"	Compact	2014-10-28
<b>IRC+10216</b>	RA: 09 <sup>h</sup> 47 <sup>m</sup> 57.381 <sup>s</sup> Dec +13°16'43.70"	Compact	2009-11-24
<b>NGC1333</b>	RA 03 <sup>h</sup> 28 <sup>m</sup> 55.580 <sup>s</sup> Dec +31°14'37.10"	Compact	2010-10-14

**Table 1.** Summary of Archival Observations Used

signal decreases after squint correction. Stokes V can be found in the average of all the visibility data as well, though the significance is  $3\text{--}5\sigma$  in that case compared to approximately  $6\text{--}10\sigma$  when the spectra is taken from the inverted maps. We present here only the map spectra. In general the peaks of Stokes I and Stokes V do not align in the maps.

In Orion KL the lines of CO ( $J = 3 \rightarrow 2$  at 345.8GHz) and SiO ( $J = 8 \rightarrow 7$  at 347.3GHz) are both bright (peak Stokes I of  $\sim 20$  Jy/beam and  $\sim 55$  Jy/beam respectively; not shown) and both show Stokes V signals. The CO Stokes V signal has an antisymmetric structure while the SiO Stokes V signal is purely negative. Figure 3 shows the peak Stokes V signal of the SiO line. We checked the average of all the visibilities (average over all antennae baselines and all time intervals) and found that the CO Stokes V signal is more intense than the SiO Stokes V signal, and the Stokes I intensities after averaging the visibilities were approximately 50 Jy and 75 Jy for CO and SiO respectively. If the Stokes V signal were purely leakage from Stokes I than we would expect to see an SiO Stokes V signal that is stronger than the CO Stokes V signal, but we do not.

In IRC+10216 we again see Stokes V in the CO transition at 345.8 GHz but also several signals in CS, SiS, and H<sup>13</sup>CN. These lines and their frequencies are listed in Table 2. We again note that the Stokes I intensity is not proportional to the Stokes V intensity. The detection in CS at 342.8 GHz is huge compared to it's I intensity.

In NGC7538 there was a strong Stokes V detection in CH<sub>2</sub>CO at 346.6 GHz that completely disappeared after correction. The Stokes V signal in CO at 345.8 GHz decreased in intensity but is still extremely prominent.


Finally Figure 2 shows no detection in NGC1333, with only a weak detection of CO in Stokes I.

When assessing the Stokes V detection we consult the map for obvious pairs of positive/negative peaks that would indicate beam offset and therefore a false Stokes V signal. The maps are taken around a very narrow frequency band of around 2 MHz so any peaks that exist shouldn't be washed out by noise in adjacent channels. In the maps for Orion KL and IRC+10216 shown in Figure 1 there are no negative peaks around the peak of Stokes V. However in NGC7538 there is quite a large negative Stokes V peak near our chosen peak that may indicate squint. The top panel of Figure 5 shows what 'squint peaks' look like (we know these peaks are from squint because they disappear after correction) and the pairs tend to resemble each other in shape. The pair of peaks around our chosen peak in NGC7538 however have distinct shapes. The worst case here is that the signal is entirely squint but on the other hand the signal may be a mixture of real and heavily affected by squint. The detections in Orion KL and IRC+10216 are more reliable.


Table 1 shows a summary of the objects presented and related information.

## 2. SQUINT CORRECTION


Here we describe spurious Stokes V that arises when using the SMA and our scheme for correcting it. This instrumental Stokes V comes from a slight pointing offset between the left- and right-handed CP beams.



**Figure 1.** Corrected maps of the CO  $J = 3 \rightarrow 2$  line (345.8GHz) and corrected spectra for Orion KL, NGC7538, IRC+10216. **Maps:** Blue contours are Stokes V and are shown at the -4, -3, -2, 2, 3, 4 $\sigma$  levels. The RMS error for each Stokes V map is found using *Miriad*'s *imstat* command. Dark red contours are Stokes I and the levels are 15%, 30%, 45%, 60%, 85% and 95% of the maximum. The value in the top left is the frequency of the mapped signal. Each map has a narrow bandwidth of  $\sim 2$  MHz around the frequency given in the top left corner. **Spectra:** *Miriad*'s *maxfit* is used on the CO map to obtain the location on the image where the Stokes V signal at 345.8GHz is maximum, and a spectra is obtained through that point. The cross on the map denotes the location of that peak. The spectrum for NGC7538 is Hanning smoothed by a length of 15.



**Figure 2.** Corrected maps of the CO  $J = 3 \rightarrow 2$  line and corrected spectrum for NGC1333 (IRAS2a). No significant Stokes V signal is detected here, probably because the object is too dim. Contours are the same levels as in Figure 1 and the spectrum is obtained the same way.




**Figure 3.** Peak Stokes V signal for the SiO ( $J = 7 \rightarrow 8$  at 347.3 GHz) transition in Orion KL. Note there is also a strong Stokes V signal in the CO ( $J = 3 \rightarrow 2$  at 345.8 GHz) transition here. The SiO signal is purely negative but the CO signal is antisymmetric.

The archival data used was in all cases observed with the goal of measuring linear polarization (i.e., the Stokes Q and Stokes U parameters). On the SMA this is done with a quarter-waveplate placed in front of the linear receivers to convert incident circular polarization (CP) to linear polarization. While this method is relatively imprecise, it avoids having to solve for the linear polarization terms of calibration objects when obtaining Stokes V (Marrone et al. 2008; Sault et al. 1996) (cite the textbook). Obtaining Stokes V from the visibilities measured with circular feeds is done as follows. Given antennae  $a$  and  $b$ , the Stokes V visibility in the circular feed case is found through (Muñoz et al. 2012):

$$\mathcal{V}_V \simeq \frac{1}{2} \left\{ \mathcal{V}_{RR} / (g_{Ra} g_{Rb}^*) - \mathcal{V}_{LL} / (g_{La} g_{Lb}^*) \right\}$$

## Stokes V Map Spectra before and after squint correction



**Figure 4.** Stokes V spectra of all objects before and after squint correction. *Miriad*'s `maxfit` is used on the CO map for each respective object to obtain the location in the image where the Stokes V signal at 345.8GHz is maximum, and a spectra is obtained through that point.

Object	Line	(GHz)	Stokes V (Jy/beam)
<b>Orion KL</b>	CO $J=3\rightarrow2$	345.78	0.65
	SiO $J=8\rightarrow7$	347.3	-0.65
<b>NGC7538</b>	CO $J=3\rightarrow2$	345.8	0.85
<b>IRC+10216</b>	CS $J=7\rightarrow6$	342.88	0.6
	SiS $J=19\rightarrow18$	344.78	0.2
	H <sup>13</sup> CN $J=4\rightarrow3$	345.34	0.8
	CO $J=3\rightarrow2$	345.8	0.4
<b>NGC1333 (IRAS2a)</b>	CO $J=3\rightarrow2$	345.8	None


**Table 2.** Summary of corrected Stokes V signals found. The beam size is determined by the configuration of the antennae array. An intensity for the Stokes V signal is only given if the peak is noticeably higher than the noise level. The intensity quoted for CO in NGC7538 is before smoothing is applied.

Here the right-handed CP and left-handed CP visibilities are  $\mathcal{V}_{RR}$  and  $\mathcal{V}_{LL}$  and are measured by orienting the quarter-waveplate that is placed in the beam of the antennae. The complex gain factors for each polarization for each antennae are  $g_{Ra}$ ,  $g_{Rb}$ ,  $g_{La}$  and  $g_{Lb}$  with 'R' and 'L' for right- and left-CP, respectively. Because the Stokes V visibility is found by taking the difference of two beams a slight offset gives rise to pairs of positive and negative peaks of Stokes V, as shown in Figure 5. This offset likely arises because of slight differences in the index of refraction of the quarter-waveplate when it is rotated, but hasn't been studied with the SMA as far as we know.

Because the visibilities are the Fourier transform of the intensity map, an offset in image space is equivalent to a complex exponential term in visibility space that can be absorbed into the gain coefficients (see Appendix). Thus it is easiest to correct for the offset in visibility-space by using *Miriad* (citation) to solve for the gain coefficients on each beam. Specifically the process is:

1. Observations are calibrated for gain and phase in the usual way using calibration observations of 'good' sources (usually quasars like 3C84, 3C454, etc.) (See *Miriad* User Guide [Sault et al. \(2008\)](#))
2. Visibilities are split into line-free continuum data and line data. These are then mapped and used to obtain CLEAN models
3. The separate continuum and line data are further split into  $LL$  and  $RR$  visibilities. *Miriad*'s `selfcal` is used on the continuum data to solve for the gain coefficients of each antenna and each polarization ( $L$  or  $R$ ). This is done by minimizing the difference between measured visibilities  $\mathcal{V}_{ij}$  of antennae  $i$  and  $j$  and model visibilities  $\hat{\mathcal{V}}_{ij}$  according to  $\epsilon^2 = \sum |\mathcal{V}_{ij} - g_i g_j^* \hat{\mathcal{V}}_{ij}|^2$  for each of the correlations  $LL$  and  $RR$  ([Sault et al. 2008](#); [Schwab 1980](#)). The model visibilities used are those found earlier.
4. The gains found from the continuum  $LL$  data is then applied to the line  $LL$  data, similarly for the  $RR$  continuum and line data.
5. The different polarizations ( $LL$ ,  $RR$ ,  $RL$ ,  $LR$ ) are recombined and inverted to produce corrected maps. Spectra can be obtained either from the corrected visibilities or the maps.

Figure 5 shows maps before and after this correction is applied.



**Figure 5.** Map of the continuum around 345GHz in Orion KL before (top) and after (bottom) squint correction. Red contours are Stokes I, blue contours are Stokes V. Dashed lines denote negative values, solid lines denote positive values. Note the multiple pairs of positive and negative Stokes V peaks. These largely disappear after correction. The contour scales on both maps are identical. Red Stokes I contours are at 15%, 35%, 55%, 75% and 95% of the peak intensity. Blue Stokes V contours are at -8, -7, -6, -5, -4, -3, -2, 2, 3, 4, 5, 6, 7 and  $8\sigma$  levels.

## REFERENCES

- |   |             |
|---|-------------|
| Muñoz, D. J., et al. 2012 ApJ, 745, 115                     | Schwab 1980 |
| Marrone et al. 2008   |             |
| Sault et al. 1996   |             |
| Sault, B., Killeen, N., 2008, Miriad Users Guide, Australia |             |
| Telescope National Facility                                 |             |