

Hypothesis Testing II

Megan Ayers

Stat 141 | Fall 2026

Friday, Week 9

Announcements

- **10am section**: Ending class a few minutes early for pigeonhole innovations (you're invited!)
- Office hour changes next week - double check Moodle!

Goals for today


- Practice framing research questions in terms of hypotheses
- Practice defining and simulating null distributions
- 1 vs 2 sided tests

Recap: Framework for Hypothesis Testing

1. Present research question and identify hypotheses
   - Null hypothesis (H_0): status quo, random chance, no effect…
   - Alternative hypothesis (H_a): the researchers' conjecture
2. Describe **Null** distribution
3. Obtain data, calculate relevant **Test Statistic**
4. Calculate the **P-value**
   - P-value = likelihood of observing the Test Statistic or something more extreme assuming the Null Hypothesis
5. Use the P-value to make a conclusion on the research question

Let's Practice

Example: Does Extrasensory Perception (ESP) exist?





Psychologists Bem and Honorton conducted extrasensory perception studies:

- A "sender" randomly chooses an object out of 4 possible objects and sends that information to a "receiver".
- The "receiver" is then given a set of 4 possible objects and they must decide which one most resembles the object sent to them.

Out of **329 reported trials**, the "receivers" correctly identified the object **106 times**.

Hypothesis Testing Steps: ESP

Psychologists Bem and Honorton conducted extrasensory perception studies:

- A "sender" randomly chooses an object out of 4 possible objects and sends that information to a "receiver".
- The "receiver" is then given a set of 4 possible objects and they must decide which one most resembles the object sent to them.

Out of **329 reported trials**, the "receivers" correctly identified the object **106 times**.

Discuss with neighbor(s):

1. What is the relevant null hypothesis (H_0) and alternative hypothesis (H_a)?

2. Describe **Null distribution**. How might we simulate it?

3. What is our relevant **Test Statistic**?

4. What does the **P-value** represent here? How might we calculate it, given step 2?

5. How would we use the P-value to make a conclusion on the research question?

05:00

Hypothesis Testing Steps: ESP

1. What is the relevant null hypothesis (H_0) and alternative hypothesis (H_a)?
   - H_0: Participants were guessing randomly ($p = 0.25$)
   - H_A: Participants were guessing better than random ($p > 0.25$)

Hypothesis Testing Steps: ESP

2. Describe "Null" distribution. How might we simulate it?
   - The distribution of $\hat{p}$ (the proportion of correct guesses) we would see if participants guess at random each time.

- **Bonus Q**: What's the lowest that $\hat{p}$ could be under the null? The highest?


Hypothesis Testing Steps: ESP

2. Describe "Null" distribution. **How might we simulate it?**
   - The distribution of $\hat{p}$ (the proportion of correct guesses) we would see if participants guess at random each time.

Steps to simulate this:

1. Sample with replacement from a vector of 0's and 1's, where we have a 25% chance of sampling 1 each time
2. Compute proportion of 1's sampled
3. Repeat 1 and 2 many times.

```
set.seed(123)
guesses <- data.frame(
  correct = c(0, 0, 0, 1)  # Like a 4-sided dice
)                          # with sides 0, 0, 0, 1
null_stats <- guesses %>%
  rep_sample_n(size = 329, replace = TRUE,
               reps = 10000) %>%
  group_by(replicate) %>%
  summarize(p_hat = sum(correct) / n())
ggplot(null_stats, aes(x = p_hat)) + geom_histogram(col
```




Hypothesis Testing Steps: ESP

2. Describe "Null" distribution. **How might we simulate it?**
   - The distribution of $\hat{p}$ (the proportion of correct guesses) we would see if participants guess at random each time.

Steps to simulate this:

1. Sample with replacement from a vector of 0's and 1's, where we have a 25% chance of sampling 1 each time
2. Compute proportion of 1's sampled
3. Repeat 1 and 2 many times.

```
set.seed(123)
guesses <- data.frame(
  correct = c(0, 0, 0, 1)  # Like a 4-sided dice
)                          # with sides 0, 0, 0, 1
null_stats <- guesses %>%
  rep_sample_n(size = 329, replace = TRUE,
               reps = 10000) %>%
  group_by(replicate) %>%
  summarize(p_hat = sum(correct) / n())
ggplot(null_stats, aes(x = p_hat)) + geom_histogram(col
```




Hypothesis Testing Steps: ESP

2. Describe "Null" distribution. **How might we simulate it?**
   - The distribution of $\hat{p}$ (the proportion of correct guesses) we would see if participants guess at random each time.

Steps to simulate this:

1. Sample with replacement from a vector of 0's and 1's, where we have a 25% chance of sampling 1 each time
2. Compute proportion of 1's sampled
3. Repeat 1 and 2 many times.

```
set.seed(123)
guesses <- data.frame(
  correct = c(0, 0, 0, 1)  # Like a 4-sided dice
)                          # with sides 0, 0, 0, 1
null_stats <- guesses %>%
  rep_sample_n(size = 329, replace = TRUE,
               reps = 10000) %>%
  group_by(replicate) %>%
  summarize(p_hat = sum(correct) / n())
ggplot(null_stats, aes(x = p_hat)) + geom_histogram(col
```




Hypothesis Testing Steps: ESP

2. Describe "Null" distribution. **How might we simulate it?**
   - The distribution of $\hat{p}$ (the proportion of correct guesses) we would see if participants guess at random each time.

Steps to simulate this:


1. Sample with replacement from a vector of **0**'s and **1**'s, where we have a 25% chance of sampling **1** each time
2. Compute proportion of **1**'s sampled
3. Repeat 1 and 2 many times.

```
set.seed(123)
guesses <- data.frame(
  correct = c(0, 0, 0, 1)  # Like a 4-sided dice
)                          # with sides 0, 0, 0, 1
null_stats <- guesses %>%
  rep_sample_n(size = 329, replace = TRUE,
               reps = 10000) %>%
  group_by(replicate) %>%
  summarize(p_hat = sum(correct) / n())
ggplot(null_stats, aes(x = p_hat)) + geom_histogram(col
```



Hypothesis Testing Steps: ESP

3. What is our relevant "Test Statistic"?
   - $\hat{p}$ = 106 / 329 = 0.32



4. How can we calculate the "P-value"?
   - Can use probability theory to find $P(\text{Correct guesses} \geq 106) = \ldots?$
   - Or our simulated null distribution

```
mean(null_stats$p_hat >= 106/329)
```
```
[1] 0.0013
```

5. How would we use the P-value to make a conclusion on the research question?
   - If the p-value is "small enough", we reject the null hypothesis of random guessing.

Hypothesis Testing: ESP

- We got a pretty small p-value (0.0013). Hooray! It is reasonable to reject the null hypothesis.
- But really, do we believe that ESP is real?
- Next lecture, we'll talk more about hypothesis testing errors.



Side Note: Reproducibility and Replicatability

- Two important words in data analysis:
  - Reproducibility
  - Replicability
- **Reproducibility**: If I give you the raw data and my write-up, you will get to the exact same final numbers that I did.
  - By using `Quarto` Documents, we are learning a **reproducible** workflow.
- **Replicability**: If you follow my study design but collect new data (i.e. repeat my study on new subjects), you will come to the same conclusions that I did.
  - Sadly, **replication** studies of Bem and Honorton's ESP trials largely failed to find evidence of ESP.

Switching gears: 1 vs 2 sided tests


- Say we're **flipping a coin 20 times**, and (for some reason) we are worried that it's not a "fair" coin
  - i.e., The probability of heads is not 0.5
- Let $p =$ **probability of getting a heads with this coin**
- **Q**: What is the null hypothesis?
- Possible alternative hypotheses:
  - $H_a: p < 0.5$
  - $H_a: p > 0.5$
  - $H_a: p \neq 0.5$
- We've been thinking about the first two cases (1 sided tests). Let's see what changes if $H_a: p \neq 0.5$ (2 sided test).

Recall the hypothesis testing steps again:

1. Present research question and identify hypotheses (H_0 and H_a)
   - $H_a: p < 0.5$, **or** $p > 0.5$, **or** $p \neq 0.5$**?**
   - $H_0: p = 0.5$
2. Describe "Null" distribution
   - **This is the same regardless of which H_a we use, depends only on H_0**
3. Obtain data, calculate relevant "Test Statistic"
4. **Calculate the "P-value"**
   - P-value = likelihood of observing the Test Statistic **or something more extreme** assuming the Null Hypothesis
5. Use the P-value to make a conclusion on the research question


1 vs 2 sided tests: what does "more extreme" mean?

- Suppose our test statistic from 20 coin flips is $\hat{p} = 0.85$.
- If we have $H_a: p > 0.5$ (**right**-tailed test)
  - We'd want $P(\hat{p} \geq 0.85)$ under the null hypothesis as our P-value




```
1 sum(null_stats$p_hat >= 0.85) / 10000
[1] 0.0351
```

- If instead we have $H_a: p < 0.5$ (**left**-tailed test)
  - We'd want $P(\hat{p} \leq 0.85)$ under the null hypothesis as our P-value





```
1 sum(null_stats$p_hat <= 0.85) / 10000
[1] 0.9649
```


1 vs 2 sided tests: what does "more extreme" mean?

- Suppose our test statistic from 20 coin flips is $\hat{p} = 0.85$.
- Suppose we have $H_a: p \neq 0.5$
  - $\hat{p} = 0.85$ is 0.35 *more* than $p = 0.5$, so we want to include $P(\hat{p} \geq 0.85)$
  - $\hat{p} = 0.15$ is 0.35 *less* than $p = 0.5$, so we'd <u>also</u> want $P(\hat{p} \leq 0.15)$ (this would be more extreme in the other direction, ex. seeing 3/20 heads)
- So our P-value would be the sum of those two things (both tails)!

```
1 (sum(null_stats$p_hat >= 0.85) + sum(null_stats$p_hat <= 0.15)) / 10000
[1] 0.0732
```



1 vs 2 sided tests: what does "more extreme" mean?

Note the effect that our choice of H_a had on our p-values!

- Our test statistic was $\hat{p} = 0.85$
- $H_a: p < 0.5$: P-value = 0.9649
- $H_a: p > 0.5$: P-value = 0.0351
- $H_a: p \neq 0.5$: P-value = 0.0732

Another Example: Mouse Frowns :(

Can you tell if a mouse is in pain by looking at its facial expression? A recent study created a "mouse grimace scale" and tested to see if there was a positive correlation between scores on that scale and the degree of pain (based on injections of a weak and mildly painful solution). The study's authors believe that if the scale applies to other mammals as well, it could help veterinarians test how well painkillers and other medications work in animals.

- **Q**: How should we write out H_0 and H_a in terms of population parameters? (*Hint: recall that we use r for correlation*)
- **Q**: How would we expect the sample statistic to behave under H_0?


Null Distribution in R via Permutation

- **Rat Pain and grimacing :(**
  - H_0: The correlation between grimacing and the level of pain is 0
  - Under the null, grimacing and level of pain are totally unrelated
  - To approximate the null distribution, we could:
    1. Randomly shuffle the `grimace` column of our data
    2. Calculate the correlation between `grimace` and `pain` in the shuffled data
    3. Repeat 1-2 many times.

```
1 head(df, 3)
```

```
      pain   grimace
1 9.391984 12.538458
2 7.001035  8.759391
3 9.793066 15.637338
```

```
1  set.seed(111)
2  null_stats <- df %>%
3    select(grimace) %>%  # Grab just grimace
4    rep_sample_n(size = 100, replace = FALSE,
5                 reps = 5000) %>%  # Sample from it
6    add_column(pain = rep(df$pain,  # Copy in pain
7                          times = 5000)) %>%
8    group_by(replicate) %>%
9    summarize(stat = cor(grimace, pain))
10
11 ggplot(null_stats, aes(x = stat)) +
12   geom_histogram(color = "white", fill = "gray60")
```



Another Example: Crocodile Smiles

Can a simple smile have an effect on punishment assigned following an infraction? In a 1995 study, Hecht and LeFrance examined the effect of a smile on the leniency of disciplinary action for wrongdoers. Participants in the experiment took on the role of members of a college disciplinary panel judging students accused of cheating. For each suspect, along with a description of the offense, a picture was provided with either a smile or neutral facial expression. A leniency score was calculated based on the disciplinary decisions made by the participants.

- **Q**: How should we write out H_0 and H_a in terms of population parameters? (*Hint, we can write the population mean leniency score for smilers as μ_S*).
- **Q**: How would we expect the sample statistic to behave under H_0?


Null Distribution in R via Permutation

- **Discipline for Smilers vs Non-Smilers**
  - H_0: We expect the difference in average leniency scores between the smiling and non-smiling groups to be zero - no relationship between the leniency score and the treatment group.
  - To approximate the null distribution, we could:
    1. Randomly shuffle the `treatment` column of our data
    2. Calculate the difference in means between the leniency scores in each group
    3. Repeat 1-2 many times.

```r
head(df, 3)
```

```
  treatment score
1     smile     6
2   neutral     4
3     smile     8
```

```r
# NOTE: This is more advanced wrangling than we've seen
set.seed(111)
null_stats <- df %>% select(treatment) %>%
  rep_sample_n(size = 60, replace = FALSE, reps = 5000) %>%
  add_column(score = rep(df$score, times = 5000)) %>%
  group_by(replicate, treatment) %>%
  summarize(ybar = mean(score)) %>%
  pivot_wider(names_from = treatment, values_from = ybar) %>%
  mutate(stat = smile - neutral)
ggplot(null_stats, aes(x = stat)) +
  geom_histogram(color = "white", fill = "gray60")
```



Next time

- Decisions in hypothesis testing
- Types of hypothesis testing errors
- Power



Have a fun and safe Halloween!