

W

DISCREPANCY MODELING OF ANKLE EXOSKELETON WALKING CAN IMPROVE RESPONSE PREDICTIONS

Megan R Ebers¹, Michael C Rosenberg¹, J Nathan Kutz², Katherine M Steele¹

¹Department of Mechanical Engineering, University of Washington, Seattle, WA

²Department of Applied Mathematics, University of Washington, Seattle WA

Exoskeletons change gait, but predicting response is hard.

Ankle exoskeletons (exos) are prescribed to **assist walking** or **improve gait** following neurological injuries.



Changes in gait kinematics and muscle activity in **response** to exo properties are **highly heterogeneous** [1].

Data-driven modeling enables subject-specific dynamics to be estimated without detailed knowledge of an individual's physiology [2].

Identifying **interpretable dynamics** governing exo response may highlight individualized drivers of response.

However, if standard measurements (kinematics and muscle activity) can **encode** the dynamics of exo responses is **unclear**.

Driving Question

To what extent can **discrepancies** between walking dynamics in zero-stiffness and high-stiffness passive exos be identified using kinematic and myoelectric data?

Response as a Discrepancy



Discrepancies are mismatches between system and model dynamics.

Discrepancy models **disambiguate** missing dynamics from noise and may quantify responses to exos.


For ankle exos:

An individual's response to exoskeleton properties, such as stiffness or torque, may be represented as a discrepancy between baseline dynamics (e.g., walking in a zero-stiffness exo) and the dynamics in a perturbed condition (e.g., walking in a high-stiffness exo)

Methods & Results



Bias analysis of model error suggests discrepancy modeling identifies missing dynamics



Discussion

Discrepancy models **encode changes in dynamics** with ankle exos using kinematic and myoelectric measurements up to the limits imposed by measurements and noise.

Unexplained myoelectric variance suggests **additional measurements are needed** to encode myoelectric response dynamics to ankle exos (e.g., ultrasound for MTU properties) [4].

Discrepancy structures identified from kinematics and muscle activity may still provide insight into different **mechanisms driving exo responses**.

Reducing model complexity and **identifying additional measurements** explaining responses represent important next steps to developing interpretable data-driven models of exo responses.

Acknowledgements

This research was supported by the NSF under award CBET 1452646 and GRFP DGE-1762114

References

- [1] E. M. McCain, T. J. Dick, T. N. Giest, R. W. Nuckols, M. D. Lewek, K. R. Saul, et al., J Neuro Rehab, vol. 16, pp. 1-12, 2019.
- [2] M. C. Rosenberg, B. S. Banjanin, S. A. Burden, and K. M. Steele, JRSI, vol. 17, p. 20200487, 2020.
- [3] S. N. Kumpati and P. Kannan, TNNLS, vol. 1, pp. 4-27, 1990.
- [4] R. W. Nuckols, T. J. Dick, O. N. Beck, and G. S. Sawicki, Scientific Reports, 10.1, pp. 1-15, 2020.