Notes on Gravity

Megha Burri

Preface

General Relativity is one of the most successful theories of Physics in history. It seems to hold up to every experimental test thrown at it, even more than a century after its conception. In undergraduate lectures this topic is often presented alongside an introduction to tensor calculus and differential geometry. In contrast, these notes will not contain this background information. The aim is to focus on the consequences of the theory rather than provide a pedagogical overview, and will therefore assume knowledge of some Riemannian geometry. One reference for these notes is the incredibly comprehensive book by Straumann [Str13], though only a small fraction of the content will be covered here. Chapter 1 will recap the theory of General Relativity in a hopefully succint manner. Chapters 2 & 3 will be devoted to stationary and rotating black hole solutions respectively. For Chapter 4 the aim is to understand the nature of Lorentzian geometry in contrast to the Riemannian case. Witten has some very approachable lectures [Wit20] as does Galloway [Gal14]. The focus will be on the causal structure of spacetime and singularity theorems. These notes are a work in progress so there will be comments spotted throughout for areas which need to be filled in. Time permitting, I aim to keep this as a long term project so that I can discuss more exotic features of gravity as my understanding improves.

Contents

1 Setup of General Relativity **3**
- 1.1 Motivation . . . 3
- 1.2 Einstein-Hilbert Action . . . 5
- 1.3 Geodesic Flow . . . 7
- 1.4 General Covariance . . . 11
- 1.5 Moving Frames . . . 11
  - 1.5.1 Klein Geometry of Minkowski Space . . . 11
  - 1.5.2 Cartan Geometry . . . 12

2 Schwarzschild Spacetime **16**
- 2.1 Derivation of the Metric . . . 16
- 2.2 Chart Problems . . . 19
- 2.3 Geodesics in Schwarzschild Spacetime . . . 20

3 Kerr Spacetime **22**
- 3.1 Metric . . . 22
- 3.2 Analysing the Geometry of Kerr Spacetime . . . 22
- 3.3 Integrability of Kerr Geodesic Flow . . . 22
  - 3.3.1 Hamilton-Jacobi equation . . . 23

4 Causality & Topology **24**
- 4.1 Obstructions to Spacetime . . . 24
- 4.2 Causality . . . 26
- 4.3 Inner Product Structure on the Tangent Space . . . 28
- 4.4 Causality & Energy Conditions . . . 30
- 4.5 Null Geometry . . . 30
- 4.6 Penrose's Theorem . . . 36
- 4.7 Instability of Kaluza-Klein Theories . . . 37

Chapter 1

Setup of General Relativity

1.1 Motivation

This section will provide an overview of the framework of Einstein's theory. The starting point is to understand the theory of special relativity. We assume that the three dimensions of space and one dimension of time can be treated as a combined space, $\mathbb{R}^4$, equipped with an inner product or metric η. The key distinction between ordinary Euclidean inner product spaces and those discussed here is that the metric is no longer positive definite. The metric can be put in diagonal form and the number of $\pm$ eigenvalues determine its signature. We say that a Lorentzian metric has signature (p, q) if there are p positive eigenvalues and q negative eigenvalues. In the context of special relativity we demand that the metric has signature $(1, 3)$ or $(3, 1)$. Both of the above choices are equivalent, but we will choose to work with the second, so the metric has diagonal form $(-1, +1, +1, +1)$. We write $\mathbb{R}^{1,3}$ or $\mathbb{R}^{3,1}$ when referring to four dimensional Cartesian space equipped with a metric of signature $(3, 1)$ or $(1, 3)$. Roughly speaking, the indefinite signature allows us to distinguish a 'time direction' in spacetime based on the signature. For example, a vector $v \in \mathbb{R}^{1,3}$ is 'pointing in the time direction' if $\eta(v, v) < 0$ in our choice of signature. Actually the vector could be pointing 'up' in the time axis or 'down', we need to prescribe which direction time 'flows' in and this is given by a time-orientation. In this case it is just choice a constant vector, T, with $\eta(T, T) < 0$.

Definition 1.1.1. $\mathbb{R}^{1,3}$ *equipped with the* ***Minkowski metric****,* $\eta(v, v) = -c^2 v_0^2 + \sum_{i=1}^{3} v_i^2$*, and a time-orientation is* ***Minkowski spacetime****.*

With any metric we can associate the orthogonal group $\mathrm{O}(p, q)$, in this case $\mathrm{O}(3, 1)$, which is the group linear transformations on $\mathbb{R}^{1,3}$ which preserve η. This is the **Lorentz group**. Not all of these transformations are considered to be physical. We restrict to the **proper orthochronous** Lorentz group $\mathrm{SO}^+(3, 1)$ which only includes those transformations which have positive determinant and do not allow time and space directions to be reflected. This is also denoted $\mathrm{SO}_0(3, 1)$ for the connected component of the identity. In special relativity observers moving at constant velocity define a so-called inertial reference frame. Any other inertial frame is related to by a transformation in the above group. The transformations can also act vectors with $\eta(v, v) = 0$. In two dimensions, $\mathbb{R}^{1,1}$, with time on the y-axis and space on the x, this relation defines two lines crossing the origin with gradient $\pm 1/c$; in other words light rays with velocity c. In this way we recover one of the postulates of special relativity that the speed of light must be constant in all frames of reference.

To get to General Relativity (GR) we suppose that spacetime is no longer described by a Cartesian space, but has the property of being curved. To enforce the equivalence principle we demand that no matter how curved spacetime is, it must locally look flat. This leads to considering that spacetime must be described by a manifold, M, instead of a vector space. It also ought to be a smooth manifold to permit the methods of calculus to be used. Moreover, we want that special relativity should hold locally at each point in space, so it should 'look like' Minkowski spacetime at any point. From Riemannian geometry we know that a manifold at a point looks like its tangent space at that point, the content of the normal neighbourhood theorem. It makes sense to then demand that our tangent spaces are actually Minkowski spaces. This idea is part of Cartan geometry.

In the physics literature, tensors are often **defined** by their transformation law. By contrast, in mathematics, one introduces tensors as arbitrary sections of the appropriate tensor tangent bundle. These two viewpoints don't agree a priori. For example, even in Minkowski space, a physicist only considers special 'four-vectors' to be vectors. To a mathematician, any old tuple of four numbers is a perfectly well-defined vector. In fact, not even all real numbers are classed as 'scalars' in the Physics literature. For example, the energy of a particle changes depending on which reference frame one chooses. These viewpoints can be reconciled by being slightly more pedantic. A postulate of special relativity is that that physical laws should not depend on the choice of inertial reference frame. This can be enforced by demanding that physical quantities should be described by representations of the appropriate group. For example, vectors in Minkowski space which transform in the Lorentz group are **four-vectors**.

In GR, we need to promote this to the manifold level. We want a metric tensor defined pointwise as, $g_p : T_pM \times T_pM \to \mathbb{R}$, which has a $(3,1)$- signature. Recall that there always exists a **frame bundle** on any manifold which consists of the disjoint union of the set of bases at each point. More precisely, this is a principal $\mathrm{GL}(n,\mathbb{R})$-bundle. If there exists a Lorentzian metric, there is a left action of Lorentz group on the frame bundle defined by a representation. We can then construct the assiocated bundle to a given representation with fibre isomorphic to $\mathbb{R}^n$. The upshot is then that tensor fields ought to sections of the associated bundle, to ensure that they transform correctly. The usual tangent bundle to a manifold can be constructed from the structure group $\mathrm{GL}(n,\mathbb{R})$. Here we have a reduction of the structure group to the proper orthochronous Lorentz group, encoding the necessary transformation properties on the associated bundle. With a Lorentzain metric, vectors can be split into three classes.

Definition 1.1.2. *At a point, $p \in M$, a vector, $X \in T_pM$, is said to be;*

1. ***timelike*** *if $g(X,X) < 0$,*
2. ***lightlike*** *if $g(X,X) = 0$,*
3. ***spacelike*** *if $g(X,X) > 0$.*

A Lorentzian metric on a spacetime also defines light cones, $\mathcal{V}_p$, in the tangent spaces at points by,

$$\mathcal{V}_p \coloneqq \{X \in T_pM : g_p(X,X) = 0\}.$$

At each point there are actually two 'components' of $\mathcal{V}_p$, and we need to distinguish one of these as future pointing. This assignment needs to smooth and defined over the whole spacetime to have a reasonable sense of the 'future'. Such an assignment is a time-orientation.

Definition 1.1.3. *A Lorentzian manifold M is time-orientable if it admits a smooth, timelike, nowhere vanishing global section $T \in \mathfrak{X}(M)$.*

Using the time-orientation we can define whether things are travelling to the future or the past. A vector $X \in T_pM$ is **future directed** at a point $p \in M$ if $g_p(X,T) < 0$. Note that any other timelike, global, nowhere vanishing section is still a valid time-orientation. If we're being proper then we should quotient out by the equivalence relation,

$$T, T' \in \mathfrak{X}(M), \quad T \sim T' \iff g(T,T') < 0,$$

so that then a choice of time-orientation is techinically an equivalence class $[T]_\sim$. Finally, orientability is imposed as a condition for all spacetimes we examine in these notes. Collecting all of this information gives a working definition.

Definition 1.1.4. ***Spacetime*** *is tuple $(M, g, \nabla, \mathrm{d}\mu_g, T)$ where;*

1. *M is a (smooth) four-dimensional manifold,*
2. *$g \in \mathrm{Sym}^2(T^\star M)$ is a Lorentzian metric with signature $(-,+,+,+)$,*
3. *∇ is the Levi-Civita connection associated to g,*
4. *$\mathrm{d}\mu_g \in \Omega^4(M)$ is a volume form on M,*
5. *and $T \in \mathfrak{X}(M)$ is a time-orientation.*

Recall that every Riemannian manifold admits a unique connection which is metric compatible and torsion free. The same is true for a Lorentzian manifold, and this is still called the Levi-Civita connection. In more general theories of gravity one considers arbitrary connections which are perhaps not torsion free etc. This is why we've made it explicit in our definition despite it not being 'additional' specified structure. On the other hand not all manifolds are orientable, this is an extra condition we impose on a valid spacetime. Furthermore, we have the following statements.

Remark 1.1.1. *A smooth manifold may not admit a Lorentzian metric, despite always admitting a Riemannian one.*

Remark 1.1.2. *An orientable manifold may not admit a time-orientation.*

The spacetimes which are allowed is restricted somewhat by the above two facts. More about causality and Lorentzian geometry is discussed in chapter 4.

1.2 Einstein-Hilbert Action

General relativity is a classical field theory where the dynamical field is the metric tensor. This is a crucial difference from other forces modelled by quantum gauge theories, which take place on a fixed background. Einstein originally deduced the equations of motion from physical principles. Another route which was taken by Hilbert —[and maybe also Einstein, check]— is to guess an appropriate Lagrangian and derive the equations of motion from the principle of least action.

Definition 1.2.1. *The* ***Einstein-Hilbert action*** *on a Spacetime M is*

$$S[g] = \int_M \star_g \left(\frac{1}{2\kappa} R + \mathcal{L}_{\mathcal{M}} \right). \tag{1.1}$$

1. $\kappa = 8\pi G c^{-4}$. *Where G is the gravitational constant and c is the speed of light;*
2. $\star_g : \Omega^k(M) \to \Omega^{n-k}(M)$ *is the Hodge dual for the metric g;*
3. *R is the Ricci scalar;*
4. *$\mathcal{L}_{\mathcal{M}}$ is a matter Lagrangian describing the fields and particle distribution.*

It's natural to ask why this specific Lagrangian and not another. One could just say this mathematical formula is the one that gets the correct equations of motion, but there are multiple ways to justify it from physical principles. One such justification is that this action is actually the unique low energy effective gravitational field theory [**?**]. From here on we will make use of local coordinates on spacetime when needed. Denote the chart around a neighbourhood $p \in U \subseteq M$ by $x^\mu \in \mathbb{R}^4$ where $\mu = \{0, 1, 2, 3\}$. Correspondingly, the tangent space basis vectors will be denoted $\{\partial_\mu\}$ and the cotangent basis $\{\mathrm{d}x^\mu\}$. The Einstein summation convention will always be assumed unless stated otherwise. The gravitational constant and the speed of light will be also be set to unity in the appropriate units.

Definition 1.2.2. *The energy momentum tensor, $T_{\mu\nu}$, is defined in terms of the matter Lagrangian by*

$$T_{\mu\nu} := \frac{-2}{\sqrt{-g}} \frac{\delta}{\delta g^{\mu\nu}} \left[\sqrt{-g} \mathcal{L}_{\mathcal{M}} \right]. \tag{1.2}$$

The energy momentum tensor encapsulates the fact matter and energy act as a source for gravity. This quantity won't be used much in these notes, but it is obviously of extreme importance since there is matter in the universe! Moreover, the energy momentum tensor also includes contributions from 'fields' such as the electromagnetic field or scalar fields as well as matter. In a certain sense everything gravitates; even gravity. This is one of the reasons that general relativity is so difficult.

Theorem 1.2.1. *The sationary action principle, $\delta S = 0$, implies Einstein's field equations in local coordinates $x^\mu(p)$:*

$$R_{\mu\nu} - \frac{1}{2} R g_{\mu\nu} = \kappa T_{\mu\nu} \tag{1.3}$$

Proof. The variation of the matter Lagrangian is almost by definition the energy momentum tensor so we'll focus on only the first term of the Lagrangian density.

$$\delta(\star_g R) = \delta\left(\sqrt{g} g^{\mu\nu} R_{\mu\nu}\right) = \delta\left(R_{\mu\nu}\right) \sqrt{g} g^{\mu\nu} + \delta\left(\sqrt{g} g^{\mu\nu}\right) R_{\mu\nu} \tag{1.4}$$

Let's compute the second term first. Consider the variation of the inverse metric $\delta(g^{\mu\nu})$. We start by using the definition of the inverse,

$$g^{\mu\sigma} g_{\sigma\nu} = \delta^\mu_\nu, \tag{1.5}$$

now we can apply the variation to both sides and rearrange:

$$\begin{aligned}
&\delta(g^{\mu\nu} g_{\nu\sigma}) = \delta(g^{\mu\nu}) g_{\nu\sigma} + g^{\mu\nu} \delta(g_{\nu\sigma}) = 0 \\
&\delta(g^{\mu\nu}) g_{\nu\sigma} = -g^{\mu\nu} \delta(g_{\nu\sigma}) \\
&\delta(g^{\mu\nu}) g_{\nu\sigma} g^{\sigma\rho} = -g^{\mu\nu} \delta(g_{\nu\sigma}) g^{\sigma\rho} \\
&\delta(g^{\mu\rho}) = -g^{\mu\nu} \delta(g_{\nu\sigma}) g^{\sigma\rho}
\end{aligned}$$

Next observe we need to work out the variation of the determinant of a metric. This follows from Jacobi's formula,

$$\delta(g) = g\mathrm{Tr}(g^{\mu\alpha}\delta(g_{\alpha\nu})), \tag{1.6}$$

where we can use the chain rule to obtain,

$$\begin{aligned}\delta\sqrt{-g} &= \frac{1}{2\sqrt{-g}}(-g)g^{\mu\nu}\delta(g_{\mu\nu})\\ \delta\sqrt{-g} &= \frac{1}{2}\sqrt{-g}g^{\mu\nu}\delta(g_{\mu\nu}).\end{aligned}$$

We can simplify the calculation of $\delta R_{\mu\nu}$ by using Riemann normal coordinates at $p \in U$, meaning the Christoffel symbols vanish, leaving only

$$\delta R_{\mu\nu} = \partial_\alpha\left(\delta\Gamma^\alpha_{\mu\nu}\right) - \partial_\nu\left(\delta\Gamma^\alpha_{\mu\alpha}\right). \tag{1.7}$$

We can now swap $\partial \to \nabla$ since $\delta\Gamma$ are tensor quantities. This result is the Platini identity.

$$\delta R_{\mu\nu} = \nabla_\alpha\delta\Gamma^\alpha_{\mu\nu} - \nabla_\nu\delta\Gamma^\alpha_{\mu\alpha} \tag{1.8}$$

—[COMPLETE, NOTE BOUNDARY TERM]— □

The field equations are incredibly difficult to solve. They are non-linear, 2nd order, coupled partial differential equations in terms of the metric. The only way to get analytic solutions is in highly symmetric toy models, as we will see in the next two chapters. Yet even then it can be cumbersome! One way to simplify the situation is to consider the case of a vanishing energy momentum tensor,

$$\begin{aligned}R_{\mu\nu} - \frac{1}{2}Rg_{\mu\nu} &= 0\\ \implies R - \frac{1}{2}R\delta^\mu_\mu &= 0\\ \implies R &= 0,\end{aligned}$$

then substitute $R = 0$ back into the vacuum field equations to get,

$$R_{\mu\nu} = 0, \tag{1.9}$$

which is the common form Einstein's vacuum field equations. These notes will be focused mostly on this restricted regime.

1.3 Geodesic Flow

We can model the motion of small objects travelling in spacetime in the absence of forces with geodesics. Since the mass is assumed small we can ignore the back reaction of the mass of the matter on the spacetime/gravitational field. Recall that a **symplectic form** is a closed non-degenerate two form. A manifold with equipped with a symplectic form is a **symplectic manifold**. When such a space also has a designated a smooth energy function, it is a Hamiltonian system. It will be useful to formalise the dynamics here as a Hamiltonian system on the phase space of spacetime,

$T^\star M$. We may also make use of the bundle isomorphism $T^\star M \cong TM$ induced by the metric as $(x^\mu, p_\mu) \mapsto (x^\mu, g^{\mu\nu} p_\nu)$. This lets us pull back the symplectic form to the tangent bundle as,

$$\omega_g = \mathrm{d}x^\mu \wedge (g_{\mu\nu} p^\nu)\,,$$

and then one can consider Lagrangian mechanics. We will stick to Hamilton's formulation for these upcoming definitions.

Definition 1.3.1 (Geodesic Flow)**.** *We have a Hamiltonian system, $(T^\star M, \omega, H)$, where in local coordinates (x^μ, p_μ) we define;*

1. *The **symplectic form**, $\omega \in \Omega^2(T^\star M)$, on $T^\star M$ in local coordinates is given by $\omega = \mathrm{d}x^\mu \wedge \mathrm{d}p_\mu$.*
2. *The Hamiltonian is $H = \frac{1}{2} g^{\mu\nu} p_\mu p_\nu$.*

Definition 1.3.2. *$\exists!\ X_H \in \mathfrak{X}(T^\star M)$ called the **Hamiltonian vector field** satisfying,*

$$\iota_{X_H} \omega = \mathrm{d}H, \tag{1.10}$$

and defining Hamilton's equations of motion.

So, for geodesic flow, the symplectic form is just the canonical one. We need a conserved energy to define a Hamiltonian system, here it is just the usual energy defined from the square of momentum, except using the induced metric. We will write trajectories in spacetime as $x^\mu(\tau)$ as a slight abuse of notation for the coordinates of the curve in a chart, $(x^\mu \circ \gamma)(\tau)$. The components of the tangent vector to the curve at a point is then written $\dot{x}^\mu \partial_\mu$.

Theorem 1.3.1. *Geodesic flow is satisfied by a trajectory $x^\mu(\tau) \iff$*

$$\nabla_{\dot{\gamma}} \dot{\gamma} = 0. \tag{1.11}$$

$$\iff \ddot{x}^\mu + \Gamma^\mu_{\sigma\nu} \dot{x}^\sigma \dot{x}^\nu = 0 \tag{1.12}$$

Proof. Start with Hamilton's equations of motion in Definition 1.3.2, which is equivalent to

$$\dot{x}^\mu = \frac{\partial H}{\partial p_\mu}, \quad \dot{p}_\mu = -\frac{\partial H}{\partial x^\mu}.$$

Then we obtain,

$$\dot{x}^\mu = g^{\mu\nu} p_\nu, \quad \dot{p}_\mu = -\frac{1}{2} \partial_\sigma (g^{\mu\nu}) p_\mu p_\nu.$$

Now take the derivative of the left-hand equation to get,

$$\begin{aligned} \ddot{x}^\mu &= g^{\mu\nu} \dot{p}_\nu + \dot{x}^\sigma \partial_\sigma g^{\mu\nu} p_\nu \\ &= g^{\mu\nu} \dot{p}_\nu + \dot{x}^\sigma \partial_\sigma g^{\mu\nu} g_{\nu\rho} \dot{x}^\rho, \end{aligned}$$

where we've used the chain rule $\frac{\mathrm{d}}{\mathrm{d}\tau} = \dot{x}^\sigma \partial_\sigma$, and the definition of four momentum $p_\nu = g_{\mu\rho} \dot{x}^\rho$. Next substitute the equation for $\dot{p}_\nu$ to get,

$$\ddot{x}^\mu = -\frac{1}{2} \partial_\nu g^{\gamma\delta} p_\gamma p_\delta g^{\mu\nu} + \dot{x}^\sigma \partial_\sigma g_{\nu\rho} \dot{x}^\rho.$$

We need to eliminate the derivative of the inverse metric so note from product rule we have :

$$\begin{aligned}
\partial_\gamma(g_{\mu\nu}g^{\nu\sigma}) &= \partial_\gamma(\delta^\sigma_\mu) = 0 \\
&= g_{\mu\nu}\partial_\gamma g^{\nu\sigma} + g^{\nu\sigma}\partial_\gamma g_{\mu\nu} = 0 \\
\implies g^{\mu\rho}g_{\mu\nu}\partial_\gamma g^{\nu\sigma} &= -g^{\mu\rho}g^{\nu\sigma}\partial_\gamma g_{\mu\nu} \\
\implies \partial_\nu g^{\gamma\delta} &= -g^{\mu\gamma}g^{\sigma\delta}\partial_\nu g_{\mu\sigma}.
\end{aligned}$$

Using this identity we obtain,

$$\begin{aligned}
\ddot{x}^\mu &= \frac{1}{2}g^{\mu\gamma}g^{\sigma\delta}\partial_\nu g_{\mu\sigma}g_{\beta\gamma}\dot{x}^\beta g_{\alpha\delta}\dot{x}^\alpha g^{\mu\nu} - g^{\alpha\mu}g^{\beta\nu}\partial_\sigma g_{\alpha\beta}g_{\nu\rho}\dot{x}^\sigma\dot{x}^\rho \\
\implies \ddot{x}^\mu &= \frac{1}{2}g^{\mu\nu}\dot{x}^\sigma\dot{x}^\alpha\partial_\nu g_{\alpha\sigma} - g^{\alpha\mu}\dot{x}^\sigma\dot{x}^\beta\partial_\sigma g_{\alpha\beta} \\
\ddot{x}^\mu &= \frac{1}{2}g^{\mu\alpha}(\partial_\alpha g_{\sigma\nu} - \partial_\sigma g_{\alpha\nu} - \partial_\nu g_{\sigma\alpha})\dot{x}^\sigma\dot{x}^\nu \\
\implies \ddot{x}^\mu &+ \Gamma^\mu_{\sigma\nu}\dot{x}^\sigma\dot{x}^\nu = 0,
\end{aligned}$$

where we've relabelled indices and recognised the formula for the Christoffel symbols of a Levi-Civita connection. On the other hand note that by definition of the induced covariant derivative on a curve,

$$\begin{aligned}
\nabla\dot{\gamma}\dot{\gamma} = 0 &\iff \dot{x}^\mu\nabla\partial_\mu\dot{x}^\nu = 0 \\
\iff \frac{\mathrm{d}x^\mu}{\mathrm{d}\tau}\frac{\partial}{\partial x^\mu}\frac{\mathrm{d}x^\nu}{\mathrm{d}\tau} &+ \frac{\mathrm{d}x^\mu}{\mathrm{d}\tau}\Gamma^\mu_{\sigma\nu}\frac{\mathrm{d}x^\sigma}{\mathrm{d}\tau} = 0,
\end{aligned}$$

where we've converted to statements for all charts. Then by chain rule we obtain,

$$\ddot{x}^\mu + \Gamma^\mu_{\sigma\nu}\dot{x}^\sigma\dot{x}^\nu = 0.$$

□

Note that we're being somewhat informal in distinguishing between geodesic and so-called co-geodesic flow, since the equations of motion are the same it won't make a difference in these notes. We will identify $g^{\mu\nu}p_\nu = p^\mu := \dot{x}^\mu$ as the **four-momentum**. For any path of a massive particle in spacetime, geodesic or not, we can define the length by integration of the metric.

Definition 1.3.3. *The **length** of curve $\gamma : [a, b] \to M$ in spacetime is defined as :*

$$L(\gamma) := \int_a^b \sqrt{|g(\dot{\gamma}(\lambda), \dot{\gamma}(\lambda))|}\mathrm{d}\lambda$$

In the case of a massive particle we can use this to define the **proper time** elapsed along a trajectory.

Definition 1.3.4. *The **proper time** τ for a massive particle is the arc length parameter associated to the length functional.*

Traditionally this is written as $ds^2 = -d\tau^2$. Recall that in Riemannian geometry geodesics are stationary points of the length functional which locally minimise distances (but not necessarily globally). In Lorentzian geometry geodesics are still stationary points of the new length functional, but now we have the interpretation of locally **maximising** elapsed proper time. Intuitively in the case of special relativity, a stationary particle relative to a frame has no time dilation or anything strange. On the other hand wiggling around causes less proper time to elapsed due to time dilation effects. A similar story is true for general relativity, infinitesimally at least. Note that the concept of things being at rest is a lot more subtle in general relativity, since inertial frames can only be defined locally.

Definition 1.3.5. *A **conserved quantity**, $f \in C^\infty(T^\star M)$, is constant along geodesic flow,*

$$\gamma_\star \partial_\tau \cdot f = \frac{\mathrm{d}}{\mathrm{d}\tau}(f \circ \gamma)(\tau) = 0, \tag{1.13}$$

where $\gamma(\tau) = (x^\mu(\tau), p_\mu(\tau))$ is now a curve in phase space, with the usual abuse of notation for coordinates on the image.

Definition 1.3.6. *A vector field $X \in \mathfrak{X}(T^\star M)$ is a symmetry for the Hamiltonian system if*

1. $X(H) = 0$
2. $\mathcal{L}_X \omega = 0$

Theorem 1.3.2 (Noether)**.** *If $X \in \mathfrak{X}(T^\star M)$ is a symmetry and $H^1(T^\star M) = 0$ then $\exists f \in C^\infty(T^\star M)$ such that*

$$\iota_X \omega = \mathrm{d}f, \tag{1.14}$$

where f is a conserved quantity.

Proof. To show such f exists we just expand the Lie derivative using Cartan's magic formula,

$$\mathcal{L}_X \omega = \iota_X \circ \mathrm{d}\omega + \mathrm{d} \circ \iota_X \omega = 0,$$

but we know that $\mathrm{d}\omega = 0$ since ω is symplectic. Then remaining is,

$$\mathrm{d} \circ \iota_X \omega = 0,$$

which implies, using the triviality of $H^1(T^\star M)$, that $\exists f \in C^\infty(M)$ satisfying $\iota_X \omega = \mathrm{d}f$. Now it remains to show that f is conserved along flows generated by X. We just have to apply the interior product successively to the symplectic form,

$$\begin{aligned} \iota_X \circ \iota_{X_H} \omega = \iota_X \circ \mathrm{d}H = X(H) = 0 \\ -\iota_{X_H} \circ \iota_X \omega = -\iota_{X_H} \circ \mathrm{d}f = 0 \implies X_H(f) = 0. \end{aligned}$$

□

The above is Noether's theorem in Hamiltonian mechanics, which we will make use of this when solving the geodesic equation. —[PERHAPS QUICKLY DEFINE INTEGRABILITY FOR LATER]—Recall that in Riemannian geometry one has the notion of Killing vector fields:

Definition 1.3.7. *A **Killing vector field**, $X \in \mathfrak{X}(M)$, satisfies $\mathcal{L}_X g = 0$.*

Equivalently, a Killing vector field can be thought of as generating infinitesimal isometries along its flow. Note that, along a geodesic, we have that a Killing vector field induces a conserved quantity, $g(\dot{\gamma}, X)$. —[PROVE]—.

The above discussion suffices to describe the path a unit mass particle travels through spacetime, in general we also want to consider massless particles such as photons. In the massive case the Hamiltonian defines the normalisation of momenta by $H = -\frac{1}{2}$, in the massless case the hamiltonian is zero since light rays are null $g(\dot{\gamma}, \dot{\gamma}) = 0$. Furthermore, the length function is identiacally zero so we must find a more suitable parameter to describe the distance travelled. We can actually define the paramater as that which satisfies the geodesic equation.

Definition 1.3.8. *An **affine parameter**, λ, is a parameter which satisfies the geodesic equation,*

$$\frac{\mathrm{d}^2 x^\mu}{\mathrm{d}\lambda^2} + \Gamma^\mu_{\sigma\nu} \frac{\mathrm{d}x^\sigma}{\mathrm{d}\lambda} \frac{\mathrm{d}x^\nu}{\mathrm{d}\lambda} = 0.$$

This parameter is unique up to an affine reparameterisation $\lambda' = a\lambda + b$ where $a, b \in \mathbb{R}$. So even along a light ray, there is still a way to track a 'length' along the geodesic.

1.4 General Covariance

In the abstract mathematical sense a manifold is a perfectly well-defined concept. The extent to which spacetime itself coincides with this definition is not immediatey obvious. When Einstein was originally formulating his theory he struggled with this concept. In this section we'll briefly talk about some of these issues and resolutions.

In general relativity, and physics as whole, we require laws to be **generally covariant**. —[DEFINE]—

The resolution to the paradox comes from realising that different field configurations have to be deemed equivalent if they are related to each other by an isometry. In other words we must quotient the space of metrics by some —[COMPLETE]—

1.5 Moving Frames

1.5.1 Klein Geometry of Minkowski Space

This section will give introduction to Cartan geometry in the context of GR. An exposition can be found in [Cat15] and [Wis10]. A full overview of the geometry is in [Sha02]. In Minkowski spacetime there is a group of symmetries which preserves the metric, the isometry group. In particular, we have translational invariance and Lorentz invariance. Combining these two together gives us the isometry group of Minkowski space; the **Poincaré group** $\mathrm{ISO}(3,1)$. We denote the connected component of the identity of the Lorentz group by $\mathrm{SO}_0(3,1)$, and likewise for the Poincaré group $\mathrm{ISO}_0(3,1)$. We may get sloppy when referring to the Lorentz group as opposed to the connected component of the identity. Both subgroups of the Poincaré group have natural right actions on the whole group itself. We can study the coset spaces of these actions to understand geometric aspects of Minkowski spacetime.

Consider the action of the three-dimensional Euclidean Poincaré group $\mathrm{ISO}(3)$ on the connected Minkowski Poinacaré group. The coset space $\mathrm{ISO}_0(3,1)/\mathrm{ISO}(3)$ is the stabiliser of this action.

Hence, this coset space characterises the set of **spacelike hyperplanes** in Minkowski spacetime. On the other hand consider the action of connected Lorentz group on the connected POincaré group. In this case we're 'modding out' by all rotations and boosts, hence the stabilisers are just points. The coset space $\mathrm{ISO}_0(3,1)/\mathrm{SO}_0(3,1)$ characterises all the **events** in Minkowski spacetime.

The above exaples are special cases of a general idea by Felix Klein.

Definition 1.5.1. *A **Klein geometry**, (H, G), consists of a Lie group G and a closed subgroup H such that the coset space is connected.*

Sometimes this definition is relaxed by allowing for disconnected Klein geometries, we will not consider these here. Note that from Lie theory we know that a closed subgroup of a Lie group is automatically a Lie subgroup. Furthermore, the coset space $(G/H = X)$ is a smooth manifold, which can be acted on transitatively on the left by G with group multiplication. In this case the pair G, X has X as a **homogenous space**. The subgroup also Hhas the natural right action on G induced from group multiplication. One can use this action to construct a bundle over the coset space $G/H$with fibres as left cosets of H. Now the right action is free and transitative, so the coset space is a **principal homogenous space** or **H-torsor**. In this case the fibre bundle is actually a **principal H-bundle** over G/H.

Recall that Lie group always has an adjoint left action on its Lie algebra. Equivalently we have a representation of H on the underlying vector space of $\mathfrak{g}/\mathfrak{h}$. Using this data we can construct the vector bundle associated to the principal H-bundle via the adjoint representation, $G \times_{\mathrm{Ad}_H} \mathfrak{g}/\mathfrak{h}$. On the other hand, since G/H is a smooth manifold, we can also consider its tangent bundle $T[G/H] \to G/H$. It turns out that these two vector bundles are canonically isomorphic [Sha02].

$$\begin{tikzcd}
T[G/H] \arrow[rr, "\sim"] \arrow[dr] & & G \times_{\mathrm{Ad}_H} \mathfrak{g}/\mathfrak{h} \arrow[dl] \\
& G/H &
\end{tikzcd}$$

1.5.2 Cartan Geometry

In Minkowski spacetime we have a global Poincaré symmetry since there is no dependence on the points of the manifold. In gauge theory a powerful principle is to take a global symmetry and turn it into a local, or gauge, symmetry. A common example is the theory of electrically charged particles. Gauging the global U(1)-symmetry imposes the condition that an electromagnetic force appears. One can play the same game with the Poincaré symmetry of Minkowski spacetime. Gauging this global symmetry also causes a new force to appear; gravity.

One way to formally describe this process is through the reduction of a structure group. To every smooth manifold one can consider the principal frame bundle over it. Specifically this comes from the fact that every smooth manifold comes equipped with a metric, so one can define an $\mathrm{O}(n)$ principal bundle over the space. This $\mathrm{O}(n)$-bundle can be thought of as arising from the reduction of the structure group of a principal $\mathrm{GL}(n, \mathbb{R})$-bundle. The same idea works for the Lorentz group in indefinite signature. Using Klein geometries we can alternatively capture the reduction of structure group, as well as the connection data, in one object. Promoting to Cartan geometries is to consider a space modelled locally as a Kelin geometry, analagous to gauging the symmetry.

Definition 1.5.2. *A $H \hookrightarrow G$ **Cartan connection** over a manifold M consists of the following data associated to a principal H-bundle $P_H \to M$.*

- *A principal H-connection, $A \in \Omega^1(P_H; \mathfrak{g})$, such that;*

1. *there is a reduction of structure groups through $H \hookrightarrow G$,*
2. *and the 1-form $A : TP_H \longrightarrow \mathfrak{g}$, induces an isomorphism $T_pP_H \cong \mathfrak{g}$, $\forall p \in P_H$.*

A space equipped with Cartan connection is a **Cartan geometry**. At each point there is a given fibre of the model Klein geometry. We can equip this space with a metric differs from a Riemannian metric on a manifold in general. Here it characterises the size of the Klein geometry, and may contain some length scale relvant to the spacetime.

Definition 1.5.3. *A **metric** for a Klein geometry, (G, H), is a potentially degenerate Ad_H-invariant metric on $\mathfrak{g}/\mathfrak{h}$.*

If Riemannian curvature measures the deviation of a manifold from being flat, then Cartan geometry measures the deviation of a geometry from the model homogenous space.

Definition 1.5.4. *The **curvature** associated to a Cartan connection is defined by,*

$$F = \mathrm{d}A + \frac{1}{2}[A \wedge A]. \tag{1.15}$$

We say that a Cartan geometry is **reductive** if the direct sum $\mathfrak{g} \cong \mathfrak{g}/\mathfrak{h} \oplus \mathfrak{h}$ is Ad_H-invariant. In our case we consider the pair $\mathrm{SO}(3,1) \hookrightarrow \mathrm{ISO}(3,1)$ as the 'model geometry'. Now the Lie algebra of the Poincaré group has a natural decomposition as,

$$\mathfrak{iso}(3,1) \cong \mathbb{R}^4 \oplus \mathfrak{so}(3,1).$$

Then the cartan connection can be decomposed as $A = \sigma + \omega$, where σ is a coframe field and ω is a principal $\mathrm{SO}(3,1)$-connection. Note that the vielbein is usually denoted by the letter, e, especially in the physics literature. Diagramatically we have:

$$\begin{tikzcd}
 & TP \arrow[d, "A"] \arrow[ddl, bend right, "\sigma"'] \arrow[ddr, bend left, "\omega"] & \\
 & \mathfrak{iso}(3,1) \arrow[dl] \arrow[dr] & \\
\mathbb{R}^4 & & \mathfrak{so}(3,1)
\end{tikzcd}$$

Assuming that the Lie algebra $\mathfrak{g}/\mathfrak{h}$ is abelian, then we can decompose the curvature as $F = \tau + \Omega$. Here Ω is the usual Riemann curvature 2-form and τ is the torsion.

$$\begin{tikzcd}
 & \bigwedge^2(TP) \arrow[d, "F"] \arrow[ddl, bend right, "\tau"'] \arrow[ddr, bend left, "\Omega"] & \\
 & \mathfrak{iso}(3,1) \arrow[dl] \arrow[dr] & \\
\mathbb{R}^4 & & \mathfrak{so}(3,1)
\end{tikzcd}$$

Then Cartan's first structure equation is the vanishing of the torsion,

$$\tau^\alpha := \mathrm{d}\sigma^\alpha + \omega^\alpha_\beta \wedge \sigma^\beta. \tag{1.16}$$

Cartan's second structure equation gives a formula for the curvature in terms of only the connection,

$$\Omega^\alpha_\beta = \mathrm{d}\omega^\alpha_\beta + \omega^\alpha_\gamma \wedge \omega^\gamma_\beta. \tag{1.17}$$

The traditional Riemann curvature tensor is related to the curvature 2-form by,

$$\Omega^\alpha_\beta = \frac{1}{2} R^\alpha_{\beta\mu\nu} \mathrm{d}x^\mu \wedge \mathrm{d}x^\nu, \tag{1.18}$$

in a coordinate basis. To simplify computations we like to stay in the orthonormal basis defined by the vielbeins as opposed to the coordinate basis. In this basis we have some useful symmetry properties. The Lie algebra $\mathfrak{so}(3,1)$ satisfies [Frè13],

$$\omega \in \mathfrak{so}(3,1) \iff \omega^T \eta + \eta\omega = 0,$$

where η is the Minkowski metric. In a basis this can be written as $\omega^\gamma_\alpha \eta_{\beta\gamma} + \omega^\gamma_\beta \eta_{\alpha\gamma} = 0$. If we make the definition, $\omega_{\alpha\beta} := \eta_{\alpha\gamma}\omega^\gamma_\beta$, then this relation is often written,

$$\omega_{\alpha\beta} = -\omega_{\beta\alpha}. \tag{1.19}$$

Since we have a Lorentzian signature ω^α_β is not antisymmetric as in the Riemannian case. It picks up minus signs when swapping 'spatial' indices, but is symmetric under the interchange of the time and space indices. Note that in particular, the spin connection has $d^2(d-1)/2$ independant components, but the Christoffel symbols have $d^2(d+1)/2$. The fact there are less independant components is one of the reasons why this formalism is easier to work with for computations. In practice one starts with the coframe field σ^α_μ, which is obtained from (and equivalent to) the metric ansatz. Then we compute derivatives and write all expressions in the orthonormal frame. Using Cartan's structure equations we can deduce the forms of the connection $\omega^\alpha_{\mu\beta}$ and the curvature form $\Omega^\alpha_{\beta\mu\nu}$. Working with the vielbein field σ^α_μ and connection form $\omega^\alpha_{\mu\beta}$ is equivalent to working with a metric. The vilebein encodes the Lorentzian metric through,

$$g = -\sigma^0 \otimes \sigma^0 + \sum_{i=1}^{3} \sigma^i \otimes \sigma^i. \tag{1.20}$$

This is usually written,

$$g_{\mu\nu} = \sigma^\alpha_\mu \sigma^\beta_\nu \eta_{\alpha\beta}, \tag{1.21}$$

in local coordinates, which is common in the physics literature. It's also common to treat the vielbein field (and their dual frame field) as an object which 'converts' spacetime indices to internal 'flat' indices. For example a four-vector could be expressed as,

$$A^\alpha = \sigma^\alpha_\mu A^\mu,$$

in a local orthonormal frame. Of course, this is just a change of basis applied to each ∂_μ in the tensor, which also works for the dual basis $\mathrm{d}x^\mu$ and frame fields. What is sometimes not emphasised in the physics literature is that these fields may not be well-defined **globally**. Indeed, it would be true

if the manifold admits a **framing**, but this need not be a requirement for an arbitrary Lorentzian manifold. Globally hyperbolic spacetimes, discussed in chapter 4, do actually enjoy this property howerver.

The above formalism is extremely insightful, but there's no new physics apparent yet, so what's the point? Beyond a better understanding of the theory, there are at least a few good reasons. The first is that it enables us to perform computations much easier, taking advantage of antisymmetry and differential forms. Another is that the universe contains fermions. We won't describe the details here, but the metric tensor alone isn't enough to describe spinor fields. In order to have a consistent unified theory, one must inevitably pass to the vielbein formalism to account for fermion fields in curved spacetime. This is especially true for supergravity and supersymmetric string theories, in which generalisations of the vielbein play important role. Even in classical gravity we can use Cartan geometry to decribe spacetimes which are not locally Minkowski. For example de Sitter space is described by the Klein geometry $\mathrm{SO}(4,1)/\mathrm{SO}(3,1)$. Then it might make sense to consider a spacetime which is locally modelled on this geometry as opposed to flat Minkowski space. This approach is useful in MacDowell-Mansouri gravity [Wis10]. For the moment, cosmological aspects of GR won't be discussed in these notes. One can assume that, when a vielbein field is introduced, we are working with the Cartan geometry modelled on $\mathrm{ISO}(3,1)/\mathrm{SO}(3,1)$.

Chapter 2

Schwarzschild Spacetime

2.1 Derivation of the Metric

The Schwarzchild solution is a static and spherically symmetric metric aiming to model the spacetime around a massive object. A derivation can be found in [Str13], we will follow this and also include explicit computations for all quantities. By definition, a static spacetime admits a timelike killing vector field, and we can decompose the manifold as a product,

$$M = \mathbb{R}_t \times \Sigma,$$

for some spatial manifold Σ. In particular the metric has the form,

$$g = -f^2 \mathrm{d}t \otimes \mathrm{d}t + h,$$

where $f \in C^\infty(\Sigma)$ and h is a Riemannian metric on Σ. It is a vacuum solution and therefore satisfies $R_{\mu\nu}$. A spacetime is spherically symmetric if there is an SO(3) action which induces an isometry on the metric. Furthermore, we require that the orbits of this action are spacelike hypersuraces. If the spacetime is also static, as in this case, then the action is trivial on $\mathbb{R}_t$. This works when we explicitly make the assumption the timelike vector field is unique, so that there is a distinguished time coordinate. If ∂_t is unique then the SO(3) action must leave it invariant. The orbits of an isomtric SO(3) action are homogenous Riemannian manifolds of two dimensions. In this case it must have constant curvature, which implies that it is geodsically complete. Furthermore, this means is must be simply-connected.

Recall that the standard simply-connected surface is either the 2-sphere, the flat Euclidean plane, or the Hyperbloic plane depending on the sign of the curvature. There is a known result that there exists a covering of our orbits by the standard surface such that the covering map is a local isometry. Since SO(3) is a subgroup of the isometry group of the orbit space, then the curvature must be strictly positive. From this we deduce the orbit space has to be the 2-sphere, S^2. Now we have a foliation of the spatial manifold as $\mathbb{R}_r \times S^2$, where r is the radial coordinate. This **defined** through the area of the orbit sphere at that point, $A = 4\pi r^2$. Now we can write an ansatz for the metric.

$$g = -e^{2a(r)} \mathrm{d}t \otimes \mathrm{d}t + e^{2b(r)} \mathrm{d}r \otimes \mathrm{d}r + r^2 \mathrm{d}\theta \otimes \mathrm{d}\theta + r^2 \sin^2(\theta)\, \mathrm{d}\varphi \otimes \mathrm{d}\varphi \tag{2.1}$$

The spherical part of the metric is the standard round metric for a two-sphere of radius r. Note that we use exponentials since the functions have to be positive, we demand that $\mathrm{d}t \neq 0 \neq \mathrm{d}r$. The

functions $a(r)$ and $b(r)$ do not depend on the other variables by the symmetry conditions. We will also impose that these functions satisfy **asymptotic flatness**, so that $r \to \infty \implies g \to \eta$. This is physically justified since we expect the gravitational field to taper off to zero at large distances. We won' t use this until near the end of the derivation, but this is enforced by demanding that $a(r) \to 0$ as $r \to \infty$ since $e^a \to 1$; and similarly for $b(r)$. Furthermore, we include a factor of two for convenience in computation. Now we need to compute the Einstein tensor for the spacetime. We can read the coframes directly from the metric, $\sigma^\alpha = \sigma^\alpha_\mu \mathrm{d}x^\mu$ as;

$$\sigma^0 = e^{a(r)}\mathrm{d}t, \quad \sigma^1 = e^{b(r)}\mathrm{d}r, \quad \sigma^2 = r\mathrm{d}\theta, \quad \sigma^3 = r\sin(\theta)\,\mathrm{d}\phi.$$

Taking derivatives we obtain;

$$\mathrm{d}\sigma^0 = a' e^{-b}\sigma^1 \wedge \sigma^0, \quad \mathrm{d}\sigma^1 = 0,$$
$$\mathrm{d}\sigma^2 = \frac{e^{-b}}{r}\sigma^1 \wedge \sigma^2, \quad \mathrm{d}\sigma^3 = \frac{1}{r}\left(e^{-b}\sigma^1 \wedge \sigma^3 + \cot(\theta)\,\sigma^2 \wedge \sigma^3\right).$$

Now we use Cartan's first structure equation, $\mathrm{d}\sigma^\alpha = -\omega^\alpha_\beta \wedge \sigma^\beta$, to determine the connection form.

$$\mathrm{d}\sigma^0 = -\omega^0_1 \wedge \sigma^1 \implies \omega^0_1 = a' e^{-b}\sigma^0 = \omega^1_0$$
$$\mathrm{d}\sigma^2 = -\omega^2_1 \wedge \sigma^1 \implies \omega^2_1 = \frac{e^{-b}}{r}\sigma^2 = -\omega^1_2$$
$$\mathrm{d}\sigma^3 = -\omega^3_1 \wedge \sigma^1 - \omega^3_2 \wedge \sigma^2 \implies \omega^3_1 = \frac{e^{-b}}{r}\sigma^3 = -\omega^1_3,\ \omega^3_2 = \frac{\cot(\theta)}{r}\sigma^3 = -\omega^2_3$$

The curvature 2-form is computed using Cartan's second structure equation, $\Omega^\alpha_\beta = \mathrm{d}\omega^\alpha_\beta + \omega^\alpha_\gamma \wedge \omega^\gamma_\beta$.

$$\begin{aligned}\Omega^0_1 &= \mathrm{d}\omega^0_1 + \omega^0_\gamma \wedge \omega^\gamma_1 = \mathrm{d}\left(a' e^{-b}\right) \wedge \sigma^0 + a' e^{-b}\mathrm{d}\sigma^0 \\ &= \left(a'' e^{-b} - a' b' e^{-b}\right) \wedge \sigma^0 + (a')^2 e^{-b}\sigma^1 \wedge \sigma^0 \\ &\implies \Omega^0_1 = -e^{-2b}\left(a'' - a' b + (a')^2\right)\sigma^0 \wedge \sigma^1\end{aligned}$$

$$\begin{aligned}\Omega^0_2 &= -\omega^0_1 \wedge \omega^1_2 = -a' e^{-b}\sigma^0 \wedge \frac{1}{r}e^{-b}\sigma^2 \\ &\implies \Omega^0_2 = \frac{a'}{r}e^{-2b}\sigma^0 \wedge \sigma^2\end{aligned}$$

$$\begin{aligned}\Omega^0_3 &= \omega^0_\gamma \wedge \omega^\gamma_3 = -a' e^{-b}\sigma^0 \wedge \frac{1}{r}e^{-b}\sigma^3 \\ &\implies \Omega^0_3 = -\frac{a'}{r}e^{-2b}\sigma^0 \wedge \sigma^3\end{aligned}$$

$$\begin{aligned}\Omega^1_2 &= -\mathrm{d}\left(\frac{e^{-b}}{r}\right) \wedge \sigma^2 - \frac{e^{-b}}{r}\mathrm{d}\sigma^2 \\ &= \left(\frac{e^{-b}}{r^2} + \frac{b'}{r}e^{-b}\right)\mathrm{d}r \wedge \sigma^2 - \frac{e^{-}2b}{r}\left(\frac{1}{r} + b'\right)\sigma^1 \wedge \sigma^2 - \frac{e^{-2b}}{r^2}\sigma^1 \wedge \sigma^2 \\ &\implies \Omega^1_2 = \frac{b'}{r}e^{-2b}\sigma^1 \wedge \sigma^2\end{aligned}$$

$$\Omega_3^2 = -\mathrm{d}\left(\frac{\cot(\theta)}{r}\right) \wedge \mathrm{d}\sigma^3 - \frac{\cot(\theta)}{r}\mathrm{d}\sigma^3 - \frac{e^{-2b}}{r^2}\sigma^2 \wedge \sigma^3$$

$$= \frac{1}{r^2}\cot(\theta)\,\mathrm{d}r \wedge \sigma^3 + \frac{1}{r\sin^2(\theta)}\mathrm{d}\theta \wedge \sigma^3 - \frac{\cot(\theta)}{r^2}\left(e^{-b}\sigma^1 \wedge \sigma^3 + \cot(\theta)\,\sigma^2 \wedge \sigma^3\right) - \frac{e^{-2b}}{r^2}\sigma^2 \wedge \sigma^3$$

$$\implies \Omega_3^2 = \frac{1-e^{2b}}{r^2}\sigma^2 \wedge \sigma^3$$

Next we can compute the components of the Ricci tensor using $R_{\alpha\beta} = \Omega_\alpha^\gamma(\pi_\gamma, \pi_\beta)$, where $\{\pi_\alpha\}_{\alpha=0}^3$ is the dual basis to the coframes. For convenience write $A = -e^{-2b}\left(a'' - a'b + (a')^2\right)$.

$$R_{00} = \Omega_0^\gamma(\pi_\gamma, \pi_0) = -A + \frac{2a'}{r}e^{-2b}$$

$$R_{11} = \Omega_1^\gamma(\pi_\gamma, \pi_1) = A + \frac{2b'}{r}e^{-2b}$$

$$R_{22} = \Omega_2^\gamma(\pi_\gamma, \pi_2) = \frac{e^{-2b}}{r}(b' - a') + \frac{1}{r^2} - \frac{e^{-2b}}{r^2} = R_{33}$$

The Ricci scalar is then just the contraction with the Minkowski metric.

$$R = 2A + 4\frac{e^{-2b}}{r}(b' - a') + \frac{2}{r^2} - \frac{2e^{-2b}}{r^2}$$

Finally, the Einstein tensor components, $G_{\alpha\beta} = R_{\alpha\beta} - \frac{1}{2}Rg_{\alpha\beta}$, are as follows:

$$G_{00} = \frac{1}{r^2} - e^{-2b}\left(\frac{1}{r^2} - \frac{2b'}{r}\right)$$

$$G_{11} = -\frac{1}{r^2} + e^{-2b}\left(\frac{1}{r^2} + \frac{2a'}{r}\right)$$

$$G_{22} = e^{-2b}\left((a')^2 - a'b' + a'' + \frac{a' - b'}{r}\right) = G_{33}$$

Now we only have two functions to solve for, namely $a(r)$ and $b(r)$, using $G_{\alpha\beta} = 0$. Note that,

$$G_{00} + G_{11} = \frac{2(b' + a')}{r} = 0 \implies b + a = C, \tag{2.2}$$

but we imposed asymptotic flatness meaning the integration constant C has to be zero to keep $a, b \to 0$. This means we can eliminate $a = -b$ and look at G_{00}.

$$\frac{1}{r^2} - e^{-2b}\left(\frac{1}{r^2} - \frac{2b'}{r}\right) = 0 \tag{2.3}$$

Multiplying through by r^2 we get,

$$1 = e^{-2b} - 2b're^{-2b}, \tag{2.4}$$

where we can recognise the right-hand side as $(re^{-2b})'$. Now we can integrate both sides and pick up a constant $2m$,

$$r = re^{-2b} + 2m \implies e^{-2b} = 1 - \frac{2m}{r},$$

leaving us finally with the full metric

$$g = -\left(1 - \frac{2m}{r}\right) \mathrm{d}t \otimes \mathrm{d}t + \left(1 - \frac{2m}{r}\right)^{-1} \mathrm{d}r \otimes \mathrm{d}r + r^2 \mathrm{d}\theta \otimes \mathrm{d}\theta + r^2 \sin^2(\theta) \mathrm{d}\phi \otimes \mathrm{d}\phi. \tag{2.5}$$

One can show that m should be equal to GM/c^2, where G is Newton's gravitational constant and M has the interpretation as the mass of the central body. The reason is that we expect GR to reduce to Newtonian gravity in a sufficient limiting scenario. This is called the weak field approximation. In partiular we consider non-relativistic velocities and suppose that the gravitational field is weak, $g \approx \eta + h$, with $|h| \ll 1$. Then the time component of the metric satisfies,

$$g_{00} \approx -\left(1 + \frac{2\Phi}{c^2}\right).$$

Here, Φ is the Newtonian gravitational potential obeying Poisson's equation,

$$\nabla^2 \Phi = 4\pi G \rho,$$

for some mass distribution ρ.

2.2 Chart Problems

Importantly, there are two singularities appearing the above metric. The first of which occurs at the so-called event horizon $r = 2m$.

$$r \to 2m \implies g_{tt} \to 0,\, g_{rr} \to \infty$$

The second singularity occurs when $r = 0$.

$$r \to 0 \implies g_{tt} \to \infty,\, g_{rr} \to 0$$

Upon close analysis it's revealed that the first is actually just a coordinate singularity, meaning our choice of chart is at fault here. In particular there exists a change of coordinates such that the metric is well-behaved at the event horizon, which will be discussed momentarily. On the other hand, the singularity at $r = 0$ is very real, no matter what chart we use this is a problematic point in spacetime and must be removed. We must modify our definition of chart to account for this, in particular the image of the r coordinate map should be:

$$\mathrm{Im}(x^r) = (0, 2m) \cup (2m, \infty) \subset \mathbb{R}$$

Now, just because our chart doesn't see the points at $r = 2m$, it doesn't mean that this is not a part of the spacetime. We can extend the chart to cover the whole manifold, excluding the singularity, using the following change of coordinates:

$$u(t, r) = \sqrt{\frac{r}{2m} - 1} e^{r/4m} \cosh\left(\frac{t}{4m}\right) \tag{2.6}$$

$$v(t, r) = \sqrt{\frac{r}{2m} - 1} e^{r/4m} \sinh\left(\frac{t}{4m}\right) \tag{2.7}$$

These are the **Kruskal** transformations. It's not immediately clear where these come from, but they can actually be derived from physical principles, as is done in [Str13]. Nothing special happens with the θ and φ coordinates so we can ignore those. Analysis of the transformed coordinates shows that u and v are in fact well-defined at $r = 2m$ and there is no longer a singularity. We can now consider the metric in the new coordinates with the points at $r = 2m$ included,

$$g = \frac{32m^3}{r} e^{-r/2m} (\mathrm{d}v \otimes \mathrm{d}v - \mathrm{d}u \otimes \mathrm{d}u) + r^2 (\mathrm{d}\theta \otimes \mathrm{d}\theta + \sin^2(\theta) \mathrm{d}\varphi \otimes \mathrm{d}\varphi). \tag{2.8}$$

The previous discussion highlights that singularities in general relativity are a delicate matter. They will be discussed more in Chapter 4.

2.3 Geodesics in Schwarzschild Spacetime

We will stick to the Lagrangian description for now, which is equivalent to the Hamiltonian description in chapter 1. We will make use of the abuse of notation, $x^\mu(\tau) \coloneqq x^\mu \circ \gamma(\tau)$, for the coordinates along a curve in spacetime, $\gamma : \mathbb{R} \to M$. The Lagrangian is then,

$$2\mathcal{L} = g_{\mu\nu} \dot{x}^\mu \dot{x}^\nu = -1, \tag{2.9}$$

where the dot denotes differentiation with respect to proper time. Beacuse there is spherical symmetry, geodesics are confined to stay on a fixed plane. Without loss of generality we can fix this plane to be $\theta = \pi/2$ to simplify the setup. We then obtain the Lagrangian from the metric as,

$$2\mathcal{L} = -\left(1 - \frac{2m}{r}\right) \dot{t}^2 + \left(1 - \frac{2m}{r}\right)^{-1} \dot{r}^2 + r^2 \dot{\varphi}^2. \tag{2.10}$$

Recall that symmetries of the metric, killing vectors, correspond to conserved quantities along geodesic flow. So, before even writing the geodesic equations, we have by definition two conserved quantities:

$$\begin{aligned} \mathcal{L}_{\partial_t} g = 0 &\implies -\left(1 - \frac{2m}{r}\right) \dot{t}^2 \coloneqq -E \\ \mathcal{L}_{\partial_\varphi} g = 0 &\implies r^2 \dot{\varphi} \coloneqq L \end{aligned}$$

These correspond to an energy and angular momentum of the particle. Now we substitute the energy back into eq. (2.10) to obtain,

$$\dot{r}^2 + \left(1 - \frac{2m}{r}\right)\left(1 + \frac{L^2}{r^2}\right) = E^2. \tag{2.11}$$

Note also that we can express $\dot{r}$ in terms of $r' \coloneqq \frac{\mathrm{d}r}{\mathrm{d}\varphi}$ with the chain rule, $\dot{r} = r'\dot{\varphi}$. Then substiuting the formula for angular momentum we have,

$$\frac{\mathrm{d}r}{\mathrm{d}\tau} = \frac{L}{r^2} \frac{\mathrm{d}r}{\mathrm{d}\varphi},$$

which can be substituted back into $2\mathcal{L} = -1$ to get:

$$\left(\frac{\mathrm{d}r}{\mathrm{d}\varphi}\right)^2 \frac{L^2}{r^4} + \left(1 + \frac{2m}{r}\right)\left(1 - \frac{L^2}{m}\right) = E^2. \tag{2.12}$$

This can be solved exactly by using the transformation $u = 1/r$, but we won't do this here. The upshot is that, for massive particles, the geodesics are ellipses when projected to a three dimesnional spatial slice. This agrees with what's expected from Newtonian gravity. The difference, however, is that the ellipses exhibts **precession** over time. One of the first predictions of general relativity was in fact the famous 'anomalous perihelion of Mercury', which can be derived with remarkable accuracy using Einstein's theory.

In preparation for the Kerr metric, it will be useful exercise to work with the Hamiltonian formalism and derive the equations of motion again. —[DO THIS]—

—[SHOW PARTICLES CAN'T ESCAPE BLACK HOLE]—.

Chapter 3

Kerr Spacetime

3.1 Metric

Kerr spacetime is again a vacuum solution to the Einstein's field equations. It models the field generated by rotating matter, and like the Schwarzschild case it contains a singularity. In the Schwarzschild case we had a spherical symmetry, in this case we loosen this to an axial one. The metric will also be static, hence there will be something similar to a timelike killing field. Note that this is slightly strange considering the mass is spinning constantly (and hence not 'static'), but we're assuming that the angular momentum is fixed for all time. —[ASYMPTOTIC TIMELIKE]—

There's not really a quick and conceptually intuitive way to derive this metric, unlike in the Schwarzschild case. There is a derivation in Straumann [Str13] which serves the purpose of being more general in that at can be extended to account for rotating stars as well, as opposed to just the vacuum surrounding a rotating black hole solution. A quicker but less conceptual derivation can be done 'analytically rotating' into complex coordinates covered in [New73]. One can also 'brute force' the calculation by considering an ellipsoidal ansatz [NP13], which is more intuitive but perhaps tedious in terms of computation. All of the above approaches result in the same metric regardless, given by the following expression:

$$ds^2 = -\frac{\Delta}{\Sigma}(dt - a\sin^2(\theta)d\phi)^2 + \frac{\sin^2(\theta)}{\Sigma^2}((r^2+a^2)d\phi - adt)^2 + \frac{\Sigma}{\Delta}dr^2 + \Sigma^2 d\theta^2 \tag{3.1}$$

Here, $\Delta = r^2 - 2Mr + a^2$, $\Sigma = r^2 + a^2\cos^2(\theta)$, and $a = \frac{J}{M}$ is the Kerr parameter.

3.2 Analysing the Geometry of Kerr Spacetime

3.3 Integrability of Kerr Geodesic Flow

Kerr geodesic flow has three obvious constants of motion. Since we have geodesic flow there is the Hamiltonian itself, or the rest mass of the particle. These arise from three obvious symmetries of the metric. Another conserved quantity arises from the axial symmetry generated by ∂_φ, implying that there is a conserved momentum $p_\varphi = g_{\mu\varphi}p^\mu$. Furthermore, there is a timelike killing field which induces a conserved p_t associated to the energy. There is in fact a third 'hidden symmetry'

associated to the total phase space $T^\star M$. We say hidden since this is not a lift of a symmetry in coniguration space, like usual translations along coordinates for example. The existence of such a transformation is usually hard to deduce, especially in 8-dimensional curved phase space.

Instead, we can go the other way and infer the existence of a global conserved quantity. This is known as Carter's constant. Then, by Noether's theorem, this does correspond to a symmetry vector field in phase space. Though, the symmetry vecror field itself won't be as important to the following discussion. Recall that the existence of four Poisson commuting and independant conserved quantities makes geodesic flow **integrable**. In particular, this means that we can now solve the geodesic equation exactly.

3.3.1 Hamilton-Jacobi equation

Chapter 4

Causality & Topology

4.1 Obstructions to Spacetime

The main goal of this chapter will be to understand the causal nature of spacetime and cover Penrose's singularity theorem. We will mainly follow Witten's [Wit20], as well as Galloway's [Gal14] lectures. Note that we won't be proving every statement carefully, such proofs can be found in [O'N83] and references therein. By spacetime, we mean as defined in Definition 1.1.4, but we allow arbitrary $d \coloneqq \dim M \geq 2$. There are a lot of subtelties involved in determing whether a manifold is/can be a spacetime. For example, not all Lorentzian manifolds are time-orientable, even if the underlying manifold is orientable. Consider the manifold $\mathbb{R} \times [0, 2\pi)$ with opposite sides indentified, $\mathbb{R} \times S^1$. In this case the cylinder is of course orientable, but a time-orientation cannot always be chosen. Consider the metric given by,

$$g = -\mathrm{d}t \otimes \mathrm{d}t + \mathrm{d}\theta \otimes \mathrm{d}\theta,$$

where $\theta \sim \theta + 2\pi$ is the standard coordinate on the circle. It's clear that the vector ∂_t defines a time-orientation since it's globally well-defined and nowhere vanishing. On the other hand consider the following counter-example.

Proposition 4.1.1. *The manifold $\mathbb{R} \times S^1$ is Lorentzian when equipped with the metric,*

$$g = -\cos(\phi)\mathrm{d}x \otimes \mathrm{d}x + \sin(\phi)(\mathrm{d}x \otimes \mathrm{d}\phi + \mathrm{d}\phi \otimes \mathrm{d}x) + \cos(\phi)\mathrm{d}\phi \otimes \mathrm{d}\phi,$$

yet $(\mathbb{R} \times S^1, g)$ is not time-orientable.

Proof. To show that $\mathbb{R} \times S^1, g$ is Lorentzian we need to show that it can be put in diagonal form with eigenvalues $(-1, +1)$. Consider the following vector fields,

$$\sigma_0 \coloneqq \cos(\varphi)\partial_x - \sin(\varphi)\partial_\phi; \quad \sigma_1 \coloneqq \sin(\varphi)\partial_x + \cos(\varphi)\partial_\phi,$$

where $\varphi \coloneqq \phi/2$ for convenience. We would like to show that these define orthonormal frames for

the metric. First let's show that $g(\sigma_0, \sigma_0) = -1$.

$$\begin{aligned} g(\sigma_0, \sigma_0) &= -\cos\phi\, \mathrm{d}x\big(\cos\varphi\partial_x - \sin\varphi\partial_\phi\big)^2 \\ +2\sin\phi\, \mathrm{d}x\big(\cos\varphi\partial_x - \sin\varphi\partial_\phi\big) &\otimes \mathrm{d}\phi\,(\cos\varphi\partial_x - \sin\varphi\partial_\phi) + \cos\phi\, \mathrm{d}\phi\big(\cos\varphi\partial_x - \sin\varphi\partial_\phi\big)^2 \\ &= -\cos\phi\cos^2\varphi - 2\sin\phi\cos\varphi\sin\varphi + \cos\phi\sin^2\varphi \\ &= -\cos\phi(\cos^2\varphi - \sin^2\varphi) - \sin^2\phi = -1 \end{aligned}$$

Similarly, for $\langle\sigma_1, \sigma_1\rangle$;

$$\begin{aligned} g(\sigma_0, \sigma_0) &= -\cos\phi\, \mathrm{d}x\big(\sin\varphi\partial_x + \cos\varphi\partial_\phi\big)^2 \\ +2\sin\phi\, \mathrm{d}x\big(\sin\varphi\partial_x + \cos\varphi\partial_\phi\big) &\otimes \mathrm{d}\phi\,(\sin\varphi\partial_x + \cos\varphi\partial_\phi) + \cos\phi\, \mathrm{d}\phi\big(\sin\varphi\partial_x + \cos\varphi\partial_\phi\big)^2 \\ &= -\cos\phi\sin^2\varphi + 2\sin\phi\cos\varphi\sin\varphi + \cos\phi\cos^2\varphi \\ &= \cos\phi(\cos^2\varphi - \sin^2\varphi) + \sin^2\phi = 1 \end{aligned}$$

Finally, the cross term $\langle\sigma_0, \sigma_1\rangle$;

$$\begin{aligned} g(\sigma_0, \sigma_1) &= -\cos\phi\, \mathrm{d}x\,(\cos\varphi\partial_x - \sin\varphi\partial_\phi) \otimes \mathrm{d}x\,(\sin\varphi\partial_x + \cos\varphi\partial_\phi) \\ &+ \sin\phi\, \mathrm{d}x\,(\cos\varphi\partial_x - \sin\varphi\partial_\phi) \otimes \mathrm{d}\phi\,(\sin\varphi\partial_x + \cos\varphi\partial_\phi) \\ &+ \sin\phi\, \mathrm{d}\phi\,(\cos\varphi\partial_x - \sin\varphi\partial_\phi) \otimes \mathrm{d}x\,(\sin\varphi\partial_x + \cos\varphi\partial_\phi) \\ &+ \cos\phi\, \mathrm{d}\phi\,(\cos\varphi\partial_x - \sin\varphi\partial_\phi) \otimes \mathrm{d}\phi\,(\sin\varphi\partial_x + \cos\varphi\partial_\phi) \\ &= -2\cos\phi\cos\varphi\sin\varphi + \sin\phi\,\big(\cos^2\varphi - \sin^2\varphi\big) = 0 \end{aligned}$$

So g is indeed a valid Lorentzian metric. Now suppose for contradiction that $(\mathbb{R}\times S^1, g)$ is time-orientable. Then there exists a nowehere vanishing global timelike section, $T \in \mathfrak{X}(M)$, such that $g(\sigma_0, T)|_p < 0$, where $x^\mu(p) = (0,0)$. Now define a function, $f(s) \coloneqq g(X_0, T)(\gamma(s))$, where $\gamma : [0,1) \to \mathbb{R}\times[0,2\pi)$ is a smooth curve. We have that $f : [0,1) \to \mathbb{R}$ is a smooth function since γ and g are both smooth. Furthermore, assume that the curve satisfies the initial condition $\gamma(0) = p$. Now we would like to show that $g(\sigma_0, T)|_\gamma < 0$ for all points on the curve. Suppose for contradiction that $\exists t \in [0,1) : f(t) < 0$. Then $g(X_0, T)(\gamma(t)) = 0$. But we also know that $g(\sigma_0, \sigma_0) = -1$, so σ_0 is timelike. If $g(\sigma_0, T)|_{\gamma(t)} = 0$ then T must be spacelike since it is orthognal to the timelike vector σ_0. Then this is a contradiction since T was assumed to be timelike everywhere. Furthermore, we have $f(0) < 0$, so by continuity $f(s) < 0\,\forall s \in [0,1)$. Now consider a curve such that $\gamma(s) = (0, 2\pi s)$. We have that $\lim_{s\to 1} g(\sigma_0, T)(\gamma(s)) = -g(\sigma_0, T)(\gamma(0))$ since $\lim_{s\to 1}\sigma_0(0, 2\pi s) = -\partial_x = -\sigma_0(0,0) = \partial_x$. Recall that $\mathbb{R}\times S^1$ is defined by identifying $\phi \sim \phi + 2\pi$, in particular $0 \sim 2\pi$. Now $f(0) = -\lim_{s\to 1} f(s) \sim -f(0)$ in the quotient, hence $f = 0$, a contradiction—[IRON THIS OUT]—. $\square$

There is a nice visualisation of the above exmaple. The frame fields precisely define a smooth rotation as ϕ varies along the horizontal of the strip $\mathbb{R}\times[0,2\pi)$. The half-angle results in the roation approaching a flip along the horizontal as $\phi \to 2\pi$. When we pass to the quotient, the two vertical edges of the strip are identified. In this case a light cone is smoothly transported around the cylinder while rotating, until it is upside down when it comes back to its starting point. Hence, there is no way to 'pick' either the top or bottom half-cone unambiguously.

Time orientability is defined as a global property, since every Lorentzian admits local timelike sections. There is already a topological obstruction to the existence of global nowhere vanishing

sections in the **Euler class** of the tangent bundle. —[REF]—. But, even if there exists an orientable Lorentzian manifold with a nowehere vanishing section, there is a further geometric obstruction from the metric itself. So time-orientability is both a topological and geometric characteristic of the space.

Proposition 4.1.2. *A Lorentzian manifold is time-orientable iff. . . .*

—[FINISH, A LOT MORE TO SAY ABOUT THIS]—

4.2 Causality

Recall that curves in spacetime can be characterised into three groups.

Definition 4.2.1. *A curve $\gamma : \mathbb{R} \longrightarrow M$ is;*

1. ***timelike*** *if* $g(\dot{\gamma}, \dot{\gamma}) < 0$,
2. ***null*** *if* $g(\dot{\gamma}, \dot{\gamma}) = 0$,
3. ***spacelike*** *if* $g(\dot{\gamma}, \dot{\gamma}) > 0$,

for all points in the image of γ.

We say that a curve is causal if it is either timelike or null. Intuitively, matter particles must travel along timelike curves and photons along null curves. There is a natural relation on events in spacetime based on whether they lie in each others futures or pasts.

Definition 4.2.2 (Causal relations)**.**

1. $p \ll q \iff \exists$ *a future directed timelike curve from p to q*
2. $p < q \iff \exists$ *a non-trivial future directed causal curve from p to q*

We then say that q is in the timelike/causal future of p. The notation $p \leq q$ if q is used if q is either in the causal future of p or $p = q$. We can define the regions of causality as follows.

Definition 4.2.3 (Causal regions)**.**

1. *The* ***timelike future*** *of a point $p \in M$ is $I^+(p) := \{q \in M : p \ll q\}$*
2. *The* ***causal future*** *of a point $p \in M$ is $J^+(p) := \{q \in M : p \leq q\}$*

We defined the case for the future regions, it can be easily extended to past regions by reversing time. For example $J^-(p)$ would be set of all q such that $q \leq p$. For the rest of the discussion we may construct definitions for only future $(+)$ events, but let it be implicit that there are also time dual past $(-)$ definitions. Note that $I^+(p)$ is open $\forall p \in M$, but $J^+(p)$ may or may not be closed. —[EXAMPLE + PICTURE]—.

Proposition 4.2.1. *Suppose $q \in J^+(p) - I^+(p)$, then any future directed causal curve from p to q must be a null geodesic.*

This is just the statement that no particle can move faster than the speed of light locally. From this proposition we have the following two statements.

Remark 4.2.1. *The causal relations are transitive, and we have the following:,*

$$(p \ll q \ \wedge \ q < r) \ \vee \ (p < q \ \wedge \ q \ll r) \implies p \ll r.$$

We can generalise the causal/timelike futures of points to arbitrary subspaces in spacetime.

Definition 4.2.4. *Consider a subset $S \subseteq M$.*

1. *The **timelike future** of S is*

$$I^+(S) \coloneqq \bigcup_{p \in S} I^+(p) = \{q \in M : p \ll q \quad \text{for some} \quad s \in S\}.$$

2. *The **causal future** of S is*

$$J^+(S) \coloneqq \bigcup_{p \in M} J^+(p) = \{q \in M : p \leq q \quad \text{for some} \quad s \in S\}.$$

It will also be useful to consider causual relations only on open subsets in the following sense.

Definition 4.2.5. *The timelike future of p **within** $U \subseteq M$ is the set of $q \in U$ such that there is a timelike curve **within** U such that $p \ll q$.*

Annoyingly, $I^+(p, U) \neq I^+(p) \cap U$ in general. Now in classical theories of physics we would like to have a sense of determinism. Traditionally we'd have initial conditions on a 'subset of spacetime' with a differential equation. We expect to be able to determine what happens, and has happened, to the future and past of this subset by solving the differential equation. To implement this in GR, where time becomes more complicated, we must restrict our definition of subset.

Definition 4.2.6. *A submanifold of spacetime, $\iota : \Sigma \hookrightarrow M$, is a **spacelike hypersurface** if $\iota^\star g$ has Riemannian signature.*

Consider Minkowski space with standard coordinates and metric,

$$g = -\mathrm{d}t \otimes \mathrm{d}t + \mathrm{d}x \otimes \mathrm{d}x.$$

Define a family of submanifolds by,

$$\Sigma_\lambda \coloneqq \{(\lambda, x) : \lambda \in \mathbb{R}\},$$

which are just horizontal lines of constant time. Clearly each Σ_λ is a spacelike since the induced metric is just $\iota^\star g = \mathrm{d}x \otimes \mathrm{d}x$. This spacelike condition is still not enough implement determinism, we need something more.

Definition 4.2.7. *A subset Σ of spacetime is **achronal** if there exists no timelike curve connecting two arbitrary points $p, q \in \Sigma$.*

To see why this definition is needed, we come back to the trusty cylinder $\mathbb{R} \times S^1$ for a non-example. A family of one-dimensional submanifolds, Σ_λ, can be defined by,

$$t = \lambda\phi,$$

where (t, ϕ) are the standard coordinates on the cylinder and $\lambda \in \mathbb{R}$ is a constant. Pictorialy this is a line wrapping around the cylinder infinitley many times. Now equip the space with the standard flat metric,

$$g = -\mathrm{d}t \otimes \mathrm{d}t + \mathrm{d}\phi \otimes \mathrm{d}\phi.$$

For small λ, Σ_λ is a spacelike surface, approaching the circle for $\lambda \to 0$. However, Σ_λ is never achronal for $\lambda > 0$. This is clear when we consider the curve paramaterised by,

$$\gamma(\tau) = (\tau, 0),$$

for $\tau \in \mathbb{R}$. It's clear that the curve connects the points $(0, 0)$ and $(2\pi\lambda, 0)$ with a straight vertical line, which is evidently a timelike geodesic. In fact, the points $\{(n2\pi\lambda, 0) : n \in \mathbb{Z}\}$ are all connected by this geodesic. This poses an issue for specifying intitial conditions over Σ_λ since data at $(0, 0)$ can influence data $(2\pi\lambda, 0)$ in the future for example. Thus, it doesn't make sense to define initial conditions over spacelike hypersufaces if they are not also achronal.

Definition 4.2.8. *An **achronal boundary** is a subset of the form $\partial I^{\pm}(S)$ for some achronal subspace S.*

Proposition 4.2.2. *If an achronal boundary is non-empty, $\partial I^{\pm}(S) \neq \varnothing$, then it is a closed achronal C^0 hypersurface in M.*

Note that by C^0 hypersurface we mean a topological submanifold of codimension one. Achronal boundaries need not be smooth in general.

Definition 4.2.9. *Suppose $S \subseteq M$ is achronal. $p \in \bar{S}$ is an **edge point** of S if every neighbourhood $U \ni p$ contains a timelike curve, γ, from $I^-(p, U)$ to $I^+(p, U)$ that does not meet S, $\gamma \cap S = \varnothing$. Denote the set of edge points* $\mathrm{edge}(S)$. *We say that S is **edgeless** if* $\mathrm{edge}(S) = \varnothing$.

—-[GIVE TWO EXAMPLES + PICTURES]—-.

Proposition 4.2.3. *An edgeless achronal set, $S : \mathrm{edge}(S) = \varnothing$, is a C^0 hypersurface.*

The proof of the above can be found in [O'N83].
—[HYPERBOLIC STUFF]—

4.3 Inner Product Structure on the Tangent Space

Now let's recall some facts about inner products with indefinite signature. For this section we fix a given metric on the tangent space, (T_pM, g).

Definition 4.3.1. *The orthognal complement of a vector subspace, $V \subseteq T_pM$, is defined by,*

$$V^\perp := \{X \in T_pM : g(X, Y) = 0, \forall Y \in V\}.$$

Proposition 4.3.1. *Let $V \subseteq T_pM$ be a subspace. The following statements are equivalent:*

$$g\big|_V \text{ is non-degenerate} \iff V \cap V^\perp = \{0\} \iff T_pM = V \oplus V^\perp.$$

There are two inequalities which always hold for Lorentzian metrics.

Proposition 4.3.2.

1. *The* ***reversed Schwarz inequality****,*

$$|g(X,Y)| \geq |X||Y|, \tag{4.1}$$

holds for all causal vectors X and Y.

2. *The* ***reversed triangle inequality****,*

$$|X+Y| \geq |X| + |Y|, \tag{4.2}$$

holds for all similary directed causal vectors X, Y.

Proof. First we will show the reversed Schwarz inequality, which then implies the reverse triangle inequality. The inequality is trivially satisfied for null vectors since all terms are zero. Assuming X is timelike, we consider the subspace,

$$\mathrm{span}_{\mathbb{R}} X = \{\lambda X : \lambda \in \mathbb{R}\} \coloneqq \mathbb{R} \cdot X,$$

and then apply the decomposition from proposition 4.3.1,

$$T_p M = \mathbb{R} \cdot X \oplus (\mathbb{R} \cdot X)^{\perp},$$

to write $Y = \lambda X + Y^{\perp}$. Here $Y^{\perp} \in (\mathbb{R} \cdot X)^{\perp}$ is spacelike since X is timelike. Now we have,

$$g(X,Y) = g(X, \lambda X + Y^{\perp}) = \lambda g(X,X) + g(X, Y^{\perp}) = \lambda g(X,X), \tag{4.3}$$

where $g(X, Y^{\perp}) = 0$ by definition. Also note that,

$$g(Y,Y) = \lambda^2 g(X,X) + 2g(X,Y^{\perp}) + g(Y^{\perp}, Y^{\perp}) = \lambda^2 g(X,X) + g(Y^{\perp}, Y^{\perp}), \tag{4.4}$$

again using the definition of $Y^{\perp}$. Now square both sides of eq. (4.3) to get,

$$g(X,Y)^2 = g(X,X)\lambda^2 g(X,X) = g(X,X)\left(g(Y,Y) - g(Y^{\perp}, Y^{\perp})\right), \tag{4.5}$$

where we substituted eq. (4.4) in the second step. Finally remember that $g(X,X) < 0$, $g(Y,Y) \leq 0$, and $g(Y^{\perp}, Y^{\perp}) > 0$. Then the inequality follows,

$$\begin{aligned} & g(X,Y)^2 \geq g(X,X)g(Y,Y) \\ \implies & |g(X,Y)| \geq |X||Y|, \end{aligned}$$

by realising absolute values and square roots can be applied to both positive sides of the inequality. For the reveresed triangle inequality we recall that if X and Y are similarly directed causal then $g(X,Y) \leq 0$. Then keeping track of signs,

$$\begin{aligned} & |X+Y|^2 = -g(X+Y, X+Y) = -g(X,X) - 2g(X,Y) + g(Y,Y) \\ & |X+Y|^2 = |X|^2 + 2|g(X,Y)| + |Y|^2, \end{aligned}$$

and then finally use the reveresed Schwarz inequality to get,

$$\begin{aligned} & |X+Y|^2 \geq |X|^2 + 2|X||Y| + |Y|^2 = (|X| + |Y|)^2 \\ \implies & |X+Y|^2 \geq |X| + |Y|. \end{aligned}$$

□

These counter-intuitive relatations are the cause of the famous 'twin paradox' from special relativity.

4.4 Causality & Energy Conditions

Definition 4.4.1. *The **chronology condition** states there are no closed **timelike** curves in a spacetime M.*

Definition 4.4.2. *The **strong causality condition** states that there a non-trivial closed **causal** curves in a spacetime M.*

Proposition 4.4.1. *Every compact spacetime contains closed timelike curves.*

Proof. —[COMPLETE]— □

Definition 4.4.3. *The **null energy condition** states that $T(X, X) \geq 0$ for all null vectors X.*

Definition 4.4.4. *The **dominant energy condition** states that $T(X, Y) \geq 0$ for all future directed vectors X, Y.*

There exist some classical fields which violate the null energy condition. —[REF]—. The dominant energy condition requires the 'energy flow' velocity to be less than the speed of light. This is more restrictive than the null energy condition, and is satisfied by most classical fields.

4.5 Null Geometry

Definition 4.5.1. *A **null hypersurace** is a smooth, connected hypersurface $\iota : S \hookrightarrow M$ such that $\iota^\star g$ is degenerate.*

In particular, $\forall p \in S, \exists 0 \neq K_p \in T_pS : (\iota^\star g)(K_p, X) = 0, \forall X \in T_pS$. It follows immediately that K_p, called the direction of degeneracy, is null $\langle K_p, K_p \rangle = 0$. Furthermore, we have $(\mathbb{R} \cdot K_p)^\perp = T_pS$. Without loss of generality we can take K_p to be future pointing with respect to the time orientation on M. Any non-zero vector that isn't a multiple of K_p is also necessarily spacelike.

$$0 \neq X_p \in T_pS : X_p \notin \mathbb{R} \cdot K_p \implies (\iota^\star g)(X_p, X_p) > 0$$

Thus these vectors can be stitched together to form global section, $K \in \mathfrak{X}(S)$, on the hypersurface. Note that this section is indeed a smooth vector field when suitably the K_p are suitably normalised. K is also unique up to pointwise multiplication by a postive function. In Minkowski space the future and past achronal boundaries, $\partial I^\pm(p)$, of a point $p \in \mathbb{R}^{3,1}$. These form the furture and past light cones at p, and neither are smooth. Away from the vertex however, $\partial I^\pm(p) - \{p\}$, are both null hypersurfaces. The notation $\langle -, - \rangle \coloneqq \iota^\star g(-, -)$ may be used when the contect is clear.

Proposition 4.5.1. *Suppose $\iota : S \hookrightarrow M$ is a null hypersurface. Then the integral curves of a null vector field, $K \in \mathfrak{X}(S)$, are null geodesics under a suitable choice of parameterisation.*

Proof. We would like to show that $\nabla_K K = \lambda K$ for some $\lambda \in \mathbb{R}$. Then the integral curves of K are pregeodesics, and hence geodesics once reparemeterised appropriately. Consider an arbitrary vector $X_p \in T_pS$. Then this can be extended as an invariant vector field, $X \in \mathfrak{X}(\gamma)$, along an integral curve γ of K which passes through the point $p \in S$.

$$\mathcal{L}_K X = \nabla_K X - \nabla_X K = 0$$

By definition of the null vector field, $\langle K, X\rangle_\gamma = 0$, since X is still tangent to S. Now differentiate,

$$\begin{aligned} K\langle K, X\rangle &= \langle \nabla_K K, X\rangle + \langle K, \nabla_K X\rangle = 0 \\ \implies \langle \nabla_K K, X\rangle &= -\langle K, \nabla_X K\rangle, \end{aligned}$$

where metric compatiblilty was used to introduce the covariant derivative. Not that on the other hand,

$$X\langle K, K\rangle = \langle \nabla_X K, K\rangle + \langle K, \nabla_X K\rangle = 2\langle K, \nabla_X K\rangle,$$

by symmetry. Then, using the defintion of the invariant extenstion of X, we obtain

$$\langle \nabla_K K, X\rangle = -\frac{1}{2} X\langle K, K\rangle = 0.$$

So we have shown that $\mathbb{R} \cdot \nabla_K K \perp T_p S$. In particular, by the uniqueness of the null vector field up to pointwise scaling,

$$\langle \nabla_K K, X\rangle\Big|_p = 0 \quad \forall X \implies \nabla_K K = \lambda K.$$

□

In Riemannian geometry it often useful to study submanifolds through the normal vector field associated to it. Similarly, we can study the geometry of null hypersrfaces through the null vector field. It's conveniant to work in a formalism where the direction of degeneracy is quotiented out.

Definition 4.5.2.

1. *Define an equivalence relation on the tangent space, $T_p S$, by*

$$X, X' \in T_p S, \quad X \sim_K X' \iff X - X' = \lambda K, \quad \lambda \in \mathbb{R}.$$

2. *Denote the equivalence class by $\bar{X} \coloneqq [X]_{\sim_K} = \{Y \in T_p S : X \sim_K Y, X \in T_p S\}$. The **mod K tangent space** at a point $p \in S$ is the quotient vector space*

$$T_p S/K \coloneqq \{\bar{X} : X \in T_p S\}.$$

3. *The **mod K tangent bundle** is the disjoint union of the mod K tangent spaces over S.*

$$TS/K \coloneqq \bigsqcup_{p \in S} T_p S/K$$

Recall we write $d = \dim M$, and the hypersurface is codimension 1, $\dim S = d - 1 \coloneqq n$. Then the mod K tangent bundle is a smooth vector bundle of rank $n - 1$. Note that, despite the name, the mod K tangent bundle does not depend on the choice of null vector field.

Proposition 4.5.2. *The mod K tangent bundle is well-defined for an arbitray choice of null vector field. In particular, for two null vector fields $K, K' \in \mathfrak{X}(S)$:*

$$TS/K \cong TS/K'$$

Proof. It suffices to show that the quotient tangent spaces are equal, $T_pS/K = T_pS/K'$. Suppose $X, X' \in T_pS/K$ then,

$$X - X' = \alpha K,\ \alpha \in \mathbb{R},$$

by the previous definition. Recall on the other hand that null vector fields on a null hypersuface are unique up to pointwise scaling by a non-zero number. So if $K \in T_pS$ is a null vector, $\exists \beta \in \mathbb{R} : K = \beta K'$. Then it's immediate that

$$X - X' = \alpha K \iff X - X' = \alpha\beta K'.$$

So $X \sim_K X' \iff X \sim_{K'} X'$, and $T_pS/K = T_pS/K'$. □

There is a natural metric induced by the quotient map.

Definition 4.5.3. *The **null induced metric**, $h : T_pS/K \otimes T_pS/K \longrightarrow \mathbb{R}$, is defined by*

$$h(\bar{X}, \bar{Y}) := \langle X, Y \rangle.$$

Proposition 4.5.3. *The induced metric well-defined mod K.*

Proof. Consider two vectors, $X, Y \in T_pS$, so that $X' = X + \alpha K$ and $Y' = Y + \beta K$ are two different representatives in the quotient. Then,

$$\begin{aligned} h(X', Y') =& \langle X + \alpha K, Y + \beta K \rangle \\ =& \langle X, Y \rangle + \langle X, \beta K \rangle + \langle \alpha K, Y \rangle + \langle \alpha K, \beta K \rangle \\ =& \langle X, Y \rangle. \end{aligned}$$

□

Definition 4.5.4. *The **null Weingarten map**, $b_K : T_pS/K \longrightarrow T_pS/K$, is defined by*

$$b_K(\overline{X}) := \overline{\nabla_X K}$$

Proposition 4.5.4.

1. *The null Weingerten map is well-defined.*
2. *$b_K = b$ is unique up to a smooth positive function.*

Proof.

1. Consider two equivalent representatives, $X, X' \in T_pS : X' = X + \alpha K$.

$$\begin{aligned} b_K(\overline{X'}) &= \overline{\nabla_{X'} K} = \overline{\nabla_{X+\alpha K} K} \\ &= \overline{\nabla_X K} + \alpha \overline{\nabla_K K} \\ &= \overline{\nabla_X K} = b_K(X) \end{aligned}$$

   In the second line we have used the earlier result that the integral curves of K are null (pre)geodesics, $\nabla_K K = \lambda K$, $\lambda \in \mathbb{R}$.

2. If $\tilde{K} \in \mathfrak{X}(S)$ is another null vector field then $\exists f \in C^\infty(S)$, a smooth positive function, such that $\tilde{K} = fK$. Then $\forall X \in T_pS$,

$$\begin{aligned} b_{\tilde{K}}(\overline{X}) &= \overline{\nabla_X(fK)} \\ &= \overline{f\nabla_X K} + \overline{X(f)\cdot K} \\ &= f b_K(\overline{X}). \end{aligned}$$

□

From now on the null vector field associated to the null Weignarten map will be omitted in the notation.

Definition 4.5.5. *The **null second fundamental form**, $B : T_pS/K \otimes T_pS/K \longrightarrow \mathbb{R}$, associated to a null vector field K on S is defined by*

$$B(\overline{X}, \overline{Y}) \coloneqq h(b(\overline{X}), \overline{Y}).$$

Proposition 4.5.5. *The null second fundamental form is symmetric. Equivalently the null Weingarten map, b, is self-adjoint with respect to the induced metric h.*

Proof. Consider two vectors $X, Y \in T_pS$ and extend them to vector fields in a neighbourhood of p. Then,

$$\begin{aligned} \langle K, Y\rangle = 0 \quad &\& \quad \langle K, X\rangle = 0 \\ \implies X\langle K, Y\rangle = 0 \quad &\& \quad Y\langle K, X\rangle = 0 \\ \iff \langle \nabla_X K, Y\rangle = -\langle K, \nabla_X Y\rangle \quad &\& \quad \langle \nabla_Y K, X\rangle = -\langle K, \nabla_Y X\rangle. \end{aligned}$$

Recall that the Levi-Civita connection is torsion free,

$$\nabla_X Y - \nabla_Y X = [X, Y].$$

Now apply the definition of the fundamental form.

$$\begin{aligned} B(\overline{X}, \overline{Y}) = h(b(\overline{X}), \overline{Y}) &= \langle \nabla_X K, Y\rangle = -\langle K, \nabla_X Y\rangle \\ &= -\langle K, \nabla_Y X\rangle - \langle K, [X, Y]\rangle \\ &= \langle \nabla_Y K, X\rangle = \langle X, \nabla_Y K\rangle \\ &= h(\overline{X}, b(\overline{Y})) = B(\overline{Y}, \overline{X}) \end{aligned}$$

□

Definition 4.5.6. *A null hypersuface is **totally geodesic** if $B \equiv 0$.*

If S is totally geodesic, then any geodesic starting tangent to S stays in S. Null hyperplanes in Minkowski spacetime are totally geodesic. Another example is in Schwarzschild spacetime. The event horizon is a null hypersurface, and light rays tangent to it must stay tangent. Studying the deviation of light rays is a useful tool to analyse the geometry of spacetimes. One can define a function which formalises this notion.

Definition 4.5.7. *The **null mean curvature** or **null expansion scalar** is the function,*

$$\vartheta := \operatorname{Tr} b.$$

Consider a spacelike hypersurface in M which is transverse to K at $p \in S$. Denote the intersection of the hypersuface with S by Σ. This is a codimension 2 spacelike submanifold such that K is orthognal on it. —[DRAW PICTURE]—. Now choose an orthonormal basis $\{\sigma_\alpha\}_{\alpha=1}^{n-1}$ of $T_p\Sigma$. Passing to the quotient we have an orthonormal basis, $\{\overline{\sigma}_{\alpha=1}^{n-1}\}$, for T_pS/K. Then the null mean curvature is,

$$\vartheta = \sum_{\alpha=1}^{n-1} h(b(\overline{\sigma_\alpha}), \overline{\sigma_\alpha}) = \sum_{\alpha=1}^{n-1} \langle \nabla_{\sigma_\alpha} K, \sigma_\alpha \rangle, \tag{4.6}$$

which is recognisable as the **divergence** of the null vector field along Σ. If ω is the volume form on Σ then we can write,

$$\mathcal{L}_K \omega = \vartheta \omega, \tag{4.7}$$

in the direction of the flow generated by K. At a point $p \in \mathbb{R}^{3,1}$ in Minkowski spacetime, consider the future null cone in Minkowski spacetime away from the vertex $\partial I^+ - \{p\}$. The null expansion on this hypersurface is greater than zero, $\vartheta > 0$, matching the fact that the null generators are diverging in time. Likewise, for the past null cone, $\partial I^- - \{p\}$, we have $\vartheta < 0$ since the null generators are converging to p.

Recall from Proposition 4.5.4 that $b_{\tilde{K}} = f b_K$ for some other null vector field, $\tilde{K}$, related by a positive smooth function $f \in C^\infty(S)$ such that $\tilde{K} = fK$. It follows immediately, from the linearity of the trace operation, that $\tilde{\vartheta} = f\vartheta$. In particular the sign of the null expansion is preserved. This is useful since inequalities such as $\vartheta > 0$ do not depend on the choice of null vector field. Now let's define these constructions over curves. Consider an affinely parametrised null geodesic generator, $s \mapsto \gamma(s)$, on S. A vector field, $s \mapsto \mathcal{Y}(s) \in T_{\gamma(s)}S/K$, is smooth if $\exists$ a smooth vector field, $s \mapsto Y \in T_{\gamma(s)}S$, such that $\mathcal{Y}(s) = \overline{Y(s)}$. Now we can differentiate the vector field on the mod K bundle by defining,

$$\mathcal{Y}'(s) := \overline{\nabla_{\dot{\gamma}} Y(s)}, \tag{4.8}$$

where $\nabla_{\dot{\gamma}} = \gamma^\star \nabla$ is the induced covariant derivative.

The null Weignarten map can now be defined along the curve in the usual way, $b(s) := b_{\dot{\gamma}(s)} : T_{\gamma(s)}S/\dot{\gamma}(s) \longrightarrow T_{\gamma(s)}S/\dot{\gamma}(s)$. THe derivative of the null Weingarten map is defined by,

$$b'(\overline{X}) := \nabla_{\dot{\gamma}} b(\overline{X}) - b(\overline{\nabla_{\dot{\gamma}} X}). \tag{4.9}$$

Proposition 4.5.6. *The null Weingarten map satisfies the Ricatti equation,*

$$b' + b \circ b + R = 0, \tag{4.10}$$

where $R : T_{\gamma(s)}S/\dot{\gamma}(s) \longrightarrow T_{\gamma(s)}S/\dot{\gamma}(s)$ is the Riemann curvature endomorphism defined by,

$$R(\overline{X}) := \overline{R(X, \dot{\gamma}(s))\dot{\gamma}(s)}. \tag{4.11}$$

Proof. For $s \in (a, b)$, let $p = \gamma(s_0)$. The null vector field, K, on S can be scaled in a neighbourhood $U \ni p$ so that K is an affinely parameterised geodesic and $K|_\gamma$ is the velocity vector field to γ. Explicitly, $K_{\gamma(s)} = \dot{\gamma}(s)$. Now for $X \in T_pS$ we can pick another neighbourhood $V \ni p$ in which X is extended to smooth vector field such that

$$[X, K] = \nabla_X K - \nabla_K X = 0.$$

Now the Riemann curvature simplifies,

$$\begin{aligned} R(X, K)K &= \nabla_X \nabla_K K - \nabla_K \nabla_X K - \nabla_{[X,K]} K \\ &= -\nabla_K \nabla_X K = -\nabla_K \nabla_K X. \end{aligned}$$

In the first line $\nabla_K K = 0$ by the null geodesic property. The Lie bracket $[X, Y]$ vanishes to eliminate the third term in the first line, and leads to the final line. So along the curve,

$$\nabla_{\dot{\gamma}} \nabla_{\dot{\gamma}} X = -R(X, \dot{\gamma})\dot{\gamma},$$

so that X is a **Jacobi** field along γ. Turning to the derivative of the Weingarten map,

$$\begin{aligned} b'(\overline{X}) &= \nabla_{\dot{\gamma}} \overline{\nabla_X K} - b(\overline{\nabla_K X}) \\ &= \nabla_{\dot{\gamma}} \overline{\nabla_K X} - b \circ b(\overline{X}) \\ &= \overline{\nabla_{\dot{\gamma}} \nabla_{\dot{\gamma}} X} - b \circ b(\overline{X}) \\ &= -\overline{R(X, \dot{\gamma})\dot{\gamma}} - b \circ b(\overline{X}). \end{aligned}$$

□

Taking the trace of the Riccati equation leads to,

$$\dot{\vartheta} = -Rc(\dot{\gamma}, \dot{\gamma}) - \sigma^2 - \frac{1}{n-1}\vartheta^2, \tag{4.12}$$

where $\sigma := \sqrt{\operatorname{Tr} \hat{b}^2}$ with $\hat{b} = b - \frac{\vartheta}{n-1}\,\mathrm{id}$. The above is a form of the **Raychaudhuri equation** for null irrotational geodesic congruence. σ is sometimes called the **shear scalar** and $\hat{b}$ is the traceless part of the Weingarten map.

Proposition 4.5.7. *Suppose a spacetime, M, obeys the null energy condition $Rc(X, X) \geq 0$. Let $S \hookrightarrow M$ be a smooth null hypersurface such that the generators of S are future geodesically complete. Then S has non-negative null expansion $\vartheta \geq 0$.*

Proof. Suppose for contradiction $\exists p \in S : \vartheta < 0$. Pick an affinely paramaterised null geodesic generator, $s \mapsto \gamma(s)$, passing through $p = \gamma(0)$. Then $\vartheta(0) < 0$. The null energy condition applied to the Raychaudhuri equation gives an inequality,

$$\frac{\mathrm{d}\vartheta}{\mathrm{d}s} \leq -\frac{1}{n-1}\vartheta^2. \tag{4.13}$$

But $\vartheta^2 > 0 \implies \dot{\vartheta} < 0, \quad \forall s > 0$. Rearranging,

$$-\frac{1}{\vartheta^2}\frac{\mathrm{d}\vartheta}{\mathrm{d}s} \geq \frac{1}{n-1}, \tag{4.14}$$

and noticing that,

$$\frac{\mathrm{d}}{\mathrm{d}s}\left(\frac{1}{\vartheta}\right) = -\frac{1}{\vartheta^2}\frac{\mathrm{d}\vartheta}{\mathrm{d}s},$$

results in

$$\frac{\mathrm{d}}{\mathrm{d}s}\left(\frac{1}{\vartheta}\right) \geq \frac{1}{n-1}. \tag{4.15}$$

So $1/\vartheta \to 0 \iff \vartheta \to -\infty$ in finite affine time. This contridacticts smoothness. □

4.6 Penrose's Theorem

—[WITTEN LECTURES BLACK HOLE TRAPPED SURFACE MOTIVATING EXAMPLE]—

Consider a closed (compact without boundary), spacelike submanifold $\iota : \Sigma \hookrightarrow M$. $\forall p \in \Sigma$, $(T_p\Sigma)^\perp$ is timelike and 2-dimensional, and so admits two futre directed null directions to Σ. —[NEED SUITABLE ORIENTATION CONDITIONS, CHECK]—. Denote the null vectors associated to these two directions by ℓ_- and ℓ_+ for ingoing and outgoing light rays. Denote the associated second fundamental forms by,

$$\chi_\pm(X,Y) \coloneqq g(\nabla_X \ell_\pm, Y), \tag{4.16}$$

and the expansions scalars are the usual traces,

$$\vartheta_\pm \coloneqq \mathrm{Tr}_{\iota^\star g}\, \chi_\pm = (\iota^\star g)^{ab}(\chi_\pm)_{ab} = \mathrm{div}_\Sigma\, \ell_\pm. \tag{4.17}$$

Definition 4.6.1. *A **trapped surface** is a closed spacelike submanifold such that both $\vartheta_+ < 0$ and $\vartheta_- < 0$.*

Now the famous result by Penrose can be formulated [Pen65].

Theorem 4.6.1 (Penrose)**.** *Let M be a globally hyperbolic spacetime satisfying the null energy condition, $Rc(X,X) \geq 0$ for all null vectors X. Suppose M has a non-compact Cauchy surface S. If M contains a trapped surface, Σ, then M is future null geodesically incomplete.*

Proof.

- Claim: $\partial I^+(\Sigma)$ is non-compact.
  Since $\partial I^+(\Sigma)$ is an achronal boundary, it is an achronal C^0 hypersurface by Proposition 4.2.2. Suppose for contradiction that $\partial I^+(\Sigma)$ is in fact compact. Then by —[NEED PROP]—, $\partial I^+(\Sigma)$ would be a compact hypersurface. However, since M is globally hyperbolic, every Cauchy surface is homeomorphic $\partial I^+(\Sigma) \cong_{\mathrm{Top}} S$. This contradicts the assumption of a non-compact Cauchy hypersurface.

- Claim: There exists a future inextendible null geodesic in $\partial I^+(\Sigma)$.
  From point set topology we have,

$$\partial I^+(\Sigma) \cong \overline{I^+(\Sigma)} - \mathrm{int}\, I^+(\Sigma).$$

Moreover, since the spacetime is globally hyperbolic,

$$\partial I^+(\Sigma) \cong J^+(\Sigma) - I^+(\Sigma).$$

From Proposition 4.2.1 each $q \in \partial I^+(\Sigma)$ must lie on a null geodesic in $\partial I^+(\Sigma)$ with the initial endpoint on $p \in \Sigma$. Furthermore, one can show that this null geodesic must meet Σ orthogonally [O'N83]. Since $\partial I^+(\Sigma)$ is closed and non-compact, $\exists \{q_k\}_{k\in\mathbb{N}} \subset \partial I^+(\Sigma)$, a sequence of points which diverge to infinity. Now for all $k \in \mathbb{N}$ there exists a null geodesic γ_k from Σ to q_k which is contained in $\partial I^+(\Sigma)$ and meets Σ orthogonally. Compactness of Σ implies that every subsequence $\{p_k\}_{k\in\mathbb{N}} \subset \Sigma$ has a convergent subsequence $p_{k_j} \to p \in \Sigma$. So for the sequence of geodesice $\{\gamma_k\}_{k\in\mathbb{N}}$ there exists a subsequence $\gamma_{k_j} \to \gamma$ convergent to an inextendible null geodesic in $\partial I^+(\Sigma)$ meeting Σ orthogonally at p.

- Claim: γ is future incomplete.
  Suppose for contradiction that γ is in fact future complete. The achronality of $\partial I^+(\Sigma)$ implies that no other future directed null geodesic starting on Σ can meet γ. Then there are no null focal points to Σ along γ, and so there exists a null hypersurface $H \subset \partial I^+(\Sigma)$ containing γ [O'N83]. Let ϑ be the null expansion of H along γ. By assumption there exists a trapped surface, Σ, and so $\vartheta(p) < 0$. From similar arguments as in Section 4.5, $\vartheta \to -\infty$ in finite affine time. This is a contradiction, hence γ must be incomplete.

□

—[INCLUDE MORE DETAILS IN PROOF]— The assumption of a non-compact Cauchy surface is motivated by the usual spacetimes considered in physics which satisfy asymptotic boundary conditions. For example, a Schwarzschild black hole has the property of being asymptotically flat.

4.7 Instability of Kaluza-Klein Theories

Following from Penrose's theorem one can deduce that spacetime prefers to be topologically trivial when possible. The result is due to [Gan75] and [Lee76].

Theorem 4.7.1. *Let M be a globally hyperbolic spaetime satisfying the null energy condition and admitting an asymptotically flat spacelike Cauchy surface Σ. If $\pi_1(\Sigma) \neq 0$ then M is future null geodesically incomplete.*

—[DISCUSS NEED FOR MODULI STABILISATION IN STRING THEORY]—

Bibliography

[Cat15] Gabriel Catren. Geometric foundations of cartan gauge gravity. *International Journal of Geometric Methods in Modern Physics*, 12(04):1530002, April 2015.

[Frè13] Pietro Giuseppe Frè. *Gravity, a Geometrical Course: Vol. 2: Black Holes, Cosmology and Introduction to Supergravity*. Springer, Dordrecht, 2013.

[Gal14] G. J. Galloway. Notes on lorentzian causality. ESI–EMS–IAMP Summer School on Mathematical Relativity, Vienna, Austria, August 2014. Unpublished lecture notes.

[Gan75] D Gannon. Singularities in nonsimply connected spacetimes. *J. Math. Phys. (N.Y.), v. 16, no. 12, pp. 2364-2367*, 12 1975.

[Lee76] C. W. Lee. A restriction on the topology of cauchy surfaces in general relativity. *Communications in Mathematical Physics*, 51:157–162, 1976.

[New73] E. T. Newman. Complex coordinate transformations and the Schwarzschild-Kerr metrics. *J. Math. Phys.*, 14(6):774, 1973.

[NP13] Branislav D. Nikolić and Milan R. Pantić. A possible intuitive derivation of the kerr metric in orthogonal form based on ellipsoidal metric ansatz, 2013.

[O'N83] Barrett O'Neill. *Semi-Riemannian Geometry: With Applications to Relativity*, volume 103 of *Pure and Applied Mathematics*. Academic Press, New York, 1983.

[Pen65] Roger Penrose. Gravitational collapse and space-time singularities. *Phys. Rev. Lett.*, 14:57–59, Jan 1965.

[Sha02] Richard Sharpe. An introduction to cartan geometries. In *Proceedings of the 21st Winter School "Geometry and Physics"*, pages [61]–75. Circolo Matematico di Palermo, 2002.

[Str13] Norbert Straumann. *General Relativity*. Springer, Dordrecht, 2013.

[Wis10] Derek K Wise. Macdowell–mansouri gravity and cartan geometry. *Classical and Quantum Gravity*, 27(15):155010, June 2010.

[Wit20] Edward Witten. Light rays, singularities, and all that. *Rev. Mod. Phys.*, 92:045004, Nov 2020.