

1 The everyday speech environments of preschoolers with
2 and without cochlear implants

3 Margaret CYCHOSZ¹, Jan R. EDWARDS², Benjamin MUNSON³, Rachel R. ROMEO^{2,4}, Jessica
4 E. KOSIE⁵, and Rochelle S. NEWMAN²

5 ¹Department of Linguistics, University of California, Los Angeles

6 ²Department of Hearing and Speech Sciences, University of Maryland-College Park

7 ³Department of Speech-Language-Hearing Sciences, University of Minnesota

8 ⁴Department of Human Development and Quantitative Methodology, University of Maryland-College Park

9 ⁵Department of Psychology, Princeton University

10 Author Note

11 Author to whom correspondence should be addressed: Margaret Cychosz, 3125 Campbell Hall,
12 University of California, Los Angeles, Los Angeles, CA, 90095. Email: mcychosz4@ucla.edu. Competing
13 interests: The authors declare none.

Abstract

Children who receive cochlear implants develop spoken language on a protracted timescale. The home environment facilitates speech-language development, yet it is relatively unknown how the environment differs between children with cochlear implants and typical hearing. We matched eighteen preschoolers with implants (31-65 months) to two groups of children with typical hearing: by chronological age and hearing age. Each child completed a long-form, naturalistic audio recording of their home environment (appx. 16 hours/child; >730 hours of observation) to measure adult speech input, child vocal productivity, and caregiver-child interaction. Results showed that children with cochlear implants and typical hearing were exposed to and engaged in similar amounts of spoken language with caregivers. However, the home environment did not reflect developmental stages as closely for children with implants, or predict their speech outcomes as strongly. Home-based speech-language interventions should focus on the unique input-outcome relationships for this group of children with hearing loss.

Keywords: cochlear implant, deafness, language input, spoken language, language interaction

Introduction

Children learn the sounds and structure of their native language(s) from the input that they receive from caregivers around them. Yet children vary in the type and quantity of their speech-language exposure, including the number and diversity of word types (Rowe, 2012), ratio of male to female speakers (Bergelson et al., 2019), amount of child-directed speech (Rowe, 2008; Rowe & Weisleder, 2020; Schwab & Lew-Williams, 2016), and acoustics of caregiver speech (Dilley, Millett, Mcauley, & Bergeson, 2014; Kalashnikova & Burnham, 2018). Crucially, these individual differences in linguistic exposure may account for some differences in children’s speech-language development. For example, 6 to 14-month-olds who engage in more contingent interactions with caregivers—typically defined as conversational exchanges that take place in quick succession (≤ 2 seconds)—produce more babbling sounds and grow larger expressive and receptive vocabularies (Ferjan Ramírez, Lytle, & Kuhl, 2020). Similarly, infants who hear more child-directed speech at seven (Newman, Rowe, & Bernstein Ratner, 2016) or nineteen months (Weisleder & Fernald, 2013) learn to process words more efficiently, helping them grow larger vocabularies by age two.

This causal link between individual differences in linguistic input and speech-language development, which persists independent of family socioeconomic status or caregiver characteristics (Romeo et al., 2018), likewise extends to non-typically developing groups such as children who receive autism spectrum disorder diagnoses (Swanson et al., 2019) and, most critically for the current work, children with hearing loss, including those who receive cochlear implants (CIs) (Ambrose, VanDam, & Moeller, 2014; Ambrose, Walker, Unflat-Berry, Oleson, & Moeller, 2015; Arjmandi, Houston, Wang, & Dilley, 2021; DesJardin & Eisenberg, 2007; Dilley et al., 2020; Wang, Jung, Bergeson, & Houston, 2020) (see Nittrouer, Lowenstein, and Antonelli (2019) for review). For example, Dilley et al. (2020) correlated acoustic-lexical properties of speech directed to a cohort of

recently-implanted infants (8-29 months at implantation) and found that more diverse word types, and more dispersed vowel spaces in infant-directed compared to adult-directed speech, predicted higher scores on standardized language assessments two years post-implantation (see also Wang, Jung, et al. 2020). Elsewhere, caregivers of children with CIs who had greater mean lengths of utterance, and used more open-ended and/or recast questions, likewise had children with higher standardized receptive and expressive language scores (DesJardin & Eisenberg, 2007). Although standardized assessments never capture the full complexity of a child's linguistic development, these results do show that individual differences in linguistic input to children with CIs can predict some of the well-acknowledged variation in implantees' developmental outcomes (Pisoni et al., 2008). As such, having a clear characterization of these children's everyday speech-language environments is fundamental to understanding their speech and language development.

How cochlear implantation might shape the early language environment

There are two components of cochlear implantation that may systematically alter child implantees' speech-language environments (Houston, 2022). First, prelingually-deafened children often do not receive one or both of their CIs until their second or third birthdays (Fitzpatrick, Ham, & Whittingham, 2015), resulting in an extended period of AUDITORY ABSENCE pre-implantation where (1) the children are not exposed to spoken language models/caregivers may direct less spoken language to the child. Second, once CIs are activated, recipients have the added challenge of compensating for SIGNAL DEGRADATION: CI electrode arrays stimulate the cochlea at discrete points, discretizing the speech envelope. There are additional issues inherent to the hardware such as electrode interaction and interaural mismatch, as well as physiological aspects such as irregular neuronal survival, that together result in a highly degraded auditory signal from which pre-lingual implantees must learn speech and language. CIs also only introduce the sensation of hearing when the devices are being worn, and children vary greatly in the number of hours

of typical device use (Ganek, Cushing, Papsin, & Gordon, 2020; Majorano et al., 2021).

How auditory absence shapes the early environment. Pediatric CI candidates must pass a series of candidacy requirements (e.g., >70dB pure-tone thresholds in both ears, limited benefit from acoustic amplification, typical anatomical cochlear development). Although the Food and Drug Administration permits implantation in children as young as 9 months of age (USFDA, 2020), recipients are often only implanted between 24 and 36 months (Fitzpatrick et al., 2015). Prior to implantation, some children are exposed to one or more forms of signed language, while others are instead exposed to more limited home sign systems especially if they are born to hearing parents (Humphries et al., 2012).¹

The absence of auditory input pre-implantation can impact the speech-language environments of children with severe to profound hearing loss. In infancy, children with TH who vocalize more receive more contingent responses from caregivers (and vice versa) resulting in a social feedback loop that spurs early speech development (Gros-Louis, West, Goldstein, & King, 2006; Warlaumont, Richards, Gilkerson, & Oller, 2014). Yet despite a common saying that “*even children who are deaf babble*” (orally), prior to intervention, children who go on to receive CIs vocalize infrequently and babble immaturely (Fagan, 2014, 2015).² In reality, it takes many months *post*-implantation for pediatric CI recipients to produce speech on par with their peers with TH and even longer to engage in socially-contingent linguistic interactions with caregivers (Fagan, Bergeson, & Morris, 2014; Kondaurova, Fagan, & Zheng, 2020; Kondaurova, Smith, Zheng, Reed, & Fagan, 2020), demonstrating how auditory absence may shape the early linguistic environment.

How signal degradation shapes the early environment. The signal transmitted by the CI is degraded in ways that may clearly implicate and shape children’s

¹ This limited exposure to signed languages pre-implantation can put infants and children at risk for language deprivation syndrome (?).

² Early evidence for mature babbling among children with hearing loss is often confounded with the degree of hearing loss; see Oller and Eilers (1988) for discussion.

speech-language environments. CI users of all ages have reduced access to temporal fine structure cues, such as those that encode the fundamental frequency (f0) of the voice and reflect speaker pitch or lexical tone (Chatterjee & Peng, 2008; Deroche et al., 2019; Lee, van Hasselt, Chiu, & Cheung, 2002; Stickney, Assmann, Chang, & Zeng, 2007). Yet children with TH use pitch cues to help differentiate between speakers in their environment (Nagels, Gaudrain, Vickers, Hendriks, & Başkent, 2021). Since children with CIs do not have the same access to these cues, certain aspects of speech processing such as normalization for speaking rate or vocal tract length, as well the ability to segregate overlapping speakers, prove more difficult for them (Cleary & Pisoni, 2002; Nittrouer, Caldwell-Tarr, Moberly, & Lowenstein, 2014).

In addition, interaural differences in CI electrode insertion depths result in different frequency-to-place mismatch interaurally, compromising sound localization cues (see evidence in middle childhood: Todd, Goupell, and Litovsky (2016)).³ Deficits in localization cues can make it more difficult for children to *locate* the source of a caregiver’s voice—and a failure to look towards a speaker could alter the feedback loop that drives caregiver speech input (Wang, Bergeson, & Houston, 2017; Wang, Shafto, & Houston, 2018). The deficits may also make it difficult for children to *attune* to audio-visual cues from the lips that may help compensate for a degraded auditory signal (Bergeson, Pisoni, & Davis, 2005).

Altogether, the degraded CI signal, including compromised f0 cues and interaural frequency-to-place mismatch, may shape children’s daily language environments and interactions with caregivers: the degraded signal affects how children attune to linguistic input, potentially explaining some of the differences between children with CIs and TH in child-caregiver vocal contingency (Kondaurova, Smith, et al., 2020), and synchronization

³ Reduced sound localization cues are also due to lack of experience with acoustic hearing (for pre-lingually deafened children) and lack of experience with binaural hearing (Grieco-Calub & Litovsky, 2012) (e.g., for single-sided deafness).

during joint attention (Chen, Castellanos, Yu, & Houston, 2019), even a year post-implantation.

Current study

The purpose of this study was to evaluate how the listening experiences of children with CIs shape their everyday speech-language environments. To evaluate this, we matched a cohort of 3- to 5-year-olds with CIs, with at least 8 months of hearing experience, separately to a group of their *hearing* age- and *chronological* age-matched peers. We densely sampled the children's naturalistic home environments using child-friendly wearable recording devices (appx. 16 hrs./child or >730 total hours of observation) allowing us to assess a battery of characteristics of the home speech-language environment. Specifically, we characterized the *quantity*, *consistency*, and experience-related *differences* in children's speech input, vocal output, and conversational interactions with caregivers. Finally, we assessed how the children's speech-language input predicted their vocal productivity. Below we outline our predictions for how each metric may differ for children with CIs and TH:

1. *Why might quantity of caregiver input differ?* Children with CIs could be exposed to more caregiver input than HA matches because they are at a more advanced cognitive developmental stage, and caregivers of children with TH use more word tokens and complex grammatical structures as children develop (Huttenlocher, Vasilyeva, Waterfall, Vevea, & Hedges, 2007; Rowe, 2012). Or, alternatively, caregivers could talk more to children with CIs in an attempt to compensate, believing the child needs more exposure. Children with CIs may receive less caregiver input than chronological matches, however, if caregivers are sensitive to the children's developing linguistic capabilities, especially in the first months and years post-implantation (DesJardin & Eisenberg, 2007).

2. *Why might child vocal output differ?* The vocal output of children with CIs is likely to be less mature than chronological age matches (less frequent, shorter duration) because the children with CIs have had less experience incorporating auditory and somatosensory feedback from their own speech production due to the time spent without access to sound pre-implantation (Fagan, 2014, 2015). Signal degradation from the CIs post-implantation may also make it harder for children with CIs to establish reliable speech-motor maps, thereby delaying progression through later stages of speech development, and resulting in less mature or error-prone vocal production several months post-implantation (Serry & Blamey, 1999). (Children's vocalizations are also part of their auditory input so less mature vocalizations mean that the child would also be receiving input that is less mature.) Yet, children with CIs would be expected to have more mature vocal output than HA matches because their speech-motor apparatuses are more mature and capable of sustaining phonation, and they are also more able to coordinate inter-articulatory movement such as the tongue and the jaw (Moore, 2004; Smith & Goffman, 1998).

3. *Why might conversational interactions differ?*

The children with CIs had less opportunity to establish vocal contingency patterns (although not necessarily other forms of contingency such as gaze (Chen, Castellanos, Yu, & Houston, 2020)) pre-implantation, in infancy (Northrup & Iverson, 2020). Among children with TH, these early contingency behaviors are highly correlated with later linguistic outcomes (Bornstein, Tamis-LeMonda, & Haynes, 1999; Donnelly & Kidd, 2021; Hirsh-Pasek et al., 2015), so the reduced interactions in infancy could set the stage for different quantities or types of conversational interactions even post-implantation. Such a result would mean that at least early after gaining access to sound, the children might pattern like the HA matches, not chronological.

One might assume that chronologically older children would engage in more

conversational interactions, because they have the cognitive and linguistic skills to support the exchanges, in which case children with CIs would engage in more conversational turns than HA matches. However, children with CIs have actually been found to engage in *fewer* turns than HA matches, perhaps because their older age grants them more bodily autonomy, distance from caregivers, and opportunity to interact with other interlocutors such as siblings. This is the result concluded by Kondaurova, Zheng, VanDam, and Kinney (2022), and indeed conversational turn development among children with TH follows this developmental pattern—initially increasing in toddlerhood but decreasing in the preschool years—through 48 months of age (Gilkerson et al., 2017).

Signal degradation of, for example, f0 cues, may also play a role. Children with CIs may be less attuned to cues for speaker location, interlocutor identity, and utterance polarity (questions are produced with rising f0 contours and statements with flat or downward sloping contours and/or a creaky voice modality), all causing children with CIs to respond or engage less readily with caregivers in their environments.

4. *Why might the relationship between speech-language input and child vocal productivity differ?* Children with TH who hear more speech and/or engage in more conversational turns have more mature speech production outcomes in infancy and early childhood (Ferjan Ramírez et al., 2020; Ruan, 2022; Wang, Williams, Dilley, & Houston, 2020). However, this relationship may be less predictive for children with CIs than children with TH. First, factors such as device and implant properties, as well as rehabilitation and therapy, may play outsize roles for the vocal maturity of children with CIs, rendering caregiver input less predictive of the children's outcomes. Second, children with CIs could have more difficulty locating caregivers in the environment due to signal degradation and device limitations (Houston, 2022; Wang et al., 2018), making it more difficult for them to separate speech from other sounds and process words (Vavatzanidis, Mürbe, Friederici, & Hahne, 2018), and potentially

making conversational interactions less predictive. Finally, signal degradation from the devices could lower the quality of speech input that children with CIs receive, so hearing more input may not be as beneficial for CI users as children with TH.

Materials and Methods

Participants

Fifty-two children participated in this study. The N=18 children with CIs were individually matched by parent-reported gender, socioeconomic status, and age to two groups of children with TH: (1) *chronological* age matches, to match for cognitive and articulatory development (N=18), and (2) *hearing* age matches, to match for auditory experience (N=16 as 2 children with CIs had <1 year of hearing experience). All children were monolingual English speakers and age matching was made within 3 months whenever possible. Table 1 presents demographic information by hearing group. There were no reliable differences in number of siblings ($F(2)=1.16$, $p > .05$) or household members ($F(2)=0.98$, $p > 0.05$) by hearing group. See Supplementary Materials I for detailed reports on family composition by hearing group. This work was approved by the relevant institutions' Institutional Review Boards.

Socioeconomic status was instantiated as the highest level of maternal education achieved. To facilitate matching, we binned education into seven levels: 1) < high school, 2) high school equivalent certificate (e.g., General Education Development [GED]), 3) high school diploma, 4) technical-associate degree, 5) some college (2+ years)/ trade school, 6) college degree, and 7) graduate degree. All maternal education matching was made within 1 degree of freedom (i.e., caregiver with level 3 education preferentially matched to caregiver with level 3, and if not then matched to level 2 or 4).

N=14 children used bilateral CIs, N=2 unilateral, and N=2 had a bimodal CI+hearing aid configuration. The children with CIs had hearing parents and were being schooled in

auditory-verbal (N=11), auditory-verbal+aural focus (N=2), aural focus (N=1), cued speech (N=1), or auditory-verbal+total communication (N=1) environments.⁴ See Appendix A for detailed, by-child audiological information. All children with TH had parent-reported typical speech-language development.

Data collection

Each child completed one daylong audio recording where they wore a small, lightweight Language ENvironment Analysis (LENA) recording device (2"x3"; 2 oz.) in a specialized vest for an entire day. Recordings were completed on a typical, non-school day in the child's life. Families were instructed to turn the device on in the morning when the child awoke and continue recording for the duration of the device battery (16 hrs.). During bathtime and other water activities, parents were told to place the recorder in a safe, dry place as close to the child as possible. All families except one completed the full 16-hour recording; the remaining family completed a 12.83-hour recording. In all cases, the device continued recording while the child napped.

Audio processing

Measures of the children's home speech-language environments were semi-automatically derived from each child's recording using LENA's diarization algorithm which assigns speaker tags and timestamps to audio clips (Xu, Yapanel, & Gray, 2009). Speech clips tagged as "Target child" (CHN), "Male adult near" (MAN), and "Female adult near" (FAN) were extracted. We filtered out CHN clips that contained cries, FAN/MAN clips that contained any non-speech, and FAN/MAN clips > 10s (574 clips, 0.07%).⁵ All audio processing scripts are included in the project's Github repository

⁴ Data on school environments were unavailable for two children.

⁵ In our experience these clips longer than 10s tend to be mislabeled. The decision to remove all FAN/MAN clips that contained *any* non-speech elements inevitably resulted in the removal of some adult

Table 1

Demographic and audiological information. Mean (SD), range.

	Chrono. age matches	Cochlear implant	Hearing age matches
Chrono. Age (mos)	46.28 (10.8), 32-66	47.72 (9.84), 31-65	35 (12.71), 17-52
Gender (F,M)	9,9	9,9	9,7
Mat. Ed.	6.22 (1), 3-7	6.11 (1.02), 3-7	6.25 (1), 3-7
Hearing Age (mos)	NA	31.28 (14.3), 8-54*	NA
Activation Age (mos)	NA	16.44 (9.7), 7-45	NA
Num. of siblings	1.61 (0.98), 1-5	1.39 (0.85), 0-3	1.12 (0.96), 0-3
Num. of household members	4.56 (1.04), 3-8	4.33 (0.97), 2-6	4.06 (1.06), 2-6
Ethnicity (N) [†]			
Hispanic	2	0	2
Race (N)			
Asian	0	1	2
Black	2	0	0
White	14	14	13
American Indian	1	0	0
Asian & white	0	0	1
Black & white	0	1	0
More than 1 race (unspecified)	1	0	0

*Includes the 2 children with cochlear implants who had hearing ages < 12 mos and were thus not included in those matched by hearing age to children with typical hearing. [†]Ethnicity information was unavailable for one child with implants and race information unavailable for two children with implants.

(*masked for review* https://github.com/megseekosh/everyday_CI).

Word token count estimates from the adult clips were likewise extracted. For simplicity, in the remainder of the manuscript, we will refer to the FAN and MAN clips as “caregiver speech,” although we stress that the clips could have contained speech from a non-caregiver adult who was speaking within 10 feet of the child. Finally, algorithmic estimates of “conversational turns” were extracted. These were defined as target child and adult utterances spoken within 5 seconds, not conversations between e.g. two adults or an adult and another child. See Figure 1).

Speech input was modeled in minutes (total duration of FAN+MAN clips) and number of words from adults. Speech output was modeled in seconds (duration of CHN clips) and the number of vocalizations from the child.⁶ Caregiver-child interaction was instantiated as the number of conversational turns. We normalized all measures by hour to account for time-of-day differences, as well as different recording lengths. To derive *quantity* estimates of each construct, we computed the average number of (words, vocalizations, turns) per hour. For *consistency* estimates, we followed King, Querdasi, Humphreys, and Gotlib (2021) in computing the percentage of units containing at least one measure (speech input: percentage of minutes in the recording containing \geq one adult word; speech output: percentage of minutes containing \geq one child vocalization). For interaction, we computed the percentage of 5-minute epochs containing at least one conversational turn. Finally, to derive experience-related *differences* in the estimates, we computed the slope of the relationship between child age (in months, either chronological or hearing) and each measure. We elaborate upon this modeling in the results. Together, this workflow allowed us to assess the quantity, consistency, and experience-related

speech near the child, but the step was maximally conservative and allowed us to be sure that the adult speech clips in the final analysis contained only speech.

⁶ Seconds were used, instead of minutes as in the adult speech, because the child vocalizations were typically much shorter than the adults’ speech.




Figure 1. Daylong audio recording processing steps for the primary analysis. Hours reflect totals after removing clips without speech/crying.

differences of various metrics of the children’s speech-language environments.

There has been substantial work evaluating the LENA system’s algorithmic performance for children with and without hearing loss learning English (Cristia, Bulgarelli, & Bergelson, 2020; Gilkerson et al., 2017; Lehet, Arjmandi, Dilley, & Houston, 2021; VanDam & Silbert, 2016). (It is beyond the scope of this paper to report on validity and reliability of all LENA metrics so we refer readers to those citations.) Crucially, our analyses relied on diarization and tags that have relatively high recall and precision for the language and age group studied (e.g., “Female adult near” > 60% for English-learning infants and preschoolers (Cristia et al., 2020)) and not those categories, such as electronic speech, that have poorer reliability. Nevertheless, as algorithmic performance is not exact for any of the analyzed categories, we interpret our results by comparing *across* hearing groups. There should be no reason why algorithmic performance would be better or worse by hearing group, and we stress that reports of exact amounts of e.g., words, vocalizations, or turns per hour, should be interpreted with caution (Ferjan Ramírez, Hippe, & Kuhl, 2021; Lehet et al., 2021).

Results

We divide the results section into the various components of each child’s daily speech-language experience: (1) caregiver input, (2) target child output (production), and (3) caregiver-child conversational turns—and evaluate the impact of hearing group upon each outcome. See Table 2 for summary statistics of the measures.

Data were analyzed in the RStudio computing environment (R version 4.2.1; RStudioTeam, 2022). All computing and statistical analyses are included in the GitHub repository affiliated with this project (*masked for review* https://github.com/megseekosh/everyday_CI). Visualizations were made using `ggplot2` (Wickham, 2016) and modeling was conducted using `lme4` and `lmerTest`

packages (Bates, Maechler, Bolker, & Walker, 2015; Kuznetsova, Brockhoff, & Christensen, 2017); see project documentation for package versions. All model fitting began with a baseline, random-effects only model. Model fit improvements were evaluated by comparing model log-likelihood values and AIC estimates. Unless noted otherwise, the predictor Hearing Group (3-levels: CI, chronological age matches, hearing age matches) was contrast-coded with ‘CI’ as the reference level so model coefficients for the chronological and hearing age match groups refer to deviance from the CI group as this is our main comparison of interest. All continuous variables were mean-centered for modeling but visualizations present the unscaled data. For all examples of repeated measures, such as child vocalization duration (i.e., a duration measure was taken from each child vocalization), we fit linear mixed effects models with random intercepts by child and a fixed effect of **Hearing Group**. Models of hourly measures (words, minutes) additionally included random intercepts by hour of recording.

Input

We quantified the children’s speech-language input by computing the average number of minutes/hour that contained speech from an adult female or male near the child. We additionally computed the average number of words/hour spoken by an adult near the child.

There were no reliable differences by **Hearing Group** for measures of input quantity (hourly words, hourly minutes of speech; log-likelihood tests all $p > .05$); thus, all groups received similar amounts of input in the environment (words and minutes).

We evaluated the consistency of speech input by hearing group by computing the percentage of minutes in each recording containing ≥ 1 word from an adult (King et al., 2021). There were no differences in speech input consistency by **Hearing Group** ($p > .05$). However, speech input became more *consistent* with **Child Age** (age coded continuously,

Table 2

Measures of the naturalistic speech environment, by hearing group. Mean(SD), range.

	Chrono. age matches	Cochlear implant	Hearing age matches
Recording duration (hrs)	15.82(0.75), 12.83-16	16(0), 16-16	16(0), 16-16
Input			
Adult speech/hr (words)	1081.49(481.29), 285.63-2250.39	1217.23(508.87), 411.36-2127.7	1105.54(433.01), 170.88-1630.86
Adult speech/hr (s)*	258.36(118.69), 72.79-533.67	288.52(121.52), 95.31-499.53	264.55(102.57), 43.67-386.22
Adult word consistency	0.52(0.13), 0.31-0.78	0.58(0.11), 0.25-0.7	0.51(0.12), 0.19-0.66
Output			
Child voc. quantity	308.03(142.81), 90.12-575.81	271.75(69.23), 48.75-381.62	254.5(108.83), 42.5-424
Voc. duration (ms)	1004.46(662.3), 80-10940	937.93(569.76), 80-13270	966.59(627.6), 80-19730
Child voc. consistency	0.55(0.15), 0.34-0.84	0.58(0.13), 0.17-0.72	0.49(0.14), 0.22-0.69
Interaction			
Convo. turn quantity	61.71(32.78), 20.69-150.94	68.17(26.47), 8.5-116.75	65.13(25.47), 11.12-92.62
Convo. turn consistency	0.58(0.14), 0.38-0.84	0.64(0.13), 0.22-0.77	0.56(0.12), 0.36-0.74

*Descriptive statistics are reported for seconds of speech/hour, but modeling was conducted on minutes of speech/hour.

in months) across the entire sample, independent of hearing status (model fit: $\beta=0.004$, $t=3.16$, $p=.003$). This result indicates that speech is more continuously present throughout the day in older children (Figure 2). Note that this measure of consistency is independent of speech quantity, or the overall *amount* of speech input (words or minutes). Speech input is more consistent—more evenly spread out and less clustered into bursts over the course of the day—in older children across the sample.




Figure 2. Cross-sectional analysis of speech input consistency across entire sample (A) and by hearing status (B). Each point represents one child. Dark, gray regression line represents local regression fit to all children; ribbons represent 95% confidence intervals. Speech input becomes more consistent, and less bursty, in older children with no interaction by hearing status.

Finally, we evaluated differences by hearing group in a cross-sectional analysis of speech input by age. For this analysis, we modeled the effect of **Child Age** (in mos) upon hourly adult word token count and minutes of adult speech/hour in the children’s environments.

Naturally, children with CIs vary along three factors: chronological age, hearing age/experience, and activation age/duration of deafness. These three measures are so highly interrelated in childhood (i.e., a child with an older activation age has less hearing experience) that here we model only by chronological and hearing age (time since activation). Effects of hearing age were only modeled for children with ≥ 12 mos of hearing experience (see Section in Methods). Additionally, there was one clear outlier in age of activation (activated at 45 mos; this child also only had 8 mos hearing experience); in Supp. materials III, we replicate all modeling results with this child removed to ensure that our effects were robust to the outlier.

Hourly word counts and minutes of speech/hour increased with child age in both groups of children with TH, but not by hearing or chronological age among the children with CIs (Table 3): for every month of development, chronological age matches (spanning 32-66 mos) received approximately 21 additional words/hour and 5 additional seconds of speech/hour while hearing age matches (17-52 mos) received an additional 16 words/hour and 4 seconds of speech/hour. Again, no such cross-sectional effect by age was seen for the children with CIs, meaning that unlike children with TH, the quantity of speech input does not reflect child age (hearing or chronological) as well among children with CIs.

Output

To assess each child’s speech output (production), we computed the average number of vocalizations from the target child spoken/hour. We additionally analyzed the impact of hearing group upon the duration of children’s vocalizations.

Table 3

*Relationship between experience (age, in mos) and measures of the naturalistic speech environment, by hearing group. β =model coefficient from linear regression, p -value from linear model parameter ($***p < .001$, $**p < .01$, $*p < .05$), $+p < .1$, r =Pearson correlation coefficient. No p -value annotation indicates $p > .1$.*

	Chrono. age matches	CI chrono. age	CI hearing age	Hearing age matches
Adult words	$\beta=20.7+$ $r=0.46$	$\beta=3.81$ $r=0.07$	$\beta=0.77$ $r=0.02$	$\beta=16.49+$ $r=0.48$
Adult speech (s)	$\beta=5.27^*$ $r=0.48$	$\beta=1.12$ $r=0.09$	$\beta=0.31$ $r=0.03$	$\beta=3.78+$ $r=0.47$
Child voc. quantity	$\beta=3.98$ $r=0.34$	$\beta=2.71$ $r=0.33$	$\beta=1.06$ $r=0.17$	$\beta=8.03^{***}$ $r=0.89$
Child voc. duration (s)	$\beta=.59$ $r=0.02$	$\beta=3.16^*$ $r=0.05$	$\beta=1.84$ $r=0.03$	$\beta=6.59^{**}$ $r=0.11$
Convo. turn	$\beta=0.83$ $r=0.29$	$\beta=0.01$ $r=0$	$\beta=-0.33$ $r=-0.16$	$\beta=2.18^{***}$ $r=0.93$

For the repeated measures (vocalization duration), we fit linear mixed effects models with random intercepts by child and a fixed effect of **Hearing Group**. Models of the hourly vocalizations additionally included random intercepts by hour of recording. There was no effect of **Hearing Group** on the number of vocalizations/hour ($p > .05$); so, hearing status did not dictate the amount of the children's speech. However, there was an effect of hearing status in the model predicting vocalization duration (comparison of models with and without **Hearing Group**: $\chi^2=6.95$, $df=2$, $p=.03$): the chronological age matches produced significantly longer vocalizations than both the children with CIs ($\beta=59.25$) and hearing age matches ($\beta=78.71$).

We additionally measured the consistency of children’s speech output which we quantified as the percentage of minutes in each recording containing at least one vocalization from the target child; there was no effect of hearing experience upon children’s vocalization output consistency.

Finally, we measured the cross-sectional differences by age in vocalization quantity and duration: there was a significant, positive effect of **Child Age** (mos) on vocalization duration among the children with CIs by chronological age, and for the hearing age matches (Figure 3). With each additional month, the duration of the children with CIs’ vocalizations increased by approximately 3.16ms, a shallower slope than for the hearing age-matched children with TH (6.59ms/month; Table 3).

Caregiver-child interactions

We next evaluated the impact of hearing group upon caregiver-target child conversational turns. There was no effect of **Hearing Group** upon the quantity or consistency of turns (both $p > .05$). The cross-sectional analysis by age showed a positive relationship between age and conversational turn quantity only for the hearing age matches (e.g. for the youngest children).

Predicting vocal productivity from input measures

For the final analysis, we examined how the speech environment predicted children’s overall speech productivity and how this relationship varied by hearing group. It is expected that children who hear more speech, and engage in more linguistic interactions with caregivers, should vocalize more (Albert, Schwade, & Goldstein, 2018; Ferjan Ramírez et al., 2020; Harbison et al., 2018; Long et al., 2020; Ruan, 2022; Warlaumont et al., 2014), but it is unclear how the strength of this relationship varies by hearing status and experience. For this analysis, the measure of input that we examined was the **Average**




Figure 3. Cross-sectional analysis by age in child vocalization duration (top) and vocalization quantity (bottom), by hearing group. Given the number of distinct data points ($N=167130$ across all children), local regression lines are fit to averages over each child; ribbons represent 95% confidence intervals. See Table 3 for exact model fit statistics for each outcome.

number of conversational turns/hr and the measure of speech productivity used was the average number of target child vocalizations/hour in each recording (Figure 4).

We fit a linear regression model, controlling for chronological age (in mos), to predict the average number of target child vocalizations per hour in each recording from the parameter **Average number of conversational turns/hour**. Then, we evaluated how the relationships between input and child vocal productivity might differ by hearing status. The interaction of **Average number of conversational turns/hour** and **Hearing Group** improved upon a model without the interaction (model fit comparison: $\chi^2=3.72$, $df=2$, $p=.03$), suggesting differences in the predictive strength of conversational turns for children with CIs. Specifically, for every additional conversational turn per hour that children with CIs engaged in, they produced approximately two additional vocalizations per hour ($\beta=2.22$, $t=5.28$, $p<.001$). However, this relationship between turns and child vocal productivity was significantly steeper for both groups of children with TH who produced approximately 3 or 4 additional vocalizations per hour for every hourly conversational turn that they engaged in (chronological matches: $\beta=1.44$, $t=2.53$, $p=.02$, or a slope of 3.66; hearing age matches: $\beta=1.31$, $t=2.18$, $p=.03$, or a slope of 3.53), suggesting differences in the predictive nature of language in the home environment for children with CIs.

Discussion

Results from this study can be distilled into two main findings. First, *the language environment does not appear to reflect development as closely for children with CIs*. Unlike children with TH, older children with CIs do not hear more speech than younger children with CIs (in hearing or chronological years). Hourly conversational turns and turns were less predictive of vocal productivity for children with CIs than either group of children with TH.

Second, *children with CIs engage in just as much caregiver-child vocal interaction as*




Figure 4. Relationships between child vocal productivity and conversational turns between adult and child. Each point represents the averaged values from one child's recording and dark lines represent the local regression around those values; ribbons represent 95% confidence intervals. The relationship between hourly conversational turns and child vocal productivity is weakest for the children with CIs.

children with TH. Despite decades of research arguing that children with hearing loss interact less with caregivers (Lederberg & Mobley, 1990; Meadow-Orlans, 1997; Meadow-Orlans & Steinberg, 1993; Spencer & Meadow-Orlans, 1996), including recent evidence from more controlled but less densely sampled lab-based work (Kondaurova, Smith, et al., 2020), we show no evidence that children with CIs vocally engage with caregivers less frequently.

In the following sections, we explore both of these points in detail and situate the results in the context of previous work, especially work involving lab-based samples.

Caregiver speech input

The amount of caregiver speech in the children’s environments was predicted to differ by hearing status. Perhaps the children with CIs would hear more speech input than the hearing age matches (due to differences in cognitive maturity) but less than the chronological age matches (due to differences in linguistic ability). The final picture was more complex: although all groups received similar *amounts* of adult speech input, this differed systematically by age. In typical development, children hear more word types, tokens, and overall amounts of speech as they progress from infancy to preschoolhood (Cychosz, Edwards, Bernstein Ratner, Torrington Eaton, & Newman, 2021; Glas, Rossi, Hamdi-Sultan, Batailler, & Bellemmouche, 2018; Rowe, 2008, 2012)—we replicated these differences by age in cross-sectional samples in both of our TH groups. However, there were no such cross-sectional age differences for the children with CIs, not by hearing age or chronological age. Again, it is not the case that children with CIs simply hear more speech in general and are thus “saturated,” with little room for differences by age; there were no differences in overall input quantity by hearing group. Instead, we take this as the first piece of evidence that the language environments of children with CIs may *reflect the individual child less* (in this case the child’s hearing or chronological age) than the environments of children with TH. Alternatively, the linguistic and cognitive development

of children with CIs may correlate less strongly with age, chronological or hearing, and more with another developmental index unique to this population, such as the combination of hearing age and CI device performance or hours of daily use. In that case, caregivers of children with CIs may attune their input to *that particular construct*, which would explain the lack of an age effect in the analyses.

Child vocal output

As predicted, the children with CIs produced shorter vocalizations than chronological age matches; crucially, however, they did not vocalize less *overall* than either TH group and had some cross-sectional differences by age in vocalization duration (albeit less than the hearing age matched group). So, the children with CIs vocalize just as frequently as TH groups and the cross-sectional comparison suggests developmental progression, although longitudinal data would be needed to confirm this finding. Nevertheless, given the differences in vocalization duration between the children with CIs and chronological matches, and the significantly weaker effect of age among the children with CIs, it appears that the children with CIs follow a different vocal pattern than either TH group. It could be that children with CIs produce shorter vocalizations but supplement them with other modalities (e.g. gestures). Note that this conclusion differs from a number of studies looking at short-term changes in vocal productivity following cochlear implant activation which have found that children produce developmentally-appropriate (for their hearing age) amounts of canonical and reduplicated babble months post-activation (Fagan, 2014; Schauwers, Gillis, Daemers, De Beukelaer, & Govaerts, 2004; Schauwers, Gillis, & Govaerts, 2008). The duration of a child's vocalizations, however, reflects a number of different components in speech development: how long can a child sustain phonation, how many sequential syllables can the child produce, how many phonemes is the child producing, etc. and as such is a different metric of speech development than babbling landmarks. The vocal development findings from Fagan, Schauwers, and other colleagues

also stemmed from shorter observations (ranging 15-80 mins/child), often collected in the lab. Going forward we plan to evaluate vocal maturity in this maximally naturalistic dataset that we have collected using a combination of vocal maturity algorithms (e.g., Räsänen, Seshadri, Lavechin, Cristia, and Casillas 2020) and hand-coding which should allow us to compare our results more directly with previous research.

At-home language interventions among children with TH often target caregiver input (caregiver-child conversational turns: Ferjan Ramírez et al. (2020); Romeo et al. (2021); caregiver words: Greenwood, Schnitz, Carta, Wallisch, and Irvin (2020); Suskind, Leffel, et al. (2016); or both: Suskind et al. (2013)). But child vocal productivity partially *drives* caregiver input (Gros-Louis et al., 2006; Warlaumont et al., 2014). Indeed, a number of interventions for groups with different developmental profiles, including less verbal profiles such as children with autism spectrum disorder diagnoses or classic galactosemia, instead target the child’s own vocal productions (Peter et al., 2021). Results here suggest that preschoolers with CIs are not necessarily lacking speech input—there were no differences in overall amounts of input by group (though quantity of input vs. intake could differ). Instead, the children may require increased opportunity for their own vocal practice in order to progress developmentally and hit speech production landmarks.

Caregiver input and child vocal output

Infants and children speak more when spoken to (Albert et al., 2018; Ferjan Ramírez et al., 2020; Harbison et al., 2018; Long et al., 2020; Ruan, 2022; Warlaumont et al., 2014). Yet children with CIs may be less likely to notice when a caregiver is speaking to them, distinguish caregiver speech input from distractors, and parse individual words because CIs compromise sound localization, speaker identity, and prosodic cues (Chatterjee & Peng, 2008; Stickney et al., 2007; Todd et al., 2016). This was the result that we found—the daily speech-language environment was *less* predictive of vocal productivity for children with CIs than either TH group.

Children with CIs consistently underperform their peers with TH on almost every measure of speech, language, and literacy (Mayer & Trezek, 2018; Morini, Golinkoff, Morlet, & Houston, 2017; Nitttrouer & Caldwell-Tarr, 2016). This speech-language gap persists into adolescence and even adulthood (Desai, Stickney, & Zeng, 2008; Schorr, Roth, & Fox, 2008). And while some children with CIs develop stronger speech-language skills than others—nearly on par with their TH peers (Niparko et al., 2010)—even 4-6 years post-implantation, more than 50% of child CI recipients perform 1-2 standard deviations below peers on many standardized measures of speech and language (James, Rajput, Brinton, & Goswami, 2009; Pisoni et al., 2008).

Since the richness of the home language environment is a strong predictor of future speech, language, and literacy outcomes for a variety of populations of children with TH (Hurtado, Marchman, & Fernald, 2008; Rowe, 2012; Swanson et al., 2019; Weisleder & Fernald, 2013), and child CI recipients are at risk of speech and language delays, providing caregiver counseling to optimize the home language environments of children with CIs could be a promising method to close the spoken language gap for these children (Suskind, Graf, et al., 2016). Many unknowns remain before such interventions could reliably be implemented, but the work we present here is an important step towards making at-home interventions a reality. Demonstrating how the home environment systematically differs by hearing status affects how clinical interventions to *shape* the home linguistic environment should be implemented. For example, our results suggest that the daily speech environment is less predictive of vocal productivity among children with CIs than children with TH. So we might not expect interventions targeting increased caregiver-child vocal interaction to elicit the same vocal productivity benefit for preschoolers with CIs as those with TH; instead, we should be evaluating intervention success based on the unique input-outcome relationships for children with CIs, such as those documented here.

Limitations

Sampling biases are a concern for developmental research in general (Draper et al., 2022; Singh, Rajendra, & Mazuka, 2022), but may be especially prevalent for methodologies such as at-home recordings since some families are unwilling to record in their homes. Marginalized groups with a history of being tracked may, understandably, be especially wary. As such, samples in developmental science that already skew white and middle to upper class may be especially biased for at-home methods. Thus, we stress that current findings about e.g. the strength of the relationship between adult input and child vocal production may not generalize to all children, even within North America.

Another limitation of this study is that recordings were only taken on one day of the child's life. Although our own work and others' suggests that differences across days *within* households are far less than differences *between* households (de Barbaro & Fausey, 2022; Havard et al., 2023), future work could more comprehensively address this concern by ensuring that all families record over two or three days, as some others have done (Romeo et al., 2018). Another way to ensure sampling consistency, but instead between families, would be to request that families record, for example, on both one weekend and week day, or for an entire weekend. Observing each child for 16 hrs. already lends insight that more limited, in-lab observations cannot; yet collecting even denser samples than those we present here—those that span longer time periods—could ensure that the observations made on a single day in the child's life are not biased by exceptional events.

Conclusion

This work evaluated the daily speech environments of preschoolers with cochlear implants in comparison to two groups of their peers with typical hearing. Using incredibly dense sampling of children's everyday environments in their homes, we assessed how a battery of everyday speech-language experiences—caregiver speech input, child vocal

production, and caregiver-child conversational turns—differed by hearing status. Take-aways are that (1) the speech-language environment reflects development less closely for children with implants than typical hearing and (2) there were minimal differences by hearing status in caregiver-child interaction, even after implementing the measure in multiple ways. The unique auditory experiences of preschoolers who receive cochlear implants, the time they spend without auditory access pre-implantation and the degraded device signal they learn from post-implantation, do shape their everyday speech and language environments.

Acknowledgements

The authors wish to thank the children and families who contributed data to this study. Additional thanks to Kim Coulter (LENA Foundation) for her support on this project. This work was funded by National Institute on Deafness and Other Communication Disorders grants F32DC019539 (M.C.), R01DC02932 (J.R.E., B.M., & Mary E. Beckman) and National Institute of Child Health and Human Development grants F31HD086957, (R.R.R.), K99/R00HD103873 (R.R.R.), and F32HD103439 (J.E.K), as well as National Science Foundation SBE Postdoctoral Research Fellowship 2004983 (J.E.K).

Data availability statement

All data and code used to generate the analyses in this paper are publicly available at: *masked for review*

Author contributions

Conception: M.C., R.S.N., and J.R.E. Data collection and funding: M.C., J.R.E., B.M., R.R.R., and J.K. Manuscript writing: M.C. Results and analyses: M.C. Manuscript editing: all authors.

Appendix

Table A1

Audiological information from children with cochlear implants

Participant	Chronological age	Age at hearing loss (mos)	Age at activation	Hearing age (since activation)	Etiology	Device configuration	Activation
300ECV1	58	0	13	45	genetic	bilateral	simultaneous
301ECV1	53	0	45	8	unknown	bilateral	R-L
302ECV1	37	0	13	24	unknown	bilateral	R-L
303ECV1	65	6	13	52	unknown	bilateral	simultaneous
304ECV1	48	0	12	36	genetic	bilateral	R-L
307ECV1	44	0	15	29	genetic	bilateral	R-L
308ECV1	39	0	13	26	genetic	bilateral	simultaneous
311ECV1	62	9	13	49	unknown	bilateral	L-R
312ECV1	44	0	24	20	genetic	unilateral	R
314ECV1	38	10	17	21	unknown	bilateral	R-L
801ECV1	39	1.5	15	24	unknown	bilateral	simultaneous
804ECV1	56	0	7	49	genetic	bilateral	simultaneous
806ECV1	45	14	34	11	genetic	unilateral	L
807ECV1	51	10	22	29	Mondini malformation	bimodal	n/a
309ECV1	61	0.5	7	54	genetic	bilateral	simultaneous
306ECV1	49	0	8	41	unknown	bilateral	R-L
605LTP1	31	0	16	15	unknown	bimodal	n/a
608LTP1	39	0.5	9	30	Connexin 26	bilateral	simultaneous

Information about the specific dB of hearing loss was not available.

1 References

- Albert, R. R., Schwade, J. A., & Goldstein, M. H. (2018). The social functions of babbling: Acoustic and contextual characteristics that facilitate maternal responsiveness. *Developmental Science*, 21(5), e12641. doi: 10.1111/desc.12641
- Ambrose, S. E., VanDam, M., & Moeller, M. P. (2014). Linguistic Input, Electronic Media, and Communication Outcomes of Toddlers with Hearing Loss. *Ear and Hearing*, 35(2), 139–147. doi: 10.1097/AUD.0b013e3182a76768
- Ambrose, S. E., Walker, E. A., Unflat-Berry, L. M., Oleson, J. J., & Moeller, M. P. (2015). Quantity and Quality of Caregivers' Linguistic Input to 18-Month and 3-Year-Old Children Who Are Hard of Hearing: *Ear and Hearing*, 36(1), 48S-59S. doi: 10.1097/AUD.0000000000000209
- Arjmandi, M., Houston, D., Wang, Y., & Dilley, L. C. (2021). Estimating the reduced benefit of infant-directed speech in cochlear implant-related speech processing. *Neuroscience Research*, S0168010221000213. doi: 10.1016/j.neures.2021.01.007
- Bates, D., Maechler, M., Bolker, B., & Walker, S. (2015). Fitting linear mixed-effects models using lme4. *Journal of Statistical Software*, 67(1), 1–48.
- Bergelson, E., Casillas, M., Soderstrom, M., Seidl, A., Warlaumont, A. S., & Amatuni, A. (2019). What Do North American Babies Hear? A large-scale cross-corpus analysis. *Developmental Science*, 22(1), e12724. doi: 10.1111/desc.12724
- Bergeson, T. R., Pisoni, D. B., & Davis, R. A. O. (2005). Development of Audiovisual Comprehension Skills in Prelingually Deaf Children With Cochlear Implants. *Ear and hearing*, 26(2), 149–164.
- Bornstein, M. H., Tamis-LeMonda, C. S., & Haynes, O. (1999). First words in the second year: Continuity, stability, and models of concurrent and predictive correspondence in vocabulary and verbal responsiveness across age and context. *Infant Behavior and Development*, 22(1), 65–85. doi: 10.1016/S0163-6383(99)80006-X
- Chatterjee, M., & Peng, S.-C. (2008). Processing F0 with cochlear implants: Modulation

frequency discrimination and speech intonation recognition. *Hearing Research*,
 235(1-2), 143–156. doi: 10.1016/j.heares.2007.11.004

Chen, C.-h., Castellanos, I., Yu, C., & Houston, D. M. (2019). Effects of children’s hearing
 loss on the synchrony between parents’ object naming and children’s attention.
Infant Behavior and Development, 57, 101322. doi: 10.1016/j.infbeh.2019.04.004

Chen, C.-h., Castellanos, I., Yu, C., & Houston, D. M. (2020). What leads to coordinated
 attention in parent–toddler interactions? Children’s hearing status matters.
Developmental Science, 23(3). doi: 10.1111/desc.12919

Cleary, M., & Pisoni, D. B. (2002). Talker Discrimination by Prelingually Deaf Children
 with Cochlear Implants: Preliminary Results. *Annals of Otology, Rhinology &*
Laryngology, 111(5_suppl), 113–118. doi: 10.1177/00034894021110S523

Cristia, A., Bulgarelli, F., & Bergelson, E. (2020). Accuracy of the Language Environment
 Analysis System Segmentation and Metrics: A Systematic Review. *Journal of*
Speech, Language, and Hearing Research, 63(4), 1093–1105. doi:
 10.1044/2020_JSLHR-19-00017

Cychosz, M., Edwards, J. R., Bernstein Ratner, N., Torrington Eaton, C., & Newman,
 R. S. (2021). Acoustic-Lexical Characteristics of Child-Directed Speech Between 7
 and 24 Months and Their Impact on Toddlers’ Phonological Processing. *Frontiers in*
Psychology, 12, 712647. doi: 10.3389/fpsyg.2021.712647

de Barbaro, K., & Fausey, C. M. (2022). Ten Lessons About Infants’ Everyday
 Experiences. *Current Directions in Psychological Science*, 31(1), 28–33. doi:
 10.1177/09637214211059536

Deroche, M. L. D., Lu, H.-P., Kulkarni, A. M., Caldwell, M., Barrett, K. C., Peng, S.-C.,
 ... Chatterjee, M. (2019). A tonal-language benefit for pitch in normally-hearing and
 cochlear-implanted children. *Scientific Reports*, 9, 109. doi:
 10.1038/s41598-018-36393-1

Desai, S., Stickney, G., & Zeng, F.-G. (2008). Auditory-visual speech perception in

- normal-hearing and cochlear-implant listeners. *The Journal of the Acoustical Society of America*, 123(1), 428–440. doi: 10.1121/1.2816573
- DesJardin, J. L., & Eisenberg, L. S. (2007). Maternal Contributions: Supporting Language Development in Young Children with Cochlear Implants. *Ear and Hearing*, 28(4), 456–469. doi: 10.1097/AUD.0b013e31806dc1ab
- Dilley, L. C., Lehet, M., Wieland, E. A., Arjmandi, M. K., Kondaurova, M., Wang, Y., . . . Bergeson, T. (2020). Individual Differences in Mothers' Spontaneous Infant-Directed Speech Predict Language Attainment in Children With Cochlear Implants. *Journal of Speech, Language, and Hearing Research*, 63(7), 2453–2467. doi: 10.1044/2020_JSLHR-19-00229
- Dilley, L. C., Millett, A. L., Mcauley, J. D., & Bergeson, T. R. (2014). Phonetic variation in consonants in infant-directed and adult-directed speech: The case of regressive place assimilation in word-final alveolar stops. *Journal of Child Language*, 41(1), 155–175. doi: 10.1017/S0305000912000670
- Donnelly, S., & Kidd, E. (2021). The Longitudinal Relationship Between Conversational Turn-Taking and Vocabulary Growth in Early Language Development. *Child Development*, 92(2), 609–625. doi: 10.1111/cdev.13511
- Draper, C. E., Barnett, L. M., Cook, C. J., Cuartas, J. A., Howard, S. J., McCoy, D. C., . . . Yousafzai, A. K. (2022). Publishing child development research from around the world: An unfair playing field resulting in most of the world's child population under-represented in research. *Infant and Child Development*, e2375. doi: 10.1002/icd.2375
- Fagan, M. K. (2014). Frequency of vocalization before and after cochlear implantation: Dynamic effect of auditory feedback on infant behavior. *Journal of experimental child psychology*, 126, 328–338. doi: 10.1016/j.jecp.2014.05.005
- Fagan, M. K. (2015). Why repetition? Repetitive babbling, auditory feedback, and cochlear implantation. *Journal of experimental child psychology*, 137, 125–136. doi:

10.1016/j.jecp.2015.04.005

Fagan, M. K., Bergeson, T. R., & Morris, K. J. (2014). Synchrony, Complexity and Directiveness in Mothers' Interactions with Infants Pre- and Post-Cochlear Implantation. *Infant behavior & development*, 37(3), 249–257. doi: 10.1016/j.infbeh.2014.04.001

Ferjan Ramírez, N., Hippe, D. S., & Kuhl, P. K. (2021). Comparing Automatic and Manual Measures of Parent–Infant Conversational Turns: A Word of Caution. *Child Development*, 92(2), 672–681. doi: 10.1111/cdev.13495

Ferjan Ramírez, N., Lytle, S. R., & Kuhl, P. K. (2020). Parent coaching increases conversational turns and advances infant language development. *Proceedings of the National Academy of Sciences*, 117(7), 3484–3491. doi: 10.1073/pnas.1921653117

Fitzpatrick, E. M., Ham, J., & Whittingham, J. (2015). Pediatric Cochlear Implantation: Why Do Children Receive Implants Late? *Ear and Hearing*, 36(6), 688–694.

Ganek, H. V., Cushing, S. L., Papsin, B. C., & Gordon, K. A. (2020). Cochlear Implant Use Remains Consistent Over Time in Children With Single-Sided Deafness. *Ear & Hearing*, 41(3), 678–685. doi: 10.1097/AUD.0000000000000797

Gilkerson, J., Richards, J. A., Warren, S. F., Montgomery, J. K., Greenwood, C. R., Kimbrough Oller, D., . . . Paul, T. D. (2017). Mapping the Early Language Environment Using All-Day Recordings and Automated Analysis. *American Journal of Speech-Language Pathology*, 26(2), 248–265. doi: 10.1044/2016_AJSLP-15-0169

Glas, L., Rossi, C., Hamdi-Sultan, R., Batailler, C., & Bellemmouche, H. (2018). Activity types and child-directed speech: A comparison between French, Tunisian Arabic and English. *Canadian Journal of Linguistics/Revue canadienne de linguistique*, 63(4), 633–666. doi: 10.1017/cnj.2018.20

Greenwood, C. R., Schnitz, A. G., Carta, J. J., Wallisch, A., & Irvin, D. W. (2020). A systematic review of language intervention research with low-income families: A word gap prevention perspective. *Early Childhood Research Quarterly*, 50, 230–245. doi:

10.1016/j.ecresq.2019.04.001

Grieco-Calub, T. M., & Litovsky, R. Y. (2012). Spatial acuity in two-to-three-year-old children with normal acoustic hearing, unilateral cochlear implants and bilateral cochlear implants. *Ear and hearing*, 33(5), 561–572. doi: 10.1097/AUD.0b013e31824c7801

Gros-Louis, J., West, M. J., Goldstein, M. H., & King, A. P. (2006). Mothers provide differential feedback to infants' prelinguistic sounds. *International Journal of Behavioral Development*, 30(6), 509–516. doi: 10.1177/0165025406071914

Harbison, A. L., Woynaroski, T. G., Tapp, J., Wade, J. W., Warlaumont, A. S., & Yoder, P. J. (2018). A new measure of child vocal reciprocity in children with autism spectrum disorder. *Autism Research*, 11(6), 903–915. doi: 10.1002/aur.1942

Havard, W., Gautheron, L., Zhang, Z., Soderstrom, M., Schuller, B., Scaff, C., . . . Cristia, A. (2023). Establishing the reliability of measures extracted from long-form recordings using LENA and the ACLEW pipeline. *submitted*.

Hirsh-Pasek, K., Adamson, L. B., Bakeman, R., Owen, M. T., Golinkoff, R. M., Pace, A., . . . Suma, K. (2015). The Contribution of Early Communication Quality to Low-Income Children's Language Success. *Psychological Science*, 26(7), 1071–1083. doi: 10.1177/0956797615581493

Houston, D. M. (2022). A framework for understanding the relation between spoken language input and outcomes for children with cochlear implants. *Child Development Perspectives*, cdep.12443. doi: 10.1111/cdep.12443

Humphries, T., Kushalnagar, P., Mathur, G., Napoli, D. J., Padden, C., Rathmann, C., & Smith, S. R. (2012). Language acquisition for deaf children: Reducing the harms of zero tolerance to the use of alternative approaches. *Harm Reduction Journal*, 9(1), 16. doi: 10.1186/1477-7517-9-16

Hurtado, N., Marchman, V. A., & Fernald, A. (2008). Does input influence uptake? Links between maternal talk, processing speed and vocabulary size in Spanish-learning

children. *Developmental Science*, 11(6), F31-F39. doi:

10.1111/j.1467-7687.2008.00768.x

Huttenlocher, J., Vasilyeva, M., Waterfall, H. R., Vevea, J. L., & Hedges, L. V. (2007).

The varieties of speech to young children. *Developmental Psychology*, 43(5),

1062–1083. doi: 10.1037/0012-1649.43.5.1062

James, D., Rajput, K., Brinton, J., & Goswami, U. (2009). Orthographic influences,

vocabulary development, and phonological awareness in deaf children who use

cochlear implants. *Applied Psycholinguistics*, 30(4), 659–684. doi:

10.1017/S0142716409990063

Kalashnikova, M., & Burnham, D. (2018). Infant-directed speech from seven to nineteen

months has similar acoustic properties but different functions. *Journal of Child*

Language, 45(5), 1035–1053. doi: 10.1017/S0305000917000629

King, L. S., Querdasi, F. R., Humphreys, K. L., & Gotlib, I. H. (2021). Dimensions of the

language environment in infancy and symptoms of psychopathology in toddlerhood.

Developmental Science, 24(5). doi: 10.1111/desc.13082

Kondaurova, M. V., Fagan, M. K., & Zheng, Q. (2020). Vocal imitation between mothers

and their children with cochlear implants. *Infancy*, infa.12363. doi:

10.1111/infa.12363

Kondaurova, M. V., Smith, N. A., Zheng, Q., Reed, J., & Fagan, M. K. (2020). Vocal

Turn-Taking Between Mothers and Their Children With Cochlear Implants:. *Ear and*

Hearing, 41(2), 362–373. doi: 10.1097/AUD.0000000000000769

Kondaurova, M. V., Zheng, Q., VanDam, M., & Kinney, K. (2022). Vocal Turn-Taking in

Families With Children With and Without Hearing Loss. *Ear & Hearing*, 43(3),

883–898. doi: 10.1097/AUD.0000000000001135

Kuznetsova, A., Brockhoff, P., & Christensen, R. (2017). lmerTest Package: Tests in linear

mixed-effects models. *Journal of Statistical Software*, 82(13), 1–26.

Lederberg, A., & Mobley, C. (1990). The Effect of Hearing Impairment on the Quality of

- Attachment and Mother-Toddler Interaction. *Child development*, 61, 1596–604. doi: 10.1111/j.1467-8624.1990.tb02886.x
- Lee, K. Y., van Hasselt, C., Chiu, S., & Cheung, D. M. (2002). Cantonese tone perception ability of cochlear implant children in comparison with normal-hearing children. *International Journal of Pediatric Otorhinolaryngology*, 63(2), 137–147. doi: 10.1016/S0165-5876(02)00005-8
- Lehet, M., Arjmandi, M. K., Dilley, L. C., & Houston, D. (2021). Circumspection in using automated measures: Talker gender and addressee affect error rates for adult speech detection in the Language ENvironment Analysis (LENA) system. *Behavior Research Methods*, 53, 113–138.
- Long, H., Bowman, D., Yoo, H., Burkhardt-Reed, M., Bene, E., & Oller, D. K. (2020). Social and endogenous infant vocalizations. *PLoS ONE*, 15(8), e0224956. doi: 10.1101/821371
- Majorano, M., Brondino, M., Guerzoni, L., Murri, A., Ferrari, R., Lavelli, M., . . . Persici, V. (2021). Do Acoustic Environment Characteristics Affect the Lexical Development of Children With Cochlear Implants? A Longitudinal Study Before and After Cochlear Implant Activation. *American Journal of Audiology*, 30(3), 602–615. doi: 10.1044/2021_AJA-20-00104
- Mayer, C., & Trezek, B. J. (2018). Literacy Outcomes in Deaf Students with Cochlear Implants: Current State of the Knowledge. *The Journal of Deaf Studies and Deaf Education*, 23(1), 1–16. doi: 10.1093/deafed/enx043
- Meadow-Orlans, K. P. (1997). Effects of Mother and Infant Hearing Status on Interactions at Twelve and Eighteen Months. *Journal of Deaf Studies and Deaf Education*, 2(1), 26–36. doi: 10.1093/oxfordjournals.deafed.a014307
- Meadow-Orlans, K. P., & Steinberg, A. (1993). Effects of infant hearing loss and maternal support on mother-infant interactions at 18 months. *Applied Developmental Psychology*, 14, 407–426.

- Moore, C. A. (2004). Physiologic development of speech production. In B. Maassen,
R. Kent, H. Peters, P. van Lieshout, & W. Hulstijn (Eds.), *Speech motor control in
normal and disordered speech* (pp. 191–209). Oxford, UK: Oxford University Press.
- Morini, G., Golinkoff, R. M., Morlet, T., & Houston, D. M. (2017). Advances in pediatric
hearing loss: A road to better language outcomes. *Translational Issues in
Psychological Science*, 3(1), 80–93. doi: 10.1037/tps0000106
- Nagels, L., Gaudrain, E., Vickers, D., Hendriks, P., & Başkent, D. (2021). School-age
children benefit from voice gender cue differences for the perception of speech in
competing speech. *The Journal of the Acoustical Society of America*, 149(5),
3328–3344. doi: 10.1121/10.0004791
- Newman, R. S., Rowe, M. L., & Bernstein Ratner, N. (2016). Input and uptake at 7
months predicts toddler vocabulary: The role of child-directed speech and infant
processing skills in language development. *Journal of Child Language*, 43(5),
1158–1173. doi: 10.1017/S0305000915000446
- Niparko, J. K., Tobey, E. A., Thal, D. J., Eisenberg, L. S., Wang, N.-Y., Quittner, A. L., &
Fink, N. E. (2010). Spoken Language Development in Children Following Cochlear
Implantation. *Journal of the American Medical Association*, 303(15), 1498–1506.
- Nittrouer, S., & Caldwell-Tarr, A. (2016). Language and Literacy Skills in Children with
Cochlear Implants: Past and Present Findings. In N. M. Young & K. Iler Kirk
(Eds.), *Pediatric Cochlear Implantation* (pp. 177–197). New York, NY: Springer New
York. doi: 10.1007/978-1-4939-2788-3_11
- Nittrouer, S., Caldwell-Tarr, A., Moberly, A. C., & Lowenstein, J. H. (2014). Perceptual
weighting strategies of children with cochlear implants and normal hearing. *Journal
of Communication Disorders*, 52, 111–133. doi: 10.1016/j.jcomdis.2014.09.003
- Nittrouer, S., Lowenstein, J. H., & Antonelli, J. (2019). Parental Language Input to
Children With Hearing Loss: Does It Matter in the End? *Journal of Speech,
Language, and Hearing Research*, 63(1), 234–258. doi:

10.1044/2019_JSLHR-19-00123

Northrup, J. B., & Iverson, J. M. (2020). The Development of Mother-Infant Coordination

Across the First Year of Life. *Developmental psychology*, 56(2), 221–236. doi:

10.1037/dev0000867

Oller, D. K., & Eilers, R. E. (1988). The Role of Audition in Infant Babbling. *Child*

Development, 59(2), 441–449.

Peter, B., Davis, J., Cotter, S., Belter, A., Williams, E., Stumpf, M., . . . Potter, N. (2021).

Toward Preventing Speech and Language Disorders of Known Genetic Origin: First

Post-Intervention Results of Babble Boot Camp in Children With Classic

Galactosemia. *American Journal of Speech-Language Pathology*, 30(6), 2616–2634.

doi: 10.1044/2021_AJSLP-21-00098

Pisoni, D. B., Conway, C. M., Kronenberger, W., Horn, D. L., Karpicke, J., & Henning, S.

(2008). Efficacy and effectiveness of cochlear implants in deaf children. In *In Deaf*

Cognition: Foundations and Outcomes, M. Marschark and P. C. Hauser, Eds., pp. 52–

101. University Press.

Räsänen, O., Seshadri, S., Lavechin, M., Cristia, A., & Casillas, M. (2020). ALICE: An

open-source tool for automatic measurement of phoneme, syllable, and word counts

from child-centered daylong recordings. *Behavior Research Methods*. doi:

10.3758/s13428-020-01460-x

Romeo, R. R., Leonard, J. A., Grotzinger, H. M., Robinson, S. T., Takada, M. E., Mackey,

A. P., . . . Gabrieli, J. D. (2021). Neuroplasticity associated with changes in

conversational turn-taking following a family-based intervention. *Developmental*

Cognitive Neuroscience, 49, 100967. doi: 10.1016/j.dcn.2021.100967

Romeo, R. R., Leonard, J. A., Robinson, S. T., West, M. R., Mackey, A. P., Rowe, M. L.,

& Gabrieli, J. D. E. (2018). Beyond the 30-Million-Word Gap: Children's

Conversational Exposure Is Associated With Language-Related Brain Function.

Psychological Science, 29(5), 700–710. doi: 10.1177/0956797617742725

- Rowe, M. L. (2008). Child-directed speech: Relation to socioeconomic status, knowledge of child development, and child vocabulary skill. *Journal of Child Language*, 35(1), 185–205.
- Rowe, M. L. (2012). A Longitudinal Investigation of the Role of Quantity and Quality of Child-Directed Speech in Vocabulary Development: Child-Directed Speech and Vocabulary. *Child Development*, 83(5), 1762–1774. doi: 10.1111/j.1467-8624.2012.01805.x
- Rowe, M. L., & Weisleder, A. (2020). Language Development in Context. *Annual Review of Developmental Psychology*, 2, 201–223.
- RStudioTeam. (2020). *RStudio: Integrated Development for R*. Boston, MA: RStudio, Inc.
- Ruan, Y. (2022). *Infant Volubility and Bilingual Input in Naturalistic Day-long Recordings* (Unpublished doctoral dissertation). McGill University, Montreal, CA.
- Schauwers, K., Gillis, S., Daemers, K., De Beukelaer, C., & Govaerts, P. J. (2004). Cochlear Implantation Between 5 and 20 Months of Age: The Onset of Babbling and the Audiologic Outcome:. *Otology & Neurotology*, 25(3), 263–270. doi: 10.1097/00129492-200405000-00011
- Schauwers, K., Gillis, S., & Govaerts, P. J. (2008). The Characteristics of Prelexical Babbling After Cochlear Implantation Between 5 and 20 Months of Age. *Ear & Hearing*, 29(4), 627–637. doi: 10.1097/AUD.0b013e318174f03c
- Schorr, E. A., Roth, F. P., & Fox, N. A. (2008). A Comparison of the Speech and Language Skills of Children With Cochlear Implants and Children With Normal Hearing. *Communication Disorders Quarterly*, 29(4), 195–210. doi: 10.1177/1525740108321217
- Schwab, J. F., & Lew-Williams, C. (2016). Language learning, socioeconomic status, and child-directed speech. *WIREs Cognitive Science*, 7(4), 264–275. doi: 10.1002/wcs.1393
- Serry, T. A., & Blamey, P. J. (1999). A 4-Year Investigation Into Phonetic Inventory

- Development in Young Cochlear Implant Users. *Journal of Speech, Language, and Hearing Research*, 42(1), 141–154. doi: 10.1044/jslhr.4201.141
- Singh, L., Rajendra, S. J., & Mazuka, R. (2022). Diversity and representation in studies of infant perceptual narrowing. *Child Development Perspectives*, cdep.12468. doi: 10.1111/cdep.12468
- Smith, A., & Goffman, L. (1998). Stability and Patterning of Speech Movement Sequences in Children and Adults. *Journal of Speech Language and Hearing Research*, 41(1), 18–30. doi: 10.1044/jslhr.4101.18
- Spencer, P. E., & Meadow-Orlans, K. P. (1996). Play, Language, and Maternal Responsiveness: A Longitudinal Study of Deaf and Hearing Infants. *Child Development*, 67(6), 3176–3191. doi: 10.1111/j.1467-8624.1996.tb01908.x
- Stickney, G. S., Assmann, P. F., Chang, J., & Zeng, F.-G. (2007). Effects of cochlear implant processing and fundamental frequency on the intelligibility of competing sentences. *The Journal of the Acoustical Society of America*, 122(2), 1069–1078. doi: 10.1121/1.2750159
- Suskind, D. L., Graf, E., Leffel, K. R., Hernandez, M. W., Suskind, E., Webber, R., . . . Nevins, M. E. (2016). Project ASPIRE: Spoken Language Intervention Curriculum for Parents of Low-socioeconomic Status and Their Deaf and Hard-of-Hearing Children. *Otology & Neurotology*, 37(2), e110-e117. doi: 10.1097/MAO.0000000000000931
- Suskind, D. L., Leffel, K. R., Graf, E., Hernandez, M. W., Gunderson, E. A., Sapolich, S. G., . . . Levine, S. C. (2016). A parent-directed language intervention for children of low socioeconomic status: A randomized controlled pilot study. *Journal of Child Language*, 43(2), 366–406. doi: 10.1017/S0305000915000033
- Suskind, D. L., Leffel, K. R., Hernandez, M. W., Sapolich, S. G., Suskind, E., Kirkham, E., & Meehan, P. (2013). An Exploratory Study of “Quantitative Linguistic Feedback”: Effect of LENA Feedback on Adult Language Production. *Communication Disorders*

- Quarterly*, 34(4), 199–209. doi: 10.1177/1525740112473146
- Swanson, M. R., Donovan, K., Paterson, S., Wolff, J. J., Parish-Morris, J., Meera, S. S., . . . Piven, J. (2019). Early language exposure supports later language skills in infants with and without autism. *Autism Research*, 12(12), 1784–1795. doi: 10.1002/aur.2163
- Todd, A. E., Goupell, M. J., & Litovsky, R. Y. (2016). Binaural release from masking with single- and multi-electrode stimulation in children with cochlear implants. *The Journal of the Acoustical Society of America*, 140(1), 59–73. doi: 10.1121/1.4954717
- USFDA. (2020). *Summary of safety and effectiveness data (SSED): Nucleus 24 Cochlear Implant System* (FDA Summary of Safety and Effectiveness Data No. PMA P970051/S172). Silver Spring, MD, USA: US Food and Drug Administration.
- VanDam, M., & Silbert, N. H. (2016). Fidelity of Automatic Speech Processing for Adult and Child Talker Classifications. *PLOS ONE*, 11(8), e0160588. doi: 10.1371/journal.pone.0160588
- Vavatzanidis, N. K., Mürbe, D., Friederici, A. D., & Hahne, A. (2018). Establishing a mental lexicon with cochlear implants: An ERP study with young children. *Scientific Reports*, 8(1), 910. doi: 10.1038/s41598-017-18852-3
- Wang, Y., Bergeson, T. R., & Houston, D. M. (2017). Infant-Directed Speech Enhances Attention to Speech in Deaf Infants With Cochlear Implants. *Journal of Speech, Language, and Hearing Research*, 60(11), 3321–3333. doi: 10.1044/2017_JSLHR-H-17-0149
- Wang, Y., Jung, J., Bergeson, T. R., & Houston, D. M. (2020). Lexical Repetition Properties of Caregiver Speech and Language Development in Children With Cochlear Implants. *Journal of Speech, Language, and Hearing Research*, 63(3), 872–884. doi: 10.1044/2019_JSLHR-19-00227
- Wang, Y., Shafto, C. L., & Houston, D. M. (2018). Attention to speech and spoken language development in deaf children with cochlear implants: A 10-year longitudinal

study. *Developmental Science*, 21(6), e12677. doi: 10.1111/desc.12677

Wang, Y., Williams, R., Dilley, L. C., & Houston, D. M. (2020). A meta-analysis of the predictability of LENA™ automated measures for child language development.

Developmental Review, 57, 100921. doi: 10.1016/j.dr.2020.100921

Warlaumont, A. S., Richards, J. A., Gilkerson, J., & Oller, D. K. (2014). A Social Feedback Loop for Speech Development and Its Reduction in Autism. *Psychological Science*, 25(7), 1314–1324. doi: 10.1177/0956797614531023

Weisleder, A., & Fernald, A. (2013). Talking to children matters: Early language experience strengthens processing and builds vocabulary. *Psychological Science*, 24(11), 2143–2152. doi: 10.1177/0956797613488145

Wickham, H. (2016). *Ggplot2: Elegant Graphics for Data Analysis*. New York: Springer-Verlag New York.

Xu, D., Yapanel, U., & Gray, S. (2009). *Reliability of the LENA Language Environment Analysis System in young children's natural home environment* (Technical Report ITR-05-2). Boulder, CO: LENA Research Foundation.