

حل تکلیف سری دو - سوال اول

```
clc;
close all;
clear all;

I=imread('pout.tif');
imshow(I);
title('Original Image');

for J=0.01:0.01:0.02
    figure;
    I0=imnoise(I,'gaussian',0,J);
    imshow(I0);
    title(strcat('Noisy Image (Gaussian,d=',num2str(J),'')'));

    figure;
    Mean_Filter(I0,3,1,2,1);
    Mean_Filter(I0,5,1,2,2);

    figure;
    Gaussian_Filter(I0,3,0.3,1,2,1);
    Gaussian_Filter(I0,5,0.6,1,2,2);

    figure;
    I1=imnoise(I,'salt & pepper',J);
    imshow(I1);
    title(strcat('Noisy Image (salt & pepper,d=',num2str(J),'')'));

    figure;
    Mean_Filter(I1,3,1,2,1);
    Mean_Filter(I1,5,1,2,2);

    figure;
    Gaussian_Filter(I1,3,0.3,1,2,1);
    Gaussian_Filter(I1,5,0.6,1,2,2);

    figure;
    Median_Filter(I1,3,1,2,1);
    Median_Filter(I1,5,1,2,2);
end

function Mean_Filter(NI,SoW,X,Y,Z)
W=ones(SoW)/(SoW*SoW);
L=imfilter(NI,W);
subplot(X,Y,Z);
imshow(L);
title(strcat('Mean Filter (',int2str(SoW),'*',int2str(SoW),'')'));
end

function Gaussian_Filter(NI,hsize,sigma,X,Y,Z)
H=fspecial('gaussian',hsize,sigma);
L=imfilter(NI,H,'replicate');
```

```
subplot(X,Y,Z);
imshow(L);
title(strcat('Gaussian Filter (',num2str(hsize),' ',num2str(sigma),'')');
end
```

```
function Median_Filter(NI,SoW,X,Y,Z)
L=medfilt2(NI,[SoW SoW]);
subplot(X,Y,Z);
imshow(L);
title(strcat('Median Filter (',int2str(SoW),'*',int2str(SoW),'')');
end
```

حل تکلیف سری دو - سوال دوم - بند الف

```
clc;
close all;
clear all;

I=imread('circuit.tif');
subplot(2,3,1);
imshow(I);
title('Original Image');

BW1=edge(I,'sobel');
subplot(2,3,2);
imshow(BW1);
title('Sobel Method');

BW2=edge(I,'prewitt');
subplot(2,3,3);
imshow(BW2);
title('Prewitt Method');

BW3=edge(I,'roberts');
subplot(2,3,4);
imshow(BW3);
title('Roberts Method');

BW4=edge(I,'canny');
subplot(2,3,5);
imshow(BW4);
title('Canny Method');
```

حل تکلیف سری دو - سوال دوم - بند ب

```
clc;
close all;
clear all;

I=imread('circuit.tif');
subplot(2,3,1);
imshow(I);
title('Original Image');

BW1=edge(I,'sobel');
subplot(2,3,2);
imshow(BW1);
title('Sobel Method');
```

```

BW2=edge(I,'prewitt');
subplot(2,3,3);
imshow(BW2);
title('Prewitt Method');

BW3=edge(I,'roberts');
subplot(2,3,4);
imshow(BW3);
title('Roberts Method');

BW4=edge(I,'canny');
subplot(2,3,5);
imshow(BW4);
title('Canny Method');

BW5=edge(I,'log');
subplot(2,3,6);
imshow(BW5);
title('Laplacian of Gaussian');

```

حل تکيف سري دو - سوال دوم - بند ج

```

clc;
close all;
clear all;

I=imread('circuit.tif');
subplot(2,4,1);
imshow(I);
title('Original Image');

J=imnoise(I,'salt & pepper',0.04);
subplot(2,4,2);
imshow(J);
title('Salt and Pepper Noise');

BW1=edge(J,'sobel');
subplot(2,4,3);
imshow(BW1);
title('Sobel Method');

BW2=edge(J,'prewitt');
subplot(2,4,4);
imshow(BW2);
title('Prewitt Method');

BW3=edge(J,'roberts');
subplot(2,4,5);
imshow(BW3);
title('Roberts Method');

BW4=edge(J,'canny');
subplot(2,4,6);
imshow(BW4);
title('Canny Method');

BW5=edge(J,'log');
subplot(2,4,7);

```

```
imshow(BW5);  
title('Laplacian of Gaussian');
```

حل تکلیف سری دو - سوال دوم - بند د

```
clc;  
close all;  
clear all;  
  
I=imread('circuit.tif');  
subplot(2,4,1);  
imshow(I);  
title('Original Image');  
  
J=imnoise(I,'gaussian',0,0.01);  
subplot(2,4,2);  
imshow(J);  
title('Gaussian Noise');  
  
BW1=edge(J,'sobel');  
subplot(2,4,3);  
imshow(BW1);  
title('Sobel Method');  
  
BW2=edge(J,'prewitt');  
subplot(2,4,4);  
imshow(BW2);  
title('Prewitt Method');  
  
BW3=edge(J,'roberts');  
subplot(2,4,5);  
imshow(BW3);  
title('Roberts Method');  
  
BW4=edge(J,'canny');  
subplot(2,4,6);  
imshow(BW4);  
title('Canny Method');  
  
BW5=edge(J,'log');  
subplot(2,4,7);  
imshow(BW5);  
title('Laplacian of Gaussian');
```