

# Exercise session on finite difference approximations of parabolic PDEs, part 3

November 26, 2021

---

In all exercises, compute the  $l^2$  norm using the formula

$$\|g\|_2 = \sqrt{\sum_{i=0}^{N_x} |g_i|^2 \Delta x}.$$

## Exercise 1

Consider the homogeneous heat equation

$$\frac{\partial c}{\partial t} = \nu \frac{\partial^2 c}{\partial x^2}$$

with periodic boundary conditions on the spatial domain  $[0, L]$  and on the time interval  $[0, T]$ . Assume  $L = 10$ ,  $T = 5$ ,  $\nu = 0.1$ . Assume as initial datum

$$c_0(x) = 10 \exp \left\{ - \left( \frac{x - L/2}{L/10} \right)^2 \right\}.$$

- Compute the exact solution by separation of variables on a uniform mesh of  $N = 100$  points.
- Compute on the same mesh a numerical solution by the implicit Euler method using  $M = 20$  time steps. Plot the absolute value of the difference between the numerical and the exact solution, and compute the  $l^2$  and  $l^\infty$  relative errors.
- Estimate empirically the convergence rate by repeating the computation using  $N = 200$  time steps and  $M = 40$  time steps.
- Compare the accuracy of the results with those of the explicit Euler method applied with a time step such that  $\nu \Delta t / \Delta x^2 = 1/5$ .

## Exercise 2

Repeat the previous exercise using the Crank-Nicolson method. Compare the accuracy of the results with those of the implicit Euler method.

### Exercise 3

Consider the non homogeneous, variable coefficients diffusion equation

$$\begin{cases} \frac{\partial c}{\partial t} = \frac{\partial}{\partial x} \left[ \frac{x^2}{100} \frac{\partial c}{\partial x} \right] + s(x, t) & x \in (0, 2), t \in (0, 2), \\ y(x, 0) = y_0(x) = x^3 & x \in (0, 2), \end{cases}$$

with  $s(x, t) = 2tx^3 - 3x^3(t^2 + 1)/25$ . Notice that the exact solution is given by  $c(x, t) = x^3(t^2 + 1)$ . Solve the equation on a mesh of  $N = 200$  nodes and  $M = 50$  time steps with the Crank Nicolson method, considering

- Dirichlet boundary conditions  $g_0(t) = 0, g_L(t) = 8(t^2 + 1)$ ;
- Neumann boundary conditions  $g_0(t) = 0, g_L(t) = 12(t^2 + 1)$ ;
- flux boundary conditions  $g_0(t) = 0, g_L(t) = 12(t^2 + 1)/25$ .

Plot the absolute value of the difference between the numerical and the exact solution, and compute the  $l^2$  and  $l^\infty$  relative errors.