Discussion of paper "Scalable Nonparametric Price Elasticity Estimation" by Wang and Huang

Discussion by Vineet Kumar (Yale School of Management)

March 2023 UTD FORMS

- Motivation: Obtain **price elasticities** from aggregate data.
- Our typical approach to demand models is parametric (e.g. logit)
- Want to do this using a nonparametric approach
 - Why?
- Use ML method, specifically bagged (bootstrap averaged) nearest neighbors
- Desiderata
 - Minimal assumptions nonparametric
 - No demand model required ever
 - Closed form for function, conditional on function values at nearest neighbors
 - Computationally light and scalable

- Motivation: Obtain **price elasticities** from aggregate data.
- Our typical approach to demand models is parametric (e.g. logit)
- Want to do this using a nonparametric approach
 - Why?
- Use ML method, specifically bagged (bootstrap averaged) nearest neighbors
- Desiderata
 - Minimal assumptions nonparametric
 - No demand model required ever
 - Closed form for function, conditional on function values at nearest neighbors.
 - Computationally light and scalable

- Motivation: Obtain **price elasticities** from aggregate data.
- Our typical approach to demand models is parametric (e.g. logit)
- Want to do this using a nonparametric approach
 - Why?
- Use ML method, specifically bagged (bootstrap averaged) nearest neighbors
- Desiderata
 - Minimal assumptions nonparametric
 - No demand model required ever
 - Closed form for function, conditional on function values at nearest neighbors
 - Computationally light and scalable

- Motivation: Obtain **price elasticities** from aggregate data.
- Our typical approach to demand models is parametric (e.g. logit)
- Want to do this using a nonparametric approach
 - Why?
- Use ML method, specifically bagged (bootstrap averaged) nearest neighbors
- Desiderata
 - Minimal assumptions nonparametric
 - No demand model required even
 - Closed form for function, conditional on function values at nearest neighbors
 - Computationally light and scalable

- Motivation: Obtain **price elasticities** from aggregate data.
- Our typical approach to demand models is parametric (e.g. logit)
- Want to do this using a nonparametric approach
 - Why?
- Use ML method, specifically bagged (bootstrap averaged) nearest neighbors
- Desiderata:
 - Minimal assumptions nonparametric
 - No demand model required even
 - Closed form for function, conditional on function values at nearest neighbors
 - Computationally light and scalable

- Motivation: Obtain **price elasticities** from aggregate data.
- Our typical approach to demand models is parametric (e.g. logit)
- Want to do this using a nonparametric approach
 - Why?
- Use ML method, specifically bagged (bootstrap averaged) nearest neighbors
- Desiderata:
 - Minimal assumptions nonparametric
 - No demand model required even
 - Closed form for function, conditional on function values at nearest neighbors
 - \implies Computationally light and scalable


- Motivation: Obtain **price elasticities** from aggregate data.
- Our typical approach to demand models is parametric (e.g. logit)
- Want to do this using a nonparametric approach
 - Why?
- Use ML method, specifically bagged (bootstrap averaged) nearest neighbors
- Desiderata:
 - Minimal assumptions nonparametric
 - No demand model required even
 - Closed form for function, conditional on function values at nearest neighbors
 - — Computationally light and scalable

- Motivation: Obtain **price elasticities** from aggregate data.
- Our typical approach to demand models is parametric (e.g. logit)
- Want to do this using a nonparametric approach
 - Why?
- Use ML method, specifically bagged (bootstrap averaged) nearest neighbors
- Desiderata:
 - Minimal assumptions nonparametric
 - No demand model required even
 - Closed form for function, conditional on function values at nearest neighbors
 - — Computationally light and scalable


- Motivation: Obtain **price elasticities** from aggregate data.
- Our typical approach to demand models is parametric (e.g. logit)
- Want to do this using a nonparametric approach
 - Why?
- Use ML method, specifically bagged (bootstrap averaged) nearest neighbors
- Desiderata:
 - Minimal assumptions nonparametric
 - No demand model required even
 - Closed form for function, conditional on function values at nearest neighbors
 - — Computationally light and scalable

- Motivation: Obtain **price elasticities** from aggregate data.
- Our typical approach to demand models is parametric (e.g. logit)
- Want to do this using a nonparametric approach
 - Why?
- Use ML method, specifically bagged (bootstrap averaged) nearest neighbors
- Desiderata:
 - Minimal assumptions nonparametric
 - No demand model required even
 - Closed form for function, conditional on function values at nearest neighbors
 - \implies Computationally light and scalable


- Consider simple case without endogeneity
- Model: $s_{jt} = f_j(p_{jt}, x_{jt}) + \varepsilon_{jt}$
- Elasticities: If you know function f, then we can easily obtain elasticities
 - Change price to $p + \Delta$, determine how sales change.
- How do you get f nonparametrically?
 - Use a ML method to obtain functional approximation
 - Universal function approximators for different classes of functions




- Consider simple case without endogeneity
- Model: $s_{jt} = f_j(p_{jt}, x_{jt}) + \varepsilon_{jt}$
- Elasticities: If you know function f, then we can easily obtain elasticities
 - Change price to $p + \Delta$, determine how sales change.
- How do you get f nonparametrically?
 - Use a ML method to obtain functional approximation
 - Universal function approximators for different classes of functions




- Consider simple case without endogeneity
- ullet Model: $s_{jt} = f_j(p_{jt}, x_{jt}) + arepsilon_{jt}$
- Elasticities: If you know function f, then we can easily obtain elasticities
 - Change price to $p + \Delta$, determine how sales change.
- How do you get f nonparametrically?
 - Use a ML method to obtain functional approximation
 - Universal function approximators for different classes of functions




- Consider simple case without endogeneity
- ullet Model: $s_{jt} = f_j(p_{jt}, x_{jt}) + arepsilon_{jt}$
- Elasticities: If you know function f, then we can easily obtain elasticities
 - Change price to $p + \Delta$, determine how sales change.
- How do you get f nonparametrically?
 - Use a ML method to obtain functional approximation
 - Universal function approximators for different classes of functions




- Consider simple case without endogeneity
- ullet Model: $s_{jt} = f_j(p_{jt}, x_{jt}) + arepsilon_{jt}$
- Elasticities: If you know function f, then we can easily obtain elasticities
 - Change price to $p + \Delta$, determine how sales change.
- How do you get f nonparametrically?
 - Use a ML method to obtain functional approximation
 - Universal function approximators for different classes of functions



- Consider simple case without endogeneity
- Model: $s_{jt} = f_j(p_{jt}, x_{jt}) + \varepsilon_{jt}$
- Elasticities: If you know function f, then we can easily obtain elasticities
 - Change price to $p + \Delta$, determine how sales change.
- How do you get f nonparametrically?
 - Use a ML method to obtain functional approximation
 - Universal function approximators for different classes of functions



- Consider simple case without endogeneity
- Model: $s_{jt} = f_j(p_{jt}, x_{jt}) + \varepsilon_{jt}$
- Elasticities: If you know function f, then we can easily obtain elasticities
 - Change price to $p + \Delta$, determine how sales change.
- How do you get f nonparametrically?
 - Use a ML method to obtain functional approximation
 - Universal function approximators for different classes of functions



- Control function approach is used to deal with endogeneity
- where z's are instruments
- Want to obtain price derivatives: $\partial_p f_j(p,x) = \partial_p \mathbf{E}[s_j|p,x] \partial_p \mathbf{E}[\varepsilon_{jt}|p,x]$
- Newey, Powell and Vella (1999) provide a way to write the selection bias to get $\partial_p \mathbf{E}[\varepsilon_{jt}|p,x] = (\partial_z g(z)'\partial_z g(z))^{-1} \partial_z g(z)'\partial_z \mathbf{E}[s_j|p,x]$
- All quantities can be easily obtained from data

- Control function approach is used to deal with endogeneity
- where z's are instruments
- Want to obtain price derivatives: $\partial_p f_j(p,x) = \partial_p \mathbf{E}[s_j|p,x] \partial_p \mathbf{E}[\varepsilon_{jt}|p,x]$
- Newey, Powell and Vella (1999) provide a way to write the selection bias to get $\partial_p \mathbf{E}[\varepsilon_{jt}|p,x] = (\partial_z g(z)'\partial_z g(z))^{-1} \partial_z g(z)'\partial_z \mathbf{E}[s_j|p,x]$
- All quantities can be easily obtained from data

- Control function approach is used to deal with endogeneity
- $\bullet \quad p = g(z) + u$
- where z's are instruments
- Want to obtain price derivatives: $\partial_p f_j(p,x) = \partial_p \mathbf{E}[s_j|p,x] \partial_p \mathbf{E}[\varepsilon_{jt}|p,x]$
- Newey, Powell and Vella (1999) provide a way to write the selection bias to get $\partial_p \mathbf{E}[\varepsilon_{jt}|p,x] = (\partial_z g(z)'\partial_z g(z))^{-1} \partial_z g(z)'\partial_z \mathbf{E}[s_j|p,x]$
- All quantities can be easily obtained from data

- Control function approach is used to deal with endogeneity
- $\bullet \quad p = g(z) + u$
- where z's are instruments
- Want to obtain price derivatives: $\partial_p f_j(p,x) = \partial_p \mathbf{E}[s_j|p,x] \partial_p \mathbf{E}[\varepsilon_{jt}|p,x]$
- Newey, Powell and Vella (1999) provide a way to write the selection bias to get $\partial_p \mathbf{E}[\varepsilon_{jt}|p,x] = (\partial_z g(z)'\partial_z g(z))^{-1} \partial_z g(z)'\partial_z \mathbf{E}[s_j|p,x]$
- All quantities can be easily obtained from data

- Control function approach is used to deal with endogeneity
- $\bullet \quad p = g(z) + u$
- where z's are instruments
- Want to obtain price derivatives: $\partial_{\rho} f_j(\rho, x) = \partial_{\rho} \mathbf{E}[s_j | \rho, x] \partial_{\rho} \mathbf{E}[\varepsilon_{jt} | \rho, x]$
- Newey, Powell and Vella (1999) provide a way to write the selection bias to get: $\partial_p \mathbf{E}[\varepsilon_{jt}|p,x] = (\partial_z g(z)'\partial_z g(z))^{-1} \partial_z g(z)'\partial_z \mathbf{E}[s_j|p,x]$
- All quantities can be easily obtained from data

- Control function approach is used to deal with endogeneity
- $\bullet \quad p = g(z) + u$
- where z's are instruments
- Want to obtain price derivatives: $\partial_p f_j(p,x) = \partial_p \mathbf{E}[s_j|p,x] \partial_p \mathbf{E}[\varepsilon_{jt}|p,x]$
- Newey, Powell and Vella (1999) provide a way to write the selection bias to get: $\partial_p \mathbf{E}[\varepsilon_{jt}|p,x] = (\partial_z g(z)'\partial_z g(z))^{-1} \partial_z g(z)'\partial_z \mathbf{E}[s_j|p,x]$
- All quantities can be easily obtained from data

- One view of ML is that it is functional approximation
- If we have $\partial_p f_i(p,x)$ we can get elasticities
- Want to approximate

$$\partial_{p} f_{j}(p, x) = \partial_{p} \mathbf{E}[s_{j}|p, x] - (\partial_{z} g(z)' \partial_{z} g(z))^{-1} \partial_{z} g(z)' \partial_{z} \mathbf{E}[s_{j}|p, x]$$

- Want a method that has "fast" asymptotic convergence (rate $\sqrt{m/T}$)
- Why the complexity:
 - Let's start with k-Nearest Neighbor (Bias-Variance Tradeoff)
 - Ensemble methods like Bagging can reduce variance (think of Random Forest)
 - ...But still biased
 - use a jackknife to correct the bias (Biau, Cerou, and Guyader, 2010)
 - corrects first order bias

- One view of ML is that it is functional approximation
- If we have $\partial_p f_i(p,x)$ we can get elasticities
- Want to approximate

$$\partial_{p} f_{j}(p,x) = \partial_{p} \mathbf{E}[s_{j}|p,x] - (\partial_{z} g(z)' \partial_{z} g(z))^{-1} \partial_{z} g(z)' \partial_{z} \mathbf{E}[s_{j}|p,x]$$

- Want a method that has "fast" asymptotic convergence (rate $\sqrt{m/T}$)
- Why the complexity:
 - Let's start with k-Nearest Neighbor (Bias-Variance Tradeoff)
 - Ensemble methods like Bagging can reduce variance (think of Random Forest)
 - ...But still biased
 - use a jackknife to correct the bias (Biau, Cerou, and Guyader, 2010)
 - corrects first order bias

- One view of ML is that it is functional approximation
- If we have $\partial_p f_i(p,x)$ we can get elasticities
- Want to approximate

$$\partial_{p}f_{j}(p,x) = \partial_{p}\mathbf{E}[s_{j}|p,x] - \left(\partial_{z}g(z)'\partial_{z}g(z)\right)^{-1}\partial_{z}g(z)'\partial_{z}\mathbf{E}[s_{j}|p,x]$$

- Want a method that has "fast" asymptotic convergence (rate $\sqrt{m/T}$)
- Why the complexity:
 - Let's start with k-Nearest Neighbor (Bias-Variance Tradeoff)
 - Ensemble methods like Bagging can reduce variance (think of Random Forest
 - ...But still biased
 - use a jackknife to correct the bias (Biau, Cerou, and Guyader, 2010)
 - corrects first order bias.

- One view of ML is that it is functional approximation
- If we have $\partial_p f_i(p,x)$ we can get elasticities
- Want to approximate

$$\partial_{p}f_{j}(p,x) = \partial_{p}\mathbf{E}[s_{j}|p,x] - \left(\partial_{z}g(z)'\partial_{z}g(z)\right)^{-1}\partial_{z}g(z)'\partial_{z}\mathbf{E}[s_{j}|p,x]$$

- Want a method that has "fast" asymptotic convergence (rate $\sqrt{m/T}$)
- Why the complexity:
 - Let's start with k-Nearest Neighbor (Bias-Variance Tradeoff)
 - Ensemble methods like Bagging can reduce variance (think of Random Forest)
 - ...But still biased
 - ullet use a jackknife to correct the bias (Biau, Cerou, and Guyader, 2010)
 - corrects first order bias.

- One view of ML is that it is functional approximation
- If we have $\partial_p f_i(p,x)$ we can get elasticities
- Want to approximate

$$\partial_{p} f_{j}(p, x) = \partial_{p} \mathbf{E}[s_{j}|p, x] - \left(\partial_{z} g(z)' \partial_{z} g(z)\right)^{-1} \partial_{z} g(z)' \partial_{z} \mathbf{E}[s_{j}|p, x]$$

- Want a method that has "fast" asymptotic convergence (rate $\sqrt{m/T}$)
- Why the complexity:
 - Let's start with k-Nearest Neighbor (Bias-Variance Tradeoff)
 - Ensemble methods like Bagging can reduce variance (think of Random Forest)
 - ...But still biased.
 - use a jackknife to correct the bias (Biau, Cerou, and Guyader, 2010)
 - corrects first order bias.

- One view of ML is that it is functional approximation
- If we have $\partial_p f_i(p,x)$ we can get elasticities
- Want to approximate

$$\partial_{p} f_{j}(p, x) = \partial_{p} \mathbf{E}[s_{j}|p, x] - \left(\partial_{z} g(z)' \partial_{z} g(z)\right)^{-1} \partial_{z} g(z)' \partial_{z} \mathbf{E}[s_{j}|p, x]$$

- Want a method that has "fast" asymptotic convergence (rate $\sqrt{m/T}$)
- Why the complexity:
 - Let's start with k-Nearest Neighbor (Bias-Variance Tradeoff)
 - Ensemble methods like Bagging can reduce variance (think of Random Forest)
 - ...But still biased.
 - use a jackknife to correct the bias (Biau, Cerou, and Guyader, 2010)
 - corrects first order bias.

- One view of ML is that it is functional approximation
- If we have $\partial_p f_j(p,x)$ we can get elasticities
- Want to approximate

$$\partial_{p} f_{j}(p, x) = \partial_{p} \mathbf{E}[s_{j}|p, x] - \left(\partial_{z} g(z)' \partial_{z} g(z)\right)^{-1} \partial_{z} g(z)' \partial_{z} \mathbf{E}[s_{j}|p, x]$$

- Want a method that has "fast" asymptotic convergence (rate $\sqrt{m/T}$)
- Why the complexity:
 - Let's start with k-Nearest Neighbor (Bias-Variance Tradeoff)
 - Ensemble methods like Bagging can reduce variance (think of Random Forest)
 - ...But still biased.
 - use a jackknife to correct the bias (Biau, Cerou, and Guyader, 2010)
 - corrects first order bias.

- One view of ML is that it is functional approximation
- If we have $\partial_p f_i(p,x)$ we can get elasticities
- Want to approximate

$$\partial_{p} f_{j}(p, x) = \partial_{p} \mathbf{E}[s_{j}|p, x] - \left(\partial_{z} g(z)' \partial_{z} g(z)\right)^{-1} \partial_{z} g(z)' \partial_{z} \mathbf{E}[s_{j}|p, x]$$

- Want a method that has "fast" asymptotic convergence (rate $\sqrt{m/T}$)
- Why the complexity:
 - Let's start with k-Nearest Neighbor (Bias-Variance Tradeoff)
 - Ensemble methods like Bagging can reduce variance (think of Random Forest)
 - ...But still biased.
 - use a jackknife to correct the bias (Biau, Cerou, and Guyader, 2010)
 - corrects first order bias.

- One view of ML is that it is functional approximation
- If we have $\partial_p f_i(p,x)$ we can get elasticities
- Want to approximate

$$\partial_{p} f_{j}(p, x) = \partial_{p} \mathbf{E}[s_{j}|p, x] - \left(\partial_{z} g(z)' \partial_{z} g(z)\right)^{-1} \partial_{z} g(z)' \partial_{z} \mathbf{E}[s_{j}|p, x]$$

- Want a method that has "fast" asymptotic convergence (rate $\sqrt{m/T}$)
- Why the complexity:
 - Let's start with k-Nearest Neighbor (Bias-Variance Tradeoff)
 - Ensemble methods like Bagging can reduce variance (think of Random Forest)
 - ...But still biased.
 - use a jackknife to correct the bias (Biau, Cerou, and Guyader, 2010)
 - corrects first order bias.

- One view of ML is that it is functional approximation
- If we have $\partial_p f_i(p,x)$ we can get elasticities
- Want to approximate

$$\partial_{p} f_{j}(p, x) = \partial_{p} \mathbf{E}[s_{j}|p, x] - \left(\partial_{z} g(z)' \partial_{z} g(z)\right)^{-1} \partial_{z} g(z)' \partial_{z} \mathbf{E}[s_{j}|p, x]$$

- Want a method that has "fast" asymptotic convergence (rate $\sqrt{m/T}$)
- Why the complexity:
 - Let's start with k-Nearest Neighbor (Bias-Variance Tradeoff)
 - Ensemble methods like Bagging can reduce variance (think of Random Forest)
 - ...But still biased.
 - use a jackknife to correct the bias (Biau, Cerou, and Guyader, 2010)
 - corrects first order bias.

Things to Like

- Paper is well written and motivated, which is not common in methodological papers
- Method is new and applicable to pretty much any demand setting.
- One of the biggest issues is endogeneity, which is accommodated here.
- Structural models can often take days, weeks or even months to estimate
- Anything that helps with obtaining realistic estimates of economic quantities like elasticity very useful
- Critical to have when real-time or timely decisions are required
- Scalable to lots of products, markets

Things to Like

- Paper is well written and motivated, which is not common in methodological papers
- Method is new and applicable to pretty much any demand setting.
- One of the biggest issues is endogeneity, which is accommodated here.
- Structural models can often take days, weeks or even months to estimate
- Anything that helps with obtaining realistic estimates of economic quantities like elasticity very useful
- Critical to have when real-time or timely decisions are required
- Scalable to lots of products, markets

- Paper is well written and motivated, which is not common in methodological papers
- Method is new and applicable to pretty much any demand setting.
- One of the biggest issues is endogeneity, which is accommodated here.
- Structural models can often take days, weeks or even months to estimate
- Anything that helps with obtaining realistic estimates of economic quantities like elasticity very useful
- Critical to have when **real-time** or timely decisions are required
- Scalable to lots of products, markets

- Paper is well written and motivated, which is not common in methodological papers
- Method is new and applicable to pretty much any demand setting.
- One of the biggest issues is endogeneity, which is accommodated here.
- Structural models can often take days, weeks or even months to estimate
- Anything that helps with obtaining realistic estimates of economic quantities like elasticity very useful
- Critical to have when **real-time** or timely decisions are required
- Scalable to lots of products, markets

- Paper is well written and motivated, which is not common in methodological papers
- Method is new and applicable to pretty much any demand setting.
- One of the biggest issues is endogeneity, which is accommodated here.
- Structural models can often take days, weeks or even months to estimate
- Anything that helps with obtaining realistic estimates of economic quantities like elasticity very useful
- Critical to have when **real-time** or timely decisions are required
- Scalable to lots of products, markets

- Paper is well written and motivated, which is not common in methodological papers
- Method is new and applicable to pretty much any demand setting.
- One of the biggest issues is endogeneity, which is accommodated here.
- Structural models can often take days, weeks or even months to estimate
- Anything that helps with obtaining realistic estimates of economic quantities like elasticity very useful
- Critical to have when **real-time** or timely decisions are required
- Scalable to lots of products, markets

- Paper is well written and motivated, which is not common in methodological papers
- Method is new and applicable to pretty much any demand setting.
- One of the biggest issues is endogeneity, which is accommodated here.
- Structural models can often take days, weeks or even months to estimate
- Anything that helps with obtaining realistic estimates of economic quantities like elasticity very useful
- Critical to have when real-time or timely decisions are required
- Scalable to lots of products, markets

- Nonparametric implies the model misspecification is not likely a huge issue
 - In parametric models, sometimes assumptions drive the result(rather than the data).
- Bagged Nearest Neighbor has attractive computational properties
- Not easy to obtain a method that has all the desirable properties
- Authors show monte carlos to show approximation quality
- Practical value in realistic applications...fast..

- Nonparametric implies the model misspecification is not likely a huge issue
 - In parametric models, sometimes assumptions drive the result(rather than the data).
- Bagged Nearest Neighbor has attractive computational properties
- Not easy to obtain a method that has all the desirable properties
- Authors show monte carlos to show approximation quality
- Practical value in realistic applications...fast..

- Nonparametric implies the model misspecification is not likely a huge issue
 - In parametric models, sometimes assumptions drive the result(rather than the data).
- Bagged Nearest Neighbor has attractive computational properties
- Not easy to obtain a method that has all the desirable properties
- Authors show monte carlos to show approximation quality
- Practical value in realistic applications...fast..

- Nonparametric implies the model misspecification is not likely a huge issue
 - In parametric models, sometimes assumptions drive the result(rather than the data).
- Bagged Nearest Neighbor has attractive computational properties
- Not easy to obtain a method that has all the desirable properties
- Authors show monte carlos to show approximation quality
- Practical value in realistic applications...fast..

- Nonparametric implies the model misspecification is not likely a huge issue
 - In parametric models, sometimes assumptions drive the result(rather than the data).
- Bagged Nearest Neighbor has attractive computational properties
- Not easy to obtain a method that has all the desirable properties
- Authors show monte carlos to show approximation quality
- Practical value in realistic applications...fast...

- Nonparametric implies the model misspecification is not likely a huge issue
 - In parametric models, sometimes assumptions drive the result(rather than the data).
- Bagged Nearest Neighbor has attractive computational properties
- Not easy to obtain a method that has all the desirable properties
- Authors show monte carlos to show approximation quality
- Practical value in realistic applications...fast..

- Let's start with profit-maximizing firm: $\pi(p) = f(x, p)(p c)$
- In equilibrium, after solving FOC, Lerner index $\frac{p-c}{p} = -\frac{1}{\varepsilon(p)}$
- So, can use this to recover marginal costs if we know $\varepsilon(p)$.

- Let's start with profit-maximizing firm: $\pi(p) = f(x,p)(p-c)$
- In equilibrium, after solving FOC, Lerner index $\frac{p-c}{p} = -\frac{1}{\varepsilon(p)}$
- So, can use this to recover marginal costs if we know $\varepsilon(p)$.

- Let's start with profit-maximizing firm: $\pi(p) = f(x,p)(p-c)$
- ullet In equilibrium, after solving FOC, Lerner index $\frac{p-c}{p}=-rac{1}{arepsilon(p)}$
- So, can use this to recover marginal costs if we know $\varepsilon(p)$.

- ullet Typically we start with a utility model $u_{ijt} = eta X_{jt} + lpha p_{jt} + \xi_{jt} + arepsilon_{ijt}$
- We recover parameters $\theta = (\alpha, \beta)$.
- Here the method cannot recover preference parameters
- However, the claim is we can still do counterfactuals. How?
- **Key Restriction**: Demand f(x,p) that is recovered through ML should not change in the counterfactual
 - Demand invariance
- What class of counterfactuals would work here?

- ullet Typically we start with a utility model $u_{ijt} = eta X_{jt} + lpha p_{jt} + \xi_{jt} + arepsilon_{ijt}$
- We recover parameters $\theta = (\alpha, \beta)$.
- Here the method cannot recover preference parameters
- However, the claim is we can still do counterfactuals. How?
- **Key Restriction**: Demand f(x,p) that is recovered through ML should not change in the counterfactual
 - Demand invariance
- What class of counterfactuals would work here?

- ullet Typically we start with a utility model $u_{ijt} = eta X_{jt} + lpha p_{jt} + \xi_{jt} + arepsilon_{ijt}$
- We recover parameters $\theta = (\alpha, \beta)$.
- Here the method cannot recover preference parameters
- However, the claim is we can still do counterfactuals. How?
- Key Restriction: Demand f(x,p) that is recovered through ML should not change in the counterfactual
 - Demand invariance
- What class of counterfactuals would work here?

- ullet Typically we start with a utility model $u_{ijt} = eta X_{jt} + lpha p_{jt} + \xi_{jt} + arepsilon_{ijt}$
- We recover parameters $\theta = (\alpha, \beta)$.
- Here the method cannot recover preference parameters
- However, the claim is we can still do counterfactuals. How?
- Key Restriction: Demand f(x,p) that is recovered through ML should not change in the counterfactual
 - Demand invariance
- What class of counterfactuals would work here?

- ullet Typically we start with a utility model $u_{ijt} = eta X_{jt} + lpha p_{jt} + \xi_{jt} + arepsilon_{ijt}$
- We recover parameters $\theta = (\alpha, \beta)$.
- Here the method cannot recover preference parameters
- However, the claim is we can still do counterfactuals. How?
- **Key Restriction**: Demand f(x,p) that is recovered through ML should not change in the counterfactual
 - Demand invariance
- What class of counterfactuals would work here?

- ullet Typically we start with a utility model $u_{ijt} = eta X_{jt} + lpha p_{jt} + \xi_{jt} + arepsilon_{ijt}$
- We recover parameters $\theta = (\alpha, \beta)$.
- Here the method cannot recover preference parameters
- However, the claim is we can still do counterfactuals. How?
- **Key Restriction**: Demand f(x,p) that is recovered through ML should not change in the counterfactual
 - Demand invariance
- What class of counterfactuals would work here?

- ullet Typically we start with a utility model $u_{ijt} = eta X_{jt} + lpha p_{jt} + \xi_{jt} + arepsilon_{ijt}$
- We recover parameters $\theta = (\alpha, \beta)$.
- Here the method cannot recover preference parameters
- However, the claim is we can still do counterfactuals. How?
- **Key Restriction**: Demand f(x,p) that is recovered through ML should not change in the counterfactual
 - Demand invariance
- What class of counterfactuals would work here?

- **Key Restriction**: Demand f(x,p) that is recovered through ML should not change in the counterfactual
 - Demand invariance
- What class of counterfactuals would work here?
 - An excise tax would be covered but a sales tax would not. Why?
- Excise tax \implies cost increases $c \rightarrow c + \delta$ but does not impact demand directly in any way.
- \Longrightarrow elasticities are not affected.
- However, what happens when we have a sales tax increase?
 - The elasticity recovered from the data $\varepsilon(p)$ is different from counterfactual $\widetilde{\varepsilon}(p)$
- What about something like Petrin (2002), minivan?

- **Key Restriction**: Demand f(x,p) that is recovered through ML should not change in the counterfactual
 - Demand invariance
- What class of counterfactuals would work here?
 - An excise tax would be covered but a sales tax would not. Why?
- Excise tax \implies cost increases $c \rightarrow c + \delta$ but does not impact demand directly in any way.
- \Longrightarrow elasticities are not affected.
- However, what happens when we have a sales tax increase?
 - The elasticity recovered from the data $\varepsilon(p)$ is different from counterfactual $\widetilde{\varepsilon}(p)$
- What about something like Petrin (2002), minivan?

- **Key Restriction**: Demand f(x,p) that is recovered through ML should not change in the counterfactual
 - Demand invariance
- What class of counterfactuals would work here?
 - An excise tax would be covered but a sales tax would not. Why?
- ullet Excise tax \Longrightarrow cost increases $c o c + \delta$ but does not impact demand directly in any way.
- => elasticities are not affected.
- However, what happens when we have a sales tax increase?
 - The elasticity recovered from the data $\varepsilon(p)$ is different from counterfactual $\widetilde{\varepsilon}(p)$
- What about something like Petrin (2002), minivan?

- **Key Restriction**: Demand f(x,p) that is recovered through ML should not change in the counterfactual
 - Demand invariance
- What class of counterfactuals would work here?
 - An excise tax would be covered but a sales tax would not. Why?
- ullet Excise tax \Longrightarrow cost increases $c o c + \delta$ but does not impact demand directly in any way.
- \Longrightarrow elasticities are not affected.
- However, what happens when we have a sales tax increase?
 - The elasticity recovered from the data $\varepsilon(p)$ is different from counterfactual $\widetilde{\varepsilon}(p)$
- What about something like Petrin (2002), minivan?

- **Key Restriction**: Demand f(x,p) that is recovered through ML should not change in the counterfactual
 - Demand invariance
- What class of counterfactuals would work here?
 - An excise tax would be covered but a sales tax would not. Why?
- Excise tax \implies cost increases $c \rightarrow c + \delta$ but does not impact demand directly in any way.
- \Longrightarrow elasticities are not affected.
- However, what happens when we have a sales tax increase?
 - The elasticity recovered from the data $\varepsilon(p)$ is different from counterfactual $\widetilde{\varepsilon}(p)$
- What about something like Petrin (2002), minivan?

- **Key Restriction**: Demand f(x,p) that is recovered through ML should not change in the counterfactual
 - Demand invariance
- What class of counterfactuals would work here?
 - An excise tax would be covered but a sales tax would not. Why?
- Excise tax \implies cost increases $c \rightarrow c + \delta$ but does not impact demand directly in any way.
- ⇒ elasticities are not affected.
- However, what happens when we have a sales tax increase?
 - ullet The elasticity recovered from the data arepsilon(
 ho) is different from counterfactual $ar{arepsilon}(
 ho)$
- What about something like Petrin (2002), minivan?

- **Key Restriction**: Demand f(x,p) that is recovered through ML should not change in the counterfactual
 - Demand invariance
- What class of counterfactuals would work here?
 - An excise tax would be covered but a sales tax would not. Why?
- ullet Excise tax \Longrightarrow cost increases $c \to c + \delta$ but does not impact demand directly in any way.
- \implies elasticities are not affected.
- However, what happens when we have a sales tax increase?
 - ullet The elasticity recovered from the data arepsilon(p) is different from counterfactual $\widetilde{arepsilon}(p)$
- What about something like Petrin (2002), minivan?

- **Key Restriction**: Demand f(x,p) that is recovered through ML should not change in the counterfactual
 - Demand invariance
- What class of counterfactuals would work here?
 - An excise tax would be covered but a sales tax would not. Why?
- ullet Excise tax \Longrightarrow cost increases $c \to c + \delta$ but does not impact demand directly in any way.
- \implies elasticities are not affected.
- However, what happens when we have a sales tax increase?
 - ullet The elasticity recovered from the data arepsilon(p) is different from counterfactual $\widetilde{arepsilon}(p)$
- What about something like Petrin (2002), minivan?

- **Key Restriction**: Demand f(x,p) that is recovered through ML should not change in the counterfactual
 - Demand invariance
- What class of counterfactuals would work here?
 - An excise tax would be covered but a sales tax would not. Why?
- ullet Excise tax \Longrightarrow cost increases $c \to c + \delta$ but does not impact demand directly in any way.
- \implies elasticities are not affected.
- However, what happens when we have a sales tax increase?
 - The elasticity recovered from the data $\varepsilon(p)$ is different from counterfactual $\widetilde{\varepsilon}(p)$
- What about something like Petrin (2002), minivan?

- How are the hyperparameters (tuning parameters) of the ML algorithm set? No theoretical guidelines for this.
- 2 What about the g function (recall p = g(z) + u). This needs to be parametrically specified.
- ② Can we use any other ML method in place of Bagged nearest neighbors?
 Random Forest, Deep nets etc.
- Oboils down to what you need from a computational perspective as well as a

- How are the hyperparameters (tuning parameters) of the ML algorithm set? No theoretical guidelines for this.
- ② What about the g function (recall p = g(z) + u). This needs to be parametrically specified.
- 3 Can we use any other ML method in place of Bagged nearest neighbors?
 Random Forest, Deep nets etc.
- Oboils down to what you need from a computational perspective as well as a

- How are the hyperparameters (tuning parameters) of the ML algorithm set? No theoretical guidelines for this.
- ② What about the g function (recall p = g(z) + u). This needs to be parametrically specified.
- Can we use any other ML method in place of Bagged nearest neighbors?
 - Random Forest, Deep nets etc.
- Boils down to what you need from a computational perspective as well as a

- How are the hyperparameters (tuning parameters) of the ML algorithm set? No theoretical guidelines for this.
- ② What about the g function (recall p = g(z) + u). This needs to be parametrically specified.
- Oan we use any other ML method in place of Bagged nearest neighbors?
 - Random Forest, Deep nets etc.
- Oboils down to what you need from a computational perspective as well as a

- How are the hyperparameters (tuning parameters) of the ML algorithm set? No theoretical guidelines for this.
- ② What about the g function (recall p = g(z) + u). This needs to be parametrically specified.
- On we use any other ML method in place of Bagged nearest neighbors?
 - Random Forest, Deep nets etc.
- Oboils down to what you need from a computational perspective as well as a

Conclusion

- Very interesting and practically valuable contribution to estimate elasticities in the presence of price endogeneity
- 2 Authors show that method can satisfy a number of desirable properties both in theory and in practice
- Need to have invariance of demand curve in the counterfactual

Conclusion

- Very interesting and practically valuable contribution to estimate elasticities in the presence of price endogeneity
- Authors show that method can satisfy a number of desirable properties both in theory and in practice
- Need to have invariance of demand curve in the counterfactual

Conclusion

- Very interesting and practically valuable contribution to estimate elasticities in the presence of price endogeneity
- Authors show that method can satisfy a number of desirable properties both in theory and in practice
- Need to have invariance of demand curve in the counterfactual