Campi Flegrei unrest: provisional observational reproduction and record

Independent reproduction of Tan et al. (2025)

13 October 2026

Main paper

Abstract

Displacement observations, labeled historical source estimates and an external routine earthquake catalog reconstruct the historical context of unrest at Campi Flegrei. Daily GNSS observations now extend through 20 March 2025. Independent waveform processing produces 1015 provisional GaMMA associations from 132733 picks, but has not established the publication's relocated fault geometry, magnitude distribution or focal mechanisms. No publication panel is fully independently reproduced. Partial counterparts: 1A, 1B, 1C, S1, S2, S4A, S5/map, S5/northing-depth, S5/easting-depth, S8A, S8B, S15C. All remaining panels are not reproduced. The scientific narrative and reproduced supplement precede the input, implementation, validation and completion records.

Campi Flegrei's unrest is recorded by ground displacement and seismicity. Tan et al. (2025) use a machine-learning-assisted earthquake catalog to resolve active structures and evaluate possible sources of unrest. Their study is observational: its central results describe the location, timing and character of earthquakes, rather than solutions of a coupled physical forward model. Here, the completed independent calculations are placed in the same scientific sequence. Published images are confined to the separate comparison PDF.

The current phase of unrest

Figure 1 reconstructs the historical context from the explicitly permitted routine catalog and displacement observations and explicitly labeled historical source estimates. The routine snapshot contains 16,046 located events from 2005 through 20 March 2025, including five events with duration magnitude at least 4.0. Monthly counts include all routine events; the cumulative series includes 7,218 events with $M_d > 0.2$. These external records are separate from the independently detected catalog.

The displacement reconstruction contains 128 historical entries (114 leveling surveys, 13 source-provided estimates and one source-provided reconstruction) and 8,743 daily GNSS observations through 20 March 2025. The original 1905–2023 interval is extended with observed data. A common vertical datum is obtained from overlapping observation records. The long sequence resolves subsidence, the major twentieth-century uplift episodes, and the recent uplift. It does not by itself distinguish magmatic from hydrothermal forcing. All three panels remain partial because the historical catalog snapshot, basemap and exact displacement-stitching convention differ or are incompletely specified.

Panel 1A–C: partially reproduced




Figure 1: Seismicity and ground displacement. Partial contextual reconstruction. A: external routine catalog, 16,046 located events and five Md>=4 stars. B: 114 leveling surveys (circles), 13 source-provided estimates (triangles), one source-provided reconstruction (cross), and daily GNSS through 20 March 2025; Historical estimates are not raw observations or reconstructions made here. C: all routine monthly counts and cumulative Md>0.2; March includes days 1–20 only. Routine events are not independent detections. Datum joining is an assumption; basemap and live snapshot differ from the publication. Depth colors: fixed 0âĂŞ5 km, red to blue; magenta >5 km, no data clipping. S5 limits/ticks are printed; 1A endpoints are unlabeled and 0âĂŞ5 km is an explicit approximation. Circle area law is an explicit display assumption, not recovered exactly. Missing magnitudes are crosses. Routine events are external context.

Improving the earthquake catalog

The expanded 1015-association checkpoint supplies 916 causally usable events and 38 calibration candidates. The explicitly authorized Tan-distributed reference grid now supplies independently averaged starting profiles. Four-layer fits converge locally, and 216 events pass the paper's numerical filters after five catalog passes. Full 32/64-bit inversion paths still differ by 0.89 km in depth; their locations remain exploratory. Both fixed-grid rounds executed; validation pending. No validated multiyear structural or mechanism result is claimed.

Figure 2 — not reproduced. Panels A and B need a validated multiyear relocated catalog and independent magnitudes. The expanded 1,015 associations now enter grid-derived calibration and two TomoDD trials, but path sensitivity and sparse coverage prevent acceptance.

A clearer view of the current phase of unrest

Figures 3 and 4 address the publication's main structural observations: the distribution of seismicity and the source mechanisms of larger events. The present independent work has no scientifically val-

idated calibrated and relocated catalog, and no independent focal-mechanism solutions. It therefore cannot determine the depth range, continuity or temporal activation of the ring fault, resolve the central fault system, or establish its slip orientations. These unreproduced questions are recorded as text; no placeholder or diagnostic plot is presented as a paper result.

Figure 3 — not reproduced. The map and both cross sections require independently calibrated and double-difference-relocated events. Provisional diagnostic locations are not substitutes.

Figure 4 — not reproduced. Panels A and B require independent SKHASH/gCAP solutions and reliable locations. No focal inversion was executed.

Implications for the physical process driving the unrest

The reconstructed hybrid-template example in Figure S15C is an independently processed waveform observation. The limited template search produced 822 unconfirmed threshold crossings; it does not establish recurring hybrid families or their evolution. Neither this result nor the initial associations can test the publication's inference about the presence or absence of upward seismic migration. The partial displacement history also does not distinguish magmatic and hydrothermal forcing. Such an interpretation would require the unresolved spatial and source constraints above.

New perspectives for improved seismic risk assessment

Fault geometry and a calibrated magnitude distribution are needed to connect this analysis to seismic risk. The pilot magnitude measurements have not established the multiyear magnitude-frequency distribution required for Figure S8. The current results therefore provide no revised estimate of event probabilities, damaging-event locations or source dimensions. Their supported contribution is restricted to the partial observational reconstructions shown here and in the supplement.

Reproduced supplement

This supplement follows the original Materials and Methods topic sequence, then Figures S1–S15 and Tables S1–S2. It reports only outcomes supported by the saved independent calculations. Detailed processing settings, input inventories, assumptions and validation are collected after this supplement.

Materials and Methods

S1 Building the Campi Flegrei high-resolution earthquake catalog

Data. The plotted station intervals describe the expanded processed checkpoint; the current grid-derived calibration uses that expanded set. Earlier magnitude and diagnostic products retain their separate event identifiers. The station inventory and intervals provide partial counterparts to Figures S1 and S2. They do not establish complete network coverage over the study period.

Phase picking. Inference with released trained weights produced 126133 merged picks (18545 P and 107588 S) in the processed observations. Figure S4A now has a partial inference comparison on the station and swarm specified in its caption. The original and retrained models return 5 P/5 S and 6 P/10 S picks respectively in the selected 120 s interval. These counts do not establish accuracy without reference labels. The exact published window is unverified. Training examples S3 and the exact S4B noise example remain unreproduced.

Phase association. GaMMA associated 7268 picks into 1003 provisional events using the explicitly labeled published alternative starting profile. These are candidates, not a validated multiyear catalog.

Initial earthquake location. The newly authorized-grid VELEST trial uses the expanded independent associations. Local convergence is achieved after documented layer merging, but full-path precision and regional-start dependence remain consequential; no location set is accepted. Numerical results and validation are reported after the supplement. Figure S5 is the separate external routine-catalog reconstruction, with 12,503 located study-period events.

Earthquake relocation. Original waveforms supply new coherency-weighted differential times for the grid-derived filtered trial. The current fixed-grid TomoDD execution and validation record follows the supplement. No validated relocated catalog or specified regional SVD analysis exists, so Figures S7 and S9–S13 remain unreproduced.

Local magnitude. The pilot measured 61681 channel amplitudes; 4705 passed the 5 dB SNR threshold. It formed 1163 relative-amplitude equations and returned 119 provisional magnitude estimates. Figure S8 displays the partial pilot comparison and distribution. This does not establish the final relocated catalog's magnitude distribution or completeness.

Focal mechanism estimation. SKHASH and gCAP inversions were not run. No independent mechanism is available for Figure 4.

S2 Validation of earthquake locations

Ring-shaped structures with confined depth distributions. The provisional associations do not establish a ring structure. Internal travel-time consistency is preserved in the documentation, but does not substitute for the regional uncertainty or common-receiver tests in Figures S7 and S12.

Depth of earthquakes in the central region. There are now 137 associated CPOZ P arrivals in the provisional sample. P-aligned CPOZ waveform gathers have been saved as a reproduction diagnostic. The relocated sector selections and manual checks on apparently deep central events remain unperformed. No conclusion about a lower depth limit follows from their absence.

S3 Template Matching to Detect Hybrid Events

The search covered 75.797 hours of available common CAAM/COLB intervals and returned 822 unconfirmed threshold crossings. It did not produce independently confirmed families for Figure S14.

Depth. No independent hybrid-event location set is available for Table S2.

Frequency index. Frequency indices were calculated for the retained three templates and the earlier four candidate windows. The expanded scan now has saved eight-second CAAM frequency-index and signal/noise measurements. These use template alignment, not reviewed onsets; coda discrimination and family confirmation remain incomplete. Figure S15C displays the corresponding third written-method template. The first two templates are different examples from the published S15A/B panels and are not inserted in their place.

Relative magnitude. Cluster-relative magnitudes were not assigned because cluster membership has not been established.

Temporal evolution of the hybrid events. The limited exposure and unconfirmed detections do not support a three-year cluster history.

Further search for HB/LP events. The provisional catalog now has station-averaged frequency-index screening on retrieved vertical channels. Selected observed windows for every negative-mean candidate in this checkpoint were visually inspected. They include persistent oscillations, impulsive packets and apparent contamination by earlier-event coda; no hybrid family was confirmed. This limited review does not supply validated onset labels. Missing observations still prevent a complete multiyear search.

Supplementary figures




Figure S1: Recorded station availability. Partial: expanded checkpoint, 868 processed files, 26 station names, 8511.520 union channel-hours. Empty intervals are neither connected nor counted. Current grid-derived calibration uses this expanded checkpoint. Earlier diagnostics retain separate identifiers. Vertical ticks show observed-interval midpoints where durations are too short to resolve; ticks do not imply continuous coverage.

Panel S2: partially reproduced





Figure S2: Station locations. Partial station inventory: 51 released identifiers audited against official study-period response epochs. Filled triangles: 42 official coordinate matches; open triangles: 9 released coordinates with unresolved epochs or channel codes. Two coincident-coordinate pairs remain distinct identifiers. No verified geographic outlier. Local WGS84 projection and clipped coastline; the original 48-instrument membership is not established. Colors distinguish broadband (red), short period (yellow) and accelerometer (white) instruments; no independent relief raster is available.

Figure S3 — not reproduced. Panels A–D require the specified training traces, annotations and augmentation examples. These were not reconstructed; downstream use of trained weights is not re-training.



S4B: exact CMIS noise-window timing and reference labels are unavailable. Other calendar-window diagnostics are not inserted as that panel.

Figure S4: PhaseNet benchmark examples. Partial S4A inference comparison on original CSOB observations, 20 May 2024 20:10:03–20:12:03 UTC. The caption's two-hour offset is interpreted literally; exact published alignment is unverified. Original weights: 5 P and 5 S; retrained weights: 6 P and 10 S. No labeled accuracy or retraining claim.

Panel S5, all views: partially reproduced




Figure S5: Routine-catalog locations. External routine-catalog data, independently selected and plotted: 12,503 located events, 21 January 2022–20 March 2025. This is 420 more than the published historical snapshot. No author-derived catalog is used to adjust that difference; these are not independent detections. Depth colors: fixed 0âĂŞ5 km, red to blue; magenta >5 km, no data clipping. S5 limits/ticks are printed; 1A endpoints are unlabeled and 0âĂŞ5 km is an explicit approximation. Circle area law is an explicit display assumption, not recovered exactly. Missing magnitudes are crosses. Routine events are external context.

Figure S6 — not reproduced. No validated independent counterpart. Authorized-grid VELEST now runs on the expanded cohort but remains path-sensitive; both fixed-grid rounds executed; validation pending. Diagnostics are retained after the supplement. No rejected locations are presented as reproduced structure.

Figure S7 — not reproduced. No validated independent counterpart. Authorized-grid VELEST now runs on the expanded cohort but remains path-sensitive; both fixed-grid rounds executed; validation pending. Diagnostics are retained after the supplement. No rejected locations are presented as reproduced structure.




Figure S8: Magnitude calibration and frequency distribution. Partial amplitude-ratio pilot: 119 provisional magnitudes from independent observations, with routine Md as the documented calibration prior. A is not an independent validation of that prior. B is a limited-sample histogram with a conditional likelihood estimate, not the full-period completeness or b-value result. Locations are initial associations, before final relocation.

Figure S9 — not reproduced. All three views require independently relocated events and cross-correlation observation counts for the published quality filter. These outputs are absent.


Figure S10 — not reproduced. The four annual groups for 2022–2025 require the independent multiyear relocated catalog. The available sample does not support them.

Figure S11 — not reproduced. The four swarm maps require the independent relocated catalog and justified swarm windows. These selections and locations remain incomplete.

Figure S12 — not reproduced. Panels A–H need relocated sector selections and manual depth review. 137 independent CPOZ P arrivals now exist, but the required sector gathers remain unfinished.

Figure S13 — not reproduced. Panels A and B require reliable independent relocated coordinates and an appropriately sourced basemap. The relocated catalog does not exist.

Figure S14 — not reproduced. The cluster histories require three-year exposure, confirmed events and relative magnitudes. The present 75.797 hour scan produced 822 unconfirmed threshold crossings, not reviewed families.



Panels A and B use different published events from the saved E1/E2 templates; D has no independently selected matching example. Those panels remain not reproduced.

Figure S15: Hybrid and shallow-event waveforms. Panel C is partial: independently processed E3 CAAM/COLB observations. Onset alignment, display filtering and spectral settings differ or are incompletely specified. A, B and D have placeholders rather than substitute waveforms.

Supplementary tables

Table S1: fitted one-dimensional velocity model — not reproduced. No independently validated regional VELEST model is accepted. Pilot fits and failed attempts, when present, are recorded after the supplement; the author's fitted velocities are not substituted.

Table S2: hybrid events in the relocated catalog — not reproduced. No independently relocated hybrid-event sample exists. The author's event parameters are not substituted.

Supplementary data products

Independent counterparts of the final catalog (Data S1), associated phases of that final catalog (Data S2), focal mechanisms (Data S3), and the relocated interactive view (Interactive Plot S1) are not complete. The saved provisional products retain their existing labels. References and the complete provenance record follow in the reproduction documentation.

Reproduction documentation

This record describes the current continuation, not the superseded nine-event run. Historical outputs remain recoverable from Git. The main paper and reproduced supplement contain only corresponding publication panels; published images occur only in the separate comparison PDF.

D1 Scientific specification and input verification

The local main paper and supplement were read before changing scientific outputs. Main captions 1–4, supplementary captions S1–S15 and the methods are readable. The two uploaded ZIP archives were listed and CRC-tested without executing their contents. Their exact filenames, sizes and checksums remain in the input inventory.

File	Bytes	Inspection
`tanmodel.pdf`	2140554	Text and captions readable
`tanmodelsupp.pdf`	62164947	Text and captions readable
`science.adw9038_data_s1_to_s3.zip`	12960255	Directory readable; CRC passed; not executed
`science.adw9038_interactive_plot_s1.zip`	8852485	Directory readable; CRC passed; not executed

Data S1 is the authors' derived catalog, S2 their associated phases, and S3 their focal mechanisms. The second archive contains an interactive derived-result visualization. None supplies original waveform observations for the independent workflow. These products are excluded from initialization, filtering, calibration, tuning and plot filling. The current workflow does not read these derived tables or compute result-table comparison statistics. Earlier comparison products are quarantined. Published reference images are used only in the separate comparison PDF.

The user now expressly permits the external routine catalog for Figures 1A, 1C and S5, and for the paper's documented duration-magnitude regularization. Those roles are implemented separately from picking and association. Released trained PhaseNet weights, generic method software, original observations and required metadata are permitted. The Tan-fitted 1D model and station corrections, event results, picks and study-specific scripts remain excluded. No figure was digitized or used as numerical input.

D2 Data access and actual coverage

The access refresh found readable miniSEED examples for seven station codes in the temporary archive: CFB1, CFB3, CROS, CRUS, V0102, V0105 and CMDP. CRC, decompression, sample headers and hashes were checked using bounded HTTP ranges; nothing was executed. The saved directory contains 319,889 miniSEED members but no CGAU, CLAC, CREM or CSOC member, although these four names appear on the index page. The examples are access/format checks, not detections or continuous exposure. Their author-event-selected windows and unresolved station epochs prevent treating this archive as an unbiased discovery record. Exact receipts are in `docs/local_waveform_receipts.json`.

Quantity	Value
GNSS observations in plotted interval	8743
GNSS extension in 2025 (days)	79

Quantity	Value
GNSS overlap RMS after offset (m)	0.000295
All historical routine records	26425
Located historical routine records	16046
Located routine study events	12503
Difference from published study count	420
Processed waveform files	825
Processed union channel-hours	7807.737
Merged picks	126133
Provisional GaMMA events	1003
Candidates on 19–20 March 2025	816
Successful segmented waveform requests	651
Pending waveform requests	73994

The last two study days account for 816 of 1003 provisional events (81.4 percent). This temporal concentration prevents inferring a multiyear seismicity history from this checkpoint. The study interval is 21 January 2022 through 20 March 2025, with the exclusive endpoint 21 March. The previous full-span HTTP 413 responses were request-size failures, not evidence of unavailable observations. The current collector queries FDSN availability and requests six-hour intervals plus half-hour calendar checks with 30-second processing margins. The smaller checks sample the full study date range without implying continuous coverage. Every successful miniSEED response is read, identified by network/station/channel, clipped to its core and hashed. Overlapping channel intervals are merged before computing channel-hours. The original full-period queue contains 74,445 six-hour requests; 229 half-hour calendar checks were added across quarter starts and the study boundaries; unattempted and failed requests never count as coverage. The portable checkpoint and per-request receipts distinguish completed downloads from the pending multiyear workload. At this checkpoint the full period has not been retrieved or processed. The raw inventory includes six original recordings selected for the written-method hybrid templates. Their waveform/spectral analysis is separate; their picks never seed the independently detected catalog. Thus raw observation coverage is not identical to inference exposure.

The collector resumes from `inputs/continuation/acquisition.sqlite`, uses six concurrent transfers and stops if shared free space falls below 20 GiB. Linear extrapolation from this download batch gives roughly 970 GB for the entire queue, against about 152 GiB free when checked. That estimate varies with compression and coverage and is not a measured full-archive size. Streaming processing with separate retention or additional storage is needed for the whole queue; pending work is labeled unfinished acquisition, not remote inaccessibility. Processing hours refer to successfully read input channels. Short fragments and missing components can prevent PhaseNet output, and these hours are not claimed as complete three-component detection exposure. The phase-classification warnings are retained in execution logs.

Fresh response-level StationXML was obtained for 22 IV stations and four 2I stations, including historical epochs. Public IT response queries returned HTTP 204. Eight additional IV identifiers missing from the XML—V0106, CDOV, CROS, CASE, V0102, V0105, V0104 and CMDP—were checked individually for full-period availability; all returned HTTP 204. A subsequent all-network StationXML query for these eight codes also returned HTTP 204. These are service-specific no-data responses, not proof that local or author-held records do not exist. RAN's login page explicitly requires registration or credentials to download SAC/ASCII waveforms; those credentials are unavailable. This is an access limitation, not a prompt restriction. Public RAN metadata were subsequently obtained directly from its station directory and XML download: all 17 candidate station files are readable, with 57 study-overlapping HN/HG response epochs. Unknown orientations on ancillary housekeeping channels are

not substituted for acceleration-channel orientations. The station directory closes POZA on 15 January 2025, leaving 16 of these stations installed at the study endpoint; this does not establish which instruments entered the paper. Waveform access remains credential-gated. Exact receipts and the archive listing are in `docs/ran_metadata_access.json` and `docs/ran_metadata_members.json`. The previously accessed local/temporary archive supplies event-selected windows; those windows do not establish continuous exposure and are preserved as such. Exact actual source files are listed in the frozen input and processing manifests. Per-channel response-epoch intersections and every unobserved interval are saved separately; a usable response requires instrument sensitivity and response stages. This verifies metadata availability, not laboratory calibration accuracy. Channel/epoch coordinates differ slightly for some station identifiers; their spread is saved in `coordinate_variability.csv`. Association uses one static station coordinate per identifier, an additional approximation that matters for precise relocation.

The original RITE daily file and format note from Zenodo record 17132077 are readable. They contain MJD, Up displacement and uncertainty in meters, with 8,664 observations through 31 December 2024. No JPG was downloaded as an analysis input. A second INGV release, Zenodo 22259337, published 2 September 2026, contains seven-column observations and format notes through 2025. Its RITE file has 9,025 observations, ending 31 December 2025. We use its 79 daily values through 20 March 2025 to complete the study-period extension. Published MD5 checksums and local SHA-256 hashes were verified. Earlier searches of the Nevada service returned HTTP 404; the public RING RITE position file was readable but ended in 2016 despite its 2025 release timestamp. Neither failure is now treated as a remaining 2025-data blocker.

Routine HTML pages for every year 2005–2025 were retrieved on 12 October 2026. The selected historical interval contains 26,425 records, including unlocated events, and 16,046 located events. The study interval contains 19,274 routine records and 12,503 with latitude, longitude and depth. All five $M_d \geq 4$ events in the Figure 1 description are present. The 420-event excess over the paper's 12,083 located study events is unresolved: archive revisions and undocumented snapshot/selection differences may contribute, but their separate effects cannot be measured without the original historical snapshot. Counts were not tuned. A compressed source snapshot preserves the exact retrieved HTML because live pages can change.

One downloaded COLB file requested for 1 January 2023 00:00–00:30 UTC contains only 0.32–1.68 seconds per component inside that core. It yields no finite picker predictions. The observation headers and hash are recorded in `data/resumption/no_prediction_file_audit.json`; neither the requested duration nor absent model scores are counted as exposure.

Station geometry and historical source classes

All 51 released station identifiers were compared with official response epochs. Forty-two identifiers have matching epochs; their coordinates replace the rounded released values, with a maximum horizontal correction of 138.8 m. Nine remain unresolved and retain open symbols: eight IV identifiers and NAP, whose official channels are HG rather than the released HN code. The source coordinates are local: the old map scale was enlarged by distant vertices in whole Natural Earth coastline geometries. Those vertices are now masked before projection. Forward/inverse coordinate transforms are checked in kilometers, and the corrected local map retains all candidate identifiers. The released inventory has 25 HH, 9 EH and 17 HN identifiers; the written methods specify 23 broadband, 9 short-period and 16 accelerometer instruments. Two HH pairs share released coordinates, but no alias or deployment record establishes that they represent the same instruments, and the additional HN identifier is unresolved. Thus 51 identifiers cannot yet be reconciled with the reported 48 instruments. The complete QA and epoch tables are `data/continuation/station_geometry_qa.csv` and `data/continuation/station_geometry_epochs.csv`.

The exploratory association uses one StationXML station coordinate per network/station/location identifier, retaining the last serialized record. It does not move the station coordinate between epochs. Channel metadata differ horizontally by up to 16.906 m at CBAC and 12.990 m at CAWE; the other inspected identifiers have no such variation. All inspected channel burial-depth fields are zero; elevation supplies the vertical coordinate. These are source-metadata conventions, not an independent survey. Epoch-specific coordinate sensitivity has not been measured and remains a location-method limitation.

The 128 historical entries comprise 114 leveling surveys, 13 source-provided estimates and one source-provided graphical reconstruction. Figure 1B distinguishes them with separate symbols. These estimates were transcribed from the source's numerical table, not extracted from figure pixels; they were not reconstructed by this implementation. Month-only dates are displayed on the fifteenth, with uncertain months retained. The source uncertainty strings and ambiguous original units are preserved without conversion. Daily GNSS observations already cover 2020–2023 and extend through the study endpoint; remaining calendar gaps are unfilled.

D3 Independent processing and explicit departures

Displacement and routine-catalog context

The original Del Gaudio et al. table supplies 128 leveling entries, March 1905–July 2009. Display dates use the middle of the stated month; uncertain months remain flagged. The original March 1905 elevation is 3.464 m. Subtracting it yields a displacement range of −1.266 to 2.541 m; adding the same observed datum back supplies the ordinate used in the new figure. This datum comes from the numerical source table, not from published image pixels. Source notes identify entries estimated historically from photographs or earlier graphical reconstructions; those notes remain in the saved table. This project digitizes no graph.

We align daily GNSS observations to leveling using the median of 23 overlapping epoch offsets, each based on GNSS within 15 days of the leveling date. The offset is 1.6115 m relative to the 1905 displacement zero, with a 0.0025 m median absolute deviation. The 2025 release is aligned using all 8,664 overlapping daily values; its offset is 1.78951 m and overlap RMS after the shift is 0.000295 m. The plotted data contain 319 missing calendar days, which are not filled. Daily continuity is broken across gaps longer than seven days. The present 2025 endpoint is 3.0504 m above the 1905 datum. This assumes comparable displacement at the leveling benchmark and GNSS site; the paper's exact joining recipe is not specified.

Routine selections follow the written Figure 1 and S5 captions. Figure 1A uses located events from 1 January 2005 through 20 March 2025; stars mark $M_d \geq 4$. Figure 1C uses all routine events for monthly counts and strictly $M_d > 0.2$ for the cumulative count. The caption specifies this cutoff only for the cumulative curve. Retaining unlocated records in the monthly bins is therefore an explicit selection assumption; the paper does not specify their treatment, and no filter was chosen to match the published bars. The cumulative curve uses actual event times, avoiding a one-month bin-position shift. March 2025 includes only days 1–20. Figure S5 uses located events from 21 January 2022 onward. There is no additional tuned quality or geographic filter. Coordinates are a WGS84 azimuthal-equidistant projection centered at 14.14°E, 40.82°N; depth is below sea level. Natural Earth coastlines are a display substitute; the original relief and inland zero-elevation contour have not been reconstructed. This is unfinished cartographic work, not a verified access failure. All 12,503 study-period routine depths occupy 51 distinct values on a 0.1 km grid. They were checked against the source HTML depth fields; the plotting code does not jitter, smooth or infer finer depths. The resulting bands are a source-precision limitation, not a location-accuracy estimate. The check is saved in `data/calibration_followup/routine_depth_precision.json`. Marker area is $0.7 + 3\,10^{M_d-2}$ points squared, an explicit display choice because

the paper does not state its precise law. The additive 0.7 point-squared offset keeps small events visible without changing their magnitudes or making the area law nonmonotonic. Missing magnitudes use fixed-size crosses. Depth uses a red-to-blue palette on a black background, with white axes and magnitude legends. The revised color limits are fixed at 0–5 km, with magenta over-range values and no modification of depths. S5 endpoints are printed; 1A endpoints are unlabeled, so its 0–5 km normalization is a disclosed approximation. Actual maxima remain 7.2 and 5.4 km. Published panels were inspected only for style. The routine catalog never enters the inference or association code.

The contextual map windows are display limits rather than catalog-selection filters. 14 selected historical events lie outside the Figure 1A map window; 14 study-period events lie outside the S5 map, and 8 lie outside the northing–depth view. The easting–depth view omits 9 selected events. These events remain in the saved numerical tables and counts. Axis limits and visible/selected counts are recorded in `data/continuation/routine_plot_scope.csv`; the maps label their off-view counts.

Velocity inputs

The exact Calò–Tramelli (2018) association profile was searched in the readable publisher article and supplement, the article's data statement, INGV's Earth-prints repository, the permitted method-artifact tree, and publications/datasets citing the model. Earth-prints lists only the article PDF in its original-file bundle. Supplement section 2 describes Vp inherited from earlier studies and a depth-dependent Vp/Vs profile revised using Wadati diagrams; Figure S1.2 is graphical, without a numerical layer table. Its data statement directs requests to INGV. No graph was digitized and no Tan-fitted model was substituted.

A numerical alternative is verified in Tramelli et al. (2021), Scientific Reports 11, 7211, Table 1. Layer tops are 0, 0.5, 1, 3 and 10 km below sea level; Vp is respectively 1.7, 2.0, 3.0, 5.5 and 5.7 km/s. Vs is Vp/1.78. Layers are piecewise constant; the top value is extended above sea level for station elevations. Near-duplicate boundary nodes preserve discontinuities when preparing the 0.1 km eikonal grid. This is a published alternative, not a verified copy of the 2018 starting profile. All affected association/location results remain partial.

The Tan-distributed reference grid is now explicitly permitted as a disclosed input, despite its dimensional discrepancy with Battaglia et al. (2008). Its preserved bytes, format interpretation, regional averaging and independent calibration are documented in the current authorized-grid section below. The older dependent models and figures remain rejected historical products; no old fit is reused. Historical transformation uncertainty limits exact equivalence, but is no longer a prohibition on running this reference input.

The exact-profile input still lacks numerical layer-top depths, Vp and Vs or depth-dependent Vp/Vs, layered/interpolated conventions, units, depth datum and confirmation of the version used by Tan's GaMMA-1D. The public-source search record is `docs/velocity_profile_search.json`. This issue is assessed after the other independent work; no contact or request to an author or institution was made.

Picking, association and calibration

The released Campi Flegrei-retrained PhaseNet weights are loaded with attribution through SeisBench. Each input is demeaned and high-pass filtered at 1 Hz with a causal four-pole Butterworth filter. Thirty-minute cores have 30-second margins. Both pick thresholds are 0.3; S picks at CASE, CFB1 and CFB3 are excluded. Arrivals within 0.25 s of a higher-probability pick on the same station/phase are merged. Gaps are split, not interpolated. The retraining demonstrations remain unperformed because the exact local annotations, splits and augmentations are unavailable.

GaMMA uses a Bayesian Gaussian mixture, 1D eikonal travel times, at least six picks, and no separate minimum P/S count. Unspecified settings are explicit assumptions: time-only likelihood, DBSCAN

epsilon 10 s, minimum samples 3, oversampling factor 5, variance prior 1 s^2, horizontal bounds ±12 km, and depth 0–8 km. Independent clusters run on six CPU workers and saved identifiers are sorted by origin time. The permitted GaMMA library otherwise advances one random stream per worker, making initialization potentially sensitive to job scheduling. A local wrapper now resets the optimizer seed to 42 plus the deterministic DBSCAN cluster label before each solve; this is an explicit implementation choice, not a published Tan seed. It does not add observations or change the likelihood equations. A serial/parallel consistency test uses two observation windows from our own saved reviews, with one versus three workers and different parent random states. Its measured agreement is reported below; this tests repeatability, not earthquake or phase accuracy. Changing this numerical initialization means differences from older catalog counts cannot be attributed solely to expanded coverage. In the pinned generic GaMMA implementation, candidate origin-time seeds come from the independently picked phases with P phases preferred; horizontal seeds perturb recording-station positions, and depth starts at the 4 km midpoint of the specified bounds. The legacy `initial_points` configuration field is not used by this version. These details are implementation settings, not recovered author settings. Association is partitioned by UTC day, so phases straddling midnight may be split; this is an additional boundary limitation. All input and coverage margins are clipped to the stated study interval. The first sequential association attempt was restarted with parallel clusters to reduce runtime; neither configuration uses a reference catalog to generate events.

No accepted whole-catalog VELEST solve exists. The current blocker is recorded in `data/continuation/velest_summary.json`; prior dependent products are quarantined. The separate alternative-profile precision experiment is recorded in `data/exploratory/velest_stability/summary.json`. No earlier locations are substituted. The pilot geometry screen requires more than six picks and azimuthal gap below 180°, following generic VELEST guidance; it is not claimed to recreate the paper's manually selected 373 events. Nonpositive or invalid initial propagation times are recorded and removed before inversion. The geometry screen is recomputed; events with fewer than four remaining picks cannot proceed to the next residual-rejection solve. These are explicit numerical departures. Controls use seven iterations, hypocenter/origin damping 0.01, velocity damping 1, station damping 0.1, and S/P observation weight 0.5. The superseded workflow attempted two regional starting models, calibration passes and fixed-model catalog passes with the paper's 0.4 s/0.3 s residual rejection and final six-pick, 0.25 s RMS, 300° gap filters. If a pass fails or produces an unusable result, it is recorded rather than replaced by the authors' fitted solution. Those historical controls, logs and candidate selections are retained in the quarantine audit trail; that earlier sequence is disabled. The historical alternative-profile follow-up failed acceptance. The new authorized-grid workflow below supersedes these completion statements.

In the superseded run, 708 candidate events exceeded the generic source's 658-event array capacity. A 1,024-event capacity and bounds checking removed that memory fault; a later 32-bit solve failed a Cholesky check. Those historical failures remain in quarantine. The current isolated experiment compares 32-bit and 64-bit arithmetic from identical observations, the same alternative profile and unchanged damping. It checks both numerical agreement and physical velocity ordering; neither precision is accepted merely because it returns. Current outcomes are reported below. No accepted whole-catalog VELEST solution exists.

The current magnitude pilot includes all retrieved stations with suitable windows, including stations without an associated phase. At those stations it uses theoretical P times from the exploratory alternative profile. Magnitude measurements use picked P arrivals where available, otherwise independent predicted arrivals, a 5 s signal window, a 2 s preceding noise window and a 5 dB power-SNR threshold. Responses are removed to velocity with prefilter corners 0.1, 0.2, 35 and 45 Hz and a 60 dB water level. Event separation must be less than one fifth of the common station distances, with at least two common stations. Channel log amplitude ratios are weighted by squared, arrival-aligned zero-lag

correlation, an assumption where the precise correlation/weight specification is absent. Weighted least squares uses unit-weight routine-Md priors only for unique post-detection temporal matches within 2 s; multiple independent events claiming the same reference are rejected as ambiguous calibration anchors, without removing detections. Unconnected or unanchored components receive no magnitude. These priors are calibration inputs, not an independent validation set. Provisional initial locations limit any pilot magnitudes. Only full-rank anchored amplitude systems return magnitudes. Formal uncertainties use a residual-scaled inverse normal matrix and remain conditional on the weighting and calibration assumptions. The saved maximum-curvature/maximum-likelihood calculation cannot establish full-period completeness from this noncontinuous sample.

The CAAM/COLB hybrid scan uses the three written-method template times, 8 s templates, 0.2–6 Hz filtering, a two-station mean normalized correlation threshold of 0.25 and 8 s peak separation. Common returned intervals are intersected; duplicates and ten-second filter-edge regions are removed. It covers 75.797 hours and produces 822 unconfirmed threshold crossings. These are not confirmed hybrid clusters. The previous four-candidate review showed persistent fluctuations and did not establish isolated hybrid onsets. The extended candidates now receive saved CAAM frequency-index and signal/noise measurements; these measurements alone do not confirm events. Cluster membership, three-year history, cluster-relative magnitudes and catalog-wide coda discrimination remain unfinished.

Historical alternative-model follow-up, 13 October 2026

This earlier checkpoint is retained to explain failed approaches. The current authorized-grid run is reported in its own section; the paragraphs here do not describe its completion status. The frozen calibration cohort contains 1,003 provisional associations. Its event identifiers and earlier magnitude, template and frequency-index products are preserved. An additional 43 successful six-hour waveform requests supplied 525,189,120 bytes and 703.785 channel-hours; 5 requests returned HTTP 204. The segmented queue now has 694 successful requests, 34 no-data responses and 73,946 unattempted requests. Pending intervals remain unfinished acquisition, not evidence of remote inaccessibility.

The separate expanded checkpoint processed 868 files from 26 station names, with 8511.520 union channel-hours. It produced 132,733 merged picks (19,530 P and 113,203 S), and 1,015 provisional associations containing 7,351 assigned picks. These identifiers are not interchanged with the earlier calibration cohort. Complete epoch intersections identify 0.000 channel-hours without a usable saved response. This checks metadata availability, not response accuracy or complete three-component detection exposure. Station coverage, every unobserved interval and request status are saved under `data/calibration_followup/expanded`. No expanded event is yet promoted to an accepted location or substituted into earlier magnitude plots.

Executed VELEST chain. The written methods identify 373 high-quality calibration events but give no numerical rule that uniquely selects them. The declared causality, more-than-six-pick and gap-below-180° screen selects 130 events: 24, 1, 1, 104 in 2022–2025. 3 have only S picks. This is an unreviewed and temporally unrepresentative alternative selection. The starting layers are Tramelli et al. (2021), Table 1, with the top extended to −0.5 km for station elevations. They are not the missing Battaglia regional averages. The reference P delay at IV.CSOB. is fixed; initial delays are zero. Two calibration rounds estimate locations/velocities and then also station delays, using seven iterations per round and the preceding explicit damping and S/P weights.

Both arithmetic precisions completed both rounds. The fitted velocity tables agree at their printed precision; maximum location differences are 11.105 m horizontally and 10.0 m vertically. Five fixed-model, fixed-correction catalog passes were executed on 905 causally consistent candidates, including the stated 0.4 s and 0.3 s residual rejection. The final six-pick, 0.25 s and 300° filters retain 238

exploratory locations. Fixed models/corrections were checked after every pass; absolute own-pick times were preserved to 0.001609 s through serialized origin-time changes.

Validation rejects acceptance. Another five seven-iteration passes do not converge. The last moves an event by 4.935 km horizontally and 2.04 km vertically; a fitted depth reaches 18.40 km. These are unstable trial estimates, not newly established deep earthquakes. With every fifth chronological event held out, unweighted phase RMS worsens from 0.314 to 0.385 s, although aggregate reported event RMS changes from 0.335 to 0.319 s. Recalculation shows that VELEST reports a sample standard deviation of normalized weighted residuals rather than unweighted phase RMS. Maximum discrepancy from the printed values is 0.005221 s, consistent with output rounding. Thus the mixed validation is not resolved by relabeling both metrics RMS.

One declared sensitivity trial requires at least four P and two S picks and excludes initial depth-bound hits. It leaves 46 events. Further iterations produce $V_p \leq V_s$; its held-out training branch also fails that physical check. Neither fit is propagated. Reviewed phase identities and well-constrained events are needed before another regional fit can be accepted. The stationary-coordinate approximation differs from sampled channel epochs by up to 12.990 m, already relevant to a claimed ten-meter precision. These checks establish reasons for rejection, not an accepted calibrated catalog.




Figure D1: Reproduction diagnostic, not a publication figure. Saved convergence trials show kilometer-scale location changes despite similar residual statistics. The model and locations are rejected for scientific interpretation.

Observed delays and exploratory HypoDD. The 238-event filtered trial supplies 1653 nearest-neighbor pairs and 6362 catalog differential phases. Original response-corrected waveform measurements supply 10,882 channel comparisons, 737 passing channels and 517 coherency-weighted station/phase delays; 13 window failures are retained. The preceding two-second window, 1–15 Hz band and correlation/coherence thresholds are explicit assumptions, not recovered author configuration. Rejected picks or published event solutions are not used to fill gaps.

Generic HypoDD 2.1 was obtained from Waldhauser's official repository at revision `fb40407` (full commit in the source receipt). Bounds checks exposed an incorrect output-vector dummy extent in `matmult2.f`; it was changed from the column count to the row count, with unchanged multiplication equations. An isolated synthetic matrix assertion checks that fix and never enters empirical data. Source hashes, dimensions, compiler flags and the exact change are recorded in `hypodd21_sources.json` and `hypodd_build.json` under the follow-up data directory.

Two alternative-profile HypoDD LSQR rounds returned 127 and 126 locations. These are not TomoDD rounds. Six shared phases link clusters; two four-iteration blocks shift weight from catalog to waveform delays, with LSQR damping 50, and a 300° gap ceiling. Magnitudes and initial errors set to zero are unused format fields, not measured quantities. LSQR error columns are not interpreted as uncertainties. Quadrants about 14.14°E, 40.82°N provide explicit alternative SVD subsets; they do not recreate the paper's four numerical selections. Smaller underconstrained components failed, so only the largest connected component in each quadrant was attempted. NW, SW and SE return 40, 16, 22 estimates; NE diverges and supplies no valid output. The largest conditional 95% depth errors are 4494.8, 662.9, 507.7 m, respectively. These conditional errors and unstable locations cannot establish the publication's spatial precision. No S7 panel is promoted.




Figure D2: Reproduction diagnostic, not Figure S7. Conditional SVD errors for three exploratory connected components; the fourth diverged. Zero waveform-difference counts are retained on a symmetric-log abscissa. Error bars cannot validate unreviewed events or the background model.

Provenance and remaining prerequisites. The authorized De Siena et al. (2017) article cites the original Battaglia model and links PANGAEA 875535. Its inspected metadata list derived locations, attenuation products, geochemistry and map material; no numerical original V_p/V_s grid is identified. Those derived products were not downloaded as analysis inputs. At that checkpoint the model was quarantined. It is now explicitly authorized as a reference input; the new grid-derived results above supersede this historical completion record. Source receipts, URL and hash are recorded; no correspondence was initiated. Exact Calò–Tramelli layer depths and velocities also remain missing, but did not prevent these explicitly alternative trials. TomoDD using the stated fixed 3D background, accepted-location magnitudes, the four specified S7 subsets and their dependent structural panels remain uncompleted. Reviewed phases, stable calibration, verified numerical grid/coordinates/depth convention and regional-averaging masks remain prerequisites. Coverage remains sparse and RAN waveforms credential-gated. Additional observations alone do not resolve the failed calibration.

The complete own-run controls, input serialization, logs and outputs, including failures, are preserved in `data/calibration_followup/run_records.zip` with member hashes. Code and commands are in `docs/CALIBRATION_FOLLOWUP.md`. The incremental rebuild executes these trials and reports scientific failures separately from successful software execution; it does not claim a full-period reproduction. Earlier magnitude, focal-mechanism and hybrid statuses are unchanged.

D4 Supporting results and validation

The frozen calibration cohort has 126133 merged picks and 1003 provisional associations; the separately numbered expanded cohort is described above. Probability bounds, S-pick exclusions, the six-pick min-

imum, unique assignment and valid pick indices are checked. Synthetic homogeneous-medium, analytic and injected-signal fixtures are isolated under `tests/results`. They test software assertions only and are excluded from empirical catalogs, magnitude measurements and paper figures. Earlier rebuilds ran the equation and layered-ray checks; their retained pass records are not observations. The current grid rebuild executes the distinct validation described below.

The present S–P check includes 1703 paired observations, of which 58 are nonpositive. There are 137 associated CPOZ P arrivals. These defects and the alternative velocity profile prevent treating raw associations as validated locations. Internal residuals from the same model do not establish absolute accuracy. The location-check tables retain failed observations. The previous generic/retrained comparison on an unlabeled 1600 s CMIS window produced 17 P/4 S and 11 P/55 S picks respectively; the 120 s CSOB window produced zero and one pick. Those counts remain supporting evidence only, not training or labeled-accuracy reproduction, and are not inserted in S4. An analytic vertical-ray check of the published alternative gives maximum P/S errors of 0.040642/0.072342 s on the 0.1 km eikonal grid; halving the spacing reduces them to 0.020321/0.036171 s. This confirms first-order interface discretization behavior, not the physical accuracy of the alternative profile. The numerical checks and refinement results are saved in `tests/results/profile_ray_checks.csv`.

The continuation adds `scripts/association_quality.py`, which flags nonpositive travel times, nonpositive S–P, S-only associations and depth-boundary solutions without changing the provisional event list. The first two flags exclude an event from magnitude equations, calibration anchors and the VELEST precision experiment because they violate arrival causality. This explicit necessary screen does not reproduce the paper's manual false-event review. Five baseline multi-station waveform examples were inspected: a strongly recorded event supports functioning detection, while other examples contain late coda picks, arrivals before the fitted origin or unclear onsets. The selection was diagnostic rather than random; it supplies no precision or recall estimate. The original arrays, plotted review and qualitative notes are retained in `data/resumption/baseline_association_review`. None of the baseline's 53 provisional magnitudes belonged to the 102 events failing causality. All five updated diagnostic pages were also inspected after the corrected association. They include clearly recorded events, late or coda-associated phase marks, and a real-looking signal whose fit reaches the lower depth bound. Passing causality therefore cannot validate depth or phase identification. Current observations and hashes are recorded in `data/resumption/current_association_visual_review.json`; no manual phase edits or event acceptance were inferred from this small diagnostic sample.

The new S4A calculation uses original CSOB waveforms starting at 20 May 2024 20:10:03 UTC, exactly two hours after the event time in the official INGV bulletin. This is a literal interpretation of the caption, not an extracted figure coordinate or an asserted exact match. The bulletin provides contextual timing only; these example picks do not seed the catalog. The trained artifacts and their hashes are in `data/continuation/benchmark_window.json`. No labels, predictions or training records from Tan's outputs are used. The released repository's file listing contains trained weights, workflow code and derived products, but no trace-level training labels, split or augmentation manifest. Only the file listing was inspected; prohibited code and output contents were not read. The archived R61 release (Zenodo 16562216, version 1.0) was also directory-listed using HTTP byte ranges. Its permitted trained weights and model configuration were decoded and are byte-identical to the current method artifacts. The archive lists no original training waveform/label, split or augmentation files. Its separately listed association-pick table was not retrieved. The directory, ranges and hashes are recorded in `docs/zenodo_release_inventory.json`.

For preliminary differential times, ten nearest neighbors are selected from causally consistent, nonboundary provisional locations. P uses Z; S uses N and E separately. Original observations are response-corrected to velocity, filtered from 1–15 Hz, and cut from -0.25 to $+1.75$ s around each own pick. An integer correlation lag within ± 0.25 s precedes a cross-spectral phase-slope correction.

With angular frequency ω, cross spectrum S_{xy} and magnitude-squared coherence γ^2, the correction minimizes $\sum_f w_f[-\arg S_{xy} - \omega\delta]^2$, where $w_f = \gamma^2/(1-\gamma^2)$ with a 10^{-6} denominator floor. Welch spectra use 64-sample Hann segments and 32-sample overlap. Correlation and mean coherence must each exceed 0.6; boundary lags and residual corrections above one sample are flagged. These explicit settings are assumptions where Tan's written description is incomplete, not recovered author settings. The differential travel time is $(p_a - o_a) - (p_b - o_b) - \delta_{b-a}$, in seconds. Saved observations, measurements and failures are in `data/continuation/differential_waveform_windows.npz`, `data/continuation/differential_times.csv` and `data/continuation/differential_time_failures.csv`. Synthetic zero-lag, known-delay and reversal tests remain isolated under `tests/results`. No inferred relocation precision or accepted event coordinates follow from this pilot.

The current amplitude/SNR stage measured 61681 channel amplitudes, with 4705 passing SNR and 1163 accepted difference equations. It returned 119 provisional magnitudes. These are actual executed outputs, not a promised future count. Candidate measurement errors and quality failures are retained in the JSON/CSV records. The old 140-measurement, 23-SNR, zero-edge result is superseded for the current sample and preserved in Git and the quarantine audit trail.

Additional observational checks

The CPOZ diagnostic contains 137 events and 274 HHZ/HHN traces, aligned on own P picks over -1 to +7 s and highpass filtered above 4 Hz. These now follow the written S12 window and band. The sector/coordinate ordering and onset validation are still absent.

Spectral measurements were calculated for 822 of 822 template crossings. None is confirmed as a hybrid event; spectral windows use correlation alignment and still require onset/coda review.

Catalog frequency-index screening measured 21199 station windows for 1003 provisional events. Station FIs are averaged; no low-frequency candidate is classified as a confirmed hybrid without onset and coda review.

This screening lacked 4878 station-event windows and recorded 1 measurement failures. The saved failure is a waveform sample-count assertion, not an established archive-access failure; it remains unfilled.

Diagnostic windows were saved for 9 negative-mean-FI candidates at 26 selected station windows. These are observations selected for review, not new event classifications.

All 9 corresponding diagnostic pages were visually inspected. Persistent oscillations, impulsive packets with positive station FI, low-SNR negative-FI records, and strong earlier arrivals decaying through later windows occur. In particular, the CMIS windows for candidates 631 and 632 show apparent coda contamination. No hybrid was confirmed. This limited agent review supplies no expert-validated onset labels or catalog-wide accuracy estimate; its notes and hashes are retained in data/resumption/catalog_frequency_visual_review.json.

The 89 retained external routine-Md calibration matches span -1.10 to 1.33; 0 reach the written focal-mechanism sample's lower Md of 2.2. These are calibration reference values for independently associated candidates, not independent magnitude measurements or a bound on events outside this matched sample. No reviewed mechanism sample has been established.

The preceding checkpoint had 53 conditional magnitudes, Mc=0.4 and b=1.01. The current checkpoint has 119, Mc=-0.5 and b=0.66. Coverage, association initialization and causality screening changed together; the separate contributions were not measured. This change prevents treating the earlier apparent agreement as a stable completeness or regional b-value result.

The observed-window repeatability test used 119 own picks and returned 6 provisional associations with 79 assigned picks. Changing from one to three workers and changing the parent random state gave maximum coordinate and origin-time differences of 0.00000000 km and 0.000000 s after the per-cluster

seed correction. This tests scheduling independence on these windows, not event or location accuracy.

Of 822 mean-score template crossings, 30 have a nonpositive score at one station. The original mean-score selection is retained, but this exposes why crossing the mean threshold cannot itself confirm a two-station hybrid event.

The alternative-profile VELEST experiment used 130 provisional events. Both real precisions were attempted with identical serialized inputs.

Maximum 32/64-bit differences were 0.000 km/s for Vp and 0.000 km/s for Vs, with 0.010 km in fitted depth.

The horizontal-coordinate check gives a maximum difference of 11.105 m in 2 locations. Identical serialized locations: False. A depth-only comparison is insufficient to claim full coordinate agreement.

Maximum serialized origin-time and event-RMS differences are 0.010 s and 0.000 s. Printed resolution does not bound unprinted internal differences.

Both runs pass the necessary positive-Vp-greater-than-Vs check. This ordering and numerical agreement do not replace reviewed phases or the documented regional calibration; no experimental fit is accepted for S6 or relocation.

The 32-bit experiment has event-RMS aggregate 0.357 s.

The 64-bit experiment has event-RMS aggregate 0.357 s.

Causality screening finds 56 events with nonpositive S-P and 51 with nonpositive propagation times; 905 of 1003 pass both necessary checks. There are 66 S-only associations and 238 depth-boundary solutions. Flags are saved for all events; they are not a manual earthquake classifier. Events failing causality cannot enter the magnitude equations, calibration priors or VELEST experiment. Other provisional associations remain unvalidated.

Finite PhaseNet-score exposure for continuous checkpoint inputs is 2591.439 P station-hours and 2364.086 S station-hours after the S exclusions. Earlier baseline files lack this telemetry. These units differ from observed channel-hours. SeisBench fills missing input components with zeros for inference; those values are not counted or plotted as observations. Short fragments without scores remain distinguishable from observed samples.

Exploratory neighboring-event processing saved 9241 observed phase windows, measured 19580 channel pairs, and retained 811 under the declared correlation/coherence criteria. These are partial timing measurements on 721 unvalidated, causally consistent nonboundary events, not TomoDD results. Initial neighbor geometry and missing author window/weight settings remain limitations.

For 226 S-wave pairs passing on both horizontal components, the median absolute delay difference is 0.0019 s and the maximum is 0.0162 s. This internal component check does not measure absolute phase accuracy or hypocentral uncertainty.

Review of low-frequency candidates

The additional low-frequency review is recorded in `data/resumption/catalog_frequency_visual_review.json`, with numerical windows and their source/hash index in `data/continuation/catalog_frequency_review.npz` and `data/continuation/catalog_frequency_review_index.json`. The diagnostic selects the two highest station SNRs and the lowest station FI per candidate, retaining distinct stations. It displays original response-corrected Z observations from -20 to $+16$ s, with longer processing context and a separate 0.2–6 Hz curve. FI and SNR labels refer to the previously saved measurement windows, not a recomputation from the display. Dotted markers show other provisional arrivals; they are not validated coda labels. No qualitative review note is used to alter the catalog, and the station measurements' reviewed-onset flags remain false. The diagnostic PDF is retained in the repository and

excluded from the paper and figure-comparison deliverable.

Magnitude statistics and comparison boundary

The conditional magnitude-frequency fit gives $M_c = -0.5$ and $b = 0.66 \pm 0.05$ from only 103 selected magnitudes at or above the pilot completeness level. The quoted Shi–Bolt standard error excludes coverage, selection and magnitude-calibration biases. This calculation does not establish population completeness or a regional b-value.

No Tan-derived result-table statistics are used in this paper. The comparison PDF contains only clearly labeled published figure references and the eligible independent counterparts.

D5 Authorized-grid calibration and fixed-grid relocation

This section supersedes the calibration completion statements in earlier checkpoints. Earlier experiments are retained separately as historical diagnostics. The present run uses the expanded cohort; integer event identifiers must not be joined to the older 1,003-event cohort.

Reference input and transformation. Tan et al.'s distribution at revision `f84c9c9d` (full revision in the source receipt) supplies `models/3d_vmod`, attributed to Battaglia et al. (2008; doi:10.1111/j.1365-2478.2007.00687.x). The current instruction explicitly permits that file as a starting/reference input. Its unchanged 122,853-byte copy is `data/grid_calibration/reference_grid.original`; SHA-256 is `dc36fe2a542fc7a3b6c473c48ec109c53ddd77a1721f509b920fea97fb0166f3`. It is not claimed to be an independently retrieved original 2008 array. The grid contains 23 longitude, 14 latitude and 27 depth nodes, ordered longitude fastest, then latitude and depth. Longitude/latitude are degrees east/north, depth is km below sea level, V_p is km/s, and the second volume is dimensionless V_p/V_s. Pointwise $V_s = V_p/(V_p/V_s)$. Depth extends from -0.5 to a 200 km padding node; the shallow model ends at 7 km. The original article describes an 18 by 18 by 9 km volume with 0.5 km spacing, unlike these irregular/padded axes. The historical transformation remains undocumented. The generic TomoDD reader and manual establish the encoding; the current 3D file is not regridded.

Regional averaging uses explicit assumed boxes: 14.01–14.21°E, 40.76–40.87°N for the caldera and 14.11–14.16°E, 40.80–40.84°N for Pozzuoli–Solfatara. Node weights are cos(latitude); V_p and pointwise V_s are averaged separately. Exact $V_p = 0.1$ km/s nodes are interpreted as nonsolid/air and excluded from the 1D average. The first entirely solid plane, 0.1 km deep, extends upward to -0.5 km. Horizontal padding and the terminal 200 km layer are excluded from regional depth integration. These masks, layer boundaries and weights are assumptions where Tan gives region names without numerical masks. They do not recover an undocumented transformation.

Independent observations and corrections to the failed pilot. The saved expanded input contains 868 processed files, 26 station names and 8511.520 union channel-hours. Independent association times span 2022-01-21T01:14:48.080 to 2025-03-20T23:58:58.807, with large gaps; this is not full-period coverage. 916 of 1015 associations have positive travel times and no observed nonpositive S–P pairs. Calibration additionally requires at least four P and two S picks, more than ten total picks, gap below 180° and no initial 0/8 km depth-bound hit, leaving 38 unreviewed candidates. This is not Tan's manually selected 373-event sample. No routine catalog or author result selects these events.

The first eight-layer attempts produce $V_p \leq V_s$ after the station-delay inversion and are rejected. The generic VELEST guide recommends avoiding low-velocity layers initially, locating events before velocity inversion, merging similar unresolved layers and damping unresolved deep velocities. Accordingly

LOWVELOCLAY is zero, prelocation uses 14 iterations, and four layer tops are $-0.5, 1, 3, 5$ km. The deepest velocity damping is 100, other layer damping is 1; deeper unconstrained velocities are not freely fitted. A failed damping-1000 attempt overflowed the fixed-width output field and is retained as a serialization failure. The corrected four-layer run preserves physical ordering. It performs two seven-iteration calibration rounds followed by three declared convergence checks, without changing observations or relaxing the paper's final filters. The reference P delay is fixed at IV.CFB3.. The converged model, not the still-moving second-round fit, supplies the catalog run.

Validation and catalog filtering. Both regional fits satisfy the stated local stability gates: every horizontal/depth change below 100 m, velocity change below 0.05 km/s, and station-delay change below 0.02 s in the final extra pass. The lower converged unweighted phase RMS is 0.218 s (caldera). These are numerical gates, not absolute accuracy. Regional starts still give maximum differences of 654.7 m horizontally and 1.04 km in depth. Complete 32/64-bit paths differ by 557.8 m horizontally, 0.89 km in depth, 0.32 km/s in velocity and 0.25 s in station delay. Starting both precisions from the same converged fit instead gives identical printed coordinates and velocities. Thus local convergence does not remove path sensitivity; printed equality does not bound unprinted differences. Reviewed phases, held-out refitting for this model and independent absolute-location validation remain undone. The fit is not accepted as a scientific catalog.

All 916 usable events enter the fixed-model run. Three passes precede $|r| > 0.4$ s rejection; another pass precedes $|r| > 0.3$ s rejection. The fifth pass and final six-pick, 0.25 s reported-RMS and 300° gap filters leave 216 exploratory locations. Fixed models and station corrections were checked after each pass; rejected phases are not replaced. The generic reported event statistic differs from unweighted phase RMS, as verified in the preceding checkpoint.

Observed differential times and TomoDD. round 1: 132 outputs, 124 gap passes; round 2: 124 outputs, 123 gap passes. both fixed-grid rounds executed; validation pending. The neighbor graph and windows belong to this expanded event set; no old-cohort waveform cache is reused. Exact measured counts, failed windows, controls and residuals are in `data/grid_calibration/merged4/tomodd`. Original absolute picks supply phase-pair times; response-corrected two-second windows supply coherency-weighted delays. Ten nearest neighbors, 1–15 Hz filtering, correlation/coherence at least 0.6, maximum lag 0.25 s, six-phase clustering links, LSQR damping 50 and two four-iteration weight blocks are explicit assumptions where configuration is unpublished. JOINT=0 fixes the 3D grid. A second round rebases travel-time differences to the independently updated origins. Generic method tables required by the solver are separated from the example observations in its distribution. No author-fitted corrections, velocity model or earthquake result is substituted. Failed or unvalidated outputs do not populate Figures 2–4, S6–S7 or S9–S13.

Display verification and scientific boundaries. Every current generated figure is rebuilt from saved numerical products by `scripts/publication_plots.py`. Input/output hashes and panel correspondence are in `data/grid_calibration/style/plot_inventory.json`. Figure S5 uses the printed 0–5 km depth limits and ticks 0,1,2,3,4,5; Figure 1A has printed ticks 0,2,4 but unlabeled endpoints, so 0–5 km is an explicitly approximate normalization rather than an exact recovery. Neither numerical table changes. Magenta denotes depths above 5 km: 4 in 1A and 2 in S5; white is reserved below zero. Missing magnitudes remain crosses. The area law $0.7 + 3\,10^{M_d - 2}$ points squared is a display assumption, not a recovered published formula. S4/S15 do not print a numerical spectral color scale; their disclosed normalized-PSD display is -80 to 0 dB, with white/black under/over colors. Counts and ranges are saved in `spectral_scales.json`. S2 uses documented instrument classes; no copied relief or elevation colorbar is manufactured. Diagnostic figures have consistent typography but are explicitly not publica-

tion figures. Figures 3 and 4A–B have no valid independent counterparts, so none is invented for style matching.

Rebuild and audit trail. Run `.venv/bin/pythonscripts/rebuild_grid_workflow.py` from the repository root. It checks the existing independent processing checkpoint, rebuilds generic solvers, regenerates grid-derived starts, executes calibration/validation and both relocation attempts where supported, plots saved data and compiles both PDFs. The saved independent waveform-window cache may be reused only with matching input/output hashes; `--remeasure` recomputes it from recordings. The current procedure does not imply that unfinished full-period acquisition or inference was executed. Controls, own serialized inputs, logs and results are preserved in `data/grid_calibration/run_records.zip`; machine-readable procedure and scientific outcomes are separate. Earlier models/results remain in Git and their labeled historical directories. The current grid authorization changes input eligibility, not numerical validation requirements. The generic solver originally overwrote earlier clusters' residual tables. The output-only patch appends every cluster; final gaps now use only stations with positive-weight retained residuals. This correction changes the trial gap-passing counts and supersedes the last-cluster-only diagnostics, which are preserved separately. All 124 final output events have residuals represented. The final per-observation RMS is 0.0430 s for waveform differences and 0.1086 s for picked-phase constraints. Neither value establishes correct event identity or absolute location accuracy.




Figure D3: Reproduction diagnostic, not a publication panel. Grid-derived starting and independently fitted layer velocities, exploratory second-round TomoDD locations and the change in depth between rounds. Fixed color limits are 0–5 km; magenta retains deeper results. Calibration path dependence and unreviewed phases prevent interpreting these patterns as recovered faults.

D6 Limitations and unfinished analyses

No publication panel is fully independently reproduced. Partial counterparts: 1A, 1B, 1C, S1, S2, S4A, S5/map, S5/northing-depth, S5/easting-depth, S8A, S8B, S15C. All remaining panels are not reproduced. Routine-context permissions and the 2025 displacement gap are resolved. The full continuous download/processing workload remains unfinished. Exact Calò–Tramelli layer values are an external-input limitation; the distributed fixed 3D artifact is now authorized, while its original transformation remains unresolved and the new calibration remains path-sensitive. RAN credentials are a specific access limitation.

The accepted regional calibrated and TomoDD-relocated catalog, four specified HypoDD SVD subsets, reliable multiyear magnitudes/completeness, manually reviewed polarities and S/P ratios, SKHASH/gCAP solutions, FK Green's functions, exact event-specific gCAP filters/windows/weights,

relocated CPOZ sector gathers, manual deep-event review and three-year hybrid clustering are not complete. The current authorized-grid section records actual TomoDD execution and validation; completion is not inferred from compilation. Training labels/splits and some inversion settings are incompletely specified. Supplied author-derived results could fill plots only by violating the independence restriction, so they are not substituted. That restriction is distinct from unfinished implementation or genuinely missing source inputs.

D7 Figure-by-figure record

The machine-readable `docs/panel_record.json` records every panel's written specification, parameters, exact input paths, executed data/plot commands, saved products and limitation. The raw-waveform manifest is `docs/continuation_processed_inputs.json`; its intervals are a frozen checkpoint, not the full queue. Context data use `continue_context.py`; all current displays use `scripts/publication_plots.py`. S1 displays the expanded processing coverage. S2 is audited by `scripts/station_geometry.py`; S15C retains its original-waveform recipe. Unmatched panels have text notices in this paper and explicit placeholders only in the comparison PDF.

Panel/view	Quantity	Status, specification and limitation
1A	INGV event map and large-event markers	partially reproduced. Main p.71, Fig.1 caption. 2005-01-01 to 2025-03-20; origin 14.14 E, 40.82 N; sea-level depth; Md>=4 markers Display: depth-colored jet_r circles, area=0.7+3*10**(Md-2) points squared; missing Md shown as crosses. Exact publication area law unspecified. Full color ranges: 0–8 km for 1A, 0–6 km for S5. Display bounds and off-view counts saved in routine_plot_scope.csv.. Current catalog snapshot differs from the historical snapshot: 12,503 located study events versus 12,083. No tuning to published counts. Natural Earth coastline is a display substitute; the original relief/inland-zero contour has not been reconstructed.
1B	Vertical displacement at Pozzuoli port	partially reproduced. Main p.71, Fig.1 caption. 128 historical entries: 114 leveling surveys, 13 source-provided estimates and one source-provided graphical reconstruction. Source classes preserved separately. Daily GNSS 2000-05-29 through 2025-03-20; median datum from 23 overlapping leveling epochs, +/-15 days. Month-only dates placed on day 15 for display.. Benchmark equivalence and datum-joining convention are assumed because the exact original stitching recipe is not specified. Calendar gaps remain unfilled; no missing recent-data blocker through 2025-03-20.
1C	Temporal seismicity and cumulative counts	partially reproduced. Main p.71, Fig.1 caption. 2005-2025; cumulative Md>0.2; study interval 2022-01-21 to 2025-03-20. Monthly counts and exact-time cumulative Md>0.2 reconstructed from the current routine snapshot; historical count differences unresolved. March 2025 is a partial month.
2A	Catalog seismicity rates by region	not reproduced. Main p.72, Fig.2 caption. Central vs ring/other regions; 54319 vs 12083 catalog events; time bin unspecified. Grid input explicitly authorized; historical transformation and averaging masks not recovered exactly. Local calibration convergence passes, full-path precision and start-model dependence do not. Unreviewed phases and sparse coverage; no accepted relocated scientific catalog. both fixed-grid rounds executed; validation pending

Panel/view	Quantity	Status, specification and limitation
2B	Depth versus time, symbol size by magnitude	not reproduced. Main p.72, Fig.2 caption. Sea-level depth; near-surface 0-0.5 km; central 0.5-2.1; ring sectors to 3.7 km. Grid input explicitly authorized; historical transformation and averaging masks not recovered exactly. Local calibration convergence passes, full-path precision and start-model dependence do not. Unreviewed phases and sparse coverage; no accepted relocated scientific catalog. both fixed-grid rounds executed; validation pending
3/map	Epicentral spatial pattern	not reproduced. Main p.73, Fig.3 caption. Relative origin 14.14 E, 40.82 N; sea-level depths. Grid input explicitly authorized; historical transformation and averaging masks not recovered exactly. Local calibration convergence passes, full-path precision and start-model dependence do not. Unreviewed phases and sparse coverage; no accepted relocated scientific catalog. both fixed-grid rounds executed; validation pending
3/northing-depth	Depth versus northing	not reproduced. Main p.73, Fig.3 caption. Relative origin 14.14 E, 40.82 N; sea-level depths. Grid input explicitly authorized; historical transformation and averaging masks not recovered exactly. Local calibration convergence passes, full-path precision and start-model dependence do not. Unreviewed phases and sparse coverage; no accepted relocated scientific catalog. both fixed-grid rounds executed; validation pending
3/ easting-depth	Depth versus easting	not reproduced. Main p.73, Fig.3 caption. Relative origin 14.14 E, 40.82 N; sea-level depths. Grid input explicitly authorized; historical transformation and averaging masks not recovered exactly. Local calibration convergence passes, full-path precision and start-model dependence do not. Unreviewed phases and sparse coverage; no accepted relocated scientific catalog. both fixed-grid rounds executed; validation pending
4A	Ring-sector focal mechanisms and seismicity	not reproduced. Main p.74, Fig.4; supplement S6-S7. High quality: SKHASH A or Kagan angle<30 deg; gCAP preference with stated exception. Grid input explicitly authorized; historical transformation and averaging masks not recovered exactly. Local calibration convergence passes, full-path precision and start-model dependence do not. Unreviewed phases and sparse coverage; no accepted relocated scientific catalog. both fixed-grid rounds executed; validation pending
4B	Central/onshore focal mechanisms and seismicity	not reproduced. Main p.74, Fig.4; supplement S6-S7. High quality: SKHASH A or Kagan angle<30 deg; gCAP preference with stated exception. Grid input explicitly authorized; historical transformation and averaging masks not recovered exactly. Local calibration convergence passes, full-path precision and start-model dependence do not. Unreviewed phases and sparse coverage; no accepted relocated scientific catalog. both fixed-grid rounds executed; validation pending
S1	Station data-availability timeline	partially reproduced. Supplement S12 caption; S2 methods. 48 stations; continuous-data availability. Expanded processed subset: 868 files, 8511.520 channel-hours; full-period queue remains unfinished. No requested-but-failed interval is counted.
S2	Station map	partially reproduced. Supplement S13 caption; S2 methods. 23 broadband, 9 short-period, 16 accelerometers. The published 48-instrument membership is not verified. Two released-coordinate pairs coincide; no deployment alias mapping justifies collapsing or removing identifiers.

Panel/view	Quantity	Status, specification and limitation
S3A	Noise example: waveform, phase labels, spectrogram	not reproduced. Supplement S14 caption; S2-S3. 30-s windows; P/S labels; D amplitude rescaling. Exact training traces and labels not identified from permissible metadata. Retraining was not repeated; permitted trained weights used for inference only.
S3B	Single-event example: waveform, phase labels, spectrogram	not reproduced. Supplement S14 caption; S2-S3. 30-s windows; P/S labels; D amplitude rescaling. Exact training traces and labels not identified from permissible metadata. Retraining was not repeated; permitted trained weights used for inference only.
S3C	Two overlapping events: waveform, phase labels, spectrogram	not reproduced. Supplement S14 caption; S2-S3. 30-s windows; P/S labels; D amplitude rescaling. Exact training traces and labels not identified from permissible metadata. Retraining was not repeated; permitted trained weights used for inference only.
S3D	Five overlapping copies of one event: waveform, phase labels, spectrogram	not reproduced. Supplement S14 caption; S2-S3. 30-s windows; P/S labels; D amplitude rescaling. Exact training traces and labels not identified from permissible metadata. Retraining was not repeated; permitted trained weights used for inference only.
S4A	Swarms at CSOB: waveform, spectrogram, original and retrained picker scores	partially reproduced. Supplement S15 caption; S2-S3. 120 s CSOB window at 2024-05-20 20:10:03 UTC, interpreting the caption two-hour offset literally; 1 Hz highpass, released model settings, thresholds 0.3.. Exact published start is not specified in the caption; no reference phase labels. Different pick counts do not establish accuracy or successful retraining.
S4B	CMIS noise: waveform, spectrogram, original and retrained picker scores	not reproduced. Supplement S15 caption; S2-S3. 1600 s; coastal noise; phase threshold 0.3. Original and retrained models both executed on independently selected CSOB/CMIS calendar windows. Exact published windows and independent reference annotations remain unidentified; these diagnostics are not the published benchmark.
S5/map	Routine-catalog spatial views; map	partially reproduced. Supplement S16 caption. 12083 events; 1D model and HYPO71 Display: depth-colored jet_r circles, area=0.7+3*10**(Md-2) points squared; missing Md shown as crosses. Exact publication area law unspecified. Full color ranges: 0–8 km for 1A, 0–6 km for S5. Display bounds and off-view counts saved in routine_plot_scope.csv.. Current catalog snapshot differs from the historical snapshot: 12,503 located study events versus 12,083. No tuning to published counts. Natural Earth coastline is a display substitute; the original relief/inland-zero contour has not been reconstructed.
S5/northing-depth	Routine-catalog spatial views; northing-depth	partially reproduced. Supplement S16 caption. 12083 events; 1D model and HYPO71 Display: depth-colored jet_r circles, area=0.7+3*10**(Md-2) points squared; missing Md shown as crosses. Exact publication area law unspecified. Full color ranges: 0–8 km for 1A, 0–6 km for S5. Display bounds and off-view counts saved in routine_plot_scope.csv.. Current catalog snapshot differs from the historical snapshot: 12,503 located study events versus 12,083. No tuning to published counts. Natural Earth coastline is a display substitute; the original relief/inland-zero contour has not been reconstructed.

Panel/view	Quantity	Status, specification and limitation
S5/ easting-depth	Routine-catalog spatial views; easting-depth	partially reproduced. Supplement S16 caption. 12083 events; 1D model and HYPO71 Display: depth-colored jet_r circles, area=0.7+3*10**(Md-2) points squared; missing Md shown as crosses. Exact publication area law unspecified. Full color ranges: 0–8 km for 1A, 0–6 km for S5. Display bounds and off-view counts saved in routine_plot_scope.csv.. Current catalog snapshot differs from the historical snapshot: 12,503 located study events versus 12,083. No tuning to published counts. Natural Earth coastline is a display substitute; the original relief/inland-zero contour has not been reconstructed.
S6/map	Initial-location spatial views; map	not reproduced. Supplement S17 caption. Requires verified fixed 2008 grid, documented regional averages and validated independent VELEST inversions.. Grid input explicitly authorized; historical transformation and averaging masks not recovered exactly. Local calibration convergence passes, full-path precision and start-model dependence do not. Unreviewed phases and sparse coverage; no accepted relocated scientific catalog. both fixed-grid rounds executed; validation pending
S6/northing-depth	Initial-location spatial views; northing-depth	not reproduced. Supplement S17 caption. Requires verified fixed 2008 grid, documented regional averages and validated independent VELEST inversions.. Grid input explicitly authorized; historical transformation and averaging masks not recovered exactly. Local calibration convergence passes, full-path precision and start-model dependence do not. Unreviewed phases and sparse coverage; no accepted relocated scientific catalog. both fixed-grid rounds executed; validation pending
S6/ easting-depth	Initial-location spatial views; easting-depth	not reproduced. Supplement S17 caption. Requires verified fixed 2008 grid, documented regional averages and validated independent VELEST inversions.. Grid input explicitly authorized; historical transformation and averaging masks not recovered exactly. Local calibration convergence passes, full-path precision and start-model dependence do not. Unreviewed phases and sparse coverage; no accepted relocated scientific catalog. both fixed-grid rounds executed; validation pending
S7A/map	Cluster locations and 2-sigma errors: map and cross-sectional views; map	not reproduced. Supplement S18 caption; S5 methods. Central174, north165, southwest175, east166; HypoDD SVD; Table S1. Grid input explicitly authorized; historical transformation and averaging masks not recovered exactly. Local calibration convergence passes, full-path precision and start-model dependence do not. Unreviewed phases and sparse coverage; no accepted relocated scientific catalog. both fixed-grid rounds executed; validation pending
S7A/ easting-depth	Cluster locations and 2-sigma errors: map and cross-sectional views; easting-depth	not reproduced. Supplement S18 caption; S5 methods. Central174, north165, southwest175, east166; HypoDD SVD; Table S1. Grid input explicitly authorized; historical transformation and averaging masks not recovered exactly. Local calibration convergence passes, full-path precision and start-model dependence do not. Unreviewed phases and sparse coverage; no accepted relocated scientific catalog. both fixed-grid rounds executed; validation pending

Panel/view	Quantity	Status, specification and limitation
S7B	EX uncertainty versus differential-pick count	not reproduced. Supplement S18 caption. Log-log axes; NCCP+NCCS. Grid input explicitly authorized; historical transformation and averaging masks not recovered exactly. Local calibration convergence passes, full-path precision and start-model dependence do not. Unreviewed phases and sparse coverage; no accepted relocated scientific catalog. both fixed-grid rounds executed; validation pending
S7D	EY uncertainty versus differential-pick count	not reproduced. Supplement S18 caption. Log-log axes; NCCP+NCCS. Grid input explicitly authorized; historical transformation and averaging masks not recovered exactly. Local calibration convergence passes, full-path precision and start-model dependence do not. Unreviewed phases and sparse coverage; no accepted relocated scientific catalog. both fixed-grid rounds executed; validation pending
S7F	EZ uncertainty versus differential-pick count	not reproduced. Supplement S18 caption. Log-log axes; NCCP+NCCS. Grid input explicitly authorized; historical transformation and averaging masks not recovered exactly. Local calibration convergence passes, full-path precision and start-model dependence do not. Unreviewed phases and sparse coverage; no accepted relocated scientific catalog. both fixed-grid rounds executed; validation pending
S7C	EX uncertainty density by cluster	not reproduced. Supplement S18 caption. Kernel density estimate; bandwidth unspecified. Grid input explicitly authorized; historical transformation and averaging masks not recovered exactly. Local calibration convergence passes, full-path precision and start-model dependence do not. Unreviewed phases and sparse coverage; no accepted relocated scientific catalog. both fixed-grid rounds executed; validation pending
S7E	EY uncertainty density by cluster	not reproduced. Supplement S18 caption. Kernel density estimate; bandwidth unspecified. Grid input explicitly authorized; historical transformation and averaging masks not recovered exactly. Local calibration convergence passes, full-path precision and start-model dependence do not. Unreviewed phases and sparse coverage; no accepted relocated scientific catalog. both fixed-grid rounds executed; validation pending
S7G	EZ uncertainty density by cluster	not reproduced. Supplement S18 caption. Kernel density estimate; bandwidth unspecified. Grid input explicitly authorized; historical transformation and averaging masks not recovered exactly. Local calibration convergence passes, full-path precision and start-model dependence do not. Unreviewed phases and sparse coverage; no accepted relocated scientific catalog. both fixed-grid rounds executed; validation pending

Panel/view	Quantity	Status, specification and limitation
S8A	Estimated local versus routine duration magnitude	partially reproduced. Supplement S19 caption; S5-S6 methods. 5 s signal, 2 s preceding noise; SNR >=5 dB; all available recorded stations; own picked or predicted P arrivals; >=2 shared stations; separation < one fifth of station distance; squared zero-lag correlation weights; Md regularization lambda=1 for unique one-to-one matches within 2 s. Initial locations. B uses 0.1 magnitude bins, maximum-curvature Mc, half-bin-corrected likelihood b and formal Shi-Bolt sampling error; no arbitrary minimum sample cutoff.. 61681 channel measurements, 4705 passing SNR, 1163 relative-amplitude equations, 119 provisional magnitudes. Final relocated locations/full exposure remain missing; routine Md is an allowed regularizer, not independent validation.
S8B	Magnitude-frequency distribution and fitted b value	partially reproduced. Supplement S19 caption; S6 methods. 5 s signal, 2 s preceding noise; SNR >=5 dB; all available recorded stations; own picked or predicted P arrivals; >=2 shared stations; separation < one fifth of station distance; squared zero-lag correlation weights; Md regularization lambda=1 for unique one-to-one matches within 2 s. Initial locations. B uses 0.1 magnitude bins, maximum-curvature Mc, half-bin-corrected likelihood b and formal Shi-Bolt sampling error; no arbitrary minimum sample cutoff.. 61681 channel measurements, 4705 passing SNR, 1163 relative-amplitude equations, 119 provisional magnitudes. Final relocated locations/full exposure remain missing; routine Md is an allowed regularizer, not independent validation.
S9/map	Filtered-catalog spatial views and feature labels A-D; map	not reproduced. Supplement S20 caption. 38103 events; NCCP+NCCS threshold 100. Unfinished full-period processing and missing calibrated/relocated catalog. Provisional GaMMA diagnostics use an alternative external profile; they are not this panel. Author-derived results are not substituted.
S9/northing-depth	Filtered-catalog spatial views and feature labels A-D; northing-depth	not reproduced. Supplement S20 caption. 38103 events; NCCP+NCCS threshold 100. Unfinished full-period processing and missing calibrated/relocated catalog. Provisional GaMMA diagnostics use an alternative external profile; they are not this panel. Author-derived results are not substituted.
S9/ easting-depth	Filtered-catalog spatial views and feature labels A-D; easting-depth	not reproduced. Supplement S20 caption. 38103 events; NCCP+NCCS threshold 100. Unfinished full-period processing and missing calibrated/relocated catalog. Provisional GaMMA diagnostics use an alternative external profile; they are not this panel. Author-derived results are not substituted.
S10/2022	Annual seismicity map	not reproduced. Supplement S21 caption. Year 2022. Unfinished full-period processing and missing calibrated/relocated catalog. Provisional GaMMA diagnostics use an alternative external profile; they are not this panel. Author-derived results are not substituted.
S10/2023	Annual seismicity map	not reproduced. Supplement S21 caption. Year 2023. Unfinished full-period processing and missing calibrated/relocated catalog. Provisional GaMMA diagnostics use an alternative external profile; they are not this panel. Author-derived results are not substituted.
S10/2024	Annual seismicity map	not reproduced. Supplement S21 caption. Year 2024. Unfinished full-period processing and missing calibrated/relocated catalog. Provisional GaMMA diagnostics use an alternative external profile; they are not this panel. Author-derived results are not substituted.

Panel/view	Quantity	Status, specification and limitation
S10/2025	Annual seismicity map	not reproduced. Supplement S21 caption. Year 2025 through March20. Unfinished full-period processing and missing calibrated/relocated catalog. Provisional GaMMA diagnostics use an alternative external profile; they are not this panel. Author-derived results are not substituted.
S11/swarm-1	Map of one of four major swarms	not reproduced. Supplement S22 caption. 2022-2025; exact swarm windows not enumerated in caption. Unfinished full-period processing and missing calibrated/relocated catalog. Provisional GaMMA diagnostics use an alternative external profile; they are not this panel. Author-derived results are not substituted.
S11/swarm-2	Map of one of four major swarms	not reproduced. Supplement S22 caption. 2022-2025; exact swarm windows not enumerated in caption. Unfinished full-period processing and missing calibrated/relocated catalog. Provisional GaMMA diagnostics use an alternative external profile; they are not this panel. Author-derived results are not substituted.
S11/swarm-3	Map of one of four major swarms	not reproduced. Supplement S22 caption. 2022-2025; exact swarm windows not enumerated in caption. Unfinished full-period processing and missing calibrated/relocated catalog. Provisional GaMMA diagnostics use an alternative external profile; they are not this panel. Author-derived results are not substituted.
S11/swarm-4	Map of one of four major swarms	not reproduced. Supplement S22 caption. 2022-2025; exact swarm windows not enumerated in caption. Unfinished full-period processing and missing calibrated/relocated catalog. Provisional GaMMA diagnostics use an alternative external profile; they are not this panel. Author-derived results are not substituted.
S12A	Ring-sector division	not reproduced. Supplement S23 caption; S8. East, southwest, north; sector boundaries by location/depth. 137 independently associated CPOZ P arrivals now exist. Relocated ring-sector selection and manual depth review remain unfinished; internal S-P checks do not establish location accuracy.
S12B	Ring-sector division in complementary spatial view	not reproduced. Supplement S23 caption; S8. East, southwest, north; sector boundaries by location/depth. 137 independently associated CPOZ P arrivals now exist. Relocated ring-sector selection and manual depth review remain unfinished; internal S-P checks do not establish location accuracy.
S12C	Common-receiver gather: east / HHZ	not reproduced. Supplement S23 caption. CPOZ; -1 to +7 s about P; >4 Hz; north sector ordered by easting, others northing. 137 independently associated CPOZ P arrivals now exist. Relocated ring-sector selection and manual depth review remain unfinished; internal S-P checks do not establish location accuracy.
S12D	Common-receiver gather: southwest / HHZ	not reproduced. Supplement S23 caption. CPOZ; -1 to +7 s about P; >4 Hz; north sector ordered by easting, others northing. 137 independently associated CPOZ P arrivals now exist. Relocated ring-sector selection and manual depth review remain unfinished; internal S-P checks do not establish location accuracy.
S12E	Common-receiver gather: north / HHZ	not reproduced. Supplement S23 caption. CPOZ; -1 to +7 s about P; >4 Hz; north sector ordered by easting, others northing. 137 independently associated CPOZ P arrivals now exist. Relocated ring-sector selection and manual depth review remain unfinished; internal S-P checks do not establish location accuracy.

Panel/view	Quantity	Status, specification and limitation
S12F	Common-receiver gather: east / HHN	not reproduced. Supplement S23 caption. CPOZ; -1 to +7 s about P; >4 Hz; north sector ordered by easting, others northing. 137 independently associated CPOZ P arrivals now exist. Relocated ring-sector selection and manual depth review remain unfinished; internal S-P checks do not establish location accuracy.
S12G	Common-receiver gather: southwest / HHN	not reproduced. Supplement S23 caption. CPOZ; -1 to +7 s about P; >4 Hz; north sector ordered by easting, others northing. 137 independently associated CPOZ P arrivals now exist. Relocated ring-sector selection and manual depth review remain unfinished; internal S-P checks do not establish location accuracy.
S12H	Common-receiver gather: north / HHN	not reproduced. Supplement S23 caption. CPOZ; -1 to +7 s about P; >4 Hz; north sector ordered by easting, others northing. 137 independently associated CPOZ P arrivals now exist. Relocated ring-sector selection and manual depth review remain unfinished; internal S-P checks do not establish location accuracy.
S13A	Full caldera seismicity on satellite reference	not reproduced. Supplement S24 caption. Depth-coded relocated events; Google Earth basemap. Unfinished full-period processing and missing calibrated/relocated catalog. Provisional GaMMA diagnostics use an alternative external profile; they are not this panel. Author-derived results are not substituted.
S13B	Onshore seismicity on satellite reference	not reproduced. Supplement S24 caption. Depth-coded relocated events; Google Earth basemap. Unfinished full-period processing and missing calibrated/relocated catalog. Provisional GaMMA diagnostics use an alternative external profile; they are not this panel. Author-derived results are not substituted.
S14/1	CAAM waveforms and event annotations	not reproduced. Supplement S25 caption; S9-S11. 1-10 Hz display; 8-s FI at 2-5 and 8-15 Hz; FI<0 colored red. 75.797 common CAAM/COLB scan hours; 822 unconfirmed threshold crossings. Three-year scan, reviewed clusters, relative magnitudes and timing histories remain unfinished.
S14/2	CAAM waveforms and event annotations	not reproduced. Supplement S25 caption; S9-S11. 1-10 Hz display; 8-s FI at 2-5 and 8-15 Hz; FI<0 colored red. 75.797 common CAAM/COLB scan hours; 822 unconfirmed threshold crossings. Three-year scan, reviewed clusters, relative magnitudes and timing histories remain unfinished.
S14/3	CAAM waveforms and event annotations	not reproduced. Supplement S25 caption; S9-S11. 1-10 Hz display; 8-s FI at 2-5 and 8-15 Hz; FI<0 colored red. 75.797 common CAAM/COLB scan hours; 822 unconfirmed threshold crossings. Three-year scan, reviewed clusters, relative magnitudes and timing histories remain unfinished.
S14/ unclustered	CAAM waveforms and event annotations	not reproduced. Supplement S25 caption; S9-S11. 1-10 Hz display; 8-s FI at 2-5 and 8-15 Hz; FI<0 colored red. 75.797 common CAAM/COLB scan hours; 822 unconfirmed threshold crossings. Three-year scan, reviewed clusters, relative magnitudes and timing histories remain unfinished.
S15A	Hybrid family 1: waveforms and spectrograms	not reproduced. Supplement S26 caption; S9-S10. CAAM/COLB HHZ/HHE; amplitudes normalized by CAAM maximum. Post-generation visual comparison shows a different published example from the written-method template used here. The generated family-template figure is related analysis, not reproduction of this panel.

Panel/view	Quantity	Status, specification and limitation
S15B	Hybrid family 2: waveforms and spectrograms	not reproduced. Supplement S26 caption; S9-S10. CAAM/COLB HHZ/HHE; amplitudes normalized by CAAM maximum. Post-generation visual comparison shows a different published example from the written-method template used here. The generated family-template figure is related analysis, not reproduction of this panel.
S15C	Hybrid family 3: waveforms and spectrograms	partially reproduced. Supplement S26 caption; S9-S10. CAAM/COLB HHZ/HHE; amplitudes normalized by CAAM maximum. Written template times used; exact published example alignment, display filtering and spectral settings incompletely specified. Sensitivity-corrected velocities and stated independent display settings used.
S15D	Shallow volcano-tectonic example: waveforms and spectrograms	not reproduced. Supplement S26 caption; S9-S10. CAAM/COLB HHZ/HHE; amplitudes normalized by CAAM maximum. Exact shallow VT example not identified by written method alone; no author catalog or plot values used to select it.

Executed partial-panel chains

The following commands generate the saved numerical files and their plotted counterparts. The full input lists and checksums are retained in the machine-readable records; no unreproduced panel has an executed numerical counterpart. The source-to-output checks cover the explicit current graph and do not prove the absence of unexamined or hidden inputs.

1A — partial data-to-plot chain. External routine-catalog context; not independent detections.

22 exact input files are listed under this panel in `docs/panel_record.json`; waveform hashes and intervals are in `docs/continuation_processed_inputs.json`, with source receipts in the input manifests.

Data command:

```
.venv/bin/pythonscripts/continue_context.py
```

Saved numerical output: `data/continuation/figure_1A.csv`.

Plot command: `.venv/bin/pythonscripts/publication_plots.py`

Output: `figures/figure_1A_routine.pdf`.

The same plotting command composes the report figure: `figures/figure_1_combined.pdf`.

1B — partial data-to-plot chain. Original GNSS/leveling observations plus explicitly labeled historical source estimates/reconstruction.

Inputs: `inputs/continuation/RITE_2000_2025.txt`, `inputs/continuation/RITE_Timeseries_Format.readme`, `inputs/continuation/RITE_daily.txt`, `inputs/continuation/Time_Series_Format.txt`, `inputs/extended/delgaudio2010_table.doc`.

Data command:

```
.venv/bin/pythonscripts/continue_context.py
```

Saved numerical output: `data/continuation/rite_gnss.csv`, `data/continuation/historical_leveling.csv`.

Plot command: `.venv/bin/pythonscripts/publication_plots.py`

Output: `figures/figure_1B_displacement.pdf`.

The same plotting command composes the report figure: `figures/figure_1_combined.pdf`.

1C — partial data-to-plot chain. External routine-catalog context; not independent detections.

21 exact input files are listed under this panel in `docs/panel_record.json`; waveform hashes and intervals are in `docs/continuation_processed_inputs.json`, with source receipts in the input manifests.

Data command:

`.venv/bin/pythonscripts/continue_context.py`

Saved numerical output: `data/continuation/figure_1C.csv`, `data/continuation/figure_1C_cumulative.csv`.

Plot command: `.venv/bin/pythonscripts/publication_plots.py`

Output: `figures/figure_1C_routine.pdf`.

The same plotting command composes the report figure: `figures/figure_1_combined.pdf`.

S1 — partial data-to-plot chain. Own processing of observations/metadata; trained weights and external Md calibration are identified in the method record.

868 exact input files are listed under this panel in `docs/panel_record.json`; waveform hashes and intervals are in `docs/continuation_processed_inputs.json`, with source receipts in the input manifests.

Data command:

`.venv/bin/pythonscripts/extend_calibration_observations.py--workers8`

Saved numerical output: `data/calibration_followup/expanded/processed_coverage_union.json`.

Plot command: `.venv/bin/pythonscripts/publication_plots.py`

Output: `figures/followup_coverage.pdf`.

S2 — partial data-to-plot chain. Own processing of observations/metadata; trained weights and external Md calibration are identified in the method record.

22 exact input files are listed under this panel in `docs/panel_record.json`; waveform hashes and intervals are in `docs/continuation_processed_inputs.json`, with source receipts in the input manifests.

Data command:

`.venv/bin/pythonscripts/station_geometry.py`

Saved numerical output: `data/continuation/figure_S2.csv`, `data/continuation/station_geometry_qa.csv`, `data/continuation/station_geometry_epochs.csv`.

Plot command: `.venv/bin/pythonscripts/publication_plots.py`

Output: `figures/continuation_stations.pdf`.

S4A — partial data-to-plot chain. Own processing of observations/metadata; trained weights and external Md calibration are identified in the method record.

Inputs: `inputs/resumption/S4_CSOB_20240520.mseed`, `inputs/resumption/s4_anchor.html`.

Data command:

`.venv/bin/pythonscripts/benchmark_window.py`

Saved numerical output: `data/continuation/figure_S4A.npz`, `data/continuation/figure_S4A_picks.json`.

Plot command: `.venv/bin/pythonscripts/benchmark_window.py`

Output: `figures/figure_S4A.pdf`.

S5/map — partial data-to-plot chain. External routine-catalog context; not independent detections.

22 exact input files are listed under this panel in `docs/panel_record.json`; waveform hashes and intervals are in `docs/continuation_processed_inputs.json`, with source receipts in the input manifests.

Data command:

`.venv/bin/pythonscripts/continue_context.py`

Saved numerical output: `data/continuation/figure_S5.csv`.

Plot command: `.venv/bin/pythonscripts/publication_plots.py`

Output: `figures/figure_S5_routine.pdf`.

S5/northing-depth — partial data-to-plot chain. External routine-catalog context; not independent detections.

22 exact input files are listed under this panel in `docs/panel_record.json`; waveform hashes and intervals are in `docs/continuation_processed_inputs.json`, with source receipts in the input manifests.

Data command:

.venv/bin/pythonscripts/continue_context.py

Saved numerical output: data/continuation/figure_S5.csv.

Plot command: .venv/bin/pythonscripts/publication_plots.py

Output: figures/figure_S5_routine.pdf.

S5/easting-depth — partial data-to-plot chain. External routine-catalog context; not independent detections.

22 exact input files are listed under this panel in docs/panel_record.json; waveform hashes and intervals are in docs/continuation_processed_inputs.json, with source receipts in the input manifests.

Data command:

.venv/bin/pythonscripts/continue_context.py

Saved numerical output: data/continuation/figure_S5.csv.

Plot command: .venv/bin/pythonscripts/publication_plots.py

Output: figures/figure_S5_routine.pdf.

S8A — partial data-to-plot chain. Own processing of observations/metadata; trained weights and external Md calibration are identified in the method record.

848 exact input files are listed under this panel in docs/panel_record.json; waveform hashes and intervals are in docs/continuation_processed_inputs.json, with source receipts in the input manifests.

Data command:

.venv/bin/pythonscripts/downstream_checks.py--datasetcontinuation

.venv/bin/pythonscripts/magnitude_statistics.py

Saved numerical output: data/continuation/figure_S8A.csv.

Plot command: .venv/bin/pythonscripts/publication_plots.py

Output: figures/figure_S8_pilot.pdf.

S8B — partial data-to-plot chain. Own processing of observations/metadata; trained weights and external Md calibration are identified in the method record.

848 exact input files are listed under this panel in docs/panel_record.json; waveform hashes and intervals are in docs/continuation_processed_inputs.json, with source receipts in the input manifests.

Data command:

.venv/bin/pythonscripts/downstream_checks.py--datasetcontinuation

.venv/bin/pythonscripts/magnitude_statistics.py

Saved numerical output: data/continuation/figure_S8B.csv, data/continuation/magnitude_statistics.json.

Plot command: .venv/bin/pythonscripts/publication_plots.py

Output: figures/figure_S8_pilot.pdf.

S15C — partial data-to-plot chain. Own processing of observations/metadata; trained weights and external Md calibration are identified in the method record.

Inputs: inputs/metadata/response.xml, inputs/waveforms/E3_CAAM.mseed, inputs/waveforms/E3_COLB.mseed.

Data command:

.venv/bin/pythonscripts/analyze_observations.py

Saved numerical output: data/observations/E3_waveforms.csv, data/observations/E3_CAAM_Z_spectrum.npz, data/observations/E3_CAAM_E_spectrum.npz, data/observations/E3_COLB_Z_spectrum.npz, data/observations/E3_COLB_E_spectrum.npz.

Plot command: .venv/bin/pythonscripts/publication_plots.py

Output: figures/hybrid_E3.pdf.

D8 Saved numerical tables and equation consistency checks

Original-waveform template frequency indices

These retained numerical measurements describe written-method templates, not confirmed hybrid families. E1 and E2 are not substituted for the different published S15A/B examples.

Template	FI	Samples
E1	-0.7862	800
E2	-0.4286	800
E3	-1.0819	800

Rejected velocity experiments

The prior grid-dependent regional calibration is archived, not accepted. The separate `scripts/velest_stability.py` experiment starts from the cited Tramelli (2021) numerical table. It runs identical serialized observations and damping with 32-bit and 64-bit real arithmetic. Raw model outputs, fitted-location files and physical-velocity checks are saved under `data/exploratory/velest_stability`. Even precision agreement cannot establish a correct physical model; any $V_p \leq V_s$ layer rejects the fit. These results do not supply S6, Table S1, or downstream relocation. The following unaccepted experimental values come directly from `model_32.csv` and `model_64.csv` in that directory, generated by `.venv/bin/pythonscripts/velest_stability.py`. Velocities are in km/s. This is a reproduction diagnostic table, not a counterpart to Table S1.

Layer top (km b.s.l.)	V_p, 32 bit	V_s, 32 bit	V_p, 64 bit	V_s, 64 bit
-0.50	1.36	0.90	1.36	0.90
0.50	2.15	1.11	2.15	1.11
1.00	2.39	1.58	2.39	1.58
3.00	5.31	3.24	5.31	3.24
10.00	5.70	3.20	5.70	3.20

D9 Rebuild and replacements

The preceding alternative-model follow-up procedure was `.venv/bin/pythonscripts/rebuild_calibration_followup.py`; its executed stages and outcome are recorded in `data/calibration_followup/rebuild_status.json`. It reuses the hash-verified expanded inference/association checkpoint by default; `--fresh-association` reruns that stage. It reruns calibration, differential measurements, exploratory HypoDD, validation, plotting and both document builds. The earlier full-subset procedure remains available below. Neither procedure establishes an accepted scientific calibration.

The current source audit re-downloaded the RITE daily text and format notes, Del Gaudio numerical table and authorized Battaglia article; their hashes match the preceding inputs. CPIS January 2022 requests returned HTTP 204 even for one-minute/one-component and thirty-minute retries, despite nominal availability. Its retrieved HH response epochs begin in November 2023. This discrepancy is recorded without counting the intervals as coverage. Public RAN login and registration pages remain accessible; authenticated waveform access is unavailable here. Source receipts and the additional source bundle are documented in `docs/resumption_sources.json` and `docs/resumption_rebuild_inputs.`

json.

Six pre-expansion hybrid scan examples were visually inspected from saved observed waveforms: high mean-correlation examples included stationary oscillations or weak counterparts at the second station; high signal/noise examples included clear impulsive body waves. None establishes a hybrid family. The selection and review are preserved in data/resumption/baseline_hybrid_review/visual_review.json. New scan tables retain individual CAAM and COLB scores. Review examples remain diagnostics and are excluded from publication-panel comparisons.

The association pass was deliberately stopped after the worker-dependent initialization issue was identified. Compatible waveform inference was retained; association and all dependent stages were rerun with deterministic cluster seeds. The interruption and hash-verified resumption are recorded, rather than counted as a successful first attempt.

The amplitude stage uses sixteen spawned workers and bounded caches of unchanged input waveforms; each measurement window is copied before processing. A software check compares 60 observed windows against direct reads and requires identical samples and timestamps. An earlier amplitude run was intentionally interrupted to avoid repeated decoding. This performance restart was not an access failure; completed inference and association were retained only after their recorded hashes were verified. A previous measurement pass reached the inversion-output step but failed when JSON serialization encountered a NumPy Boolean. The acceptance flag was converted explicitly to a Python Boolean, and the full failed stage was rerun from observations. The failed attempt's saved numerical hashes are preserved in docs/magnitude_serialization_failure.json; it is not counted as a passed build. Six downstream CSVs, including the magnitude equations and estimates, are byte-identical between that failed pass and the successful rerun. The amplitude CSV hash differs. Because its pre-failure bytes were not retained, the numerical or ordering cause of that difference cannot be resolved from the hash alone; no claim of byte-for-byte amplitude reproducibility is made. That successful historical rerun was the preceding checkpoint; current measurements are regenerated from the expanded frozen input set. The current authorized-grid rebuild is scripts/rebuild_grid_workflow.py; it regenerates the new trials and the styled deliverables from the verified expanded processing checkpoint. The following paragraph describes the preceding frozen-checkpoint rebuild, retained as history. That historical continuation procedure is documented in docs/CONTINUATION.md. Restore the frozen inputs with scripts/restore_inputs.py--continuation; run scripts/rebuild_continuation.py using the repository's Python environment. The optional --fresh flag also recomputes all earlier baseline inference. The rebuild regenerates the current association, downstream attempts, context panels and both PDFs; only compatible per-file inference caches and the verified baseline may be reused. It is a rebuild of the frozen observed subset, not proof of complete multiyear reproduction. Its actual outcome and executed commands are saved in data/continuation/rebuild_status.json and data/continuation/rebuild_log.json. The preceding fresh rebuild stopped because this Python installation lacks SQLite's in-memory deserialization method. The ledger reader was changed to a temporary project-local database; missing request segment fields also required explicit null handling. Passed stages were hash-verified before resumption. Earlier worker and study-boundary failures belong to superseded runs and are not evidence of unavailable observations. The current rebuild mode and outcome are stated in the saved status file; matching inference caches may be reused, and subsequent numerical stages execute again. The document build verifies main-text sections, the eligible generated figures in original numbering order, supplement topic order, and all nineteen comparison pages in original order, and confirms that published images occur only in the comparison PDF.

This continuation replaces the prior report/comparison PDFs and generated figure fragments, panel records and obsolete context figure usage. The old displacement display ending in 2019 is replaced by observed daily coverage through March 2025. The old context plots are superseded by the corresponding Figure 1 and S5 outputs; their scripts/data remain traceable, while their obsolete PDFs/PNGs are

removed after the current counterparts exist. New seismic outputs are in `data/continuation`. Eligible baseline observations and picks remain in `data/extended`; grid-dependent baseline results and obsolete comparison statistics are in `quarantine`. Synthetic fixtures are isolated in `tests/results`. Exact changes are documented in `docs/REPLACEMENTS.md`. Project-specific caches remain under `tmp`; pip's normal cache is `/home/mehffin/.cache/pip`. Large waveforms and model/software artifacts remain ignored under `inputs`, with retrieval URLs, file sizes and hashes.

D10 Sources

The primary specification is Tan et al. (2025), Science 390, 70–75, local `tanmodel.pdf` and `tanmodelsupp.pdf`. Observation/metadata services: https://webservices.ingv.it/fdsnws/; public RAN station metadata: https://ran.protezionecivile.it/IT/stazioni.php; temporary waveform archive: https://dataserver.ov.ingv.it/users/anna.tramelli/seismic_records/index.html; routine catalog: https://terremoti.ov.ingv.it/gossip/flegrei/. GNSS: De Martino and colleagues, https://zenodo.org/records/17132077 and https://zenodo.org/records/22259337, with the requested citation to De Martino et al. (2021), doi:10.3390/rs13142725. Historical leveling: Del Gaudio et al. (2010), doi:10.1016/j.jvolgeores.2010.05.014. Published alternative profile: Tramelli et al. (2021), doi:10.1038/s41598-021-86506-6, Table 1. Unavailable exact profile: Calò and Tramelli (2018), doi:10.1038/s41598-018-34456-x. Quarantined, unverified grid attribution: Battaglia et al. (2008), doi:10.1111/j.1365-2478.2007.00687.x.

Tan and colleagues' released trained PhaseNet artifact is attributed to https://github.com/iamharrison2000/Campi-Flegrei-Earthquake-Monitoring. Only classified weights, configuration and station metadata enter the current workflow. The permitted artifact revision is

`f84c9c9dc6bb1501a95a833f6634199a63d4f3b3`.

GaMMA revision is

`80394dd4a4c29450da599f507916ecf1db6203dd`.