

Operations on Fuzzy Sets

Murat Osmanoglu

Fuzzy Operations

Standart fuzzy operations

- Complement, $\mu_C(x) = 1 - \mu_A(x)$ where $C = \neg A$

Intersection, $\mu_C(x) = \min\{\mu_A(x), \mu_B(x)\}$ where $C = A \cap B$

Union, $\mu_C(x) = \max\{\mu_A(x), \mu_B(x)\}$ where $C = A \cup B$

Fuzzy Operations

Standart fuzzy operations

- Complement, $\mu_C(x) = 1 - \mu_A(x)$ where $C = \neg A$
Intersection, $\mu_C(x) = \min\{\mu_A(x), \mu_B(x)\}$ where $C = A \cap B$
Union, $\mu_C(x) = \max\{\mu_A(x), \mu_B(x)\}$ where $C = A \cup B$
- standart fuzzy operations are generalizations of the corresponding classical set operations

Fuzzy Operations

Standart fuzzy operations

- Complement, $\mu_C(x) = 1 - \mu_A(x)$ where $C = \neg A$
Intersection, $\mu_C(x) = \min\{\mu_A(x), \mu_B(x)\}$ where $C = A \cap B$
Union, $\mu_C(x) = \max\{\mu_A(x), \mu_B(x)\}$ where $C = A \cup B$
- standart fuzzy operations are generalizations of the corresponding classical set operations
- a desirable feature of the standard operations is their inherent prevention of the compounding of errors of operands
if e is the error associated with $\mu_A(x)$ and $\mu_B(x)$, then maximum error associated with $\mu_C(x)$ remains e where $C = \neg A$, or $A \cap B$, or $A \cup B$

Fuzzy Operations

Standart fuzzy operations

- Complement, $\mu_C(x) = 1 - \mu_A(x)$ where $C = \neg A$
Intersection, $\mu_C(x) = \min\{\mu_A(x), \mu_B(x)\}$ where $C = A \cap B$
Union, $\mu_C(x) = \max\{\mu_A(x), \mu_B(x)\}$ where $C = A \cup B$
- standart fuzzy operations are generalizations of the corresponding classical set operations
- a desirable feature of the standard operations is their inherent prevention of the compounding of errors of operands
if e is the error associated with $\mu_A(x)$ and $\mu_B(x)$, then maximum error associated with $\mu_C(x)$ remains e where $C = \neg A$, or $A \cap B$, or $A \cup B$
- for each operation, there exist a broad class of functions where each of them considered a class of operations and **characterized by justified axioms**

Fuzzy Operations

Standart fuzzy operations

- Complement, $\mu_C(x) = 1 - \mu_A(x)$ where $C = \neg A$
Intersection, $\mu_C(x) = \min\{\mu_A(x), \mu_B(x)\}$ where $C = A \cap B$
Union, $\mu_C(x) = \max\{\mu_A(x), \mu_B(x)\}$ where $C = A \cup B$
- standart fuzzy operations are generalizations of the corresponding classical set operations
- a desirable feature of the standard operations is their inherent prevention of the compounding of errors of operands
if e is the error associated with $\mu_A(x)$ and $\mu_B(x)$, then maximum error associated with $\mu_C(x)$ remains e where $C = \neg A$, or $A \cap B$, or $A \cup B$
- for each operation, there exist a broad class of functions where each of them considered a class of operations and **characterized by justified axioms**
- different functions can be used in different contexts

Fuzzy Operations

Fuzzy Complement

- Let a fuzzy complement operation be defined as a function

$$c : [0,1] \rightarrow [0,1]$$

$$\mu_A(x) \rightarrow c(\mu_A(x))$$

- to produce fuzzy sets that qualify as meaningful complements of given fuzzy sets, the functions c must satisfy certain properties

Fuzzy Operations

Fuzzy Complement

- Let a fuzzy complement operation be defined as a function

$$c : [0,1] \rightarrow [0,1]$$

$$\mu_A(x) \rightarrow c(\mu_A(x))$$

- to produce fuzzy sets that qualify as meaningful complements of given fuzzy sets, the functions c must satisfy certain properties

$\mu_A(x)$ specifies the degree of which x belongs to A ,
 $\mu_C(x)$ specifies the degree of which x does not belong to A or
 x belongs to C where $C = \neg A$

Fuzzy Operations

Fuzzy Complement

- Let a fuzzy complement operation be defined as a function

$$c : [0,1] \rightarrow [0,1]$$

$$\mu_A(x) \rightarrow c(\mu_A(x))$$

- to produce fuzzy sets that qualify as meaningful complements of given fuzzy sets, the functions c must satisfy certain properties
- **Axiom 1** : $c(0) = 1, c(1) = 0$ (boundary condition)

Fuzzy Operations

Fuzzy Complement

- Let a fuzzy complement operation be defined as a function

$$c : [0,1] \rightarrow [0,1]$$

$$\mu_A(x) \rightarrow c(\mu_A(x))$$

- to produce fuzzy sets that qualify as meaningful complements of given fuzzy sets, the functions c must satisfy certain properties
- **Axiom 1** : $c(0) = 1, c(1) = 0$ (boundary condition)
- **Axiom 2** : for all a, b in $[0,1]$, if $a \leq b$, then $c(a) \geq c(b)$ (monotonicity)

Fuzzy Operations

Fuzzy Complement

- Let a fuzzy complement operation be defined as a function

$$c : [0,1] \rightarrow [0,1]$$

$$\mu_A(x) \rightarrow c(\mu_A(x))$$

- to produce fuzzy sets that qualify as meaningful complements of given fuzzy sets, the functions c must satisfy certain properties
- **Axiom 1** : $c(0) = 1, c(1) = 0$ (boundary condition)
- **Axiom 2** : for all a, b in $[0,1]$, if $a \leq b$, then $c(a) \geq c(b)$ (monotonicity)
- Axiom 1 required to get correct complements for crisp sets (since we stated that the fuzzy operations are generalizations of corresponding crisp operations)

Fuzzy Operations

Fuzzy Complement

- Let a fuzzy complement operation be defined as a function

$$c : [0,1] \rightarrow [0,1]$$

$$\mu_A(x) \rightarrow c(\mu_A(x))$$

- to produce fuzzy sets that qualify as meaningful complements of given fuzzy sets, the functions c must satisfy certain properties
- Axiom 1** : $c(0) = 1, c(1) = 0$ (boundary condition)
- Axiom 2** : for all a, b in $[0,1]$, if $a \leq b$, then $c(a) \geq c(b)$ (monotonicity)
- Axiom 1 required to get correct complements for crisp sets (since we stated that the fuzzy operations are generalizations of corresponding crisp operations)
- Axiom 2 required to be monotonic decreasing, i.e. when μ_A increases, the complement $c(\mu_A)$ must not increase (it may decrease or, at least remain same).

Fuzzy Operations

Fuzzy Complement

- to consider a function as a fuzzy complement operation, it must satisfy the first two axioms (axiomatic skeleton)

Fuzzy Operations

Fuzzy Complement

- to consider a function as a fuzzy complement operation, it must satisfy the first two axioms (axiomatic skeleton)
- each additional axiom reduces the general class to a special subclass

Fuzzy Operations

Fuzzy Complement

- to consider a function as a fuzzy complement operation, it must satisfy the first two axioms (axiomatic skeleton)
- each additional axiom reduces the general class to a special subclass
- **Axiom 3** : c is a continuous function

Fuzzy Operations

Fuzzy Complement

- to consider a function as a fuzzy complement operation, it must satisfy the first two axioms (axiomatic skeleton)
- each additional axiom reduces the general class to a special subclass
- **Axiom 3** : c is a continuous function

- a function is continuous at some x_0 in D , if f is defined at x_0 , the limit of the function exist x_0 , and equals to $f(x_0)$
- the graph of a continuous function is a single unbroken curve with no holes or jumps

Fuzzy Operations

Fuzzy Complement

- to consider a function as a fuzzy complement operation, it must satisfy the first two axioms (axiomatic skeleton)
- each additional axiom reduces the general class to a special subclass
- **Axiom 3** : c is a continuous function
- **Axiom 4** : c is involutive, i.e. $c(c(a)) = a$ for each a in $[0,1]$

Fuzzy Operations

Fuzzy Complement

- to consider a function as a fuzzy complement operation, it must satisfy the first two axioms (axiomatic skeleton)
- each additional axiom reduces the general class to a special subclass
- **Axiom 3** : c is a continuous function
- **Axiom 4** : c is involutive, i.e. $c(c(a)) = a$ for each a in $[0,1]$
- Standart Complement ; $c(a) = 1 - a$ where $a = \mu_A(x)$ in $[0,1]$

Fuzzy Operations

Fuzzy Complement

- to consider a function as a fuzzy complement operation, it must satisfy the first two axioms (axiomatic skeleton)
- each additional axiom reduces the general class to a special subclass
- **Axiom 3** : c is a continuous function
- **Axiom 4** : c is involutive, i.e. $c(c(a)) = a$ for each a in $[0,1]$
- Standart Complement ; $c(a) = 1 - a$ where $a = \mu_A(x)$ in $[0,1]$
 - $c(0) = 1 - 0 = 1$ and $c(1) = 1 - 1 = 0$

Fuzzy Operations


Fuzzy Complement

- to consider a function as a fuzzy complement operation, it must satisfy the first two axioms (axiomatic skeleton)
- each additional axiom reduces the general class to a special subclass
- **Axiom 3** : c is a continuous function
- **Axiom 4** : c is involutive, i.e. $c(c(a)) = a$ for each a in $[0,1]$
- Standart Complement ; $c(a) = 1 - a$ where $a = \mu_A(x)$ in $[0,1]$
 - $c(0) = 1 - 0 = 1$ and $c(1) = 1 - 1 = 0$
 - assume $a \leq b$, then $1 - a \geq 1 - b$, and $c(a) \geq c(b)$

Fuzzy Operations

Fuzzy Complement


- to consider a function as a fuzzy complement operation, it must satisfy the first two axioms (axiomatic skeleton)
- each additional axiom reduces the general class to a special subclass
- **Axiom 3** : c is a continuous function
- **Axiom 4** : c is involutive, i.e. $c(c(a)) = a$ for each a in $[0,1]$
- Standart Complement ; $c(a) = 1 - a$ where $a = \mu_A(x)$ in $[0,1]$
 - $c(0) = 1 - 0 = 1$ and $c(1) = 1 - 1 = 0$
 - assume $a \leq b$, then $1 - a \geq 1 - b$, and $c(a) \geq c(b)$



Fuzzy Operations

Fuzzy Complement


- to consider a function as a fuzzy complement operation, it must satisfy the first two axioms (axiomatic skeleton)
- each additional axiom reduces the general class to a special subclass
- **Axiom 3** : c is a continuous function
- **Axiom 4** : c is involutive, i.e. $c(c(a)) = a$ for each a in $[0,1]$
- Standart Complement ; $c(a) = 1 - a$ where $a = \mu_A(x)$ in $[0,1]$
 - $c(0) = 1 - 0 = 1$ and $c(1) = 1 - 1 = 0$
 - assume $a \leq b$, then $1 - a \geq 1 - b$, and $c(a) \geq c(b)$



Fuzzy Operations

Fuzzy Complement


- to consider a function as a fuzzy complement operation, it must satisfy the first two axioms (axiomatic skeleton)
- each additional axiom reduces the general class to a special subclass
- **Axiom 3** : c is a continuous function
- **Axiom 4** : c is involutive, i.e. $c(c(a)) = a$ for each a in $[0,1]$
- Standart Complement ; $c(a) = 1 - a$ where $a = \mu_A(x)$ in $[0,1]$
 - $c(0) = 1 - 0 = 1$ and $c(1) = 1 - 1 = 0$
 - assume $a \leq b$, then $1 - a \geq 1 - b$, and $c(a) \geq c(b)$
 - c is continuous



Fuzzy Operations


Fuzzy Complement

- to consider a function as a fuzzy complement operation, it must satisfy the first two axioms (axiomatic skeleton)
- each additional axiom reduces the general class to a special subclass
- **Axiom 3** : c is a continuous function
- **Axiom 4** : c is involutive, i.e. $c(c(a)) = a$ for each a in $[0,1]$
- Standart Complement ; $c(a) = 1 - a$ where $a = \mu_A(x)$ in $[0,1]$
 - $c(0) = 1 - 0 = 1$ and $c(1) = 1 - 1 = 0$
 - assume $a \leq b$, then $1 - a \geq 1 - b$, and $c(a) \geq c(b)$
 - c is continuous
 - $c(c(a)) = c(1 - a) = 1 - (1 - a) = a$



Fuzzy Operations


Fuzzy Complement



$$c(a) = \begin{cases} 1 & \text{if } a \leq z \\ 0 & \text{if } a > z \end{cases}$$

Fuzzy Operations


Fuzzy Complement



$$c(a) = \begin{cases} 1 & \text{if } a \leq z \\ 0 & \text{if } a > z \end{cases}$$

Fuzzy Operations

Fuzzy Complement




$$c(a) = \begin{cases} 1 & \text{if } a \leq z \\ 0 & \text{if } a > z \end{cases}$$

– $c(0) = 1$ and $c(1) = 0$

Fuzzy Operations

Fuzzy Complement




$$c(a) = \begin{cases} 1 & \text{if } a \leq z \\ 0 & \text{if } a > z \end{cases}$$

- $c(0) = 1$ and $c(1) = 0$
- assume $a \leq b \leq z$, then $c(a) \geq c(b)$; or $z \leq a \leq b$, then $c(a) \geq c(b)$;
or $a \leq z \leq b$, then $c(a) = 1 \geq c(b) = 0$

Fuzzy Operations

Fuzzy Complement




$$c(a) = \begin{cases} 1 & \text{if } a \leq z \\ 0 & \text{if } a > z \end{cases}$$

- $c(0) = 1$ and $c(1) = 0$
- assume $a \leq b \leq z$, then $c(a) \geq c(b)$; or $z \leq a \leq b$, then $c(a) \geq c(b)$;
or $a \leq z \leq b$, then $c(a) = 1 \geq c(b) = 0$
- c is not continuous

Fuzzy Operations

Fuzzy Complement



$$c(a) = \begin{cases} 1 & \text{if } a \leq z \\ 0 & \text{if } a > z \end{cases}$$

- $c(0) = 1$ and $c(1) = 0$
- assume $a \leq b \leq z$, then $c(a) \geq c(b)$; or $z \leq a \leq b$, then $c(a) \geq c(b)$;
or $a \leq z \leq b$, then $c(a) = 1 \geq c(b) = 0$
- c is not continuous
- assume a in $(0,1)$, then $c(c(a)) \neq a$


Fuzzy Operations

Fuzzy Complement

Yager Class

$$c_w(a) = (1 - a^w)^{1/w}$$


where w in $(0, \infty)$



Sugeno Class

$$c(a) = (1 - a) / (1 + \lambda a)$$

where λ in $(-1, \infty)$



Fuzzy Operations

Fuzzy Intersection (t-norm)

- Let a fuzzy intersection operation be defined as a function

$$i : [0,1] \times [0,1] \rightarrow [0,1]$$

$$(\mu_A(x), \mu_B(x)) \rightarrow i(\mu_A(x), \mu_B(x))$$

Fuzzy Operations

Fuzzy Intersection (t-norm)

- Let a fuzzy intersection operation be defined as a function

$$i : [0,1] \times [0,1] \rightarrow [0,1]$$

$$(\mu_A(x), \mu_B(x)) \rightarrow i(\mu_A(x), \mu_B(x))$$

- Axiom 1** : $i(1,1) = 1, i(1,0) = 0, i(0,1) = 0, i(0,0) = 0$ (boundary condition)
- Axiom 2** : $i(a,b) = i(b,a)$ (comutativity)
- Axiom 3** : if $b \leq d$, then $i(a,b) \leq i(a,d)$ (monotonicity)
- Axiom 4** : $i(i(a,b),c) = i(a,i(b,c))$ (associativity)

Fuzzy Operations

Fuzzy Intersection (t-norm)

- Let a fuzzy intersection operation be defined as a function

$$i : [0,1] \times [0,1] \rightarrow [0,1]$$

$$(\mu_A(x), \mu_B(x)) \rightarrow i(\mu_A(x), \mu_B(x))$$

- Axiom 1** : $i(1,1) = 1, i(1,0) = 0, i(0,1) = 0, i(0,0) = 0$ (boundary condition)
- Axiom 2** : $i(a,b) = i(b,a)$ (comutativity)
- Axiom 3** : if $b \leq d$, then $i(a,b) \leq i(a,d)$ (monotonicity)
- Axiom 4** : $i(i(a,b),c) = i(a,i(b,c))$ (associativity)
- these axioms ensure that the fuzzy intersection becomes the classical intersection when A and B are crisp

Fuzzy Operations

Fuzzy Intersection (t-norm)

- Let a fuzzy intersection operation be defined as a function

$$i : [0,1] \times [0,1] \rightarrow [0,1]$$

$$(\mu_A(x), \mu_B(x)) \rightarrow i(\mu_A(x), \mu_B(x))$$

- Axiom 1** : $i(1,1) = 1, i(1,0) = 0, i(0,1) = 0, i(0,0) = 0$ (boundary condition)
- Axiom 2** : $i(a,b) = i(b,a)$ (commutativity)
- Axiom 3** : if $b \leq d$, then $i(a,b) \leq i(a,d)$ (monotonicity)
- Axiom 4** : $i(i(a,b),c) = i(a,i(b,c))$ (associativity)
- these axioms ensure that the fuzzy intersection becomes the classical intersection when A and B are crisp
- monotonicity and commutativity express that a decrease in the degree of membership in sets A and B cannot produce an increase in the degree of membership of intersection

Fuzzy Operations

Fuzzy Intersection (t-norm)

- Let a fuzzy intersection operation be defined as a function

$$i : [0,1] \times [0,1] \rightarrow [0,1]$$

$$(\mu_A(x), \mu_B(x)) \rightarrow i(\mu_A(x), \mu_B(x))$$

- **Axiom 5** : i is a continuous function (continuity)
- **Axiom 6** : $i(a,a) = a$ (idempotency)

Fuzzy Operations

Fuzzy Intersection (t-norm)

- Let a fuzzy intersection operation be defined as a function

$$i : [0,1] \times [0,1] \rightarrow [0,1]$$

$$(\mu_A(x), \mu_B(x)) \rightarrow i(\mu_A(x), \mu_B(x))$$

- **Axiom 5** : i is a continuous function (continuity)
- **Axiom 6** : $i(a,a) = a$ (idempotency)
- Standart Intersection ; $i(a,b) = \min\{a,b\}$ where $a=\mu_A(x)$ and $b=\mu_B(x)$

Fuzzy Operations

Fuzzy Intersection (t-norm)

- Let a fuzzy intersection operation be defined as a function

$$i : [0,1] \times [0,1] \rightarrow [0,1]$$

$$(\mu_A(x), \mu_B(x)) \rightarrow i(\mu_A(x), \mu_B(x))$$

- **Axiom 5** : i is a continuous function (continuity)
- **Axiom 6** : $i(a,a) = a$ (idempotency)
- Standart Intersection ; $i(a,b) = \min\{a,b\}$ where $a=\mu_A(x)$ and $b=\mu_B(x)$
- $i(0,a) = \min\{0,a\} = 0$ and $i(1,1) = \min\{1,1\} = 1$

Fuzzy Operations

Fuzzy Intersection (t-norm)

- Let a fuzzy intersection operation be defined as a function

$$i : [0,1] \times [0,1] \rightarrow [0,1]$$

$$(\mu_A(x), \mu_B(x)) \rightarrow i(\mu_A(x), \mu_B(x))$$

- **Axiom 5** : i is a continuous function (continuity)
- **Axiom 6** : $i(a,a) = a$ (idempotency)
- Standart Intersection ; $i(a,b) = \min\{a,b\}$ where $a=\mu_A(x)$ and $b=\mu_B(x)$
 - $i(0,a) = \min\{0,a\} = 0$ and $i(1,1) = \min\{1,1\} = 1$
 - $i(a,b) = i(b,a)$

Fuzzy Operations

Fuzzy Intersection (t-norm)

- Let a fuzzy intersection operation be defined as a function

$$i : [0,1] \times [0,1] \rightarrow [0,1]$$

$$(\mu_A(x), \mu_B(x)) \rightarrow i(\mu_A(x), \mu_B(x))$$

- **Axiom 5** : i is a continuous function (continuity)
- **Axiom 6** : $i(a,a) = a$ (idempotency)
- Standart Intersection ; $i(a,b) = \min\{a,b\}$ where $a=\mu_A(x)$ and $b=\mu_B(x)$
 - $i(0,a) = \min\{0,a\} = 0$ and $i(1,1) = \min\{1,1\} = 1$
 - $i(a,b) = i(b,a)$
 - assume $b \leq d$; if $a < b$, then $i(a,b) = a \leq i(a,d) = a$

Fuzzy Operations

Fuzzy Intersection (t-norm)

- Let a fuzzy intersection operation be defined as a function

$$i : [0,1] \times [0,1] \rightarrow [0,1]$$

$$(\mu_A(x), \mu_B(x)) \rightarrow i(\mu_A(x), \mu_B(x))$$

- **Axiom 5** : i is a continuous function (continuity)
- **Axiom 6** : $i(a,a) = a$ (idempotency)
- Standart Intersection ; $i(a,b) = \min\{a,b\}$ where $a=\mu_A(x)$ and $b=\mu_B(x)$
 - $i(0,a) = \min\{0,a\} = 0$ and $i(1,1) = \min\{1,1\} = 1$
 - $i(a,b) = i(b,a)$
 - assume $b \leq d$; if $a < b$, then $i(a,b) = a \leq i(a,d) = a$
if $b < a$, then $i(a,b) = b \leq i(a,d)$

Fuzzy Operations

Fuzzy Intersection (t-norm)

- Let a fuzzy intersection operation be defined as a function

$$i : [0,1] \times [0,1] \rightarrow [0,1]$$

$$(\mu_A(x), \mu_B(x)) \rightarrow i(\mu_A(x), \mu_B(x))$$

- **Axiom 5** : i is a continuous function (continuity)
- **Axiom 6** : $i(a,a) = a$ (idempotency)
- Standart Intersection ; $i(a,b) = \min\{a,b\}$ where $a=\mu_A(x)$ and $b=\mu_B(x)$
 - $i(0,a) = \min\{0,a\} = 0$ and $i(1,1) = \min\{1,1\} = 1$
 - $i(a,b) = i(b,a)$
 - assume $b \leq d$; if $a < b$, then $i(a,b) = a \leq i(a,d) = a$
if $b < a$, then $i(a,b) = b \leq i(a,d)$
 - $i(i(a,b),c) = i(a,i(b,c))$

Fuzzy Operations

Fuzzy Intersection (t-norm)

- Let a fuzzy intersection operation be defined as a function

$$i : [0,1] \times [0,1] \rightarrow [0,1]$$

$$(\mu_A(x), \mu_A(x)) \rightarrow i(\mu_A(x), \mu_B(x))$$

- **Axiom 5** : i is a continuous function (continuity)
- **Axiom 6** : $i(a,a) = a$ (idempotency)
- Standart Intersection ; $i(a,b) = \min\{a,b\}$ where $a=\mu_A(x)$ and $b=\mu_B(x)$
 - $i(0,a) = \min\{0,a\} = 0$ and $i(1,1) = \min\{1,1\} = 1$
 - $i(a,b) = i(b,a)$
 - assume $b \leq d$; if $a < b$, then $i(a,b) = a \leq i(a,d) = a$
if $b < a$, then $i(a,b) = b \leq i(a,d)$
 - $i(i(a,b),c) = i(a,i(b,c))$
- i is continuous function
- $i(a,a) = a$

Fuzzy Operations

Fuzzy Intersection (t-norm)

- Algebraic Product ($A \bullet B$); $i(a,b) = a.b$

Fuzzy Operations

Fuzzy Intersection (t-norm)

- Algebraic Product ($A \bullet B$); $i(a,b) = a.b$
 - $i(0,a) = 0$ and $i(1,1) = 1$

Fuzzy Operations

Fuzzy Intersection (t-norm)

- Algebraic Product ($A \bullet B$); $i(a,b) = a.b$
 - $i(0,a) = 0$ and $i(1,1) = 1$
 - $i(a,b) = i(b,a)$

Fuzzy Operations

Fuzzy Intersection (t-norm)

- Algebraic Product ($A \bullet B$); $i(a,b) = a.b$
 - $i(0,a) = 0$ and $i(1,1) = 1$
 - $i(a,b) = i(b,a)$
 - assume $b \leq d$, then $i(a,b) = a.b \leq i(a,d) = a.d$

Fuzzy Operations

Fuzzy Intersection (t-norm)

- Algebraic Product ($A \bullet B$); $i(a,b) = a.b$
 - $i(0,a) = 0$ and $i(1,1) = 1$
 - $i(a,b) = i(b,a)$
 - assume $b \leq d$, then $i(a,b) = a.b \leq i(a,d) = a.d$
 - $i(i(a,b),c) = i(a,i(b,c)) \rightarrow a.b.c = a.b.c$

Fuzzy Operations

Fuzzy Intersection (t-norm)

- Algebraic Product ($A \bullet B$); $i(a,b) = a.b$
 - $i(0,a) = 0$ and $i(1,1) = 1$
 - $i(a,b) = i(b,a)$
 - assume $b \leq d$, then $i(a,b) = a.b \leq i(a,d) = a.d$
 - $i(i(a,b),c) = i(a,i(b,c)) \rightarrow a.b.c = a.b.c$
 - $i(a,a) = a^2 \neq a$

Fuzzy Operations

Fuzzy Intersection (t-norm)

- Algebraic Product ($A \bullet B$); $i(a,b) = a.b$
 - $i(0,a) = 0$ and $i(1,1) = 1$
 - $i(a,b) = i(b,a)$
 - assume $b \leq d$, then $i(a,b) = a.b \leq i(a,d) = a.d$
 - $i(i(a,b),c) = i(a,i(b,c)) \rightarrow a.b.c = a.b.c$
 - $i(a,a) = a^2 \neq a$
- Bounded Product ($A \odot B$); $i(a,b) = \max\{0, a+b-1\}$

Fuzzy Operations

Fuzzy Intersection (t-norm)

- Algebraic Product ($A \bullet B$); $i(a,b) = a.b$
 - $i(0,a) = 0$ and $i(1,1) = 1$
 - $i(a,b) = i(b,a)$
 - assume $b \leq d$, then $i(a,b) = a.b \leq i(a,d) = a.d$
 - $i(i(a,b),c) = i(a,i(b,c)) \rightarrow a.b.c = a.b.c$
 - $i(a,a) = a^2 \neq a$
- Bounded Product ($A \odot B$); $i(a,b) = \max\{0, a+b-1\}$
- Yager Class ; $i_w(a,b) = 1 - \min\{1, ((1-a)^w + (1-b)^w)^{1/w}\}$
where w in $(0, \infty)$

Fuzzy Operations

Fuzzy Union (s-norm)

- Let a fuzzy union operation be defined as a function

$$u : [0,1] \times [0,1] \rightarrow [0,1]$$

$$(\mu_A(x), \mu_B(x)) \rightarrow u(\mu_A(x), \mu_B(x))$$

Fuzzy Operations

Fuzzy Union (s-norm)

- Let a fuzzy union operation be defined as a function

$$u : [0,1] \times [0,1] \rightarrow [0,1]$$

$$(\mu_A(x), \mu_B(x)) \rightarrow u(\mu_A(x), \mu_B(x))$$

- Axiom 1** : $u(1,1) = 1, u(1,0) = 1, u(0,1) = 1, u(0,0) = 0$ (boundary condition)
- Axiom 2** : $u(a,b) = u(b,a)$ (comutativity)
- Axiom 3** : if $b \leq d$, then $u(a,b) \leq u(a,d)$ (monotonicity)
- Axiom 4** : $u(u(a,b),c) = u(a,u(b,c))$ (associativity)

Fuzzy Operations

Fuzzy Union (s-norm)

- Let a fuzzy union operation be defined as a function

$$u : [0,1] \times [0,1] \rightarrow [0,1]$$

$$(\mu_A(x), \mu_B(x)) \rightarrow u(\mu_A(x), \mu_B(x))$$

- **Axiom 1** : $u(1,1) = 1, u(1,0) = 1, u(0,1) = 1, u(0,0) = 0$ (boundary condition)
- **Axiom 2** : $u(a,b) = u(b,a)$ (commutativity)
- **Axiom 3** : if $b \leq d$, then $u(a,b) \leq u(a,d)$ (monotonicity)
- **Axiom 4** : $u(u(a,b),c) = u(a,u(b,c))$ (associativity)
- these axioms ensure that the fuzzy union becomes the classical union when A and B are crisp
- monotonicity and commutativity express that a decrease in the degree of membership in sets A and B cannot produce an increase in the degree of membership of union

Fuzzy Operations

Fuzzy Union (s-norm)

- Let a fuzzy intersection operation be defined as a function

$$u : [0,1] \times [0,1] \rightarrow [0,1]$$

$$(\mu_A(x), \mu_A(x)) \rightarrow u(\mu_A(x), \mu_B(x))$$

- **Axiom 5** : u is a continuous function (continuity)
- **Axiom 6** : $u(a,a) = a$ (idempotency)

Fuzzy Operations

Fuzzy Union (s-norm)

- Let a fuzzy intersection operation be defined as a function

$$u : [0,1] \times [0,1] \rightarrow [0,1]$$

$$(\mu_A(x), \mu_B(x)) \rightarrow u(\mu_A(x), \mu_B(x))$$

- **Axiom 5** : u is a continuous function (continuity)
- **Axiom 6** : $u(a,a) = a$ (idempotency)
- Standart Union ; $u(a,b) = \max\{a,b\}$ where $a=\mu_A(x)$ and $b=\mu_B(x)$

Fuzzy Operations

Fuzzy Union (s-norm)

- Let a fuzzy intersection operation be defined as a function

$$u : [0,1] \times [0,1] \rightarrow [0,1]$$

$$(\mu_A(x), \mu_B(x)) \rightarrow u(\mu_A(x), \mu_B(x))$$

- **Axiom 5** : u is a continuous function (continuity)
- **Axiom 6** : $u(a,a) = a$ (idempotency)
- Standart Union ; $u(a,b) = \max\{a,b\}$ where $a=\mu_A(x)$ and $b=\mu_B(x)$
- $u(1,a) = \max\{1,a\} = 1$ and $u(0,0) = \max\{0,0\} = 0$

Fuzzy Operations

Fuzzy Union (s-norm)

- Let a fuzzy intersection operation be defined as a function

$$u : [0,1] \times [0,1] \rightarrow [0,1]$$

$$(\mu_A(x), \mu_B(x)) \rightarrow u(\mu_A(x), \mu_B(x))$$

- **Axiom 5** : u is a continuous function (continuity)
- **Axiom 6** : $u(a,a) = a$ (idempotency)
- Standart Union ; $u(a,b) = \max\{a,b\}$ where $a=\mu_A(x)$ and $b=\mu_B(x)$
 - $u(1,a) = \max\{1,a\} = 1$ and $u(0,0) = \max\{0,0\} = 0$
 - $u(a,b) = u(b,a)$

Fuzzy Operations

Fuzzy Union (s-norm)

- Let a fuzzy intersection operation be defined as a function

$$u : [0,1] \times [0,1] \rightarrow [0,1]$$

$$(\mu_A(x), \mu_B(x)) \rightarrow u(\mu_A(x), \mu_B(x))$$

- **Axiom 5** : u is a continuous function (continuity)
- **Axiom 6** : $u(a,a) = a$ (idempotency)
- Standart Union ; $u(a,b) = \max\{a,b\}$ where $a=\mu_A(x)$ and $b=\mu_B(x)$
 - $u(1,a) = \max\{1,a\} = 1$ and $u(0,0) = \max\{0,0\} = 0$
 - $u(a,b) = u(b,a)$
 - assume $b \leq d$; if $d < a$, then $u(a,b) = a \leq u(a,d) = a$

Fuzzy Operations

Fuzzy Union (s-norm)

- Let a fuzzy intersection operation be defined as a function

$$u : [0,1] \times [0,1] \rightarrow [0,1]$$

$$(\mu_A(x), \mu_B(x)) \rightarrow u(\mu_A(x), \mu_B(x))$$

- **Axiom 5** : u is a continuous function (continuity)
- **Axiom 6** : $u(a,a) = a$ (idempotency)
- Standart Union ; $u(a,b) = \max\{a,b\}$ where $a=\mu_A(x)$ and $b=\mu_B(x)$
 - $u(1,a) = \max\{1,a\} = 1$ and $u(0,0) = \max\{0,0\} = 0$
 - $u(a,b) = u(b,a)$
 - assume $b \leq d$; if $d < a$, then $u(a,b) = a \leq u(a,d) = a$
if $a < d$, then $u(a,d) = d \geq u(a,b)$

Fuzzy Operations

Fuzzy Union (s-norm)

- Let a fuzzy intersection operation be defined as a function

$$u : [0,1] \times [0,1] \rightarrow [0,1]$$

$$(\mu_A(x), \mu_B(x)) \rightarrow u(\mu_A(x), \mu_B(x))$$

- **Axiom 5** : u is a continuous function (continuity)
- **Axiom 6** : $u(a,a) = a$ (idempotency)
- Standart Union ; $u(a,b) = \max\{a,b\}$ where $a=\mu_A(x)$ and $b=\mu_B(x)$
 - $u(1,a) = \max\{1,a\} = 1$ and $u(0,0) = \max\{0,0\} = 0$
 - $u(a,b) = u(b,a)$
 - assume $b \leq d$; if $d < a$, then $u(a,b) = a \leq u(a,d) = a$
if $a < d$, then $u(a,d) = d \geq u(a,b)$
 - $u(u(a,b),c) = u(a,u(b,c))$

Fuzzy Operations

Fuzzy Union (s-norm)

- Let a fuzzy intersection operation be defined as a function

$$u : [0,1] \times [0,1] \rightarrow [0,1]$$

$$(\mu_A(x), \mu_B(x)) \rightarrow u(\mu_A(x), \mu_B(x))$$

- **Axiom 5** : u is a continuous function (continuity)
- **Axiom 6** : $u(a,a) = a$ (idempotency)
- Standart Union ; $u(a,b) = \max\{a,b\}$ where $a=\mu_A(x)$ and $b=\mu_B(x)$
 - $u(1,a) = \max\{1,a\} = 1$ and $u(0,0) = \max\{0,0\} = 0$
 - $u(a,b) = u(b,a)$
 - assume $b \leq d$; if $d < a$, then $u(a,b) = a \leq u(a,d) = a$
if $a < d$, then $u(a,d) = d \geq u(a,b)$
 - $u(u(a,b),c) = u(a,u(b,c))$
- u is continuous function
- $u(a,a) = a$

Fuzzy Operations

Fuzzy Union (s-norm)

- Probabilistic Sum ; $u(a,b) = a + b - a.b$

Fuzzy Operations

Fuzzy Union (s-norm)

- Probabilistic Sum ; $u(a,b) = a + b - a.b$
 - $u(1,a) = 1$ and $u(0,0) = 0$

Fuzzy Operations

Fuzzy Union (s-norm)

- Probabilistic Sum ; $u(a,b) = a + b - a.b$
 - $u(1,a) = 1$ and $u(0,0) = 0$
 - $u(a,b) = u(b,a)$

Fuzzy Operations

Fuzzy Union (s-norm)

- Probabilistic Sum ; $u(a,b) = a + b - a.b$
 - $u(1,a) = 1$ and $u(0,0) = 0$
 - $u(a,b) = u(b,a)$
 - assume $b \leq d$, then $u(a,b) = a + b(1 - a) \leq u(a,d) = a + d(1 - a)$

Fuzzy Operations

Fuzzy Union (s-norm)

- Probabilistic Sum ; $u(a,b) = a + b - a.b$
 - $u(1,a) = 1$ and $u(0,0) = 0$
 - $u(a,b) = u(b,a)$
 - assume $b \leq d$, then $u(a,b) = a + b(1 - a) \leq u(a,d) = a + d(1 - a)$
 - $u(u(a,b),c) = u(a,u(b,c)) \rightarrow a+b+c-ac-ab+abc = a+b+c-ab-ac+abc$

Fuzzy Operations

Fuzzy Union (s-norm)

- Probabilistic Sum ; $u(a,b) = a + b - a.b$
 - $u(1,a) = 1$ and $u(0,0) = 0$
 - $u(a,b) = u(b,a)$
 - assume $b \leq d$, then $u(a,b) = a + b(1 - a) \leq u(a,d) = a + d(1 - a)$
 - $u(u(a,b),c) = u(a,u(b,c)) \rightarrow a+b+c-ac-ab+abc = a+b+c-ab-ac+abc$
 - $u(a,a) = 2a - a^2 \neq a$

Fuzzy Operations

Fuzzy Union (s-norm)

- Probabilistic Sum ; $u(a,b) = a + b - a.b$
 - $u(1,a) = 1$ and $u(0,0) = 0$
 - $u(a,b) = u(b,a)$
 - assume $b \leq d$, then $u(a,b) = a + b(1 - a) \leq u(a,d) = a + d(1 - a)$
 - $u(u(a,b),c) = u(a,u(b,c)) \rightarrow a+b+c-ac-ab+abc = a+b+c-ab-ac+abc$
 - $u(a,a) = 2a - a^2 \neq a$
- Bounded Sum ($A \oplus B$); $u(a,b) = \min\{1, a+b\}$

Fuzzy Operations

Fuzzy Union (s-norm)

- Probabilistic Sum ; $u(a,b) = a + b - a.b$
 - $u(1,a) = 1$ and $u(0,0) = 0$
 - $u(a,b) = u(b,a)$
 - assume $b \leq d$, then $u(a,b) = a + b(1 - a) \leq u(a,d) = a + d(1 - a)$
 - $u(u(a,b),c) = u(a,u(b,c)) \rightarrow a+b+c-ac-ab+abc = a+b+c-ab-ac+abc$
 - $u(a,a) = 2a - a^2 \neq a$
- Bounded Sum ($A \oplus B$); $u(a,b) = \min\{1, a+b\}$
- Yager Class ; $u_w(a,b) = \min\{1, (a^w + b^w)^{1/w}\}$
where w in $(0, \infty)$

Fuzzy Operations

Other Operations on Fuzzy Sets

- Disjunctive Sum ; $C = (A \cap \neg B) \cup (\neg A \cap B)$

$$\mu_C(x) = \max\{\min[\mu_A(x), 1 - \mu_B(x)], \min[\mu_B(x), 1 - \mu_A(x)]\}$$

Fuzzy Operations

Other Operations on Fuzzy Sets

- Disjunctive Sum ; $C = (A \cap \neg B) \cup (\neg A \cap B)$

$$\mu_C(x) = \max\{\min[\mu_A(x), 1 - \mu_B(x)], \min[\mu_B(x), 1 - \mu_A(x)]\}$$

- Disjoint Sum ; $C = A \Delta B$

$$\mu_C(x) = |\mu_A(x) - \mu_B(x)|$$

Fuzzy Operations

Other Operations on Fuzzy Sets

- Disjunctive Sum ; $C = (A \cap \neg B) \cup (\neg A \cap B)$

$$\mu_C(x) = \max\{\min[\mu_A(x), 1 - \mu_B(x)], \min[\mu_B(x), 1 - \mu_A(x)]\}$$

- Disjoint Sum ; $C = A \Delta B$

$$\mu_C(x) = |\mu_A(x) - \mu_B(x)|$$

- Set Difference ; $C = A - B = A \cap \neg B$

$$\mu_C(x) = \min\{\mu_A(x), 1 - \mu_B(x)\}$$

Fuzzy Operations

Other Operations on Fuzzy Sets

- Disjunctive Sum ; $C = (A \cap \neg B) \cup (\neg A \cap B)$

$$\mu_C(x) = \max\{\min[\mu_A(x), 1 - \mu_B(x)], \min[\mu_B(x), 1 - \mu_A(x)]\}$$

- Disjoint Sum ; $C = A \Delta B$

$$\mu_C(x) = |\mu_A(x) - \mu_B(x)|$$

- Set Difference ; $C = A - B = A \cap \neg B$

$$\mu_C(x) = \min\{\mu_A(x), 1 - \mu_B(x)\}$$

- Distance in Fuzzy Set ;

$$\text{Minkowski distance, } d_w(A, B) = (\sum_{x \text{ in } X} |\mu_A(x) - \mu_B(x)|^w)^{1/w}$$

Fuzzy Operations


- $\mu_A(x) = 1/(x^2+1)$ and $\mu_B(x) = x^2/(x^2+1)$

$C = \neg A, D = \neg B, E = A \cap B, F = A \cup B, \mu_C, \mu_D, \mu_E, \mu_F = ?$

Fuzzy Operations

- $\mu_A(x) = 1/(x^2+1)$ and $\mu_B(x) = x^2/(x^2+1)$


$C = \neg A, D = \neg B, E = A \cap B, F = A \cup B, \mu_C, \mu_D, \mu_E, \mu_F = ?$



Fuzzy Operations

- $\mu_A(x) = 1/(x^2+1)$ and $\mu_B(x) = x^2/(x^2+1)$


$C = \neg A, D = \neg B, E = A \cap B, F = A \cup B, \mu_C, \mu_D, \mu_E, \mu_F = ?$



Fuzzy Operations

- $\mu_A(x) = 1/(x^2+1)$ and $\mu_B(x) = x^2/(x^2+1)$


$C = \neg A, D = \neg B, E = A \cap B, F = A \cup B, \mu_C, \mu_D, \mu_E, \mu_F = ?$



Fuzzy Operations

- $\mu_A(x) = 1/(x^2+1)$ and $\mu_B(x) = x^2/(x^2+1)$

$C = \neg A, D = \neg B, E = A \cap B, F = A \cup B, \mu_C, \mu_D, \mu_E, \mu_F = ?$



$$\mu_C(x) = 1 - \mu_A(x)$$


$$\mu_C(x) = 1 - 1/(x^2+1)$$

$$\mu_C(x) = x^2/(x^2+1) = \mu_B(x)$$

Fuzzy Operations

- $\mu_A(x) = 1/(x^2+1)$ and $\mu_B(x) = x^2/(x^2+1)$

$C = \neg A, D = \neg B, E = A \cap B, F = A \cup B, \mu_C, \mu_D, \mu_E, \mu_F = ?$



$$\mu_C(x) = 1 - \mu_A(x)$$

$$\mu_C(x) = 1 - 1/(x^2+1)$$

$$\mu_C(x) = x^2/(x^2+1) = \mu_B(x)$$

$$\mu_D(x) = 1 - \mu_B(x)$$


$$\mu_D(x) = 1 - x^2/(x^2+1)$$

$$\mu_D(x) = 1/(x^2+1) = \mu_A(x)$$

Fuzzy Operations

- $\mu_A(x) = 1/(x^2+1)$ and $\mu_B(x) = x^2/(x^2+1)$

$C = \neg A, D = \neg B, E = A \cap B, F = A \cup B, \mu_C, \mu_D, \mu_E, \mu_F = ?$



$$\mu_C(x) = 1 - \mu_A(x)$$

$$\mu_C(x) = 1 - 1/(x^2+1)$$

$$\mu_C(x) = x^2/(x^2+1) = \mu_B(x)$$

$$\mu_D(x) = 1 - \mu_B(x)$$

$$\mu_D(x) = 1 - x^2/(x^2+1)$$

$$\mu_D(x) = 1/(x^2+1) = \mu_A(x)$$


$$\mu_E(x) = \min\{\mu_A(x), \mu_B(x)\}$$

$$\mu_F(x) = \max\{\mu_A(x), \mu_B(x)\}$$

Fuzzy Operations

- $\mu_A(x) = 1/(x^2+1)$ and $\mu_B(x) = x^2/(x^2+1)$

$C = \neg A, D = \neg B, E = A \cap B, F = A \cup B, \mu_C, \mu_D, \mu_E, \mu_F = ?$



$$\mu_C(x) = 1 - \mu_A(x)$$

$$\mu_C(x) = 1 - 1/(x^2+1)$$

$$\mu_C(x) = x^2/(x^2+1) = \mu_B(x)$$

$$\mu_D(x) = 1 - \mu_B(x)$$

$$\mu_D(x) = 1 - x^2/(x^2+1)$$

$$\mu_D(x) = 1/(x^2+1) = \mu_A(x)$$

$$\mu_E(x) = \min\{\mu_A(x), \mu_B(x)\}$$


$$\mu_F(x) = \max\{\mu_A(x), \mu_B(x)\}$$

$$1/(x^2+1) = x^2/(x^2+1)$$

Fuzzy Operations

- $\mu_A(x) = 1/(x^2+1)$ and $\mu_B(x) = x^2/(x^2+1)$

$C = \neg A, D = \neg B, E = A \cap B, F = A \cup B, \mu_C, \mu_D, \mu_E, \mu_F = ?$



$$\mu_C(x) = 1 - \mu_A(x)$$

$$\mu_C(x) = 1 - 1/(x^2+1)$$

$$\mu_C(x) = x^2/(x^2+1) = \mu_B(x)$$

$$\mu_D(x) = 1 - \mu_B(x)$$

$$\mu_D(x) = 1 - x^2/(x^2+1)$$

$$\mu_D(x) = 1/(x^2+1) = \mu_A(x)$$

$$\mu_E(x) = \min\{\mu_A(x), \mu_B(x)\}$$

$$\mu_F(x) = \max\{\mu_A(x), \mu_B(x)\}$$


$$1/(x^2+1) = x^2/(x^2+1)$$

$$x = \pm 1$$

Fuzzy Operations

- $\mu_A(x) = 1/(x^2+1)$ and $\mu_B(x) = x^2/(x^2+1)$

$C = \neg A, D = \neg B, E = A \cap B, F = A \cup B, \mu_C, \mu_D, \mu_E, \mu_F = ?$



$$\mu_C(x) = 1 - \mu_A(x)$$

$$\mu_C(x) = 1 - 1/(x^2+1)$$

$$\mu_C(x) = x^2/(x^2+1) = \mu_B(x)$$

$$\mu_D(x) = 1 - \mu_B(x)$$

$$\mu_D(x) = 1 - x^2/(x^2+1)$$

$$\mu_D(x) = 1/(x^2+1) = \mu_A(x)$$


$$\mu_E(x) = \min\{\mu_A(x), \mu_B(x)\}$$

$$\mu_F(x) = \max\{\mu_A(x), \mu_B(x)\}$$

Fuzzy Operations

- $\mu_A(x) = 1/(x^2+1)$ and $\mu_B(x) = x^2/(x^2+1)$

$C = \neg A, D = \neg B, E = A \cap B, F = A \cup B, \mu_C, \mu_D, \mu_E, \mu_F = ?$



$$\mu_C(x) = 1 - \mu_A(x)$$

$$\mu_C(x) = 1 - 1/(x^2+1)$$

$$\mu_C(x) = x^2/(x^2+1) = \mu_B(x)$$

$$\mu_D(x) = 1 - \mu_B(x)$$

$$\mu_D(x) = 1 - x^2/(x^2+1)$$

$$\mu_D(x) = 1/(x^2+1) = \mu_A(x)$$


$$\mu_E(x) = \min\{\mu_A(x), \mu_B(x)\}$$

$$\mu_F(x) = \max\{\mu_A(x), \mu_B(x)\}$$

Fuzzy Operations

- $\mu_A(x) = 1/(x^2+1)$ and $\mu_B(x) = x^2/(x^2+1)$

$C = \neg A, D = \neg B, E = A \cap B, F = A \cup B, \mu_C, \mu_D, \mu_E, \mu_F = ?$



$$\mu_C(x) = 1 - \mu_A(x)$$

$$\mu_C(x) = 1 - 1/(x^2+1)$$

$$\mu_C(x) = x^2/(x^2+1) = \mu_B(x)$$

$$\mu_D(x) = 1 - \mu_B(x)$$

$$\mu_D(x) = 1 - x^2/(x^2+1)$$

$$\mu_D(x) = 1/(x^2+1) = \mu_A(x)$$


$$\mu_E(x) = \min\{\mu_A(x), \mu_B(x)\}$$

$$\mu_F(x) = \max\{\mu_A(x), \mu_B(x)\}$$

Fuzzy Operations

- $\mu_A(x) = 1/(x^2+1)$ and $\mu_B(x) = x^2/(x^2+1)$

$C = \neg A, D = \neg B, E = A \cap B, F = A \cup B, \mu_C, \mu_D, \mu_E, \mu_F = ?$



$$\mu_C(x) = 1 - \mu_A(x)$$

$$\mu_C(x) = 1 - 1/(x^2+1)$$

$$\mu_C(x) = x^2/(x^2+1) = \mu_B(x)$$

$$\mu_D(x) = 1 - \mu_B(x)$$

$$\mu_D(x) = 1 - x^2/(x^2+1)$$

$$\mu_D(x) = 1/(x^2+1) = \mu_A(x)$$

$$\mu_E(x) = \min\{\mu_A(x), \mu_B(x)\}$$


$$\mu_F(x) = \max\{\mu_A(x), \mu_B(x)\}$$

$$\mu_E(x) = \begin{cases} \mu_A(x) & \text{if } x < -1 \\ \mu_B(x) & \text{if } -1 \leq x \leq 1 \\ \mu_A(x) & \text{if } x > 1 \end{cases}$$

Fuzzy Operations

- $\mu_A(x) = 1/(x^2+1)$ and $\mu_B(x) = x^2/(x^2+1)$

$$C = \neg A, D = \neg B, E = A \cap B, F = A \cup B, \mu_C, \mu_D, \mu_E, \mu_F = ?$$



$$\mu_C(x) = 1 - \mu_A(x)$$

$$\mu_C(x) = 1 - 1/(x^2+1)$$

$$\mu_C(x) = x^2/(x^2+1) = \mu_B(x)$$

$$\mu_D(x) = 1 - \mu_B(x)$$

$$\mu_D(x) = 1 - x^2/(x^2+1)$$

$$\mu_D(x) = 1/(x^2+1) = \mu_A(x)$$

$$\mu_E(x) = \min\{\mu_A(x), \mu_B(x)\}$$

$$\mu_F(x) = \max\{\mu_A(x), \mu_B(x)\}$$

$$\mu_E(x) = \begin{cases} \mu_A(x) & \text{if } x < -1 \\ \mu_B(x) & \text{if } -1 \leq x \leq 1 \\ \mu_A(x) & \text{if } x > 1 \end{cases}$$

$$\mu_F(x) = \begin{cases} \mu_B(x) & \text{if } x < -1 \\ \mu_A(x) & \text{if } -1 \leq x \leq 1 \\ \mu_B(x) & \text{if } x > 1 \end{cases}$$