

# Fuzzy Sets

Murat Osmanoglu

# Crisp Sets

## (Classical or Nonfuzzy)


- $Z = \{\dots, -2, -1, 0, 1, 2, \dots\}$ ,  $N = \{0, 1, 2, \dots\}$

**Definition :** collection of well-defined objects

# Crisp Sets (Classical or Nonfuzzy)

- $Z = \{\dots, -2, -1, 0, 1, 2, \dots\}$ ,  $N = \{0, 1, 2, \dots\}$

**Definition :** collection of well-defined objects




- $a \in A, b \in A, c \notin A$

# Crisp Sets (Classical or Nonfuzzy)

- $Z = \{\dots, -2, -1, 0, 1, 2, \dots\}$ ,  $N = \{0, 1, 2, \dots\}$

**Definition :** collection of well-defined objects




- $a \in A, b \in A, c \notin A$
- $\mu_A : X \rightarrow \{0, 1\}$   
membership function

# Crisp Sets (Classical or Nonfuzzy)

- $Z = \{\dots, -2, -1, 0, 1, 2, \dots\}$ ,  $N = \{0, 1, 2, \dots\}$

**Definition :** collection of well-defined objects




- $a \in A, b \in A, c \notin A$
- $\mu_A : X \rightarrow \{0, 1\}$   
membership function

$$\mu_A(x) = \begin{cases} 1 & \text{if } x \in A \\ 0 & \text{if } x \notin A \end{cases}$$

# Crisp Sets (Classical or Nonfuzzy)

- $Z = \{\dots, -2, -1, 0, 1, 2, \dots\}$ ,  $N = \{0, 1, 2, \dots\}$

**Definition :** collection of well-defined objects



- $a \in A, b \in A, c \notin A$
- $\mu_A : X \rightarrow \{0, 1\}$   
membership function


$$\mu_A(x) = \begin{cases} 1 & \text{if } x \in A \\ 0 & \text{if } x \notin A \end{cases}$$

- $X$  denotes the universal set, contains all possible elements under the consideration

# Crisp Sets (Classical or Nonfuzzy)

- $Z = \{\dots, -2, -1, 0, 1, 2, \dots\}$ ,  $N = \{0, 1, 2, \dots\}$

**Definition :** collection of well-defined objects



- $a \in A, b \in A, c \notin A$
- $\mu_A : X \rightarrow \{0, 1\}$   
membership function

$$\mu_A(x) = \begin{cases} 1 & \text{if } x \in A \\ 0 & \text{if } x \notin A \end{cases}$$

- $X$  denotes the universal set, contains all possible elements under the consideration
- $\emptyset$  denotes the empty set, contains no element

# Crisp Sets

## (Classical or Nonfuzzy)

- $A = \{A_1, A_2, \dots, A_n\}$ , a family of sets, that contains sets as elements  
 $A = \{A_i \mid i \in I\}$  where  $I$  is called the index set that reference the corresponding set  $A_i$



# Crisp Sets

## (Classical or Nonfuzzy)

- $A = \{A_1, A_2, \dots, A_n\}$ , a family of sets, that contains sets as elements  
 $A = \{A_i \mid i \in I\}$  where  $I$  is called the index set that reference the corresponding set  $A_i$
- If every element of set  $A$  is also an element of set  $B$ , then  $A$  is called a subset of  $B$ ,

$$A \subseteq B$$

# Crisp Sets

## (Classical or Nonfuzzy)

- $A = \{A_1, A_2, \dots, A_n\}$ , a family of sets, that contains sets as elements

$A = \{A_i \mid i \in I\}$  where  $I$  is called the index set that reference the corresponding set  $A_i$

- If every element of set  $A$  is also an element of set  $B$ , then  $A$  is called a subset of  $B$ ,

$$A \subseteq B$$

- If  $A \subseteq B$  and  $B \subseteq A$ , then  $A = B$

# Crisp Sets

## (Classical or Nonfuzzy)

- $A = \{A_1, A_2, \dots, A_n\}$ , a family of sets, that contains sets as elements  
 $A = \{A_i \mid i \in I\}$  where  $I$  is called the index set that reference the corresponding set  $A_i$
- If every element of set  $A$  is also an element of set  $B$ , then  $A$  is called a subset of  $B$ ,  
$$A \subseteq B$$
- If  $A \subseteq B$  and  $B \subseteq A$ , then  $A = B$
- $P(A)$ , the power set of  $A$ , that contains all subsets of  $A$

# Crisp Sets

## (Classical or Nonfuzzy)

- $A = \{A_1, A_2, \dots, A_n\}$ , a family of sets, that contains sets as elements  
 $A = \{A_i \mid i \in I\}$  where  $I$  is called the index set that reference the corresponding set  $A_i$
- If every element of set  $A$  is also an element of set  $B$ , then  $A$  is called a subset of  $B$ ,

$$A \subseteq B$$

- If  $A \subseteq B$  and  $B \subseteq A$ , then  $A = B$
- $P(A)$ , the power set of  $A$ , that contains all subsets of  $A$
- The number of members of a finite set  $A$  is called cardinality of  $A$ ,

$$|A|$$

# Crisp Sets

## (Classical or Nonfuzzy)

- $A = \{A_1, A_2, \dots, A_n\}$ , a family of sets, that contains sets as elements  
 $A = \{A_i \mid i \in I\}$  where  $I$  is called the index set that reference the corresponding set  $A_i$
- If every element of set  $A$  is also an element of set  $B$ , then  $A$  is called a subset of  $B$ ,

$$A \subseteq B$$

- If  $A \subseteq B$  and  $B \subseteq A$ , then  $A = B$
- $P(A)$ , the power set of  $A$ , that contains all subsets of  $A$
- The number of members of a finite set  $A$  is called cardinality of  $A$ ,

$$|A|$$

- The cardinality of power set of a given set  $A$ ,

$$|P(A)| = 2^{|A|}$$

# Crisp Sets

(Classical or Nonfuzzy)

## Operations on Sets

# Crisp Sets

## (Classical or Nonfuzzy)

### Operations on Sets

- Complement,  $\neg A = \{x \in X \mid x \notin A\}$ ,

# Crisp Sets

## (Classical or Nonfuzzy)

### Operations on Sets

- Complement,  $\neg A = \{x \in X \mid x \notin A\}$ ,
- Intersection,  $A \cap B = \{x \in X \mid x \in A \text{ and } x \in B\}$ ,



# Crisp Sets

## (Classical or Nonfuzzy)

### Operations on Sets

- Complement,  $\neg A = \{x \in X \mid x \notin A\}$ ,
- Intersection,  $A \cap B = \{x \in X \mid x \in A \text{ and } x \in B\}$ ,
- Union,  $A \cup B = \{x \in X \mid x \in A \text{ or } x \in B\}$ ,

# Crisp Sets

## (Classical or Nonfuzzy)

### Operations on Sets

- Complement,  $\neg A = \{x \in X \mid x \notin A\}$ ,
- Intersection,  $A \cap B = \{x \in X \mid x \in A \text{ and } x \in B\}$ ,
- Union,  $A \cup B = \{x \in X \mid x \in A \text{ or } x \in B\}$ ,
- Set Difference,  $A - B = \{x \in X \mid x \in A \text{ and } x \notin B\}$ ,

# Crisp Sets

## (Classical or Nonfuzzy)

### Operations on Sets

- Complement,  $\neg A = \{x \in X \mid x \notin A\}$ ,
- Intersection,  $A \cap B = \{x \in X \mid x \in A \text{ and } x \in B\}$ ,
- Union,  $A \cup B = \{x \in X \mid x \in A \text{ or } x \in B\}$ ,
- Set Difference,  $A - B = \{x \in X \mid x \in A \text{ and } x \notin B\}$ ,
- $\bigcup_{i \in I} A_i = \{x \in X \mid x \in A_i \text{ for some } i \in I\}$

# Crisp Sets

## (Classical or Nonfuzzy)

### Operations on Sets

- Complement,  $\neg A = \{x \in X \mid x \notin A\}$ ,
- Intersection,  $A \cap B = \{x \in X \mid x \in A \text{ and } x \in B\}$ ,
- Union,  $A \cup B = \{x \in X \mid x \in A \text{ or } x \in B\}$ ,
- Set Difference,  $A - B = \{x \in X \mid x \in A \text{ and } x \notin B\}$ ,
- $\bigcup_{i \in I} A_i = \{x \in X \mid x \in A_i \text{ for some } i \in I\}$
- $\bigcap_{i \in I} A_i = \{x \in X \mid x \in A_i \text{ for all } i \in I\}$

# Crisp Sets

## (Classical or Nonfuzzy)

### Operations on Sets

- Complement,  $\neg A = \{x \in X \mid x \notin A\}$ ,
- Intersection,  $A \cap B = \{x \in X \mid x \in A \text{ and } x \in B\}$ ,
- Union,  $A \cup B = \{x \in X \mid x \in A \text{ or } x \in B\}$ ,
- Set Difference,  $A - B = \{x \in X \mid x \in A \text{ and } x \notin B\}$ ,
- $\bigcup_{i \in I} A_i = \{x \in X \mid x \in A_i \text{ for some } i \in I\}$
- $\bigcap_{i \in I} A_i = \{x \in X \mid x \in A_i \text{ for all } i \in I\}$
- Fundamental characteristics of set operations,

# Crisp Sets

## (Classical or Nonfuzzy)

### Operations on Sets

- Complement,  $\neg A = \{x \in X \mid x \notin A\}$ ,
- Intersection,  $A \cap B = \{x \in X \mid x \in A \text{ and } x \in B\}$ ,
- Union,  $A \cup B = \{x \in X \mid x \in A \text{ or } x \in B\}$ ,
- Set Difference,  $A - B = \{x \in X \mid x \in A \text{ and } x \notin B\}$ ,
- $\bigcup_{i \in I} A_i = \{x \in X \mid x \in A_i \text{ for some } i \in I\}$
- $\bigcap_{i \in I} A_i = \{x \in X \mid x \in A_i \text{ for all } i \in I\}$
- Fundamental characteristics of set operations,  
 $\neg(\neg A) = A, \quad \neg X = \emptyset, \quad \neg \emptyset = X, \quad A \cup \neg A = X, \quad A \cap \neg A = \emptyset,$

# Crisp Sets

## (Classical or Nonfuzzy)

### Operations on Sets

- Complement,  $\neg A = \{x \in X \mid x \notin A\}$ ,
- Intersection,  $A \cap B = \{x \in X \mid x \in A \text{ and } x \in B\}$ ,
- Union,  $A \cup B = \{x \in X \mid x \in A \text{ or } x \in B\}$ ,
- Set Difference,  $A - B = \{x \in X \mid x \in A \text{ and } x \notin B\}$ ,
- $\bigcup_{i \in I} A_i = \{x \in X \mid x \in A_i \text{ for some } i \in I\}$
- $\bigcap_{i \in I} A_i = \{x \in X \mid x \in A_i \text{ for all } i \in I\}$
- Fundamental characteristics of set operations,  
 $\neg(\neg A) = A, \quad \neg X = \emptyset, \quad \neg \emptyset = X, \quad A \cup \neg A = X, \quad A \cap \neg A = \emptyset,$   
 $A \cup X = X, \quad A \cup \emptyset = A, \quad A \cap X = A, \quad A \cap \emptyset = \emptyset,$

# Crisp Sets

## (Classical or Nonfuzzy)

### Operations on Sets

- Complement,  $\neg A = \{x \in X \mid x \notin A\}$ ,
- Intersection,  $A \cap B = \{x \in X \mid x \in A \text{ and } x \in B\}$ ,
- Union,  $A \cup B = \{x \in X \mid x \in A \text{ or } x \in B\}$ ,
- Set Difference,  $A - B = \{x \in X \mid x \in A \text{ and } x \notin B\}$ ,
- $\bigcup_{i \in I} A_i = \{x \in X \mid x \in A_i \text{ for some } i \in I\}$
- $\bigcap_{i \in I} A_i = \{x \in X \mid x \in A_i \text{ for all } i \in I\}$
- Fundamental characteristics of set operations,  
 $\neg(\neg A) = A$ ,  $\neg X = \emptyset$ ,  $\neg \emptyset = X$ ,  $A \cup \neg A = X$ ,  $A \cap \neg A = \emptyset$ ,  
 $A \cup X = X$ ,  $A \cup \emptyset = A$ ,  $A \cap X = A$ ,  $A \cap \emptyset = \emptyset$ ,  
 $A \cup B = B \cup A$ ,  $A \cap B = B \cap A$ ,



# Crisp Sets

## (Classical or Nonfuzzy)

### Operations on Sets

- Complement,  $\neg A = \{x \in X \mid x \notin A\}$ ,
- Intersection,  $A \cap B = \{x \in X \mid x \in A \text{ and } x \in B\}$ ,
- Union,  $A \cup B = \{x \in X \mid x \in A \text{ or } x \in B\}$ ,
- Set Difference,  $A - B = \{x \in X \mid x \in A \text{ and } x \notin B\}$ ,
- $\bigcup_{i \in I} A_i = \{x \in X \mid x \in A_i \text{ for some } i \in I\}$
- $\bigcap_{i \in I} A_i = \{x \in X \mid x \in A_i \text{ for all } i \in I\}$
- Fundamental characteristics of set operations,  
 $\neg(\neg A) = A$ ,  $\neg X = \emptyset$ ,  $\neg \emptyset = X$ ,  $A \cup \neg A = X$ ,  $A \cap \neg A = \emptyset$ ,  
 $A \cup X = X$ ,  $A \cup \emptyset = A$ ,  $A \cap X = A$ ,  $A \cap \emptyset = \emptyset$ ,  
 $A \cup B = B \cup A$ ,  $A \cap B = B \cap A$ ,  
 $\neg(A \cup B) = \neg A \cap \neg B$ ,  $\neg(A \cap B) = \neg A \cup \neg B$

# Crisp Sets (Classical or Nonfuzzy)

- The cartesian product set of A and B

$$A \times B = \{(a,b) \mid a \in A \text{ and } b \in B\}$$

# Crisp Sets

## (Classical or Nonfuzzy)

- The cartesian product set of  $A$  and  $B$

$$A \times B = \{(a,b) \mid a \in A \text{ and } b \in B\}$$

- $|A \times B| = |A| \cdot |B|$

# Crisp Sets

## (Classical or Nonfuzzy)

- The cartesian product set of  $A$  and  $B$

$$A \times B = \{(a,b) \mid a \in A \text{ and } b \in B\}$$

- $|A \times B| = |A| \cdot |B|$

### Convex Set

- a set  $A$  in  $\mathbb{R}^n$  is convex iff, for every pair of points  $x$  and  $y$  in  $A$ , all the points lie in the line segment connecting  $x$  and  $y$  are also in  $A$

# Crisp Sets (Classical or Nonfuzzy)



- The cartesian product set of A and B

$$A \times B = \{(a,b) \mid a \in A \text{ and } b \in B\}$$

- $|A \times B| = |A| \cdot |B|$

## Convex Set

- a set A in  $\mathbb{R}^n$  is convex iff, for every pair of points x and y in A, all the points lie in the line segment connecting x and y are also in A



# Crisp Sets (Classical or Nonfuzzy)



- The cartesian product set of A and B

$$A \times B = \{(a,b) \mid a \in A \text{ and } b \in B\}$$

- $|A \times B| = |A| \cdot |B|$

## Convex Set

- a set A in  $\mathbb{R}^n$  is convex iff, for every pair of points x and y in A, all the points lie in the line segment connecting x and y are also in A



# Crisp Sets (Classical or Nonfuzzy)

- The cartesian product set of A and B



$$A \times B = \{(a,b) \mid a \in A \text{ and } b \in B\}$$

- $|A \times B| = |A| \cdot |B|$

## Convex Set

- a set A in  $\mathbb{R}^n$  is convex iff, for every pair of points x and y in A, all the points lie in the line segment connecting x and y are also in A

convex



# Crisp Sets (Classical or Nonfuzzy)

- The cartesian product set of A and B



$$A \times B = \{(a,b) \mid a \in A \text{ and } b \in B\}$$

- $|A \times B| = |A| \cdot |B|$

## Convex Set

- a set A in  $\mathbb{R}^n$  is convex iff, for every pair of points x and y in A, all the points lie in the line segment connecting x and y are also in A

convex





# Crisp Sets (Classical or Nonfuzzy)



- The cartesian product set of A and B

$$A \times B = \{(a,b) \mid a \in A \text{ and } b \in B\}$$

- $|A \times B| = |A| \cdot |B|$

## Convex Set

- a set A in  $\mathbb{R}^n$  is convex iff, for every pair of points x and y in A, all the points lie in the line segment connecting x and y are also in A



# Crisp Sets (Classical or Nonfuzzy)



- The cartesian product set of  $A$  and  $B$

$$A \times B = \{(a,b) \mid a \in A \text{ and } b \in B\}$$

- $|A \times B| = |A| \cdot |B|$


## Convex Set

- a set  $A$  in  $\mathbb{R}^n$  is convex iff, for every pair of points  $x$  and  $y$  in  $A$ , all the points lie in the line segment connecting  $x$  and  $y$  are also in  $A$




- $A$  is convex iff  $\forall a, b \in A, \forall \lambda \in [0,1], \lambda a + (1-\lambda)b \in A$

# Fuzzy Sets




# Fuzzy Sets




- $\mu_A : X \rightarrow [0,1]$   
membership function
- indicates the degree of membership

# Fuzzy Sets




- $\mu_A : X \rightarrow [0,1]$   
membership function
- indicates the degree of membership
- $\mu_A(a) = 1, \mu_A(b) = 1, \mu_A(c) = 0$

# Fuzzy Sets




- $\mu_A : X \rightarrow [0,1]$   
membership function
- indicates the degree of membership
- $\mu_A(a) = 1, \mu_A(b) = 1, \mu_A(c) = 0$
- $\mu_A(d) = 0.3, \mu_A(e) = 0.7$

# Fuzzy Sets



- $\mu_A : X \rightarrow [0,1]$   
membership function
  - indicates the degree of membership
  - $\mu_A(a) = 1, \mu_A(b) = 1, \mu_A(c) = 0$
  - $\mu_A(d) = 0.3, \mu_A(e) = 0.7$
- 
- $A = \{(a,1), (b,1), (d,0.3), (e,0.7)\}$

# Fuzzy Sets




- $\mu_A : X \rightarrow [0,1]$   
membership function
- indicates the degree of membership
- $\mu_A(a) = 1, \mu_A(b) = 1, \mu_A(c) = 0$
- $\mu_A(d) = 0.3, \mu_A(e) = 0.7$

- $A = \{(a,1), (b,1), (d,0.3), (e,0.7)\}$
- $A = 1/a + 1/b + 0.3/d + 0.7/e$




# Fuzzy Sets

- $A = \text{'real numbers close to 0'}$




# Fuzzy Sets

- $A = \text{'real numbers close to 0'}$



# Fuzzy Sets


- $A$  = 'real numbers close to 0'



$$\mu_A(x) = \begin{cases} x/4 + 1 & \text{if } x \text{ in } [-4, 0] \\ 1 - x/4 & \text{if } x \text{ in } [0, 4] \end{cases}$$

# Fuzzy Sets

- $A$  = 'real numbers close to 0'




$$\mu_A(x) = \begin{cases} x/4 + 1 & \text{if } x \text{ in } [-4, 0] \\ 1 - x/4 & \text{if } x \text{ in } [0, 4] \end{cases}$$

# Fuzzy Sets


- $A$  = 'real numbers close to 0'

$$\mu_A(x) = \begin{cases} x/4 + 1 & \text{if } x \text{ in } [-4, 0] \\ 1 - x/4 & \text{if } x \text{ in } [0, 4] \end{cases}$$



- $B$  = 'real numbers very close to 0'

# Fuzzy Sets




- $A = \text{'real numbers close to } 0\text{'}$

$$\mu_A(x) = \begin{cases} x/4 + 1 & \text{if } x \text{ in } [-4, 0] \\ 1 - x/4 & \text{if } x \text{ in } [0, 4] \end{cases}$$

- $B = \text{'real numbers very close to } 0\text{'}$

$$\mu_B(x) = 1 / (1 + x^2)$$

# Fuzzy Sets




- A = 'real numbers close to 0'

$$\mu_A(x) = \begin{cases} x/4 + 1 & \text{if } x \text{ in } [-4, 0] \\ 1 - x/4 & \text{if } x \text{ in } [0, 4] \end{cases}$$

- B = 'real numbers very close to 0'

$$\mu_B(x) = 1 / (1 + x^2)$$

# Fuzzy Sets



- A = 'real numbers close to 0'


$$\mu_A(x) = \begin{cases} x/4 + 1 & \text{if } x \text{ in } [-4, 0] \\ 1 - x/4 & \text{if } x \text{ in } [0, 4] \end{cases}$$

- B = 'real numbers very close to 0'

$$\mu_B(x) = 1 / (1 + x^2)$$



# Fuzzy Sets



- A = 'real numbers close to 0'


$$\mu_A(x) = \begin{cases} x/4 + 1 & \text{if } x \text{ in } [-4, 0] \\ 1 - x/4 & \text{if } x \text{ in } [0, 4] \end{cases}$$

- B = 'real numbers very close to 0'

$$\mu_B(x) = 1 / (1 + x^2)$$

- C = 'real numbers very very close to 0'

# Fuzzy Sets



- A = 'real numbers close to 0'

$$\mu_A(x) = \begin{cases} x/4 + 1 & \text{if } x \text{ in } [-4, 0] \\ 1 - x/4 & \text{if } x \text{ in } [0, 4] \end{cases}$$


- B = 'real numbers very close to 0'

$$\mu_B(x) = 1 / (1 + x^2)$$

- C = 'real numbers very very close to 0'

$$\mu_C(x) = (1 / (1 + x^2))^2$$

# Fuzzy Sets



- A = 'real numbers close to 0'

$$\mu_A(x) = \begin{cases} x/4 + 1 & \text{if } x \text{ in } [-4, 0] \\ 1 - x/4 & \text{if } x \text{ in } [0, 4] \end{cases}$$


- B = 'real numbers very close to 0'

$$\mu_B(x) = 1 / (1 + x^2)$$

- C = 'real numbers very very close to 0'

$$\mu_C(x) = (1 / (1 + x^2))^2$$

# Fuzzy Sets



- A = 'real numbers close to 0'

$$\mu_A(x) = \begin{cases} x/4 + 1 & \text{if } x \in [-4, 0] \\ 1 - x/4 & \text{if } x \in [0, 4] \end{cases}$$

- B = 'real numbers very close to 0'


$$\mu_B(x) = 1 / (1 + x^2)$$

- C = 'real numbers very very close to 0'

$$\mu_C(x) = (1 / (1 + x^2))^2$$


# Fuzzy Sets

- $X = \{5, 10, 15, 20, 25, 30, 35, 40, 45, 50\}$



# Fuzzy Sets


- $X = \{5, 10, 15, 20, 25, 30, 35, 40, 45, 50\}$



- $\text{Support}(A) = \{x \in X \mid \mu_A(x) > 0\}$

# Fuzzy Sets


- $X = \{5, 10, 15, 20, 25, 30, 35, 40, 45, 50\}$



- $\text{Support}(A) = \{x \in X \mid \mu_A(x) > 0\}$   
Support(Cold) =  $\{5, 10, 15\}$   
Support(Mild) =  $\{15, 20, 25\}$   
Support(Hot) =  $\{25, 30, 35, 40, 45, 50\}$

# Fuzzy Sets

- $X = \{5, 10, 15, 20, 25, 30, 35, 40, 45, 50\}$




- $\text{Support}(A) = \{x \in X \mid \mu_A(x) > 0\}$   
 $\text{Support}(\text{Cold}) = \{5, 10, 15\}$   
 $\text{Support}(\text{Mild}) = \{15, 20, 25\}$   
 $\text{Support}(\text{Hot}) = \{25, 30, 35, 40, 45, 50\}$

- $A_\alpha = \{x \in X \mid \mu_A(x) \geq \alpha\}$ ,  $\alpha$ -cut set



# Fuzzy Sets

- $X = \{5, 10, 15, 20, 25, 30, 35, 40, 45, 50\}$



- $\text{Support}(A) = \{x \in X \mid \mu_A(x) > 0\}$   
 $\text{Support}(\text{Cold}) = \{5, 10, 15\}$   
 $\text{Support}(\text{Mild}) = \{15, 20, 25\}$   
 $\text{Support}(\text{Hot}) = \{25, 30, 35, 40, 45, 50\}$


- $A_\alpha = \{x \in X \mid \mu_A(x) \geq \alpha\}$ ,  $\alpha$ -cut set

$$\text{Cold}_{0.5} = \{5, 10, 15\}$$

$$\text{Cold}_1 = \{5, 10\}$$

# Fuzzy Sets

- $X = \{5, 10, 15, 20, 25, 30, 35, 40, 45, 50\}$



- $\text{Support}(A) = \{x \in X \mid \mu_A(x) > 0\}$   
 $\text{Support}(\text{Cold}) = \{5, 10, 15\}$   
 $\text{Support}(\text{Mild}) = \{15, 20, 25\}$   
 $\text{Support}(\text{Hot}) = \{25, 30, 35, 40, 45, 50\}$

- $A_\alpha = \{x \in X \mid \mu_A(x) \geq \alpha\}$ ,  $\alpha$ -cut set

$$\text{Cold}_{0.5} = \{5, 10, 15\}$$


$$\text{Cold}_1 = \{5, 10\}$$

$$\text{Mild}_{0.5} = \{15, 20, 25\}$$

$$\text{Mild}_1 = \{20\}$$

# Fuzzy Sets

- $X = \{5, 10, 15, 20, 25, 30, 35, 40, 45, 50\}$



- $\text{Support}(A) = \{x \in X \mid \mu_A(x) > 0\}$   
 $\text{Support}(\text{Cold}) = \{5, 10, 15\}$   
 $\text{Support}(\text{Mild}) = \{15, 20, 25\}$   
 $\text{Support}(\text{Hot}) = \{25, 30, 35, 40, 45, 50\}$

- $A_\alpha = \{x \in X \mid \mu_A(x) \geq \alpha\}$ ,  $\alpha$ -cut set

$$\text{Cold}_{0.5} = \{5, 10, 15\}$$

$$\text{Hot}_{0.33} = \{25, 30, 35, 40, 45, 50\}$$

$$\text{Cold}_1 = \{5, 10\}$$

$$\text{Hot}_{0.66} = \{30, 35, 40, 45, 50\}$$


$$\text{Mild}_{0.5} = \{15, 20, 25\}$$

$$\text{Hot}_1 = \{35, 40, 45, 50\}$$

$$\text{Mild}_1 = \{20\}$$

# Fuzzy Sets

- $X = \{5, 10, 15, 20, 25, 30, 35, 40, 45, 50\}$



- $\text{Support}(A) = \{x \in X \mid \mu_A(x) > 0\}$   
 $\text{Support}(\text{Cold}) = \{5, 10, 15\}$   
 $\text{Support}(\text{Mild}) = \{15, 20, 25\}$   
 $\text{Support}(\text{Hot}) = \{25, 30, 35, 40, 45, 50\}$

- $A_\alpha = \{x \in X \mid \mu_A(x) \geq \alpha\}$ ,  $\alpha$ -cut set

$$\text{Cold}_{0.5} = \{5, 10, 15\}$$

$$\text{Hot}_{0.33} = \{25, 30, 35, 40, 45, 50\}$$

$$\text{Cold}_1 = \{5, 10\}$$

$$\text{Hot}_{0.66} = \{30, 35, 40, 45, 50\}$$

$$\text{Mild}_{0.5} = \{15, 20, 25\}$$


$$\text{Hot}_1 = \{35, 40, 45, 50\}$$

$$\text{Mild}_1 = \{20\}$$

- if  $u > v$ , then  $A_u \subseteq A_v$

# Fuzzy Sets

- $X = \{5, 10, 15, 20, 25, 30, 35, 40, 45, 50\}$



- $\text{Support}(A) = \{x \in X \mid \mu_A(x) > 0\}$   
 $\text{Support}(\text{Cold}) = \{5, 10, 15\}$   
 $\text{Support}(\text{Mild}) = \{15, 20, 25\}$   
 $\text{Support}(\text{Hot}) = \{25, 30, 35, 40, 45, 50\}$

- $A_\alpha = \{x \in X \mid \mu_A(x) \geq \alpha\}$ ,  $\alpha$ -cut set

$\text{Cold}_{0.5}$

$\text{Cold}_1 =$

- The maximum membership value is called the **height** of the fuzzy set.

$\text{Mild}_{0.5} = \{15, 20, 25\}$


$\text{Hot}_1 = \{35, 40, 45, 50\}$

$\text{Mild}_1 = \{20\}$

- if  $u > v$ , then  $A_u \subseteq A_v$

# Fuzzy Sets

- $X = \{5, 10, 15, 20, 25, 30, 35, 40, 45, 50\}$



- $\text{Support}(A) = \{x \in X \mid \mu_A(x) > 0\}$   
 $\text{Support}(\text{Cold}) = \{5, 10, 15\}$   
 $\text{Support}(\text{Mild}) = \{15, 20, 25\}$   
 $\text{Support}(\text{Hot}) = \{25, 30, 35, 40, 45, 50\}$

- $A_\alpha = \{x \in X \mid \mu_A(x) \geq \alpha\}$ ,  $\alpha$ -cut set

Cold<sub>0.5</sub>

Cold<sub>1</sub> =

Mild<sub>0.5</sub>

Mild<sub>1</sub> =

- The maximum membership value is called the **height** of the fuzzy set.

- If the height of a fuzzy set is 1, then it is called **normalized**.

- If  $u > v$ , then  $A_u \subseteq A_v$

# Fuzzy Sets

- $A$  is a subset of  $B$ ,  $A \subseteq B$ , if  $\mu_A(x) \leq \mu_B(x)$  for all  $x$  in  $X$

# Fuzzy Sets

- $A$  is a subset of  $B$ ,  $A \subseteq B$ , if  $\mu_A(x) \leq \mu_B(x)$  for all  $x$  in  $X$

## Convex Fuzzy Set

- Let  $A$  be a fuzzy set.  $\forall a, b$  in  $A$ , if  $\mu_A(\lambda a + (1-\lambda)b) \geq \min \{\mu_A(a), \mu_A(b)\}$  where  $\lambda$  in  $[0,1]$ , then  $A$  is a convex fuzzy set





# Fuzzy Sets

- $A$  is a subset of  $B$ ,  $A \subseteq B$ , if  $\mu_A(x) \leq \mu_B(x)$  for all  $x$  in  $X$

## Convex Fuzzy Set

- Let  $A$  be a fuzzy set.  $\forall a, b$  in  $A$ , if  $\mu_A(\lambda a + (1-\lambda)b) \geq \min \{\mu_A(a), \mu_A(b)\}$  where  $\lambda$  in  $[0,1]$ , then  $A$  is a convex fuzzy set





# Fuzzy Sets

- $A$  is a subset of  $B$ ,  $A \subseteq B$ , if  $\mu_A(x) \leq \mu_B(x)$  for all  $x$  in  $X$

## Convex Fuzzy Set

- Let  $A$  be a fuzzy set.  $\forall a, b$  in  $A$ , if  $\mu_A(\lambda a + (1-\lambda)b) \geq \min \{\mu_A(a), \mu_A(b)\}$  where  $\lambda$  in  $[0,1]$ , then  $A$  is a convex fuzzy set





# Fuzzy Sets

- $A$  is a subset of  $B$ ,  $A \subseteq B$ , if  $\mu_A(x) \leq \mu_B(x)$  for all  $x$  in  $X$

## Convex Fuzzy Set

- Let  $A$  be a fuzzy set.  $\forall a, b$  in  $A$ , if  $\mu_A(\lambda a + (1-\lambda)b) \geq \min \{\mu_A(a), \mu_A(b)\}$  where  $\lambda$  in  $[0,1]$ , then  $A$  is a convex fuzzy set





# Fuzzy Sets

- $A$  is a subset of  $B$ ,  $A \subseteq B$ , if  $\mu_A(x) \leq \mu_B(x)$  for all  $x$  in  $X$

## Convex Fuzzy Set

- Let  $A$  be a fuzzy set.  $\forall a, b$  in  $A$ , if  $\mu_A(\lambda a + (1-\lambda)b) \geq \min \{\mu_A(a), \mu_A(b)\}$  where  $\lambda$  in  $[0,1]$ , then  $A$  is a convex fuzzy set





# Fuzzy Sets

- $A$  is a subset of  $B$ ,  $A \subseteq B$ , if  $\mu_A(x) \leq \mu_B(x)$  for all  $x$  in  $X$

## Convex Fuzzy Set

- Let  $A$  be a fuzzy set.  $\forall a, b$  in  $A$ , if  $\mu_A(\lambda a + (1-\lambda)b) \geq \min \{\mu_A(a), \mu_A(b)\}$  where  $\lambda$  in  $[0,1]$ , then  $A$  is a convex fuzzy set



## Magnitude of Fuzzy Set



- Scalar Cardinality,  $|A| = \sum \mu_A(x)$ ,

# Fuzzy Sets

- $A$  is a subset of  $B$ ,  $A \subseteq B$ , if  $\mu_A(x) \leq \mu_B(x)$  for all  $x$  in  $X$

## Convex Fuzzy Set

- Let  $A$  be a fuzzy set.  $\forall a, b$  in  $A$ , if  $\mu_A(\lambda a + (1-\lambda)b) \geq \min \{\mu_A(a), \mu_A(b)\}$  where  $\lambda$  in  $[0,1]$ , then  $A$  is a convex fuzzy set



## Magnitude of Fuzzy Set



- Scalar Cardinality,  $|A| = \sum \mu_A(x)$ ,  
 $|Hot| = 0.33 + 0.66 + 1 + 1 + 1 + 1 = 5$

# Fuzzy Sets

- $A$  is a subset of  $B$ ,  $A \subseteq B$ , if  $\mu_A(x) \leq \mu_B(x)$  for all  $x$  in  $X$

## Convex Fuzzy Set

- Let  $A$  be a fuzzy set.  $\forall a, b$  in  $A$ , if  $\mu_A(\lambda a + (1-\lambda)b) \geq \min \{\mu_A(a), \mu_A(b)\}$  where  $\lambda$  in  $[0,1]$ , then  $A$  is a convex fuzzy set



## Magnitude of Fuzzy Set



- Scalar Cardinality,  $|A| = \sum \mu_A(x)$ ,  
 $|Hot| = 0.33 + 0.66 + 1 + 1 + 1 + 1 = 5$
- Relative Cardinality,  $\|A\| = |A| / |X|$

# Fuzzy Sets

- $A$  is a subset of  $B$ ,  $A \subseteq B$ , if  $\mu_A(x) \leq \mu_B(x)$  for all  $x$  in  $X$

## Convex Fuzzy Set

- Let  $A$  be a fuzzy set.  $\forall a, b$  in  $A$ , if  $\mu_A(\lambda a + (1-\lambda)b) \geq \min \{\mu_A(a), \mu_A(b)\}$  where  $\lambda$  in  $[0,1]$ , then  $A$  is a convex fuzzy set



## Magnitude of Fuzzy Set

- Scalar Cardinality,  $|A| = \sum \mu_A(x)$ ,  
 $|Hot| = 0.33 + 0.66 + 1 + 1 + 1 + 1 = 5$
- Relative Cardinality,  $\|A\| = |A| / |X|$   
 $\|Hot\| = 5/10 = 0.5$





# Fuzzy Sets

- $A$  is a subset of  $B$ ,  $A \subseteq B$ , if  $\mu_A(x) \leq \mu_B(x)$  for all  $x$  in  $X$

## Convex Fuzzy Set

- Let  $A$  be a fuzzy set.  $\forall a, b$  in  $A$ , if  $\mu_A(\lambda a + (1-\lambda)b) \geq \min \{\mu_A(a), \mu_A(b)\}$  where  $\lambda$  in  $[0,1]$ , then  $A$  is a convex fuzzy set



## Magnitude of Fuzzy Set



- Scalar Cardinality,  $|A| = \sum \mu_A(x)$ ,  
 $|Hot| = 0.33 + 0.66 + 1 + 1 + 1 + 1 = 5$
- Relative Cardinality,  $\|A\| = |A| / |X|$   
 $\|Hot\| = 5/10 = 0.5$
- Fuzzy Cardinality,  $|A|_F = \{|A_u|, u\}$ ,

# Fuzzy Sets

- $A$  is a subset of  $B$ ,  $A \subseteq B$ , if  $\mu_A(x) \leq \mu_B(x)$  for all  $x$  in  $X$

## Convex Fuzzy Set

- Let  $A$  be a fuzzy set.  $\forall a, b$  in  $A$ , if  $\mu_A(\lambda a + (1-\lambda)b) \geq \min \{\mu_A(a), \mu_A(b)\}$  where  $\lambda$  in  $[0,1]$ , then  $A$  is a convex fuzzy set



## Magnitude of Fuzzy Set

- Scalar Cardinality,  $|A| = \sum \mu_A(x)$ ,  
 $|Hot| = 0.33 + 0.66 + 1 + 1 + 1 + 1 = 5$
- Relative Cardinality,  $\|A\| = |A| / |X|$   
 $\|Hot\| = 5/10 = 0.5$
- Fuzzy Cardinality,  $|A|_F = \{(|A_u|, u)\}$ ,  
 $|Hot|_F = \{(6, 0.33), (5, 0.66), (4, 1)\}$