

several ionizing energies in the same measurement. Combined with a forthcoming purely circular ($S_3 = 1$) synchrotron radiation calibration of the molecular chiroptical response at harmonic photon energies, this PECD measurement will allow the disentangling of Stokes parameters S_3 (circular) and S_4 (unpolarized radiation) of the harmonic emission. In a first approximation, assuming a fully polarized elliptical radiation with vertical ellipse axis, and a zero anisotropy parameter β , we can extract a value for the first-order Legendre polynomial coefficient $b_1 = \text{PECD}/(2S_3)$ by using the ellipticities measured by optical polarimetry (Fig. 2). For the HOMO this gives $b_1 \approx 4 \pm 2 \times 10^{-3}$ at a photon energy of 15.5 eV (harmonic 5) and $b_1 \approx 1.1 \pm 0.5 \times 10^{-2}$ at a photon energy of 21.7 eV (harmonic 7). This is in decent agreement with synchrotron radiation measurements performed at neighbouring energies: $b_1 \approx 7 \pm 5 \times 10^{-3}$ at 14 and 16 eV, and $b_1 \approx 2 \pm 1 \times 10^{-2}$ at 22 eV (ref. 34).

By enabling the extension of the current frequency-resolved experiments carried out with synchrotron radiation to the time domain via pump-probe experiments, resonant HHG sources enable a very large avenue of research for 'femtochirality'. In particular, the transition states of enantioselective photochemical processes could be directly addressed by applying standard pump-probe femtochemistry methods. Other femtosecond probes of chirality in the gas phase have been demonstrated recently. Coulomb explosion imaging enables the absolute configuration of molecules to be determined³⁵, but is restricted to small systems. Compared to multi-photon ionization^{36,37}, XUV PECD has the advantage of being a universal direct ionization probe, involving no intermediate resonant state. Many photochemical processes could therefore be mapped out in the time domain by means of PECD, which is known to be extremely sensitive to static² and dynamical³ molecular structures. Elliptical HHG also opens the path to attosecond metrology of circular dichroism. The temporal profile associated with the superposition of consecutive harmonics is an attosecond pulse train. Using an additional probe laser field enables resolving the phase of the molecular photoionization process¹⁵, giving access to attosecond time delays of the photoelectrons¹⁴. It will thus be possible to resolve the different photoionization times of photoelectrons emitted from a chiral molecule forward or backwards relative to the direction of the light propagation, thereby obtaining a unique insight into the fundamental dynamics of electrons in chiral potentials.

Methods

The experiments were performed using the Aurore laser system at CELIA, which delivers 8 mJ, 25 fs, 800 nm pulses at 1 kHz. Frequency doubling was achieved using a 200- μm -thick type I BBO crystal (4 mJ of 800 nm gives 1 mJ of 400 nm light). The polarization state of the fundamental radiation was shaped by rotating a broadband zero-order half-waveplate in front of a quarter-waveplate, which enabled the laser ellipticity ε_0 to be varied while keeping the major axis of the ellipse fixed. The laser was focused by a thin 50 cm SiO_2 lens into a vacuum chamber and interacted with a continuous jet (300 μm nozzle) at 100 mbar or a static gas cell (3 mm long) filled at 5 mbar. Optical polarimetry using the two sources gave similar results. Note that SF_6 has the advantage of being non-toxic, non-corrosive and gaseous at room temperature, with a high vapour pressure. The harmonic polarization state was characterized using an XUV polarizer consisting of three bare gold mirrors under 75° – 60° – 75° incidence, leading to a contrast of 5–20 between S and P polarizations for harmonics 5 to 9 of the 400 nm light. A second half-waveplate was used to rotate the laser polarization, which is equivalent to rotating the XUV polarizer, to record the Malus' law.

For the PECD experiment, a reservoir of enantiopure fenchone (Sigma Aldrich) at room temperature was connected through a microleak to a heated metallic nozzle (120 $^\circ\text{C}$) of 300 μm diameter located 7 cm away from the interaction zone. The pressure in the interaction region was 2×10^{-6} mbar (5×10^{-8} mbar background pressure).

Received 11 March 2014; accepted 27 November 2014;
published online 22 December 2014

References

- Böwering, N. *et al.* Asymmetry in photoelectron emission from chiral molecules induced by circularly polarized light. *Phys. Rev. Lett.* **86**, 1187–1190 (2001).
- Powis, I. Photoelectron circular dichroism in chiral molecules. *Adv. Chem. Phys.* **138**, 267–329 (2008).
- Garcia, G. A., Nahon, L., Daly, S. & Powis, I. Vibrationally induced inversion of photoelectron forward-backward asymmetry in chiral molecule photoionization by circularly polarized light. *Nature Commun.* **4**, 2132 (2013).
- Stohr, J. *et al.* Element-specific magnetic microscopy with circularly polarized X-rays. *Science* **259**, 658–661 (1993).
- Boeglin, C. *et al.* Distinguishing the ultrafast dynamics of spin and orbital moments in solids. *Nature* **465**, 458–461 (2010).
- Schoenlein, R. W. *et al.* Generation of femtosecond pulses of synchrotron radiation. *Science* **287**, 2237–2240 (2000).
- Čutić, N. *et al.* Vacuum ultraviolet circularly polarized coherent femtosecond pulses from laser seeded relativistic electrons. *Phys. Rev. ST Accel. Beams* **14**, 030706 (2011).
- Allaria, E. *et al.* Highly coherent and stable pulses from the FERMI seeded free-electron laser in the extreme ultraviolet. *Nature Photon.* **6**, 699–704 (2012).
- Le Déroff, L., Salières, P., Carré, B., Joyeux, D. & Phalippou, D. Measurement of the degree of spatial coherence of high-order harmonics using a Fresnel-mirror interferometer. *Phys. Rev. A* **61**, 043802 (2000).
- Hergott, J.-F. *et al.* Extreme-ultraviolet high-order harmonic pulses in the microjoule range. *Phys. Rev. A* **66**, 021801 (2002).
- Mahieu, B. *et al.* Full tunability of laser femtosecond high-order harmonics in the ultraviolet spectral range. *Appl. Phys. B* **108**, 43–49 (2012).
- Mairesse, Y. *et al.* High harmonic XUV spectral phase interferometry for direct electric-field reconstruction. *Phys. Rev. Lett.* **94**, 173903 (2005).
- Chini, M., Zhao, K. & Chang, Z. The generation, characterization and applications of broadband isolated attosecond pulses. *Nature Photon.* **8**, 178–186 (2014).
- Klünder, K. *et al.* Probing single-photon ionization on the attosecond time scale. *Phys. Rev. Lett.* **106**, 143002 (2011).
- Haessler, S. *et al.* Phase-resolved attosecond near-threshold photoionization of molecular nitrogen. *Phys. Rev. A* **80**, 011404(R) (2009).
- Lépine, F., Ivanov, M. Y. & Vrakking, M. J. J. Attosecond molecular dynamics: fact or fiction? *Nature Photon.* **8**, 195–204 (2014).
- Bauer, M. Femtosecond ultraviolet photoelectron spectroscopy of ultra-fast surface processes. *J. Phys. D* **38**, R253–R267 (2005).
- Vodungbo, B. *et al.* Polarization control of high order harmonics in the EUV photon energy range. *Opt. Express* **19**, 4346–4356 (2011).
- Budil, K. S., Salières, P., Perry, M. D. & L'Huillier, A. Influence of ellipticity on harmonic generation. *Phys. Rev. A* **48**, R3437–R3440 (1993).
- Antoine, P., Carré, B., L'Huillier, A. & Lewenstein, M. Polarization of high-order harmonics. *Phys. Rev. A* **55**, 1314–1324 (1997).
- Milšević, D. B., Becker, W. & Kopold, R. Generation of circularly polarized high-order harmonics by two-color coplanar field mixing. *Phys. Rev. A* **61**, 063403 (2000).
- Yuan, K.-J. & Bandrauk, A. D. Single circularly polarized attosecond pulse generation by intense few cycle elliptically polarized laser pulses and terahertz fields from molecular media. *Phys. Rev. Lett.* **110**, 023003 (2013).
- Fleischer, A., Kfir, O., Diskin, T., Sidorenko, P. & Cohen, O. Spin angular momentum and tunable polarization in high-harmonic generation. *Nature Photon.* **8**, 543–549 (2014).
- Smirnova, O. *et al.* Attosecond circular dichroism spectroscopy of polyatomic molecules. *Phys. Rev. Lett.* **102**, 063601 (2009).
- Zhou, X. *et al.* Elliptically polarized high-order harmonic emission from molecules in linearly polarized laser fields. *Phys. Rev. Lett.* **102**, 073902 (2009).
- Mairesse, Y. *et al.* High harmonic spectroscopy of multichannel dynamics in strong-field ionization. *Phys. Rev. Lett.* **104**, 213601 (2010).
- Yang, L. *et al.* Energy-dependent valence photoelectron spectra of SF_6 : *ab initio* calculations and measurements. *J. Electron Spectrosc. Rel. Phenom.* **94**, 163–179 (1998).
- Ferré, A. *et al.* Multi-channel static and dynamical resonant high-order harmonic generation. *Nature Commun.* **5**, 5952 (2014).
- Chini, M. *et al.* Coherent phase-matched VUV generation by field-controlled bound states. *Nature Photon.* **8**, 437–441 (2014).
- Rabinovitch, K., Canfield, L. R. & Madden, R. P. A method for measuring polarization in the vacuum ultraviolet. *Appl. Opt.* **4**, 1005–1010 (1965).
- Ferré, A. *et al.* High-harmonic transient grating spectroscopy of SF_6 molecular vibrations. *J. Phys. B* **47**, 124023 (2014).
- Falcão-Filho, E. L. *et al.* Scaling of high-order harmonic efficiencies with visible wavelength drivers: a route to efficient extreme ultraviolet sources. *Appl. Phys. Lett.* **97**, 061107 (2010).
- Nahon, L., Garcia, G. A., Harding, C. J., Mikajlo, E. & Powis, I. Determination of chiral asymmetries in the valence photoionization of camphor enantiomers by photoelectron imaging using tunable circularly polarized light. *J. Chem. Phys.* **125**, 114309 (2006).
- Powis, I., Harding, C. J., Garcia, G. A. & Nahon, L. A valence photoelectron imaging investigation of chiral asymmetry in the photoionization of fenchone and camphor. *Chem. Phys. Chem.* **9**, 475–483 (2008).